

RESEARCH ARTICLE

Multilocus Bayesian Estimates of Intra-Oceanic Genetic Differentiation, Connectivity, and Admixture in Atlantic Swordfish (*Xiphias gladius* L.)

Brad L. Smith^{1,2*}, Ching-Ping Lu², Blanca García-Cortés³, Jordi Viñas⁴, Shean-Ya Yeh⁵, Jaime R. Alvarado Bremer^{1,2}

1 Department of Marine Biology, Texas A&M University at Galveston, OCSB 3029, Galveston, TX 77553, United States of America, **2** Department of Wildlife and Fisheries Sciences, Texas A&M University, 210 Nagle Hall, TAMU 2258, College Station, TX, 75044, United States of America, **3** Centro Oceanográfico A Coruña, Instituto Español de Oceanografía, Muelle de las Animas s/n P.O. Box 130, 15080 A Coruña, Spain, **4** Department de Biologia, Universitat de Girona, Campus Montilivi, E-17071 Girona, Spain, **5** Institute of Oceanography, National Taiwan University, Taipei 106, Taiwan

* brad.smith@byuh.edu



OPEN ACCESS

Citation: Smith BL, Lu C-P, García-Cortés B, Viñas J, Yeh S-Y, Alvarado Bremer JR (2015) Multilocus Bayesian Estimates of Intra-Oceanic Genetic Differentiation, Connectivity, and Admixture in Atlantic Swordfish (*Xiphias gladius* L.). PLoS ONE 10(6): e0127979. doi:10.1371/journal.pone.0127979

Academic Editor: Daniele Canestrelli, Tuscia University, ITALY

Received: January 17, 2015

Accepted: April 21, 2015

Published: June 9, 2015

Copyright: © 2015 Smith et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](http://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: DNA sequences of marker loci: Genbank accessions: JX042446-JX04251, AF069912-AF069913, FJ890938-FJ890941, FJ911901-FJ911904. SNP allelic data for 774 swordfish used for analyses are included in Supporting Information. All other relevant data are within the paper and its Supporting Information files.

Funding: Funding and support was provided by Texas Institute of Oceanography (<http://www.tamug.edu/research/TIO/TIO.html>), The McDaniel Charitable Foundation Small Grant Program (<http://mcdanielcharitablefoundation.org/research-grant>)

Abstract

Previous genetic studies of Atlantic swordfish (*Xiphias gladius* L.) revealed significant differentiation among Mediterranean, North Atlantic and South Atlantic populations using both mitochondrial and nuclear DNA data. However, limitations in geographic sampling coverage, and the use of single loci, precluded an accurate placement of boundaries and of estimates of admixture. In this study, we present multilocus analyses of 26 single nucleotide polymorphisms (SNPs) within 10 nuclear genes to estimate population differentiation and admixture based on the characterization of 774 individuals representing North Atlantic, South Atlantic, and Mediterranean swordfish populations. Pairwise F_{ST} values, AMOVA, PCoA, and Bayesian individual assignments support the differentiation of swordfish inhabiting these three basins, but not the current placement of the boundaries that separate them. Specifically, the range of the South Atlantic population extends beyond 5°N management boundary to 20°N-25°N from 45°W. Likewise the Mediterranean population extends beyond the current management boundary at the Strait of Gibraltar to approximately 10°W. Further, admixture zones, characterized by asymmetric contributions of adjacent populations within samples, are confined to the Northeast Atlantic. While South Atlantic and Mediterranean migrants were identified within these Northeast Atlantic admixture zones no North Atlantic migrants were identified respectively in these two neighboring basins. Owing to both, the characterization of larger number of loci and a more ample spatial sampling coverage, it was possible to provide a finer resolution of the boundaries separating Atlantic swordfish populations than previous studies. Finally, the patterns of population structure and admixture are discussed in the light of the reproductive biology, the known patterns of dispersal, and oceanographic features that may act as barriers to gene flow to Atlantic swordfish.

[program/small-grant-program/](#)) to JRAB, TAMUG-Graduate Research Office and Marine Biology Department, the Erma Lee and Luke Mooney Graduate Student Travel Grant, and the Erma Lee and Luke Mooney Graduate Student Research Excellence Award. The open access publishing fees for this manuscript have been covered by the Texas A&M University Online Access to Knowledge (OAK) Fund, supported by the University Libraries and the Office of the Vice President for Research. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

Introduction

The epipelagic realm of the world's oceans is a relatively homogeneous and contiguous environment characterized by the near absence of physical barriers. These features when coupled with the high dispersal behavior and demographic characteristics of many marine species result in high levels of gene flow [1]. Specifically, many epipelagic species of fishes are characterized by long distance potential dispersal of pelagic eggs and larvae facilitated by currents, high vagility of juveniles and adults, opportunistic feeding behaviors, eurythermal physiologies of adults, and large effective population sizes. Consequently, the levels of genetic differentiation in marine fishes are typically substantially lower compared to populations of anadromous and freshwater fishes [2]. It is thus not surprising that the distributions of many species of tunas, billfishes, dolphinfishes, lamnid sharks, are cosmopolitan [3, 4], with some species, such as skipjack tuna (*Katsuwonus pelamis*) displaying no genetic differentiation among-oceans [5], and others, like wahoo (*Acanthocybium solandri*) showing only incipient levels inter-oceanic differentiation [6]. Significant inter-oceanic differentiation, however, has been documented in many highly migratory fishes, including bigeye tuna [7], Atlantic bluefin tuna [8, 9], albacore tuna [10, 11], and many species of billfishes [12] and swordfish [13]. Similarly, genetic investigations in several species of billfish have shown intra-oceanic genetic differentiation within the Pacific Ocean, including sailfish [14], white marlin, and blue marlin [12]. Within the Atlantic, no significant population substructure was detected in several epipelagic fishes; including shortfin mako (*Isurus oxyrinchus*) [15], blue marlin (*Makaira nigricans*) and sailfish (*Istiophorus platypterus*) [12], bigeye tuna (*Thunnus obesus*) [16], and in both yellowfin tuna (*Thunnus albacares*) and skipjack tuna [5]. By contrast, swordfish (*Xiphias gladius* L.) shows significant genetic differentiation within the Atlantic Ocean [13, 17, 18] and also within the Mediterranean Sea [19].

Swordfish is a large epipelagic monotypic cosmopolitan species capable of long distance movements, as documented using both conventional tags [20] and pop-up satellite tags (PSATs) [21–24]. Swordfish are subject to intensive commercial exploitation worldwide, and in the Atlantic Ocean they are managed by the International Commission for the Conservation of Tunas (ICCAT) as two separate stocks: the North Atlantic and the South Atlantic separated at 5°N. In turn, North Atlantic swordfish is managed separately from Mediterranean swordfish with a boundary at the Strait of Gibraltar. Previous genetic studies on Atlantic swordfish employing mitochondrial DNA (mtDNA) (see [25] for summary), single copy nuclear DNA (scnDNA) [18, 26, 27], and to lesser extent microsatellite loci [28–30], confirm the differentiation among North Atlantic, South Atlantic and Mediterranean populations. Some of these studies also suggested that the range of these swordfish populations extend beyond current management boundaries [26, 31]. However, limitations in geographic sampling coverage, and the use of single loci, precluded an accurate placement of boundaries and of estimates of admixture. Multilocus analyses of microsatellite data [29, 30] attempted to refine these estimates by expanding sampling coverage and by utilizing Bayesian individual assignment to identify zones of admixture. While, microsatellite data confirmed the pronounced differences between Atlantic and Mediterranean swordfish [29], it failed to delineate the separation of North Atlantic and South Atlantic swordfish populations.

In recent years there has been an increase in the adoption of single nucleotide polymorphisms (SNPs) as an alternative to microsatellite data in Bayesian analyses of population admixture. In addition to a comparable differentiation power, SNPs have lower associated genotyping errors, simpler mutation models, higher amenability to automation and high throughput technologies, and increased amplification success with low-quality and ancient DNA as compared to microsatellites [32–34]. Further, the levels of genetic differentiation

among Atlantic swordfish populations using SNPs was an entire order of magnitude greater compared to microsatellites [35, 36]. Accordingly, in this study we evaluate the patterns of population differentiation and admixture of Atlantic and Mediterranean swordfish based on Bayesian analyses of multilocus nuclear SNPs using a more extensive and representative geographic sampling coverage compared to previous studies. We describe the patterns of differentiation and admixture among North Atlantic, South Atlantic and Mediterranean swordfish populations.

Materials and Methods

Sampling

A total of 774 swordfish specimens from 18 localities in the Atlantic and western Mediterranean were collected from 1991–2006 (Table 1 and Fig 1). Muscle, liver, and heart tissue samples of were collected by observers on commercial longline vessels or obtained from collaborating researchers. Ethical approval was not required for the collection of adult swordfish, as these were collected during routine commercial operations. Swordfish adults were sacrificed by the fishermen by decapitation. In all instances the location of capture for each fish was recorded. Larval swordfish (n = 52), collected as part of ichthyoplankton surveys in the northern Gulf of Mexico during the summers of 2005–2006 [37], were also characterized. Permits for fish larvae collections in the Gulf of Mexico were issued by the Highly Migratory Species Management Division of the National Oceanic and Atmospheric Administration.

(permits: Billfish-SRP-06-01, Billfish-EFP-07-03, and Billfish-EFP-08-03) to Dr. J.R. Rooker. The management boundaries at 5°N, separating the North Atlantic stock from the South

Table 1. Sampling localities for 774 swordfish subdivided into three stocks (North Atlantic, Mediterranean, South Atlantic) by ICATT management boundaries of 5°N and Strait of Gibraltar.

Population	Sample Locality	Latitude	Longitude	Sampling Date(s)	n
North Atlantic					
1	Gulf of Mexico	26-28N	86-93W	06/06-07/06, 06/07	52
2	Florida	21-32N	78-85W	1/93-8/93	49
3	Northeast US	35-42N	67-74W	8/90-11/90, 1/91-3/91	42
4	Lesser Antilles	12-20N	49-64W	10/92-2/93, 2/94, 1/95	48
5	East of Flemish Cap	43-47N	37-41W	9/04-11/04	17
6	Central North Atlantic	32-47N	37-43W	11/91, 12/95, 9/04-11/04	20
7	Iberian	35-40N	10-25W	11/91, 12/92, 3/93-3/93, 12/95-4/96	51
8	Strait of Gibraltar	33-35N	5-10W	8/92, 6/93-9/93, 5/96-11/96	83
10	Morocco	30-33N	12-15W	4/93-5/93, 10/96-11/96, 6/02-8/02	20
11	Western Sahara	20-28N	12-20W	2/92, 5/92, 6/93, 12/95, 4/96, 7/96-11/96	37
12	Cape Verde	5N-17N	21-32W	8/04-12/04, 9/05-10/05	56
Mediterranean					
9	West Mediterranean	35-39N	0-8E	5/92-7/92, 8/03-12/03	59
South Atlantic					
13	Equatorial Brazil	1S-3N	23-35W	11/91, 3/00, 9/04	42
14	Gulf of Guinea	5N	4W	11/91, 7/98-8/98	45
15	Central South Atlantic	14-16S	10-17W	10/04-12/04	28
16	Brazil	15-23S	22-27W	3/96, 5/96, 4/01, 12/04	35
17	Brazil-Uruguay	27-37S	32-42W	8/95-10/95, 4/01, 6/03-7/03	44
18	Namibia	26-27S	10-11E	7/99-8/99	46

doi:10.1371/journal.pone.0127979.t001




Fig 1. The geographical sampling of 774 swordfish (blue dots) and the locality groupings (numbered black circles) used for analyses as outlined in Table 1.

doi:10.1371/journal.pone.0127979.g001

Atlantic stock, and at the Strait of Gibraltar, separating the North Atlantic stock from the Mediterranean stock, were initially used to assign localities to the corresponding three management units or stocks currently recognized by ICCAT [38].

DNA extraction and nuclear loci genotyping

Total genomic DNA was isolated from tissue with a modified TENS and Proteinase K (20 mg/μL) digestion followed by ethanol precipitation without organic extraction [39]. Ten nuclear loci were amplified and genotyped for genetic population structure analyses: acidic ribosomal phosphoprotein P0 (ARP); adenine nucleotide translocator (ANT); aldolase-B (AldB); alpha-skeletal actin (Act2α); ATP synthase beta-subunit (ATPsβ); Calmodulin (CaM); Golgi pH regulator (GpHr); lactose dehydrogenase A (LdhA); myosin light chain (Mlc2); and signal

recognition particle 54 (SRP54). Details on loci and corresponding SNPs selection, the minimization of ascertainment bias as well as the criteria to design primers and probes that target these, are given in Smith et al. [35]. Briefly, to minimize ascertainment bias potential loci and SNPs were selected after conducting a preliminary characterization with HRMA of 30 swordfish, 10 from each of three geographically disjunct swordfish populations (NW Atlantic, Mediterranean and Pacific). Those amplicons that generated distinct melting profiles, were sequenced, and electropherogram inspections were then used to validate the SNPs responsible for the polymorphic melting curves (see [35] for details). Sequence alignments were also used to evaluate the placement and design of new HRMA-genotyping primers and probes, when required. A total of 85 SNPs were initially identified but only those differing at frequencies >5% among reference samples were targeted for further HRMA primer and or probe design. PCR conditions of selected potential loci were further optimized and validated using additional samples ($n = 40$) from each of the three reference populations. This process ultimately yielded a total of 26 SNPs among 10 loci that we targeted for genotyping (see Table 2 in [35]). Procedures for the amplification and subsequent genotyping of SNPs for short amplicon high-resolution melting analysis (SA-HRMA) of ARP, ATPs β , *CaM*, and SRP54 and unlabeled probe high-resolution melting analysis (UP-HRMA) of ANT, GpHR, *ldhA*, and *Mlc2* are described in Smith et al. [35].

Two additional loci, *AldB* and *Act2 α* , were scored as size polymorphisms as follows. A simple sequence repeat (SSR) identified in *AldB* [39] was amplified with the following primers; 5'-VIC-TGTGCCCAGTATAAGAAGGATGG-3' and 5'-CTGTGGAGAATCAGGGCTCC-3' (JX042447). Polymerase chain reactions (PCR) were conducted in 12.5 μ L reactions containing 10 ng of genomic DNA, 1X Econotaq Green Master Mix (Lucigen), and 0.20 μ M of each primer. Thermocycling was performed on an Eppendorf Mastercycler (Eppendorf) with an initial denaturation of 10 min at 95°C followed by 35 cycles denaturing for 1 min at 94°C, annealing for 1 min at 54°C, and extension for 1 min at 72°C. PCR products were diluted 1:10 and 1 μ L of diluted PCR template and 0.2 μ L of ROX size standard (Life Technologies) in 10 μ L of Hi-Di formamide were run on an ABI 3130 Genetic Analyzer (Applied Biosystems). SSRs were scored by size and analyzed using GeneMapper v.4.0 (Applied Biosystems).

Locus *Act2 α* contains an SNP that corresponds to a restriction site of endonuclease Hpy8I and was amplified using primers; 5'-GTCACCGGAGTCCAGGACG-3' and 5'-ATCTGGCAC CACACCTTCTACAA-3' (JX042448) using the same chemistry and thermocycling profile used in *AldB* amplification. Prior to digestion, the quality and quantity of PCR products were visualized in 1% Tris-acetate (TA) agarose gels via gel electrophoresis prior to digests. Restriction digests were conducted in 10 μ L volumes containing 2–4 μ L PCR product, 1X Buffer TANGO (Fermentas), and 0.4 U of Hpy8I (Fermentas), that were incubated for 16 hours at 37°C, followed by thermal inactivation at 80°C for 5 min. Restriction Fragment Length Polymorphisms (RFLPs) were separated in 2% TA agarose gels run at 100V for 30 min, and were visualized and captured on a Gel Doc XR (Bio-Rad) UV transilluminator using the Quality One v.4.6 (Bio-Rad) software.

Data analysis

Genetic diversity. The number of alleles, observed (H_o) and expected (H_e) heterozygosities under Hardy-Weinberg equilibrium (HWE), and inbreeding coefficients (F_{IS}) were calculated using GENALEX version 6.41 [40]. Departures from HWE were estimated using an exact probability test [41, 42] and global linkage disequilibrium using a maximum likelihood-ratio test [43] in GENEPOP version 4.1 (Markov chain parameters: 10000 dememorization steps, 1000 batches, 10000 iterations per batch) [44, 45]. Sequential Bonferroni corrections were

applied to adjust the levels of statistical significance for multiple comparisons [46]. An F_{ST} -outlier detection method, in which observed locus F_{ST} values are compared to calculated global F_{ST} values expected under neutrality using coalescent simulations [47] was performed in LOSITAN [48]. A total of 50,000 simulations were conducted to detect putative loci under selection. To remove potential selective loci when computing the initial mean F_{ST} , the first simulation used the 'neutral' mean F_{ST} option with a 0.995 confidence interval, a false discovery rate set at 0.1, and by pooling the sampling localities into the corresponding three populations ($K = 3$) identified by Bayesian clustering analyses (see below) as follows, North Atlantic (1–7, 10–11), South Atlantic (12–18), and Mediterranean (8–9). The assignment power of each locus was ranked *a posteriori* to Bayesian analysis using the critical population method in WHICHLOCI v1.0 [49] using samples from localities identified as non-mixing areas.

Population differentiation. A hierarchical analysis of molecular variance (AMOVA) [50] was implemented in Arlequin version 3.5 [51] to estimate levels of population subdivision. Population differentiation was also evaluated using both global and pairwise F_{ST} tests [52]. Initially, AMOVA groupings of two (MED and Atlantic) and three (MED, NA, and SA) adhering to the current management regions were evaluated. Alternative groupings in AMOVA were evaluated *a posteriori* to F_{ST} and Bayesian analyses (see below). Slatkin's linearized F_{ST} [53] was also calculated in Arlequin. Principal Coordinates Analysis (PCoA) [54], based on a standardized covariance matrix of genetic distances between each locality pair, was carried in GenAlEx version 6.4 [40].

Genetic clustering analysis. Genetic population structure and patterns of inter-population gene flow were assessed by Bayesian inference implemented in STRUCTURE version 2.3 [55]. Previous genetic studies using mtDNA and a single nuclear locus identified comparatively low levels of gene flow in between North and South Atlantic swordfish [17, 26, 56]. Accordingly, a no admixture ancestry with correlated allele frequencies model, as outlined in Falush et al. [57] was adopted. Compared to freshwater and anadromous fishes, the populations of marine fishes display weak levels of differentiation ($F_{ST} < 0.20$) [58]. Accordingly, the LOCPRIOR option was implemented to detect weak signals of population structure using an *a priori* grouping as outlined in Hubisz et al. [59]. The number of clusters (K) was estimated using an *ad hoc* approach [55] by obtaining the mean posterior probability of the data ($L(K)$) and the ΔK approach of Evanno et al. [60] using STRUCTURE HARVESTER v0.6.92 [61]. Twenty independent runs for each K value (1–10) were performed using 100,000 Markov chain Monte Carlo (MCMC) iterations with a burn-in period of 100,000. Results from STRUCTURE were compared with the results from GENELAND [62], which incorporates individual spatial and genetic data to infer population structure and spatial boundaries between clusters. Individual spatial coordinates were set to the corresponding latitude and longitude of sampling. An uncertainty value on the spatial coordinates of each sample was set to 30 decimal degrees corresponding to the documented displacement of North Atlantic swordfish from tagging experiments [22, 23]. Correlated allele and spatial models were implemented for twenty independent runs with 100,000 MCMC iterations and a burn-in of 10,000 iterations for each K value (1–8). After estimating K , 25 independent runs with the optimal K value were run in GENELAND, 50 independent runs in STRUCTURE, and the optimal alignment of replicates were completed using CLUMPP version 1.1.12 [63].

Results

Hardy-Weinberg, linkage disequilibrium, and F_{ST} outlier analysis

A total of 774 swordfish were genotyped successfully at the ten loci (see supporting S5 Table). Five of these loci contained a single SNP and were bi-allelic and the rest contained both

multiple SNPs and alleles. The number of alleles per locus ranged from 2 to 6, and in total 26 SNPs were characterized (see supporting [S1 Table](#)). Average (H_o) and expected (H_e) heterozygosities, and inbreeding coefficients (F_{IS}) are given in [Table 2](#). None of the Fisher's exact probability tests for departure from HWE were significant ($P < 0.05$) after Bonferroni correction. Genotypic linkage was not significant among the 44 possible pairs of nuclear loci across all samples with ($P < 0.01$) and without ($P < 0.05$) Bonferroni correction. Likewise exact tests for genotypic linkage disequilibrium within each sample were also non-significant after Bonferroni correction ($P < 0.05$) (not shown). Initial F_{ST} outlier analysis of the ten loci of all individuals from the 18 localities identified *CaM* as a potentially under selection ($P < 0.05$) ([Fig 2A](#)). However, since both STRUCTURE and GENELAND identified several localities (6–7, 10–11) in the North Atlantic as admixture zones there was the possibility that heterozygote deficits biased the outlier analysis. Consequently, this analysis was conducted excluding admixed samples resulting in no locus identified as an outlier ($P < 0.05$). This procedure also yielded higher F_{ST} values for all loci as well as higher H_e values for *Act2 α* , *AldB*, *ATPs β* , *CaM*, and *Mlc2*. ([Fig 2B](#)). The power assignment of each loci using WHICHLOCI v1.0 identified *CaM* as the locus with the highest power of discrimination for North and South Atlantic populations followed by *Mlc2*, *SRP54*, *AldB*, *ARP*, *Act2 α* , *ldhA*, *GpHR*, *ATPs β* , and *ANT* respectively (see supporting [S2 Table](#)). Interestingly, *CaM* displayed the lowest power of discrimination between North Atlantic and Mediterranean populations, with *Mlc2* yielding the highest power of discrimination between them followed by *ATPs β* , *Act2 α* , *ldhA*, *ANT*, *AldB*, *SRP54*, *ARP*, and *GpHR*, respectively (see supporting [S3 Table](#)).

Population differentiation

Pairwise F_{ST} . Multilocus pairwise F_{ST} values among the 18 populations ranged considerably (<0.001 to 0.121) ([Table 3](#)). The comparison between Mediterranean and South Atlantic populations yielded the highest indices of differentiation ($F_{ST} > 0.081 - 0.121$). Several sampling localities identified as admixture zones (6–7, 10–11) were excluded when estimating gene flow among North Atlantic (1–5), Mediterranean (8–9) and South Atlantic (12–18) populations. The lowest level of gene flow of about 5 migrants per generation (Nm) was estimated between Mediterranean and South Atlantic populations, compared to about 9 migrants per generation between North Atlantic and South Atlantic swordfish ([Table 4](#)). Slatkin's linearized F_{ST} values ([Fig 3](#)) highlights the homogeneity between the Mediterranean sample collected to the east of the Alboran Sea (8) and the adjacent Northeast Atlantic area west of the Strait of Gibraltar (9). The magnitude of these fixation indices also underscore the high degree of differentiation of Mediterranean swordfish with respect to all the Atlantic localities (1–7, 10–18) including those inhabiting the adjacent NE Atlantic waters off both the Iberian Peninsula and the NW African coast. In addition, [Fig 2](#) illustrates that the North Atlantic localities (1–5) are different from the South Atlantic localities (13–18) but also from locality 12 which corresponds geographically to the North Atlantic stock since it is located north of the 5°N management boundary. Linearized F_{ST} values depict a geographic gradient of differentiation from the Northwest Atlantic to the Southeast Atlantic indicative of mixing between the South Atlantic and North Atlantic populations, primarily in the eastern localities of Iberian, Morocco, and Western Sahara (7, 10–11) and to a lesser extent in the central North Atlantic (6) locality.

AMOVA. In all alternative hierarchical arrangements tested (two- and three-group; with all localities and excluding admixture areas) with AMOVA ([Table 5](#)) the majority of the variation ($>90\%$) in swordfish was contained within individuals. In all arrangements, among-groups (F_{CT}) differentiation was highly significant ($P < 0.01$). However, the proportions of variation explained by among-groups were slightly larger in two-group AMOVAs, with 6.63%

Table 2. Average genetic diversity of 10 nuclear loci within 18 swordfish populations.

Population		Locus									
		ARP	ALDB	GpHR	Act2α	ATPsβ	CaM	Mlc2	SRP54	LDHA	ANT
1	A	2	4	5	2	2	2	3	2	3	5
	H_o	0.288	0.288	0.615	0.231	0.500	0.346	0.173	0.327	0.385	0.442
	H_E	0.447	0.287	0.581	0.473	0.453	0.497	0.206	0.322	0.450	0.567
	F_{IS}	0.354	-0.005	-0.059	0.513	-0.105	0.304	0.161	-0.014	0.145	0.220
2	A	2	4	5	2	2	2	3	2	3	5
	H_o	0.347	0.224	0.571	0.592	0.469	0.510	0.265	0.388	0.633	0.571
	H_E	0.479	0.205	0.512	0.500	0.493	0.497	0.320	0.359	0.468	0.522
	F_{IS}	0.276	-0.097	-0.115	-0.184	0.047	-0.027	0.170	-0.079	-0.352	-0.094
3	A	2	4	5	2	2	2	3	2	3	5
	H_o	0.381	0.238	0.571	0.429	0.429	0.476	0.357	0.310	0.476	0.405
	H_E	0.472	0.218	0.521	0.459	0.427	0.499	0.381	0.262	0.447	0.504
	F_{IS}	0.192	-0.094	-0.097	0.067	-0.003	0.045	0.062	-0.183	-0.065	0.197
4	A	2	4	5	2	2	2	3	2	3	6
	H_o	0.417	0.188	0.521	0.458	0.458	0.500	0.458	0.292	0.583	0.542
	H_E	0.444	0.209	0.513	0.413	0.457	0.499	0.400	0.330	0.476	0.545
	F_{IS}	0.062	0.101	-0.015	-0.109	-0.002	-0.002	-0.147	0.116	-0.225	0.007
5	A	2	3	4	2	2	2	2	2	2	5
	H_o	0.471	0.176	0.353	0.353	0.412	0.471	0.353	0.235	0.588	0.412
	H_E	0.415	0.164	0.424	0.415	0.389	0.498	0.291	0.291	0.457	0.356
	F_{IS}	-0.133	-0.074	0.167	0.150	-0.058	0.056	-0.214	0.190	-0.288	-0.155
6	A	2	3	3	2	2	2	2	2	3	4
	H_o	0.350	0.150	0.300	0.200	0.350	0.400	0.350	0.250	0.700	0.500
	H_E	0.489	0.141	0.339	0.255	0.349	0.480	0.349	0.219	0.540	0.529
	F_{IS}	0.284	-0.062	0.114	0.216	-0.004	0.167	-0.004	-0.143	-0.296	0.054
7	A	2	3	3	2	2	2	2	2	3	4
	H_o	0.350	0.150	0.300	0.200	0.350	0.400	0.350	0.250	0.700	0.500
	H_E	0.489	0.141	0.339	0.255	0.349	0.480	0.349	0.219	0.540	0.529
	F_{IS}	0.284	-0.062	0.114	0.216	-0.004	0.167	-0.004	-0.143	-0.296	0.054
8	A	2	4	5	2	2	2	3	2	3	5
	H_o	0.301	0.301	0.590	0.349	0.422	0.470	0.494	0.108	0.494	0.518
	H_E	0.384	0.294	0.503	0.319	0.451	0.497	0.506	0.164	0.511	0.532
	F_{IS}	0.215	-0.024	-0.173	-0.097	0.065	0.055	0.023	0.340	0.034	0.026
9	A	2	2	4	2	2	2	2	2	3	3
	H_o	0.339	0.390	0.525	0.356	0.475	0.424	0.559	0.051	0.424	0.627
	H_E	0.324	0.403	0.505	0.314	0.493	0.479	0.500	0.050	0.514	0.539
	F_{IS}	-0.046	0.032	-0.041	-0.134	0.037	0.116	-0.119	-0.026	0.176	-0.163
10	A	2	4	5	2	2	2	3	2	3	4
	H_o	0.400	0.200	0.400	0.300	0.600	0.350	0.200	0.200	0.400	0.400
	H_E	0.480	0.186	0.505	0.375	0.495	0.439	0.395	0.180	0.499	0.468
	F_{IS}	0.167	-0.074	0.208	0.200	-0.212	0.202	0.494	-0.111	0.198	0.144
11	A	2	4	5	2	2	2	3	2	3	6
	H_o	0.432	0.108	0.568	0.216	0.541	0.459	0.297	0.297	0.486	0.622
	H_E	0.456	0.104	0.581	0.307	0.482	0.407	0.360	0.290	0.481	0.555
	F_{IS}	0.051	-0.039	0.024	0.295	-0.121	-0.130	0.174	-0.026	-0.011	-0.120
12	A	2	6	5	2	2	2	3	2	3	6
	H_o	0.536	0.232	0.500	0.554	0.375	0.232	0.411	0.268	0.536	0.464

(Continued)

Table 2. (Continued)

Population		Locus									
		ARP	ALDB	GpHR	Act2α	ATPsβ	CaM	Mlc2	SRP54	LDHA	ANT
13	H_E	0.497	0.213	0.464	0.481	0.500	0.205	0.406	0.305	0.526	0.439
	F_{IS}	-0.077	-0.090	-0.078	-0.152	0.250	-0.131	-0.012	0.121	-0.018	-0.057
	A	2	3	5	2	2	2	3	2	3	5
14	H_o	0.405	0.262	0.524	0.452	0.429	0.190	0.286	0.452	0.429	0.476
	H_E	0.477	0.234	0.535	0.350	0.459	0.172	0.338	0.350	0.489	0.458
	F_{IS}	0.152	-0.117	0.021	-0.292	0.067	-0.105	0.154	-0.292	0.123	-0.040
15	A	2	5	5	2	2	2	3	2	3	6
	H_o	0.511	0.222	0.556	0.422	0.489	0.156	0.400	0.333	0.533	0.600
	H_E	0.475	0.206	0.573	0.437	0.429	0.143	0.336	0.278	0.529	0.552
16	F_{IS}	-0.075	-0.080	0.030	0.033	-0.141	-0.084	-0.190	-0.200	-0.009	-0.088
	A	2	3	5	2	2	2	2	2	3	5
	H_o	0.571	0.393	0.393	0.286	0.643	0.179	0.250	0.357	0.571	0.321
17	H_E	0.500	0.327	0.596	0.337	0.459	0.163	0.270	0.337	0.548	0.524
	F_{IS}	-0.143	-0.201	0.341	0.152	-0.400	-0.098	0.073	-0.061	-0.043	0.387
	A	2	3	5	2	2	2	3	2	3	4
18	H_o	0.514	0.257	0.429	0.486	0.543	0.200	0.571	0.457	0.629	0.514
	H_E	0.467	0.232	0.487	0.420	0.441	0.180	0.454	0.382	0.551	0.449
	F_{IS}	-0.101	-0.107	0.121	-0.156	-0.230	-0.111	-0.258	-0.197	-0.140	-0.144
19	A	2	6	5	2	2	2	3	2	3	4
	H_o	0.409	0.364	0.545	0.318	0.432	0.114	0.341	0.386	0.500	0.568
	H_E	0.474	0.339	0.549	0.397	0.456	0.146	0.334	0.407	0.468	0.493
20	F_{IS}	0.137	-0.073	0.006	0.198	0.054	0.224	-0.022	0.050	-0.069	-0.154
	A	2	4	5	2	2	2	3	2	3	5
	H_o	0.457	0.152	0.457	0.304	0.435	0.217	0.326	0.348	0.565	0.543
21	H_E	0.471	0.182	0.473	0.364	0.440	0.194	0.368	0.315	0.512	0.534
	F_{IS}	0.032	0.165	0.034	0.164	0.011	-0.122	0.115	-0.105	-0.104	-0.018

A: The number of alleles; H_o : observed heterozygosity; H_E : expected heterozygosity; F_{IS} : inbreeding coefficient

doi:10.1371/journal.pone.0127979.t002

with all localities and 7.05%, excluding admixture areas (localities 6, 10–11), than in three-group arrangements that yielded 5.17% and 6.45% with and without admixture areas, respectively. In addition, the proportion of variance explained by differences among populations within groups (F_{SC}) was four to ten times larger in the two-groups arrangements than in the three-groups arrangements, with the lowest proportion of variation (0.19%) for the three-groups arrangement without admixture areas ($F_{SC} = 0.00207$, $P > 0.05$; North Atlantic: South Atlantic: Mediterranean).

Principal coordinate analysis. The PCoA of the standardized covariance of pairwise population genetic distances provide concordant evidence of locality assignment to a minimum of three separate swordfish populations (Fig 4). The first three axes accounted for ~93.86% of the variation among populations with Eigen values of 0.847 and 0.328 for the first and second axes, respectively. Based on the clustering patterns, three groups can be identified corresponding to North Atlantic, South Atlantic, and Mediterranean populations. Loadings in the first axis contrast the Mediterranean from the South Atlantic, whereas the second axis contrasts the Mediterranean with the North Atlantic. The North Atlantic cluster, is not as tightly grouped as other populations, and is comprised of the Gulf of Mexico (1), Southeast U.S. Coast (2),




Fig 2. Identification of candidate loci under selection inferred from F_{ST} outlier analysis ($P < 0.05$) [47, 48] of ten nuclear markers using the individuals (A) from all 18 localities and (B) excluding admixture localities and including only localities that correspond to areas of known spawning. F_{ST} values for all the loci were notably greater and H_e for Act2 α , AldB, ATPs β , CaM, and Mlc2 increased when admixed samples were excluded (B) compared to the analysis with all samples (A).

doi:10.1371/journal.pone.0127979.g002

Northeast U.S. Coast (3), Lesser Antilles (4), and east of Flemish Cap (5) samples. The sample collected west of the Strait of Gibraltar (8), which for management purposes corresponds to the North Atlantic stock, cluster tightly with the Mediterranean sample (9), and together they form the Mediterranean group. Finally, the South Atlantic group included the samples from Cape Verde (12), collected north of the current boundary between northern and southern Atlantic swordfish stocks, which cluster tightly with the samples from Equatorial Brazil (13), Gulf of Guinea (14), Central South Atlantic (15), Brazil (16), Brazil-Uruguay (17), and Namibia

Table 3. Pairwise F_{ST} (lower diagonal) and corresponding significance values (upper diagonal) among the 18 swordfish localities representative of Atlantic and Mediterranean populations characterized in this study (see Table 1).

	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18
1		0.401	0.319	0.438	0.272	*	0.161	***	***	0.112	**	***	***	***	***	***	***	***
2	0.000		0.108	0.106	*	**	0.190	***	***	0.118	**	**	***	***	***	***	***	***
3	0.001	0.006		0.470	0.419	0.226	0.074	***	***	0.393	*	***	***	***	***	**	***	***
4	0.000	0.006	0.000		0.429	0.250	0.380	***	***	0.385	0.055	***	***	***	***	**	***	***
5	0.003	0.018	0.000	0.000		0.411	0.188	***	***	0.410	0.089	**	*	***	*	*	**	*
6	0.018	0.033	0.006	0.004	0.000		0.078	***	***	0.428	0.325	**	*	**	*	*	**	0.080
7	0.005	0.003	0.007	0.000	0.007	0.013		***	***	0.423	0.279	***	**	**	**	*	**	**
8	0.066	0.063	0.064	0.042	0.074	0.070	0.038		0.158	**	***	***	***	***	***	***	***	***
9	0.069	0.077	0.074	0.049	0.085	0.076	0.050	0.003		***	***	***	***	***	***	***	***	***
10	0.009	0.010	0.001	0.001	0.001	0.000	0.000	0.036	0.052		0.421	0.226	0.318	0.138	0.224	0.179	0.090	0.411
11	0.019	0.025	0.014	0.010	0.015	0.003	0.002	0.054	0.069	0.000		**	0.203	0.084	0.104	0.063	0.103	0.291
12	0.053	0.031	0.037	0.045	0.042	0.044	0.019	0.086	0.114	0.005	0.021		0.079	*	0.064	0.134	**	*
13	0.047	0.045	0.035	0.039	0.031	0.023	0.020	0.097	0.117	0.002	0.004	0.008		0.417	0.443	0.420	0.415	0.412
14	0.044	0.040	0.036	0.040	0.039	0.026	0.023	0.105	0.121	0.008	0.009	0.011	0.000		0.382	0.409	0.394	0.421
15	0.049	0.049	0.043	0.044	0.043	0.023	0.025	0.100	0.117	0.005	0.009	0.010	0.000	0.000		0.301	0.413	0.429
16	0.050	0.040	0.032	0.035	0.031	0.025	0.018	0.085	0.106	0.007	0.012	0.005	0.000	0.000	0.002		0.405	0.412
17	0.045	0.039	0.041	0.041	0.045	0.031	0.022	0.100	0.117	0.011	0.008	0.015	0.000	0.000	0.000	0.000		0.399
18	0.044	0.042	0.034	0.034	0.033	0.014	0.017	0.089	0.106	0.001	0.002	0.012	0.000	0.000	0.000	0.000	0.000	

Note: Significant F_{st} in bold

* = $P < 0.05$

** = $P < 0.01$

*** = $P < 0.001$

doi:10.1371/journal.pone.0127979.t003

Table 4. Matrix of migration (Nm) for North Atlantic (NA) (n = 155), Mediterranean (MED) (n = 142) swordfish, and South Atlantic (SA) (n = 256) as calculated in Arlequin v3.5.

	NA	MED
MED	7.2144	
SA	8.5915	4.8516

doi:10.1371/journal.pone.0127979.t004

(18). The remaining samples occupy an intermediate position, with central North Atlantic (6) and Iberian (7) samples associating more closely with the North Atlantic group, whereas Morocco (10) and Western Sahara (11) samples more closely with the South Atlantic group.

Bayesian genetic clustering analyses

STRUCTURE analysis. Evaluation of the mean posterior probabilities from multiple STRUCTURE analyses ($K = 1-10$) revealed that the mean $\ln P(D)$ increased from $K = 1-3$ and then sharply decreased when $K \geq 4$ (see supporting [S1 Fig](#)) indicating that $K = 3$ reflects the majority of the genetic structure in the dataset [55]. Estimated ΔK 's revealed two high ΔK peaks of similar magnitude corresponding to $K = 2$ ($\Delta K_2 = 35.887$) and $K = 3$ ($\Delta K_3 = 33.6548$), followed by very low ΔK (< 5.0) values (see supporting [S1 Fig](#)). Mediterranean localities (8–9) were removed from the analysis to test whether the strong differentiation of the Mediterranean from the Atlantic was causing an underestimation of K . The evaluation including only North Atlantic and South Atlantic localities identified a single ΔK peak at $K = 2$ (not shown) congruent with the genetic heterogeneity between the North and South Atlantic populations of swordfish. While Evanno's ΔK method seeks to detect the uppermost hierarchical level of population structure the method is less reliable at lower levels of genetic differentiation and may underestimate K [64]. Therefore, the estimation of K using the *ad hoc* evaluation of mean posterior probabilities from multiple analyses of K [55] appears to be more appropriate in this instance than the ΔK approach [60].

Average population posterior probability memberships (\bar{Q}) of the 18 localities (see supporting [S4 Table](#)) and individual assignments at $K = 3$ were concordant with results from PCoA, AMOVA, and F_{ST} analyses ([Fig 5](#)). Localities 1–5 had average values of ancestry probabilities (\bar{Q}) belonging to the North Atlantic > 0.90 with no individuals of South Atlantic or Mediterranean origin. Swordfish collected in the central North Atlantic (6) clustered between the North Atlantic and the South Atlantic indicating a potential mixing area, however sample size was small ($n = 20$). The Iberian (7) sample clustered slightly closer with the North Atlantic ($\bar{Q} = 0.55$) and represented an area of mixing with at least one individual belonging to the Mediterranean with the rest either to the North Atlantic or the South Atlantic. No North Atlantic or South Atlantic swordfish were identified within the Mediterranean sample (9) and the area west of the Strait of Gibraltar (8) clustered ($\bar{Q} = 0.85$) with the Mediterranean (9). In fact, 77 out of 83 individuals sampled (93%) immediately to the west of the Strait of Gibraltar (8) were identified by STRUCTURE as Mediterranean. Of the remaining individuals, three swordfish (3.6%) were identified as North Atlantic and another three (3.6%) as South Atlantic. Four Mediterranean swordfish were also identified in the Iberian sample (7) as follows: three at 37.5°N 12.5°W and one at 37.5°N 22.5°W, accounting for roughly 11% and 5% of the total number of individuals sampled respectively in these two locations.

The area off the Atlantic coast of Morocco (10) was also identified as an area of mixing with individuals of North Atlantic and South Atlantic ($\bar{Q} = 0.52$) origin. Although the sample of Western Sahara (11) contained a majority of individuals assigned to the South Atlantic ($\bar{Q} = 0.57$), it is a zone of admixture containing also individuals of North Atlantic origin. Swordfish




Fig 3. Slatkin's [53] linearized F_{ST} values of 18 localities of Atlantic and Mediterranean swordfish. Population numbers correspond to the sampled localities in Fig 1.

doi:10.1371/journal.pone.0127979.g003

from Cape Verde (12) along with all the individuals sampled south of 5°N (13–18) were assigned to the South Atlantic ($\bar{Q} > 0.95$), with no North Atlantic or Mediterranean migrants. STRUCTURE results using only *CaM* revealed that while this locus drives the heterogeneity between South Atlantic and North Atlantic populations, alone it fails to differentiate between North Atlantic and Mediterranean populations. Conversely, Bayesian analysis using all the loci except *CaM* differentiated between North Atlantic and Mediterranean populations. Further, this analysis also discriminates North Atlantic from South Atlantic swordfish, although posterior probabilities were higher when *CaM* was included (see supporting S2 Fig).

GENELAND analysis. The K value with the greatest density on the MCMC chain after a burn-in of 10,000 was $K = 4$. However, inspection of map of posterior probabilities identified one cluster as a 'ghost' with no associated samples (not shown). Further, this ghost cluster was unstable, as its placing or presence differed among replicate runs. The existence of 'ghost'

Table 5. Hierarchical analysis of molecular variance (AMOVA) of SNP data for 18 samples of swordfish.

Genetic Structure	All localities			Excluding admixture areas		
	Variance component	% of total	Fixation index	Variance component	% of total	Fixation index
Two Groups						
	G1: Atlantic (1–7,10–18)			G1: Atlantic (1–5, 12–18)		
	G2: Mediterranean (8–9)			G2: Mediterranean (8–9)		
Among groups	0.14991	6.63	$F_{CT} = 0.06634^{**}$	0.16037	7.05	$F_{CT} = 0.07046^{**}$
Among populations within groups	0.03691	1.63	$F_{SC} = 0.01749^{***}$	0.04373	1.92	$F_{SC} = 0.02067^{***}$
Among individuals within populations	0.03735	1.65	$F_{IS} = 0.01802^{ns}$	0.02306	1.01	$F_{IS} = 0.01113^{ns}$
Within individuals	2.03553	90.08	$F_{IT} = 0.09920^{***}$	2.04876	90.02	$F_{IT} = 0.09981^{***}$
Three Groups						
	G1: North Atlantic (1–7,10–11)			G1: North Atlantic (1–5)		
	G2: Mediterranean (8–9)			G2: Mediterranean (8–9)		
	G3: South Atlantic (12–18)			G3: South Atlantic (12–18)		
Among groups	0.11349	5.17	$F_{CT} = 0.05171^{***}$	0.14314	6.45	$F_{CT} = 0.06450^{***}$
Among populations within groups	0.00834	0.38	$F_{SC} = 0.00401^{**}$	0.00430	0.19	$F_{SC} = 0.00207^{ns}$
Among individuals within populations	0.03735	1.70	$F_{IS} = 0.05171^{ns}$	0.02306	1.04	$F_{IS} = 0.1113^{ns}$
Within individuals	2.03553	92.75	$F_{IT} = 0.07253^{***}$	2.04876	92.32	$F_{IT} = 0.07683^{***}$

ns = $P > 0.05$

* = $P < 0.05$

** = $P < 0.01$

*** = $P < 0.001$

Samples (in parentheses) were grouped into alternative groups (G1-G3) according to region of capture, with all localities, or excluding localities identified as areas of admixture (see text for explanation)

doi:10.1371/journal.pone.0127979.t005

clusters has been reported as an artifact of GENELAND which tends to overestimate K in a comparison to other Bayesian genetic clustering algorithms [65]. Accordingly, GENELAND was run with the optimal value ($K = 3$) identified by STRUCTURE. Localities 1–5 and part of 7 comprised the North Atlantic ($\bar{Q} > 0.90$), localities 8 and 9 the Mediterranean ($\bar{Q} > 0.90$), and localities 12–18 the South Atlantic ($\bar{Q} > 0.90$). The individual posterior probabilities assigned in GENELAND (Fig 6) identified the region corresponding to the central North Atlantic (6) and Western Sahara (11) samples as areas of mixing between the North Atlantic and South Atlantic populations. The samples corresponding to the North Atlantic area west of the Strait of Gibraltar (8) have the genetic signature of the western Mediterranean (9), whereas swordfish from the Atlantic locality of Morocco (10) were identified as either Mediterranean or North Atlantic. The Iberian sample (7) is comprised of three subsamples located at 37.5°N 22.5°W ($n = 19$), 37.5°N 17.5°W ($n = 5$), and 37.55°N 12.5°W ($n = 27$), with a North Atlantic origin assigned by GENELAND of $> 80\%$, $> 70\%$, and $< 50\%$, respectively. Since the precise geographic coordinates of capture for the individuals in these Atlantic subsamples were recorded only with a precision of 2.5° latitude by 2.5° longitude, and because the subsample sizes are small, these estimates are imprecise.

Discussion

Nuclear loci evaluation

Genetic population structure studies using nuclear DNA have traditionally excluded adaptive markers to avoid introducing bias to estimates of migration rates and effective population sizes




Fig 4. Principal coordinate analysis (PCoA) [54] of 18 localities of Atlantic and Mediterranean swordfish. The numbered population means, corresponding to the sampling localities in Fig 1, are identified by their *current management stock* (yellow triangle = Mediterranean, blue diamond = North Atlantic, red circle = South Atlantic) as defined by the ICCAT management boundaries at 5°N and the Strait of Gibraltar. Colored circles of equal diameter are drawn to display population groupings, red = South Atlantic, blue = North Atlantic, yellow = Mediterranean.

doi:10.1371/journal.pone.0127979.g004

[66]. In this study we attempted to identify variation in non-coding nuclear introns to maximize the potential of characterizing neutral loci. Of the 26 SNPs assayed 25 were contained in the introns of 10 nuclear genes, with the exception of a SNP corresponding to a third codon synonymous substitution in one exon of the ATPsβ gene. Although introns are generally






Fig 5. Bayesian individual assignment of Atlantic and Mediterranean swordfish in STRUCTURE v2.3 [55] using no admixture, correlated alleles, and LOCPRIOR models and inferred from 50 independent runs of $K = 3$ using 100,000 MCMC iterations and a burn-in period of 100,000. Estimated individual membership coefficients (\hat{q}) are sorted by sampling locality and correspond to the numbered localities in Fig 1.

doi:10.1371/journal.pone.0127979.g005

A) Map of posterior probability to belong to SATl



B) Map of posterior probability to belong to NWAtl



C) Map of posterior probability to belong to MED




Fig 6. Posterior probability contour maps of Atlantic and Mediterranean swordfish calculated in GENELAND [62] with an uncertainty on coordinate value = 30°, correlated allele frequency and spatial models, and inferred from 20 independent runs of $K = 3$ using 100,000 MCMC iterations and a thinning of 100. Posterior probability contours range from 1.0 in light yellow to 0.1 in red for membership to the (A) South Atlantic population vs. all other populations, (B) North Atlantic vs. all other populations, and (C) Mediterranean vs. all other populations. Black circles correspond to sampling locations and may indicate more than one individual. Posterior probability contours in areas of limited and/or no sample coverage are extrapolations and may therefore contain error (e.g. no northern Portugal sampling extends the Mediterranean posterior probability of membership).

doi:10.1371/journal.pone.0127979.g006

assumed to be neutral, some may contain transcription regulation or splicing control elements that alter gene expression [67]. Introns may also experience purifying and/or positive selection [68] or the effects of hitch-hiking (i.e. linkage between a non-coding variant and a gene under selection) as reported in Atlantic cod [69]. Therefore deviations from HWE, linkage disequilibrium tests, and outlier analysis are commonly applied to detect potentially adaptive loci. In our study exact HWE [42] and linkage equilibrium [43] tests were not significant after Bonferroni corrections. Similarly the the F_{ST} outlier analysis failed to identify any loci characterized as subject to selection. Given that the number of loci analyzed in the current study is small ($n = 10$), the associated error in the calculation of global neutral F_{ST} could be large. Accordingly, future studies utilizing genomic technologies will certainly yield greater numbers of loci to conduct more conclusive outlier tests.

Alternatively, outlier and non-neutral SNPs can provide insight into adaptive genetic diversity within and among populations [70, 71] which may be useful for resolving fine-scale population connectivity, admixture, mixed stock analysis, and identifying local adaptation for conservation or resource management [72–74]. Studies utilizing adaptive loci in Atlantic cod [75], Atlantic herring [74, 76], Atlantic salmon [77], and Pacific lamprey [78] reported that fewer loci and smaller sample sizes were required to detect population differentiation in marine fishes when using outlier loci as compared to neutral loci. Muths, Le Couls (79) hypothesized that the inclusion of adaptive loci in future genetic analyses of large pelagic fishes, specifically swordfish, may identify additional evolutionary significant population units than analyses relying on only neutral loci. However, since the inclusion of outlier loci may violate model assumptions used for individual assignment, it is important that outlier loci included in such analyses are carefully evaluated and presented with comparative results using neutral loci [74]. While the ranking analysis identified *CaM* as the most informative locus for distinguishing North and South Atlantic swordfish populations, this locus alone fails to differentiate between North Atlantic and Mediterranean swordfish populations. Further, STRUCTURE results without *CaM* (see supporting S2 Fig) show that the remaining loci were capable to discriminate between North and South Atlantic swordfish. Accordingly, the high resolution of population differentiation reported in this study among North Atlantic, South Atlantic and Mediterranean swordfish cannot be attributed to *CaM* alone. In fact, we demonstrate that the inclusion of the additional loci increases the power of population discrimination.

Atlantic swordfish population differentiation

Analyses of multilocus nuclear SNP data indicate that Atlantic swordfish can be subdivided into at least three populations, namely Mediterranean, North Atlantic, and South Atlantic. The presence of these populations is supported by linearized pairwise F_{ST} [52, 53], AMOVA [50], PCoA [54], and Bayesian genetic clustering using both in STRUCTURE [55, 57, 59] and GENELAND [62]. Though no geographic barriers prevent gene flow among North Atlantic, South Atlantic and Mediterranean swordfish populations, the number of migrants among populations is small (Table 4). Additionally, the observed temporal stability in allele frequencies

within the Northwest Atlantic, South Atlantic, and Mediterranean populations also suggests limited gene flow among these swordfish populations. This temporal stability is perhaps best evident in the parity of genetic signature between the larval Gulf of Mexico samples and the samples of adult Northwest Atlantic swordfish collected over a decade prior. Based on these results the North Atlantic population extends from the tropical spawning areas in northwestern Atlantic waters [80, 81] to 50°W. In the feeding areas north of 40°N the range of the North Atlantic population extends to the Azores and may extend at that latitude as far east as 15°W. The South Atlantic population extends from 50°S through the South Atlantic tropical and equatorial spawning areas [17] to 20°N at 40°W and 25°N along the African coast.

The Mediterranean population extends past the Strait of Gibraltar, possibly as far as 8°W, with little evidence of Atlantic swordfish entering the Mediterranean Sea. These results are concordant with the phylogeographical association of mtDNA lineages previously reported [7, 13, 17, 56] and also with evidence derived from fisheries data [82]. The Mediterranean population is identified here as the most divergent of the three populations, and these findings agree with other studies using mtDNA, scnDNA (see [25] for summary), and microsatellite loci [29, 30].

The Bayesian clustering analyses based on multilocus SNPs presented here are not consistent with analyses of global swordfish population structure using microsatellites [28, 30]. Those studies reported extremely low levels of differentiation between North and South Atlantic populations ($F_{ST} \approx 0.0015$), precluding individual cluster assignment. By contrast, Kasapidis et al. [29] argued that the considerably higher estimates of differentiation reported with nuclear SNPs, and specifically with *CaM*, is strong adaptive selection. Such explanation, however, fails to explain the concordance between mtDNA and nuclear SNPs data (reviewed in [25]), including the current study. Instead, such agreement suggests that the differentiation between North Atlantic and South Atlantic swordfish is the consequence of population subdivision and low levels of gene flow.

The shallow levels of differentiation between North and South Atlantic using microsatellites could be due to allele size homoplasy in combination with large effective sizes [83] of swordfish populations and the high mutation rates characteristic of these type of loci [84]. Alternatively, the failure to find differences using microsatellite analyses within the Atlantic [28–30] could be due to the small number of loci (4–10) characterized in those studies, or because the samples employed to represent the North Atlantic came from areas identified here as admixture zones within the Northeast Atlantic. Accordingly, a stronger signal of differentiation between South Atlantic and North Atlantic swordfish may be obtained by increasing the number of microsatellite loci aimed to characterize representative samples from reproductive areas in the Northwest Atlantic. Yet, in other highly migratory marine fishes the strong differentiation revealed with mtDNA has not been obtained with microsatellites, and such discordance was interpreted as evidence of male mediated gene flow in bigeye tuna (*Thunnus obesus*) [16], bluefin tuna (*T. thynnus*) and yellowfin tuna (*T. albacares*) [85], sandbar shark (*Carcharhinus plumbeus*) [86], and shortfin mako (*Isurus oxyrinchus*) [15], without considering alternative hypotheses. In blue marlin (*Makaira nigricans*) mitochondrial cytochrome b gene sequences [87] and RFLP data of the mtDNA molecule [88] revealed a pronounced signal of inter-oceanic differentiation that was considerably higher compared to allozyme and scnDNA data [89]. Such disparity was attributed to the enhanced effect of genetic drift under migration-drift equilibrium resulting from the fourfold lower effective population size of mitochondrial than nuclear molecules, and while the effect of male biased inter-ocean exchange could not be rejected, it was not required to account for the observed disparity. Computer simulations and power analyses conducted later on the blue marlin allozyme and scnDNA data together with newly generated microsatellite and whole molecule mtDNA RFLP data [90], indicated that the large range of F_{ST} values among marker classes and among loci within a marker class were consistent with the same

population level $N_e m$, given differences in both mode of inheritance and genomic sampling variance among estimates. Further, one microsatellite locus (Mn8) yielded a pattern concordant with mtDNA regarding both the strong inter-oceanic differentiation, and inferred unidirectional gene flow of Indo-Pacific blue marlin into the South Atlantic. In swordfish, the similarity in levels of differentiation between North and South Atlantic populations using mtDNA ($F_{ST} = 0.0244$; [17]) and multilocus nuclear SNPs ($F_{ST} = 0.0393$; this study), is not consistent with gender-biased gene flow. Considering that both swordfish and blue marlin females attain larger sizes than males and are more abundant at higher latitudes [81, 91, 92] where inter-oceanic gene flow would be facilitated (e.g., around southern Africa), it would be highly unlikely that male biased dispersal explains the observed inter-oceanic allele frequency discrepancies among nuclear and mitochondrial markers in both species. Further, in the Atlantic the differences between the sex ratios at-size seem to support the hypothesis that mature swordfish females are more mobile, tending to cover (in terms of probability) broader geographic areas to sustain the energetic requirements required to support the high production of eggs, whereas the migration of males would be generally more restricted and conditioned by the lower average body biomass and the lower energy requirements in comparison to the females to produce spawning products [92]. Thus in spite of the higher mobility and gene flow potential of females no gender-biased gene flow was detected in Atlantic swordfish.

Atlantic swordfish population subdivision and admixture

Previous studies of Atlantic swordfish population structure only broadly identified potential areas of admixture between the North Atlantic and Mediterranean populations [7, 31] and between the North Atlantic and South Atlantic populations [17, 18, 26] and offered no estimates of the contribution of each population. The Strait of Gibraltar is the current management boundary utilized by ICCAT to separate the Mediterranean and the North Atlantic populations. Multilocus analyses of SNPs with STRUCTURE (Fig 5) and GENELAND (Fig 6C) suggests that the Mediterranean stock extends past the management boundary to around 10°W into North Atlantic waters. However, given that the mapping algorithm of GENELAND extrapolates posterior probabilities into areas of limited or no sample coverage some of the posterior probability curves should be viewed with caution. Specifically, since no samples off the northern coasts of Portugal or Spain were characterized, the posterior probability curves in this region should not be interpreted as evidence of the Mediterranean population extending around the Iberian Peninsula into the Bay of Biscay and the Celtic Sea (Fig 6C). Taken all together, Mediterranean swordfish exiting through the Strait of Gibraltar appear to remain primarily in the shallower Atlantic shelf waters off the northwest African and Iberian coasts, with occasional migrants venturing farther into North Atlantic waters, perhaps as far as the Azores [30, this study]. Movements of mature swordfish inhabiting the shelf area west of the Strait of Gibraltar into the Mediterranean, putatively associated with spawning behavior around the Balearic Islands, occur from May to the beginning of July [82]. Later in the year, between August-November a migration associated with feeding behavior in the opposite direction (east to west) across the Strait takes place with no reproductively active swordfish and with 15% of the catch corresponding to juveniles [82]. Given that our samples are limited to summer and fall when migrations are believed to take place, the expected reduction in frequency of Mediterranean swordfish in this Atlantic region during the winter and spring cannot be confirmed at this time. In a previous study using mtDNA CR-I sequence data, Alvarado Bremer et al. [93] documented no Atlantic swordfish east of the Strait of Gibraltar. Similarly, in this study, no Atlantic swordfish were identified within the Mediterranean Sea ($n = 59$). It should be noted that in this study Mediterranean swordfish were only sampled in the western Mediterranean,

specifically in the area east of the Alboran Sea and south of the Balearic Islands, where the presence of swordfish juveniles has been documented [94] and reproductive behavior is expected to occur [82]. If North Atlantic swordfish were to enter the Mediterranean, their presence would be best documented in this region. Further, the low frequency of movements of Atlantic swordfish into the Mediterranean is also supported by the Bayesian analysis of microsatellite data from 602 western and eastern Mediterranean swordfish in which only three individuals of putative Atlantic origin were identified [30]. Altogether it appears that oceanographic features and or behavioral cues limit the extent of admixture of Atlantic and Mediterranean swordfish populations, with the majority of the contact confined to the shallower waters of both the northwest African shelf and the Gulf of Cadiz.

While the Strait of Gibraltar and associated oceanographic and geophysical features may constrain the admixture between North Atlantic and Mediterranean populations, no such physical barrier separates the populations of North Atlantic and South Atlantic swordfish with admixture occurring over a considerably broader geographic area. A proposal to move the current management boundary separating the North Atlantic from the South Atlantic populations from 5°N to about 15°N was suggested by Chow et al. [26] on the basis of significant differences in allele frequency of *CaM*. Our analyses with PCoA (Fig 4), STRUCTURE (Fig 5), and GENELAND (Fig 6) are in agreement with the extension of the South Atlantic swordfish stock past the current 5°N boundary but suggest an extension even farther north, past Cape Verde (12) to around 20°N. This boundary does not conform to a horizontal line extending from the Caribbean Sea to the African coast. Instead, the 0.90 isoline of GENELAND is more akin to a step extending from about 0°N 45°W north to 25°N 45°W, and from there almost horizontally towards the African coast (Fig 6). To the north and west of this boundary, an admixture zone begins around Western Sahara (11) and continues north to Morocco (10) and the Iberian sea (7) extending west towards the central North Atlantic (6) and then south towards the northern coast of Brazil. An interesting feature of the pattern of admixture between northern and southern stocks is that the relative proportion contribution to neighboring populations is asymmetrical, with South Atlantic swordfish moving past this boundary towards the north, but with no swordfish of North Atlantic origin moving into the range of the South Atlantic population, reminiscent of the asymmetric contact between the North Atlantic and Mediterranean swordfish populations.

Posterior probability membership contours generated by GENELAND (Fig 6) could be used for mixed stock allocation in the identified areas of stock admixture, with the caveat that the sensitivity of spatial Bayesian clustering, particularly when identifying linear boundaries, is contingent upon sampling coverage [95] as well as initial model assumptions. However, altering initial model assumptions did not alter the general pattern of genetic differentiation reported (not shown). Seasonal displacements of North and South Atlantic swordfish, as well as sampling in regions not properly represented here (e.g., 20–25°N and 30–40°W), may change the extent and geographical placement of admixture, but are not expected to alter the general pattern of differentiation separating these two populations (see notes below regarding reduced abundance in these areas derived from catches).

Potential barriers to gene flow

In the absence of a geographic barrier to explain the asymmetric pattern of admixture separating north and south populations of Atlantic swordfish, biological, behavioral, and oceanographic impediments to gene flow need to be examined. Conventional tag-recapture data [20, 96], though biased in its coverage to primarily the Northwest Atlantic fishery, indicates that swordfish tagged west of the Azores generally follow similar patterns of directional migration,

west by southwest, back to the primary breeding grounds of the North Atlantic indicative of spawning and feeding ground fidelity, whereas swordfish tagged east of the Azores (of which there are far fewer records) tend to move east by southeast. Interestingly, of the few swordfish tagged between 15°N and 30°N, movement generally leads east by northeast. While conventional tagging data may mirror some of the same molecular distribution trends inferred in this study, these data are heavily biased to areas of high fishing effort [20] and the reporting tag-recapture rates among fleets. Although limited by their relative short duration, Pop-up satellite archival tag (PSAT) data has yielded similar evidence of seasonal spawning and feeding ground migrations in the Northwest Atlantic [22, 23].

Recently, Abascal et al. [24] reported the PSAT tracks for 10 Atlantic swordfish tagged in the North Atlantic as part of a tagging study aimed to characterize diel vertical diel behaviors and horizontal seasonal migrations and habitat preferences of this species. Seven tracks corresponded to swordfish tagged around 35°N–45°N in latitude and 40°–50°W in longitude. The other three tracks were of swordfish tagged in a southeastern area of the North Atlantic to the north of Cape Verde archipelago, around 20°N–23°N in latitude and 20°W–25°W in longitude. The movements of these 10 swordfish are concordant with the results obtained with STRUCTURE, with four tracks indicating movements within areas (two tracks each) identified here as North Atlantic (localities 3 and 5) or South Atlantic (locality 12) populations, whereas six swordfish moved in and out of sampling areas where the posterior probabilities suggest admixture zones (i.e., locality 6, for the fish tagged in the northern area, and 6, 7 and 11, for fish tagged in the southern area). However, one PSAT deployed in the northern area after spending several months in the central North Atlantic admixture zone, spent the last six months of a year at liberty within locality 12 (Cape Verde), identified here as an area belonging to the South Atlantic population. Altogether, only one of 10 swordfish tracked were in conflict with the probability assignments (>0.95) of STRUCTURE. By contrast, concordance of the PSAT tracks with GENELAND's North-South Atlantic population boundaries (0.10–0.90 confidence limit) occurred in seven out of ten swordfish. Two swordfish tagged in the northern area, crossed the 0.90 isoline of assignment to the South Atlantic population. Conversely, one of three tracks of swordfish tagged in the southern area crossed the 0.90 confidence isoline delimiting the North Atlantic population. However, none of the swordfish tagged either in the northern or southern areas of the North Atlantic crossed into the respective 1.0 probability isolines delimiting the North Atlantic and South Atlantic swordfish populations, respectively.

Within-ocean differences in patterns of genetic differentiation

Strong natal fidelity, with admixture primarily confined to feeding grounds, has been proposed as an explanation for the unparalleled intra-oceanic differentiation separating North Atlantic and South Atlantic swordfish populations [17]. By contrast, within the Pacific Ocean [97] and along the Mediterranean Sea [31] differentiation of mtDNA appears to conform to isolation by distance (IBD), although such a pattern in the Pacific Ocean is not supported by the multi-locus analysis of nDNA SNP data [98]. Conversely, in the Indian Ocean the presence of at least two potential genetic populations was suggested [18, 99, 100], but this interpretation was rejected recently [79] in favor of panmixia since the frequency distribution of mitochondrial and microsatellite markers using a more ample sampling coverage of that basin including multiple seasons was homogeneous. Further Muths et al. [79] propose that the presence of distinct swordfish populations respectively within the Atlantic and Pacific Oceans could be partly explained by the existence in those basins of two separate hemispheric oceanographic systems with corresponding discrete feeding areas for the respective populations with limited or no admixture. The association of genetic differentiation with the existence of separate feeding areas

as proposed by Muths et al. [79] contrasts dramatically with the interpretation of spawning site fidelity as the driving force behind the observed genetic heterogeneity of swordfish populations. Accordingly, maximum levels of genetic differentiation using both mtDNA and nuclear SNP data were obtained when separate breeding areas were compared [17]. In the current study restricting the analysis to spawning areas yielded higher F_{ST} values for all loci while at the same time increased the H_e values for several loci (e.g. *Act2 α* , *CaM*, *Mlc2*). This result supports the hypothesis that samples from feeding grounds in both the NW Atlantic and the South Atlantic Ocean (i.e., Gulf of Guinea) may be more prone to population admixture as suggested by Alvarado Bremer et al. [17]. Similarly, spawning site fidelity is also supported by the evidence of the admixture of North Atlantic and Mediterranean swordfish populations in the feeding area to the west of the Strait of Gibraltar by both mtDNA [25] and nDNA data (this study). The concordance of mtDNA and nDNA data was also reported for the separation of South Atlantic from Indian Ocean swordfish [79]. Such concordance of nuclear and mtDNA data in the separation of swordfish populations inhabiting the Mediterranean, North Atlantic, South Atlantic and Indian Ocean basins, is particularly relevant given reported differences in rates of dispersal of swordfish males and females.

While fidelity to spawning and feeding grounds may explain low levels of gene flow between North and South Atlantic swordfish, it doesn't fully explain why admixture only seems to occur in the Northeast Atlantic waters east of about 50°W and south of 30°N. Chow et al. [26] suggested that changes in oceanographic parameters (e.g. salinity, dissolved oxygen, temperature, etc.) in deep waters might explain the strong genetic break that their *CaM* data revealed since swordfish are deep divers. Nutrient upwelling, caused by prevailing winds off the western coast of Africa, creates a large hypoxic zone in the eastern tropical Atlantic [101] that apparently correlates with the inferred area of population differentiation identified in this study (see supporting S3 Fig). Sensitivity to oceanographic conditions in deep waters has been documented in other billfish [102] and swordfish are known to dive substantially deeper than other billfish [21], and thus may be more sensitive to changes in concentrations in DO at depth, which might be used as a cue by North Atlantic swordfish not to move south to the areas occupied by South Atlantic swordfish.

Other oceanographic features are known to act as barriers to dispersal in proximity to coastal areas. For instance while recognizing that the Amazon discharge would not be expected to affect the movements of adult highly migratory pelagic fish, Chow et al. [26] hypothesized it may act as a freshwater barrier to passive larval dispersal between the Caribbean and Brazilian provinces. This feature geographically coincides with the posterior probability break between North Atlantic and South Atlantic populations identified by STRUCTURE and GENELAND, as well as an inferred reported break in potential spawning areas [17] (Fig 7). Further, the same region coincides with areas where historical catches of adult swordfish are low, and if catches properly reflect abundance (Fig 6), then the low density of adults in this region together with the Amazon discharge preventing the admixture of larvae from northern and southern breeding areas, as hypothesized by Chow et al. [26] could together promote the observed patterns of genetic differentiation between northern and southern populations of swordfish. Finally, the population structure of Atlantic swordfish may be influenced by sea surface currents as reported in other pelagic species (e.g. blue marlin [103]). The North Atlantic gyre, the Canary current, and the Atlantic equatorial counter current all associate with the population differentiation of North Atlantic and South Atlantic swordfish, and may explain why population admixture is primarily confined to Northeast Atlantic waters.




Fig 7. GENELAND map of posterior probability of membership to the South Atlantic population overlaid the regions of reproduction of Atlantic swordfish summarized by Alvarado Bremer et al. [17]. Shaded in dark blue are the Northwest Atlantic and South Atlantic areas where eggs and or early life history stages have been collected. The areas shaded in lighter blue correspond to areas where females with high-gonadal indices and/or hydrated oocytes have been collected that coincide to areas of biased male to female sex-ratios [104].

doi:10.1371/journal.pone.0127979.g007

Conclusions and Future Studies

The findings of this study can be summarized as follows: first, significant differentiation among Mediterranean, North Atlantic, and South Atlantic swordfish populations is confirmed with very low rates of gene flow, particularly when compared to other highly migratory fishes. Second, population admixture between the North Atlantic and the South Atlantic population, and to a much lesser extent with the Mediterranean, is primarily confined to the Northeast Atlantic and in both cases appears to be unidirectional, from the Mediterranean to the NE Atlantic and from the South Atlantic to the NE Atlantic. Third, the genetic partitioning of these populations differs from currently recognized management boundaries. Fourth, and lastly, while the oceanographic and/or behavioral barriers to gene flow are not yet identified, future tagging, genetic, oceanographic, or physiological studies should focus efforts to areas of admixture in the Northeast Atlantic and areas where individuals were assigned with high posterior probabilities, i.e. North and South Atlantic spawning and feeding grounds. Additionally, future genomic studies utilizing large numbers of SNPs may provide insight into whether adaptive genetic variation is responsible for the fine-scale population structure here reported.

Supporting Information

S1 Fig. Bayesian cluster (K) estimation. The estimation of the number of clusters (K) in the STRUCTURE v2.3 [55] analysis of Atlantic and Mediterranean swordfish using (A) the ad hoc approach of Prichard et al. [55] and the mean posterior probability of the data (L(K)) and (B) ΔK approach of Evanno et al. [60]. Twenty independent runs for each K value (1–10) were performed using 100,000 MCMC iterations with a burn in period of 100,000.
(TIF)

S2 Fig. Relative importance of CaM in STRUCTURE analysis. The STRUCTURE v2.3 individual assignments of 774 swordfish from 18 localities using the no admixture and correlated allele model and the location prior with A) only the CaM locus, B) all loci except the CaM locus, and C) all loci including the CaM locus.
(TIF)

S3 Fig. Genetic partitioning in relation to the South Atlantic OMZ. The oxygen minimum zone (OMZ) in the South Atlantic swordfish feeding grounds at 100 m depth. The annual mean dissolved oxygen (DO) at 100 m depths in mL L⁻¹ are from the World Ocean Atlas 2009 (Garcia et al. 2009)¹. The OMZ depicts a similar boundary as the posterior probability of membership contour maps in Fig 5. ¹Garcia HE, Locarnini RA, Boyer TP, Antonov JI, Baranova MM, et al., editors (2010) World Ocean Atlas 2009, volume 3: dissolved oxygen, apparent oxygen utilization, and oxygen saturation. Washington, D.C. 344 p.
(TIF)

S1 Table. Allele Frequency Distribution. The allele frequency distribution within the eighteen sampled localities for each of the ten nuclear loci.
(XLS)

S2 Table. Locus power of assignment. A consensus power of assignment for each locus using WHICHLOCI v.1.0 between North Atlantic (n = 207) and Mediterranean (n = 142) swordfish inferred in STRUCTURE v2.3.
(XLS)

S3 Table. Locus power of assignment for the South Atlantic population. A consensus critical population (South Atlantic) power of assignment for each locus using WHICHLOCI v.1.0 between North Atlantic (n = 207) and South Atlantic (n = 297) swordfish inferred in STRUCTURE v2.3.
(XLS)

S4 Table. Structure population posterior probability memberships. The average population posterior probability membership (Q) of 18 localities in the Atlantic and Mediterranean inferred in STRUCTURE v2.3.
(XLS)

S5 Table. SNP allelic data. The alleles for the ten loci and all 774 swordfish analyzed in this study. Allelic numbering corresponds to reported GenBank alleles for all loci except ALDB. Allelic numbering for ALDB corresponds to STR amplicon size scoring.
(XLSX)

Acknowledgments

The authors would like to dedicate this study to the memory of Captain Langille K. Dixon of Wood's Harbour, Nova Scotia without whom much of the sampling would not have been

possible. Additional special thanks to L. Janson, Dennis Lee, Jaime Mejuto, Jay Rooker, Bernard Stequert, Marc Griffiths, Favio Hazin, Carla Marques, Carles Pla, NMFS-SEFSC, and Instituto Español de Oceanografía for providing swordfish tissue samples for analysis.

Author Contributions

Conceived and designed the experiments: BLS C-PL JRAB. Performed the experiments: BLS C-PL. Analyzed the data: BLS. Contributed reagents/materials/analysis tools: JV BG-C S-YY JRAB. Wrote the paper: BLS JRAB.

References

1. Avise JC. Conservation genetics in the marine realm. *J Hered*. 1998; 89(5):377–82. doi: [10.1093/jhered/89.5.377](https://doi.org/10.1093/jhered/89.5.377) PMID: [ISI:000076113200001](https://pubmed.ncbi.nlm.nih.gov/13200001/).
2. Ward RD, Woodwark M, Skibinski DOF. A Comparison of Genetic Diversity Levels in Marine, Fresh-Water, and Anadromous Fishes. *J Fish Biol*. 1994; 44(2):213–32. doi: [10.1111/j.1095-8649.1994.tb01200.x](https://doi.org/10.1111/j.1095-8649.1994.tb01200.x) PMID: [ISI:A1994NC51900004](https://pubmed.ncbi.nlm.nih.gov/1994NC51900004/).
3. Avise JC. *Phylogeography: the history and formation of species*. Cambridge, Mass.: Harvard University Press; 2000. viii, 447 p. p.
4. Graves JE. Molecular insights into the population structures of cosmopolitan marine fishes. *J Hered*. 1998; 89(5):427–37. PubMed PMID: [ISI:000076113200006](https://pubmed.ncbi.nlm.nih.gov/13200006/).
5. Ely B, Viñas J, Alvarado Bremer JR, Black D, Lucas L, Covello K, et al. Consequences of the historical demography on the global population structure of two highly migratory cosmopolitan marine fishes: the yellowfin tuna (*Thunnus albacares*) and the skipjack tuna (*Katsuwonus pelamis*). *Bmc Evol Biol*. 2005; 5. Artn 19 doi: [10.1186/1471-2148-5-19](https://doi.org/10.1186/1471-2148-5-19) PMID: [ISI:000227791300002](https://pubmed.ncbi.nlm.nih.gov/13000002/).
6. Díaz-Jaimes P, Uribe-Alcocer M, Ortega-García S, Durand JD. Spatial and temporal mitochondrial DNA genetic homogeneity of dolphin populations (*Coryphaena hippurus*) in the eastern central Pacific. *Fish Res*. 2006; 80(2–3):333–8. doi: [10.1016/j.fishres.2006.04.015](https://doi.org/10.1016/j.fishres.2006.04.015) PMID: [ISI:000239850600023](https://pubmed.ncbi.nlm.nih.gov/16000023/).
7. Alvarado Bremer JR, Stequert B, Robertson NW, Ely B. Genetic evidence for inter-oceanic subdivision of bigeye tuna (*Thunnus obesus*) populations. *Mar Biol*. 1998; 132(4):547–57. PubMed PMID: [ISI:000077657100001](https://pubmed.ncbi.nlm.nih.gov/1000001/).
8. Alvarado Bremer JR, Naseri I, Ely B. A provisional study of northern bluefin tuna populations. *Col Vol Sci Pap ICCAT*. 1999; 49(1):127–9.
9. Carlsson J, McDowell JR, Carlsson JEL, Graves JE. Genetic identity of YOY bluefin tuna from the eastern and Western Atlantic spawning areas. *J Hered*. 2007; 98(1):23–8. doi: [10.1093/jhered/esl046](https://doi.org/10.1093/jhered/esl046) PMID: [ISI:000243585900003](https://pubmed.ncbi.nlm.nih.gov/16000003/).
10. Montes I, Iriondo M, Manzano C, Arrizabalaga H, Jimenez E, Pardo MA, et al. Worldwide genetic structure of albacore *Thunnus alalunga* revealed by microsatellite DNA markers. *Mar Ecol Prog Ser*. 2012; 471:183–U202. doi: [10.3354/Meps09991](https://doi.org/10.3354/Meps09991) PMID: [ISI:000312782700015](https://pubmed.ncbi.nlm.nih.gov/12782700015/).
11. Viñas J, Alvarado Bremer JR, Pla C. Inter-oceanic genetic differentiation among albacore (*Thunnus alalunga*) populations. *Mar Biol*. 2004; 145(2):225–32. doi: [10.1007/s00227-004-1319-5](https://doi.org/10.1007/s00227-004-1319-5) PMID: [ISI:000223098200002](https://pubmed.ncbi.nlm.nih.gov/123098200002/).
12. Graves JE, McDowell JR. Stock structure of the world's istiophorid billfishes: a genetic perspective. *Mar Freshwater Res*. 2003; 54(4):287–98. doi: [10.1071/Mf01290](https://doi.org/10.1071/Mf01290) PMID: [ISI:000184843100002](https://pubmed.ncbi.nlm.nih.gov/184843100002/).
13. Alvarado Bremer JR, Mejuto J, Greig TW, Ely B. Global population structure of the swordfish (*Xiphias gladius* L.) as revealed by analysis of the mitochondrial DNA control region. *J Exp Mar Biol Ecol*. 1996; 197(2):295–310. PubMed PMID: [ISI:A1996UU63500010](https://pubmed.ncbi.nlm.nih.gov/1996UU63500010/).
14. McDowell JR. Genetic stock structure of the sailfin, *Istiophorus platypterus*, based on nuclear and mitochondrial DNA. [Dissertation]: Virginia Institute of Marine Science: William and Mary; 2002.
15. Schrey AW, Heist EJ. Microsatellite analysis of population structure in the shortfin mako (*Isurus oxyrinchus*). *Can J Fish Aquat Sci*. 2003; 60(6):670–5. doi: [10.1139/F03-064](https://doi.org/10.1139/F03-064) PMID: [ISI:000184735200004](https://pubmed.ncbi.nlm.nih.gov/184735200004/).
16. Gonzalez EG, Beerli P, Zardoya R. Genetic structuring and migration patterns of Atlantic bigeye tuna, *Thunnus obesus* (Lowe, 1839). *Bmc Evol Biol*. 2008; 8. Artn 252 doi: [10.1186/1471-2148-8-252](https://doi.org/10.1186/1471-2148-8-252) PMID: [ISI:000259778400001](https://pubmed.ncbi.nlm.nih.gov/159778400001/).
17. Alvarado Bremer JR, Mejuto J, Gómez-Márquez J, Boán F, Carpintero P, Rodríguez JM, et al. Hierarchical analyses of genetic variation of samples from breeding and feeding grounds confirm the genetic

- partitioning of northwest Atlantic and South Atlantic populations of swordfish (*Xiphias gladius* L.). *J Exp Mar Biol Ecol.* 2005; 327(2):167–82. doi: [10.1016/j.jembe.2005.06.022](https://doi.org/10.1016/j.jembe.2005.06.022) PMID: [ISI:000233941400005](https://pubmed.ncbi.nlm.nih.gov/16100005/).
18. Chow S, Takeyama H. Nuclear and mitochondrial DNA analyses reveal four genetically separated breeding units of the swordfish. *J Fish Biol.* 2000; 56(5):1087–98. PubMed PMID: [ISI:000087281500005](https://pubmed.ncbi.nlm.nih.gov/1500005/).
 19. Viñas J, Perez-Serra A, Vidal O, Bremer JRA, Pla C. Genetic differentiation between eastern and western Mediterranean swordfish revealed by phylogeographic analysis of the mitochondrial DNA control region. *Ices J Mar Sci.* 2010; 67(6):1222–9. doi: [10.1093/icesjms/fsq031](https://doi.org/10.1093/icesjms/fsq031) PMID: [ISI:000280919600013](https://pubmed.ncbi.nlm.nih.gov/19600013/).
 20. Neilson JD, Paul SD, Smith SC. Stock structure of swordfish (*Xiphias gladius*) in the Atlantic: a review of the non-genetic evidence. *ICCAT Coll Vol Sci Pap.* 2007; 61:25–60.
 21. Dewar H, Prince ED, Musyl MK, Brill RW, Sepulveda C, Luo JG, et al. Movements and behaviors of swordfish in the Atlantic and Pacific Oceans examined using pop-up satellite archival tags. *Fish Oceanogr.* 2011; 20(3):219–41. doi: [10.1111/j.1365-2419.2011.00581.x](https://doi.org/10.1111/j.1365-2419.2011.00581.x) PMID: [ISI:000289259700005](https://pubmed.ncbi.nlm.nih.gov/200289259700005/).
 22. Neilson JD, Smith S, Royer F, Paul SD, Porter JM, Lutcavage M. Investigations of horizontal movements of Atlantic swordfish using pop-up satellite archival tags. 2009. In: *Rev M T Fis [Internet]*. New York: SpringerReviews: Methods and Technologies in Fish Biology and Fisheries; [145–59, 452].
 23. Sedberry GR, Loefer JK. Satellite telemetry tracking of swordfish, *Xiphias gladius* off the eastern United States. *Mar Biol.* 2001; 139(2):355–60. PubMed PMID: [ISI:000170744200015](https://pubmed.ncbi.nlm.nih.gov/170744200015/).
 24. Abascal AJ, Mejuto J, Quintans M, García-Cortés B, Ramos-Cartelle A. Horizontal tracking of the broadbill swordfish, *Xiphias gladius*, in the central and eastern North Atlantic using Pop-up satellite tags. *Coll Vol Sci Pap ICCAT.* in press;SCRS/2013/151.
 25. Alvarado Bremer JR, Mejuto J, Gómez-Márquez J, Pla-Zanuy C, Viñas J, Marques C, et al. Genetic population structure of Atlantic swordfish: current status and future directions. *ICCAT Coll Vol Sci Pap.* 2007; 61:107–18.
 26. Chow S, Clarke S, Nakadate M, Okazaki M. Boundary between the north and south Atlantic populations of the swordfish (*Xiphias gladius*) inferred by a single nucleotide polymorphism at calmodulin gene intron. *Mar Biol.* 2007; 152(1):87–93. doi: [10.1007/s00227-007-0662-8](https://doi.org/10.1007/s00227-007-0662-8) PMID: [ISI:000247471700009](https://pubmed.ncbi.nlm.nih.gov/1700009/).
 27. Grieg TW, Alvarado Bremer JR, Ely B. Nuclear markers provide additional evidence for population subdivision among Atlantic swordfish. *ICCAT Coll Vol Sci Pap.* 2000; 51:1637–41.
 28. Kasapidis P, Valeiras X, Garcia-Cortes B, Mejuto J. Genetic and growth profiles of several specimens of swordfish (*Xiphias gladius*) tagged and recaptured in the Atlantic, Indian and Pacific oceans. *ICCAT Coll Vol Sci Pap.* 2008; 62(4):1142–51.
 29. Kasapidis P, Mejuto J, Tserpes G, Antoniou A, Garcia-Cortes B, Peristeraki P, et al. Genetic structure of the swordfish (*Xiphias gladius*) stocks in the Atlantic using microsatellite DNA analysis. *ICCAT Coll Vol Sci Pap.* 2007; 61:89–98.
 30. Kotoulas G, Mejuto J, Antoniou A, Kasapidis P, Tserpes G, Piccinetti C, et al. Global genetic structure of swordfish (*Xiphias gladius*) as revealed by microsatellite DNA markers. *ICCAT Coll Vol Sci Pap.* 2007; 61:79–88.
 31. Viñas J, Alvarado Bremer JR, Mejuto J, de la Serna JM, García-Cortés B, Pla C. Swordfish genetic population structure in the North Atlantic and Mediterranean. *ICCAT Coll Vol Sci Pap.* 2007; 61:99–106.
 32. Morin PA, Martien KK, Taylor BL. Assessing statistical power of SNPs for population structure and conservation studies. *Mol Ecol Resour.* 2009; 9(1):66–73. doi: [10.1111/j.1755-0998.2008.02392.x](https://doi.org/10.1111/j.1755-0998.2008.02392.x) PMID: [ISI:000262678900009](https://pubmed.ncbi.nlm.nih.gov/18900009/).
 33. Morin PA, Luikart G, Wayne RK. SNPs in ecology, evolution and conservation. *Trends Ecol Evol.* 2004; 19(4):208–16. doi: [10.1016/j.tree.2004.01.009](https://doi.org/10.1016/j.tree.2004.01.009) PMID: [ISI:000220842400008](https://pubmed.ncbi.nlm.nih.gov/220842400008/).
 34. Ryman N, Palm S, Andre C, Carvalho GR, Dahlgren TG, Jorde PE, et al. Power for detecting genetic divergence: differences between statistical methods and marker loci. *Mol Ecol.* 2006; 15(8):2031–45. doi: [10.1111/j.1365-294X.2006.02839.x](https://doi.org/10.1111/j.1365-294X.2006.02839.x) PMID: [ISI:000238143900002](https://pubmed.ncbi.nlm.nih.gov/143900002/).
 35. Smith BL, Lu CP, Alvarado Bremer JR. Methodological streamlining of SNP discovery and genotyping via high-resolution melting analysis (HRMA) in non-model species. *Mar Genom.* 2013; 9:39–49. doi: [10.1016/j.margen.2012.10.001](https://doi.org/10.1016/j.margen.2012.10.001) PMID: [ISI:000316041500006](https://pubmed.ncbi.nlm.nih.gov/216041500006/).
 36. Smith BL, Lu CP, Alvarado Bremer JR. High-resolution melting analysis (HRMA): a highly sensitive inexpensive genotyping alternative for population studies. *Mol Ecol Resour.* 2010; 10(1):193–6. doi: [10.1111/j.1755-0998.2009.02726.x](https://doi.org/10.1111/j.1755-0998.2009.02726.x) PMID: [ISI:000272866500024](https://pubmed.ncbi.nlm.nih.gov/272866500024/).

37. Rooker JR, Simms JR, Wells RJD, Holt SA, Holt GJ, Graves JE, et al. Distribution and Habitat Associations of Billfish and Swordfish Larvae across Mesoscale Features in the Gulf of Mexico. *Plos One*. 2012; 7(4). ARTN e34180 doi: [10.1371/journal.pone.0034180](https://doi.org/10.1371/journal.pone.0034180) PMID: [ISI:000305336600022](https://pubmed.ncbi.nlm.nih.gov/22533660/).
38. ICCAT. Report of the 2009 Atlantic swordfish stock assessment session. *Coll Vol Sci Pap ICCAT*. 2010; 65(1):1–123.
39. Grieg TW. Partitioning genetic variation in swordfish (*Xiphias gladius L.*); analysis of sample variance and population structure. [Doctoral Dissertation]. Columbia: University of South Carolina; 2000.
40. Peakall R, Smouse PE. GENALEX 6: genetic analysis in Excel. Population genetic software for teaching and research. *Mol Ecol Notes*. 2006; 6(1):288–95. doi: [10.1111/j.1471-8286.2005.01155.x](https://doi.org/10.1111/j.1471-8286.2005.01155.x) PMID: [ISI:000235725600089](https://pubmed.ncbi.nlm.nih.gov/16235725600089/).
41. Fisher RA. The logic of inductive inference. *J R Stat Soc*. 1935; 98:39–82. PubMed PMID: [ISI:000189071900002](https://pubmed.ncbi.nlm.nih.gov/189071900002/).
42. Guo SW, Thompson EA. Performing the Exact Test of Hardy-Weinberg Proportion for Multiple Alleles. *Biometrics*. 1992; 48(2):361–72. PubMed PMID: [ISI:A1992HY93800002](https://pubmed.ncbi.nlm.nih.gov/1992HY93800002/).
43. Raymond M, Rousset F. An exact test for population differentiation. *Evolution*. 1995; 49(6):1280–3. PubMed PMID: [ISI:A1995TR69500027](https://pubmed.ncbi.nlm.nih.gov/1995TR69500027/).
44. Raymond M, Rousset F. Genepop (Version-1.2)—Population-genetics software for exact tests and ecumenicism. *J Hered*. 1995; 86(3):248–9. PubMed PMID: [ISI:A1995RB30200017](https://pubmed.ncbi.nlm.nih.gov/1995RB30200017/).
45. Rousset F. GENEPOP '007: a complete re-implementation of the GENEPOP software for Windows and Linux. *Mol Ecol Resour*. 2008; 8(1):103–6. doi: [10.1111/j.1471-8286.2007.01931.x](https://doi.org/10.1111/j.1471-8286.2007.01931.x) PMID: [ISI:000253827100016](https://pubmed.ncbi.nlm.nih.gov/253827100016/).
46. Rice WR. Analyzing Tables of Statistical Tests. *Evolution*. 1989; 43(1):223–5. PubMed PMID: [ISI:A1989R828900018](https://pubmed.ncbi.nlm.nih.gov/1989R828900018/).
47. Beaumont MA, Nichols RA. Evaluating loci for use in the genetic analysis of population structure. *P Roy Soc Lond B Bio*. 1996; 263(1377):1619–26. PubMed PMID: [ISI:A1996WD19600001](https://pubmed.ncbi.nlm.nih.gov/1996WD19600001/).
48. Antao T, Lopes A, Lopes RJ, Beja-Pereira A, Luikart G. LOSITAN: A workbench to detect molecular adaptation based on a F(st)-outlier method. *Bmc Bioinformatics*. 2008; 9:323. Artn 323 doi: [10.1186/1471-2105-9-323](https://doi.org/10.1186/1471-2105-9-323) PMID: [ISI:000258517000001](https://pubmed.ncbi.nlm.nih.gov/258517000001/).
49. Banks MA, Eichert W, Olsen JB. Which genetic loci have greater population assignment power? *Bioinformatics*. 2003; 19(11):1436–8. doi: [10.1093/bioinformatics/btg172](https://doi.org/10.1093/bioinformatics/btg172) PMID: [ISI:000184491500019](https://pubmed.ncbi.nlm.nih.gov/2003184491500019/).
50. Excoffier L, Smouse PE, Quattro JM. Analysis of molecular variance inferred from metric distances among DNA haplotypes—application to human mitochondrial-DNA restriction data. *Genetics*. 1992; 131(2):479–91. PubMed PMID: [ISI:A1992HW75900021](https://pubmed.ncbi.nlm.nih.gov/1992HW75900021/).
51. Excoffier L, Laval G, Schneider S. Arlequin (version 3.0): An integrated software package for population genetics data analysis. *Evol Bioinform*. 2005; 1:47–50. PubMed PMID: [ISI:000207065900004](https://pubmed.ncbi.nlm.nih.gov/207065900004/).
52. Weir BS, Cockerham CC. Estimating F-Statistics for the Analysis of Population-Structure. *Evolution*. 1984; 38(6):1358–70. PubMed PMID: [ISI:A1984TY40400017](https://pubmed.ncbi.nlm.nih.gov/1984TY40400017/).
53. Slatkin M. Isolation by Distance in Equilibrium and Nonequilibrium Populations. *Evolution*. 1993; 47(1):264–79. PubMed PMID: [ISI:A1993KR96500022](https://pubmed.ncbi.nlm.nih.gov/1993KR96500022/).
54. Orłóci L. Multivariate analysis in vegetation research. 2d ed. The Hague; Boston: Junk; 1978. ix, 451 p. p.
55. Pritchard JK, Stephens M, Donnelly P. Inference of population structure using multilocus genotype data. *Genetics*. 2000; 155(2):945–59. PubMed PMID: [ISI:000087475100039](https://pubmed.ncbi.nlm.nih.gov/200087475100039/).
56. Chow S, Okamoto H, Uozumi Y, Takeuchi Y, Takeyama H. Genetic stock structure of the swordfish (*Xiphias gladius*) inferred by PCR-RFLP analysis of the mitochondrial DNA control region. *Mar Biol*. 1997; 127(3):359–67. PubMed PMID: [ISI:A1997WJ44700001](https://pubmed.ncbi.nlm.nih.gov/1997WJ44700001/).
57. Falush D, Stephens M, Pritchard JK. Inference of population structure using multilocus genotype data: linked loci and correlated allele frequencies. *Genetics*. 2003; 164(4):1567–87. PubMed PMID: [ISI:000185248000029](https://pubmed.ncbi.nlm.nih.gov/2003185248000029/).
58. Waples RS. Separating the wheat from the chaff: Patterns of genetic differentiation in high gene flow species. *J Hered*. 1998; 89(5):438–50. PubMed PMID: [ISI:000076113200007](https://pubmed.ncbi.nlm.nih.gov/2000076113200007/).
59. Hubisz MJ, Falush D, Stephens M, Pritchard JK. Inferring weak population structure with the assistance of sample group information. *Mol Ecol Resour*. 2009; 9(5):1322–32. doi: [10.1111/j.1755-0998.2009.02591.x](https://doi.org/10.1111/j.1755-0998.2009.02591.x) PMID: [ISI:000268855000004](https://pubmed.ncbi.nlm.nih.gov/2009025915000004/).
60. Evanno G, Regnaut S, Goudet J. Detecting the number of clusters of individuals using the software STRUCTURE: a simulation study. *Mol Ecol*. 2005; 14(8):2611–20. doi: [10.1111/j.1365-294X.2005.02553.x](https://doi.org/10.1111/j.1365-294X.2005.02553.x) PMID: [ISI:000229961500029](https://pubmed.ncbi.nlm.nih.gov/229961500029/).

61. Earl DA, Vonholdt BM. STRUCTURE HARVESTER: a website and program for visualizing STRUCTURE output and implementing the Evanno method. *Conserv Genet Resour.* 2012; 4(2):359–61. doi: [10.1007/s12686-011-9548-7](https://doi.org/10.1007/s12686-011-9548-7) PMID: [ISI:000303536400036](https://pubmed.ncbi.nlm.nih.gov/21586627/).
62. Guillot G, Mortier F, Estoup A. GENELAND: a computer package for landscape genetics. *Mol Ecol Notes.* 2005; 5(3):712–5. doi: [10.1111/j.1471-8286.2005.01031.x](https://doi.org/10.1111/j.1471-8286.2005.01031.x) PMID: [ISI:000231529400078](https://pubmed.ncbi.nlm.nih.gov/15821529400078/).
63. Jakobsson M, Rosenberg NA. CLUMPP: a cluster matching and permutation program for dealing with label switching and multimodality in analysis of population structure. *Bioinformatics.* 2007; 23(14):1801–6. doi: [10.1093/bioinformatics/btm233](https://doi.org/10.1093/bioinformatics/btm233) PMID: [ISI:000249248300012](https://pubmed.ncbi.nlm.nih.gov/17309804/).
64. Waples RS, Gaggiotti O. What is a population? An empirical evaluation of some genetic methods for identifying the number of gene pools and their degree of connectivity. *Mol Ecol.* 2006; 15(6):1419–39. doi: [10.1111/j.1365-294X.2006.02890.x](https://doi.org/10.1111/j.1365-294X.2006.02890.x) PMID: [ISI:000236798200001](https://pubmed.ncbi.nlm.nih.gov/165236798200001/).
65. Chen C, Durand E, Forbes F, Francois O. Bayesian clustering algorithms ascertaining spatial population structure: a new computer program and a comparison study. *Mol Ecol Notes.* 2007; 7(5):747–56. doi: [10.1111/j.1471-8286.2007.01769.x](https://doi.org/10.1111/j.1471-8286.2007.01769.x) PMID: [ISI:000249193000004](https://pubmed.ncbi.nlm.nih.gov/17309804/).
66. Avise JC. *Molecular markers, natural history, and evolution.* 2nd ed. Sunderland, Mass.: Sinauer Associates; 2004. 684 p.
67. Nott A, Meislin SH, Moore MJ. A quantitative analysis of intron effects on mammalian gene expression. *RNA.* 2003; 9(5):607–17. Epub 2003/04/19. PubMed PMID: [12702819](https://pubmed.ncbi.nlm.nih.gov/12702819/); PubMed Central PMCID: PMC1370426.
68. Gazave E, Marques-Bonet T, Fernando O, Charlesworth B, Navarro A. Patterns and rates of intron divergence between humans and chimpanzees. *Genome Biol.* 2007; 8(2):R21. Epub 2007/02/21. doi: [10.1186/gb-2007-8-2-r21](https://doi.org/10.1186/gb-2007-8-2-r21) [pii] doi: [10.1186/gb-2007-8-2-r21](https://pubmed.ncbi.nlm.nih.gov/17309804/) PMID: [17309804](https://pubmed.ncbi.nlm.nih.gov/17309804/); PubMed Central PMCID: PMC1852421.
69. Nielsen EE, Hansen MM, Meldrup D. Evidence of microsatellite hitch-hiking selection in Atlantic cod (*Gadus morhua* L.): implications for inferring population structure in nonmodel organisms. *Mol Ecol.* 2006; 15(11):3219–29. doi: [10.1111/j.1365-294X.2006.03025.x](https://doi.org/10.1111/j.1365-294X.2006.03025.x) PMID: [ISI:000240440800010](https://pubmed.ncbi.nlm.nih.gov/165236798200010/).
70. Kirk H, Freeland JR. Applications and Implications of Neutral versus Non-neutral Markers in Molecular Ecology. *International journal of molecular sciences.* 2011; 12(6):3966–88. doi: [10.3390/ijms12063966](https://doi.org/10.3390/ijms12063966) PMID: [21747718](https://pubmed.ncbi.nlm.nih.gov/21747718/); PubMed Central PMCID: PMC3131602.
71. Helyar SJ, Hemmer-Hansen J, Bekkevold D, Taylor MI, Ogden R, Limborg MT, et al. Application of SNPs for population genetics of nonmodel organisms: new opportunities and challenges. *Mol Ecol Resour.* 2011; 11 Suppl 1:123–36. doi: [10.1111/j.1755-0998.2010.02943.x](https://doi.org/10.1111/j.1755-0998.2010.02943.x) PMID: [21429169](https://pubmed.ncbi.nlm.nih.gov/21429169/).
72. Allendorf FW, Hohenlohe PA, Luikart G. Genomics and the future of conservation genetics. *Nat Rev Genet.* 2010; 11(10):697–709. doi: [10.1038/Nrg2844](https://doi.org/10.1038/Nrg2844) PMID: [ISI:000281911300010](https://pubmed.ncbi.nlm.nih.gov/21429169/).
73. Nielsen EE, Hemmer-Hansen J, Larsen PF, Bekkevold D. Population genomics of marine fishes: identifying adaptive variation in space and time. *Mol Ecol.* 2009; 18(15):3128–50. doi: [10.1111/j.1365-294X.2009.04272.x](https://doi.org/10.1111/j.1365-294X.2009.04272.x) PMID: [ISI:000268457200003](https://pubmed.ncbi.nlm.nih.gov/193098003/).
74. Limborg MT, Helyar SJ, de Bruyn M, Taylor MI, Nielsen EE, Ogden R, et al. Environmental selection on transcriptome-derived SNPs in a high gene flow marine fish, the Atlantic herring (*Clupea harengus*). *Mol Ecol.* 2012; 21(15):3686–703. doi: [10.1111/j.1365-294X.2012.05639.x](https://doi.org/10.1111/j.1365-294X.2012.05639.x) PMID: [WOS:000306478800007](https://pubmed.ncbi.nlm.nih.gov/21429169/).
75. Skarstein TH, Westgaard JI, Fevolden SE. Comparing microsatellite variation in north-east Atlantic cod (*Gadus morhua* L.) to genetic structuring as revealed by the pantophysin (Pan I) locus. *J Fish Biol.* 2007; 70:271–90. doi: [10.1111/j.1095-8649.2007.01456.x](https://doi.org/10.1111/j.1095-8649.2007.01456.x) PMID: [ISI:000246788100001](https://pubmed.ncbi.nlm.nih.gov/17309804/).
76. Andre C, Larsson LC, Laikre L, Bekkevold D, Brigham J, Carvalho GR, et al. Detecting population structure in a high gene-flow species, Atlantic herring (*Clupea harengus*): direct, simultaneous evaluation of neutral vs putatively selected loci. *Heredity.* 2011; 106(2):270–80. doi: [10.1038/Hdy.2010.71](https://doi.org/10.1038/Hdy.2010.71) PMID: [ISI:000286375300009](https://pubmed.ncbi.nlm.nih.gov/21429169/).
77. Freamo H, O'Reilly P, Berg PR, Lien S, Boulding EG. Outlier SNPs show more genetic structure between two Bay of Fundy metapopulations of Atlantic salmon than do neutral SNPs. *Mol Ecol Resour.* 2011; 11:254–67. doi: [10.1111/j.1755-0998.2010.02952.x](https://doi.org/10.1111/j.1755-0998.2010.02952.x) PMID: [ISI:000287485100022](https://pubmed.ncbi.nlm.nih.gov/21429169/).
78. Hess JE, Campbell NR, Close DA, Docker MF, Narum SR. Population genomics of Pacific lamprey: adaptive variation in a highly dispersive species. *Mol Ecol.* 2013; 22(11):2898–916. doi: [10.1111/Mec.12150](https://doi.org/10.1111/Mec.12150) PMID: [WOS:000319675300005](https://pubmed.ncbi.nlm.nih.gov/248519675300005/).
79. Muths D, Le Couls S, Evano H, Grewe P, Bourjea J. Multi-Genetic Marker Approach and Spatio-Temporal Analysis Suggest There Is a Single Panmictic Population of Swordfish *Xiphias gladius* in the Indian Ocean. *Plos One.* 2013; 8(5). ARTN e63558 doi: [10.1371/journal.pone.0063558](https://doi.org/10.1371/journal.pone.0063558) PMID: [WOS:000320362700035](https://pubmed.ncbi.nlm.nih.gov/248519675300035/).

80. Govoni JJ, Stender BW, Pashuk O. Distribution of larval swordfish, *Xiphias gladius*, and probable spawning off the southeastern United States. *Fish B-Noaa*. 2000; 98(1):64–74. PubMed PMID: [ISI:000085512900005](#).
81. Palko BJ, Beardsley GL, Richards WJ. Synopsis of the biology of the swordfish, *Xiphias gladius* Linnaeus. Seattle, Wash.: U.S. Dept. of Commerce, National Oceanic and Atmospheric Administration, National Marine Fisheries Service; 1981. iv, 21 p. p.
82. de la Serna JM, Alot E. Consideraciones relativas a los desplazamientos efectuados por el pez espada (*Xiphias gladius*) en el área del estrecho de Gibraltar y otras observaciones relacionadas con la biología de la reproducción. *ICCAT Coll Vol Sci Pap*. 1990; 32(2):353–9.
83. O'Reilly PT, Canino MF, Bailey KM, Bentzen P. Inverse relationship between F-ST and microsatellite polymorphism in the marine fish, walleye pollock (*Theragra chalcogramma*): implications for resolving weak population structure. *Mol Ecol*. 2004; 13(7):1799–814. doi: [10.1111/j.1365-294X.2004.02214.x](#) PMID: [ISI:000221933000007](#).
84. DeWoody JA, Avise JC. Microsatellite variation in marine, freshwater and anadromous fishes compared with other animals. *J Fish Biol*. 2000; 56(3):461–73. PubMed PMID: [ISI:000085915900001](#).
85. Qiu F, Kitchen A, Beerli P, Miyamoto MM. A possible explanation for the population size discrepancy in tuna (genus *Thunnus*) estimated from mitochondrial DNA and microsatellite data. *Mol Phylogenet Evol*. 2013; 66(2):463–8. doi: [10.1016/j.ympev.2012.05.002](#) PMID: [ISI:000315553300004](#).
86. Portnoy DS, McDowell JR, Heist EJ, Musick JA, Graves JE. World phylogeography and male-mediated gene flow in the sandbar shark, *Carcharhinus plumbeus*. *Mol Ecol*. 2010; 19(10):1994–2010. doi: [10.1111/j.1365-294X.2010.04626.x](#) PMID: [ISI:000277297700002](#).
87. Finnerty JR, Block BA. Direct sequencing of mitochondrial DNA detects highly divergent haplotypes in blue marlin (*Makaira nigricans*). *Mol Mar Biol Biotechnol*. 1992; 1(3):206–14. Epub 1992/06/01. PubMed PMID: [1308204](#).
88. Graves JE, McDowell JR. Inter-ocean genetic divergence of istiophorid billfishes. *Mar Biol*. 1995; 122(2):193–203. PubMed PMID: [ISI:A1995QX95300004](#).
89. Buonaccorsi VP, Reece KS, Morgan LW, Graves JE. Geographic distribution of molecular variance within the blue marlin (*Makaira nigricans*): A hierarchical analysis of allozyme, single-copy nuclear DNA, and mitochondrial DNA markers. *Evolution*. 1999; 53(2):568–79. PubMed PMID: [ISI:000080227100026](#).
90. Buonaccorsi VP, McDowell JR, Graves JE. Reconciling patterns of inter-ocean molecular variance from four classes of molecular markers in blue marlin (*Makaira nigricans*). *Mol Ecol*. 2001; 10(5):1179–96. PubMed PMID: [ISI:000168692900010](#).
91. Nakamura Y. FAO species catalogue vol. 5: billfishes of the world: an annotated and illustrated catalogue of marlins, sailfishes, spearfishes, and swordfishes known to date. *FAO Fish Synop*. 1985; 125:48–51.
92. Neilson J, Arocha F, Cass-Calay S, Mejuto J, Ortiz M, Scott G, et al. The recovery of Atlantic swordfish: the comparative roles of the regional fisheries management organization and species biology. *Reviews in Fisheries Science*. 2013; 21(2):59–97.
93. Alvarado Bremer JR, Mejuto J, Gómez-Márquez J, Viñas J, Boán F. Hierarchical analysis of nucleotide diversity reveals extremely low levels of mitochondrial DNA gene flow between northeast Atlantic and Mediterranean swordfish populations. *ICCAT Coll Vol Sci Pap*. 1999; 49(1):457–66.
94. Rey JCA, E. Captura de estadios postlarvarius de pex espada (*Xiphias gladius*) en el oeste del Mediterraneo occidental. *Col Vol Sci Pap ICCAT*. 1988; 27:178–9.
95. Blair C, Weigel DE, Balazik M, Keeley ATH, Walker FM, Landguth E, et al. A simulation-based evaluation of methods for inferring linear barriers to gene flow. *Mol Ecol Resour*. 2012; 12(5):822–33. doi: [10.1111/j.1755-0998.2012.03151.x](#) PMID: [ISI:000307927700006](#).
96. García-Cortés B, Mejuto J, Quintans M. Summary of swordfish (*Xiphias gladius*) recaptures carried out by the Spanish surface longline fleet in the Atlantic Ocean: 1984–2002. *ICCAT Coll Vol Sci Pap*. 2003; 55(4):1476–84.
97. Reeb CA. Genetic discontinuity of big fish in a small sea. *P Natl Acad Sci USA*. 2010; 107(6):2377–8. doi: [10.1073/pnas.0914639107](#) PMID: [ISI:000274408100005](#).
98. Lu C-P. The development of high resolution melting assays for species identification and potential sex-linked genetic markers survey in istiophorid billfish and bayesian analyses of genetic variation and population differentiation in Pacific swordfish (*Xiphias Gladius* L.) [Dissertation]: Texas A&M University; 2014.
99. Lu C-P, Chen CA, Hui CF, Tzeng TD, Yeh SY. Population genetic structure of the swordfish, *Xiphias gladius* (Linnaeus, 1758), in the Indian Ocean and West Pacific inferred from the complete DNA

- sequence of the mitochondrial control region. *Zool Stud.* 2006; 45(2):269–79. PubMed PMID: [WOS:000239194200013](#).
100. Muths D, Grewe P, Jean C, Bourjea J. Genetic population structure of the Swordfish (*Xiphias gladius*) in the southwest Indian Ocean: Sex-biased differentiation, congruency between markers and its incidence in a way of stock assessment. *Fish Res.* 2009; 97(3):263–9. doi: [10.1016/j.fishres.2009.03.004](#) PMID: [WOS:000266543700011](#).
 101. Diaz RJ. Overview of hypoxia around the world. *J Environ Qual.* 2001; 30(2):275–81. PubMed PMID: [ISI:000169455400001](#).
 102. Prince ED, Luo JG, Goodyear CP, Hoolihan JP, Snodgrass D, Orbesen ES, et al. Ocean scale hypoxia-based habitat compression of Atlantic istiophorid billfishes. *Fish Oceanogr.* 2010; 19(6):448–62. doi: [10.1111/j.1365-2419.2010.00556.x](#) PMID: [ISI:000283075500003](#).
 103. Seki MP, Lumpkin R, Flament P. Hawaii cyclonic eddies and blue marlin catches: The case study of the 1995 Hawaiian International Billfish Tournament. *J Oceanogr.* 2002; 58(5):739–45. PubMed PMID: [ISI:000178241000012](#).
 104. Mejuto J, de la Serna JM, García B. Some considerations on the spatial and temporal variability in the sex-ratio at size of the swordfish (*Xiphias gladius* L.). *ICCAT Coll Vol Sci Pap.* 1998; 48(1):205–15.