

XXXVII Congreso SEBBM - Granada 2014  
Congreso anual de la Sociedad Española de  
Bioquímica y Biología Molecular  
Granada, del 9 al 12 de septiembre 2014

P15-12

**Assessment of environmental pollution applying  
oxidative stress biomarkers in the mussel *Mytilus  
galloprovincialis***

**Silvia Tejada<sup>1</sup>, Antoni Sureda<sup>2</sup>, Ainhoa Zuazu<sup>3</sup>, Miguela Méndez<sup>2</sup>,  
Antonino Natalotto<sup>4</sup>, Salvatore Fasulo<sup>4</sup>, Salud Deudero<sup>3</sup>**

<sup>1</sup>University of the Balearic Islands / Experimental Laboratory, Research Unit, Son Llätzer Hospital, IUNICS, Palma de Mallorca, ES, <sup>2</sup>Research Group on Community Nutrition and Oxidative Stress, IUNICS, University of the Balearic Islands, Palma de Mallorca, Spain and CIBER: CB12/03/30038 Fisiopatología de la Obesidad y la Nutrición, Instituto de Salud Carlos III (ISCIII), Palma de Mallorca, ES, <sup>3</sup>Spanish Institute of Oceanography, Palma de Mallorca, ES, <sup>4</sup>Dep. of Animal Biology and Marine Ecology, University of Messina, Messina, IT

In polluted environments, and especially in coastal waters, living organisms are most often exposed to complex mixtures of chemical contaminants. Mussels are desirable as bioindicators since they are likely to reflect changes in the environment pollution status. Transplanting mussels from a reference site to different locations allows biomonitoring the alterations due to environmental pollutants. The aim of the present study was to analyse the biochemical response of the mussels (*Mytilus galloprovincialis*) exposed to petrol refinery. Mussels purchased from a mussel farm in Messina (Italy) were transplanted for two months in a control clean site (Brucoli) and a polluted site (Priolo refinery). A third group was maintained in the same conditions in the farm for two months. Gills from each specimen (n=8 for each station) were dissected and used for biochemical analysis. Mussels from the three studied areas presented the same length and width. Mussels from the polluted site presented higher levels of malondialdehyde respect to the clean site. The activities of catalase and glutathione-S transferase (GST) were significantly higher in the polluted site, whereas the activity of acetylcholinesterase was significantly lower respect non-polluted sites. A significant increase in the gene expression of metallothionein (MT)-10, MT-20 and cytochrome P450 were reported in the polluted site respect to the other areas, whereas the increase in catalase and GST was not statistically significant. In conclusion, pollutants induce oxidative stress and increase the antioxidant and detoxifying system in mussels. The use of biomarkers in gills of the mussel *M. galloprovincialis* is a good tool to categorize differences between polluted and non-polluted areas.

**Acknowledgments:** National Parks Program 024/2010 and CIBERobn CB12/03/30038.

**Keywords:** Antioxidants, Biomarkers, Oxidative damage, Pollution.