

Descriptive Epidemiology of Collegiate Men's Football Injuries: National Collegiate Athletic Association Injury Surveillance System, 1988–1989 Through 2003–2004

Randall Dick, MS, FACSM*; Michael S. Ferrara, PhD, ATC†; Julie Agel, MA, ATC‡; Ron Courson, ATC, PT, NREMT-I, CSCS†; Stephen W. Marshall, PhD§; Michael J. Hanley, MS, ATC||; Fred Reifsteck, MD†

*National Collegiate Athletic Association, Indianapolis, IN; †University of Georgia, Athens, GA; ‡University of Minnesota, Minneapolis, MN; §University of North Carolina at Chapel Hill, Chapel Hill, NC; ||East Carolina University, Greenville, NC

Objective: To review 16 years of National Collegiate Athletic Association (NCAA) injury surveillance data for men's football and identify potential areas for injury prevention initiatives.

Background: Football is a high-velocity collision sport in which injuries are expected. Football tends to have one of the highest injury rates in sports. Epidemiologic data helps certified athletic trainers and other clinicians identify injury trends and patterns to appropriately design and institute injury prevention protocols and then measure their effects.

Main Results: During the 16-year reporting period, about 19% of the Division I, II, and III NCAA institutions sponsoring football participated in the Injury Surveillance System. The results from the 16-year study period show little variation in the injury rates over time: games averaged 36 injuries per 1000 athlete-exposures (A-Es); fall practice, approximately 4 injuries per 1000 A-Es; and spring practice, about 10 injuries per 1000 A-Es. The game injury rate was more than 9 times higher than the in-season practice injury rate (35.90 versus 3.80 injuries per 1000 A-Es, rate ratio = 9.1, 95% confidence interval = 9.0, 9.2), and the spring practice injury rate was more than 2 times

higher than the fall practice injury rate (9.62 versus 3.80 injuries per 1000 A-Es, rate ratio = 2.5, 95% confidence interval = 2.5, 2.6). The rate ratio for games versus fall practices was greatest for upper leg contusions (18.1 per 1000 A-Es), acromioclavicular joint sprains (14.0 per 1000 A-Es), knee internal derangements (13.4 per 1000 A-Es), ankle ligament sprains (12.0 per 1000 A-Es), and concussions (11.1 per 1000 A-Es).

Recommendations: Football is a complex sport that requires a range of skills performed by athletes with a wide variety of body shapes and types. Injury risks are greatest during games. Thus, injury prevention measures should focus on position-specific activities to reduce the injury rate. As equipment technology improves for the helmet, shoulder pads, and other protective devices, appropriate injury surveillance procedures should be performed to determine the effect of the new equipment on injury rates. A consistent evaluation of injury trends and patterns will assist decision makers in designing injury prevention techniques in areas that warrant the greatest attention and suggesting rule changes and modifications based on the data.

Key Words: athletic injuries, injury prevention, concussions, knee injuries, ankle injuries, heat illness

The game of football has been played competitively at the collegiate level in the United States for more than 100 years. In fact, it was safety concerns about the early game of football, and its notorious “flying wedge” formation, that led to the establishment of the National Collegiate Athletic Association (NCAA) during the first decade of the 20th century. At the urging of President Theodore Roosevelt, colleges banded together with the goal of reforming football to limit the injuries (and fatalities) in the young sport.

The NCAA conducted its first football championship in 1978. In the 1988–1989 academic year, 524 schools were sponsoring varsity football teams, with approximately 47 942 participants. By 2003–2004, the number of varsity teams had increased 18% to 621, involving 59 980 participants.¹ Participation growth during this time was apparent in all 3 divisions but particularly in Divisions I and II. In addition to traditional practice sessions during the fall, the NCAA also sanctions a

short spring practice season for Divisions I and II. Spring football is a maximum 15-day, formal, off-season team practice. No official competitive games are played in the spring season.

SAMPLING AND METHODS

During the 16-year period from 1988–1989 through 2003–2004, an average of 18.8% of schools sponsoring varsity football programs participated in annual fall NCAA Injury Surveillance System (ISS) data collection (Table 1). A similar percentage (18.0%) of Division I and II schools participated in spring football data collection during the same period. The sampling process, data collection methods, injury and exposure definitions, inclusion criteria, and data analysis methods are described in detail in the “Introduction and Methods” article in this special issue.²

Table 1. School Participation Frequency (in Total Numbers) by Year and National Collegiate Athletic Association (NCAA) Division, Men's Football, 1988–1989 Through 2003–2004*

Year	Division I Schools		Division II Schools		Division III Schools†		All Divisions		
	Participating	Sponsoring	Participating	Sponsoring	Participating	Sponsoring	Participating	Sponsoring	Percentage
Fall football									
1988–1989	33	193	16	116	36	213	85	524	16.2
1989–1990	31	195	16	116	32	213	79	524	15.1
1990–1991	39	193	19	120	39	221	97	535	18.1
1991–1992	39	195	24	127	43	224	106	547	19.4
1992–1993	39	195	22	129	34	229	95	553	17.2
1993–1994	49	221	23	142	34	198	106	561	18.9
1994–1995	56	224	30	141	31	201	117	568	20.6
1995–1996	52	230	27	154	34	216	113	601	18.8
1996–1997	47	229	32	154	43	217	122	600	20.3
1997–1998	52	231	26	151	23	217	101	599	16.9
1998–1999	42	231	17	157	44	217	103	605	17.0
1999–2000	68	235	30	155	57	220	155	610	25.4
2000–2001	42	236	24	157	47	231	113	624	18.1
2001–2002	53	238	29	153	46	226	128	617	20.7
2002–2003	45	240	25	150	52	229	122	619	19.7
2003–2004	37	238	25	151	52	228	114	621	18.4
Average	45	220	24	142	40	219	110	582	18.8
Spring football									
1988–1989	24	193	22	116	N/A‡	N/A	46	309	14.9
1989–1990	24	195	10	116	N/A	N/A	34	311	10.9
1990–1991	35	193	21	120	N/A	N/A	56	313	17.9
1991–1992	28	195	12	127	N/A	N/A	40	322	12.4
1992–1993	27	195	16	129	N/A	N/A	43	324	13.3
1993–1994	37	221	17	142	N/A	N/A	54	363	14.9
1994–1995	38	224	22	141	N/A	N/A	60	365	16.4
1995–1996	53	230	23	154	N/A	N/A	76	384	19.8
1996–1997	71	229	35	154	N/A	N/A	106	383	27.7
1997–1998	75	231	37	151	N/A	N/A	112	382	29.3
1998–1999	30	231	17	157	N/A	N/A	47	388	12.1
1999–2000	66	235	28	155	N/A	N/A	94	390	24.1
2000–2001	59	236	25	157	N/A	N/A	84	393	21.4
2001–2002	43	238	21	153	N/A	N/A	64	391	16.4
2002–2003	50	240	21	150	N/A	N/A	71	390	18.2
2003–2004	43	238	26	151	N/A	N/A	69	391	17.7
Average	44	220	22	142	N/A	N/A	66	362	18.0

*"Participating" refers to schools that provided appropriate data to the NCAA Injury Surveillance System; "Sponsoring" refers to the total number of schools offering the sport within the NCAA divisions.

†Division III schools do not participate in spring football.

‡N/A indicates not applicable.

RESULTS

Game and Practice Athlete-Exposures

The average annual numbers of games, fall practices, spring practices, and athletes participating for each NCAA division, condensed over the study period, are shown in Table 2. For fall football, Division I averaged 7 more practices than Division II and 13 more than Division III. Division I also had a larger practice squad size than either of the other 2 divisions. Division I averaged 1 to 2 more games and 4 more game participants than Divisions II and III. In spring practices, both divisions averaged 15 practices, and Division I averaged 12 more participants per practice than Division II.

Injury Rate by Activity, Division, and Season

Fall game, fall practice, and spring practice injury rates over time, combined across divisions, with 95% confidence inter-

Table 2. Average Annual Games, Practices, and Athletes Participating by National Collegiate Athletic Association Division per School, Men's Football, 1988–1989 Through 2003–2004

Division	Games	Athletes per Game	Practices	Athletes per Practice
Fall football				
I	11	52	81	95
II	10	48	74	81
III	9	48	68	78
Spring football				
I	N/A*	N/A	15	77
II	N/A	N/A	15	65

*N/A indicates not applicable.




Figure 1. Injury rates and 95% confidence intervals per 1000 athlete-exposures by games, practices, and academic years, men's football, 1988–1989 through 2003–2004 ($n = 30\,797$ game injuries, 42 355 fall practice injuries, and 10 943 spring practice injuries). Practice injury counts for 1996–1997 were imputed from adjacent years. Game time trend $P = .69$. Average annual change = -0.2% (95% confidence interval = $-1.1, 0.7$). Fall practice time trend $P = .82$. Average annual change = -0.4% (95% confidence interval = $-3.5, 2.8$). Spring practice time trend $P = .03$. Average annual change = -1.1% (95% confidence interval = $-2.1, -0.1$).

vals (CIs), are displayed in Figure 1. Over the 16 years of the study, the rate of injury in games was 9 times higher than the practice injury rate (35.90 versus 3.80 injuries per 1000 athlete-exposures [A-Es], rate ratio = 9.4, 95% CI = 9.3, 9.5), and the risk of injury in spring practice was more than twice that in fall practice (9.62 versus 3.80 injuries per 1000 A-Es, rate ratio = 2.5, 95% CI = 2.5, 2.6; data not shown). No statistically significant decreases were identified in the fall game (average annual change of -0.2% , $P = .69$) or the fall practice (average annual change of -0.4% , $P = .82$). The spring practice (average annual change of -1.1% , $P = .03$) injury rates decreased during the sample period.

The total number of fall games, fall practices, and spring practices with associated injury rates condensed over years by division and season (preseason, in season, and postseason) are presented in Tables 3 and 4. During the 16-year period, 30 797 injuries from almost 18 000 games, 42 355 injuries from more than 128 000 fall practices, and 10 943 injuries from more than 15 000 spring practices were reported. Fall game and practice injury rates did not differ significantly across divisions. However, fall preseason practice injury rates were more than 3 times as high as in season or postseason practice injury rates (fall preseason versus regular season: 7.24 versus 2.09 injuries per 1000 A-Es, rate ratio = 3.5, 95% CI = 3.4, 3.5, $P < .01$; fall preseason versus postseason: 7.24 versus 1.35 injuries per 1000 A-Es, rate ratio = 5.4, 95% CI = 4.8, 6.0, $P < .01$), whereas in-season game injury rates were approximately 50% higher than postseason game injury rates (36.11 versus 23.71 injuries per 1000 A-Es, rate ratio = 1.5, 95% CI = 1.4, 1.7, $P < .01$). Spring practice injury rates were one third higher than preseason fall practice rates (9.62 versus 7.24 injuries per 1000 A-Es, rate ratio = 1.3, 95% CI = 1.3, 1.4, $P < .01$) and more than 2 times higher than the overall fall practice injury rate (9.62 versus 3.80 injuries per 1000 A-Es, rate ratio = 2.5, 95% confidence interval = 2.5, 2.6, $P < .01$).

Body Parts Injured Most Often and Specific Injuries

The frequency of injury to 5 general body parts (head/neck, upper extremity, trunk/back, lower extremity, and other/system) for fall games, fall practices, and spring practices, with years and divisions combined, is shown in Table 5. The breakdown of injury frequency for each body region was similar for fall and spring practices and regular-season games. More than 50% of all injuries in the 3 activities were to the lower extremity. Systemic injuries, primarily heat illness, accounted for 6% of fall practice time-loss events.

The most common injured body part and injury type combinations for games and practices with years and divisions combined are displayed in Table 6. All injuries that accounted for at least 1% of reported injuries during the 16-year sampling period were included. In fall games, knee internal derangements (17.8%), ankle ligament sprains (15.6%), and concussions (6.8%) accounted for the majority of injuries. A variety of shoulder problems also was reported. In fall practices, knee internal derangements (12.0%), ankle ligament sprains (11.8%), upper leg muscle-tendon strains (10.7%), and concussions (5.5%) were the most frequently reported. Heat illness totaled 3.9% of fall practice injuries. Spring practice injuries followed the fall practice injury pattern, with knee internal derangements (16.4%), ankle ligament sprains (13.9%), upper leg muscle-tendon strains (10.8%), and concussions (5.6%) accounting for 46.7% of all reported injuries. Heat illness totaled less than 0.5% of spring practice injuries. Compared with a fall practice, a participant in an in-season game was 18 times more likely to sustain an upper leg contusion (1.27 versus 0.07 injuries per 1000 A-Es, rate ratio = 18.1, 95% CI = 16.6, 19.9, $P < .01$), 14 times as likely to sustain an acromioclavicular joint sprain (0.98 versus 0.07 injuries per 1000 A-Es, rate ratio = 14.0, 95% CI = 12.7, 15.4, $P < .01$), 13 times as likely to sustain knee internal derange-

Table 3. Games and Practices With Associated Injury Rates by National Collegiate Athletic Association Division and Season, Men's Football, 1988–1989 Through 2003–2004*

	Total No. of Games Reported	Game Injury Rate per 1000 Athlete-Exposures	95% Confidence Interval	Total No. of Practices Reported	Practice Injury Rate per 1000 Athlete-Exposures	95% Confidence Interval
Division I						
Preseason	298	7.44	6.09, 8.80	19 325	7.05	6.92, 7.17
In season	7369	37.79	37.17, 38.40	36 800	2.02	1.97, 2.06
Postseason	169	22.31	19.19, 25.44	1569	1.70	1.50, 1.91
Total Division I	7836	36.27	35.68, 36.85	56 694	3.66	3.61, 3.71
Division II						
Preseason	172	5.20	3.66, 6.73	9217	6.77	6.59, 6.96
In season	3749	35.23	34.36, 36.10	17 811	1.90	1.83, 1.97
Postseason	98	25.43	20.76, 30.10	482	0.75	0.48, 1.03
Total Division II	4019	33.64	32.82, 34.46	27 510	3.57	3.49, 3.65
Division III						
Preseason	281	9.14	7.53, 10.74	13 872	7.85	7.69, 8.01
In season	5602	34.31	33.61, 35.01	28 377	2.32	2.26, 2.38
Postseason	137	24.44	20.52, 28.37	691	0.75	0.51, 0.98
Total Division III	6020	32.89	32.23, 33.55	42 940	4.18	4.11, 4.25
All Divisions						
Preseason	751	7.55	6.68, 8.43	42 414	7.24	7.15, 7.32
In season	16 720	36.11	35.70, 36.52	82 988	2.09	2.05, 2.12
Postseason	404	23.71	21.54, 25.87	2742	1.35	1.20, 1.49
Total	17 911	35.90	35.50, 36.30	128 395	3.80	3.80, 3.80

*Wald χ^2 statistics from negative binomial model: game injury rates did not differ among divisions ($P = .49$) but did differ within seasons ($P < .01$). Practice injury rates did not differ among divisions ($P = .26$) but did differ within seasons ($P < .01$). Postseason sample sizes are much smaller (and have a higher variability) than preseason and in season sample sizes because only a small percentage of schools participated in the postseason tournaments in any sport and not all of those were a part of the Injury Surveillance System sample. Numbers do not always sum to totals because of missing division or season information.

Table 4. Practice Injury Rates, Men's Spring Football, 1988–1989 Through 2003–2004*

	Total No. of Practices Reported	Practice Injury Rate per 1000 Athlete-Exposures	95% Confidence Interval
Division I	10 427	10.05	9.83, 10.27
Division II	5127	8.58	8.26, 8.89
Both divisions	15 554	9.62	9.44, 9.80

*Injury rates differ between divisions ($P < .01$).

Table 5. Percentage of Game and Practice Injuries by Major Body Part, Men's Football, 1988–1989 Through 2003–2004

Body Part	Fall Games	Fall Practices	Spring Practices
Head/neck	11.5	10.1	9.8
Upper extremity	22.6	20.1	22.9
Trunk/back	9.9	13.2	9.9
Lower extremity	54.7	50.8	55.7
Other/system	1.4	5.9	1.6

ment (6.17 versus 0.46 injuries per 1000 A-Es, rate ratio = 13.4, 95% CI = 12.9, 13.9, $P < .01$), 12 times more likely to sustain an ankle ligament sprain (5.39 versus 0.45, rate ratio = 12.0, 95% CI = 11.5, 12.5, $P < .01$), and 11 times as likely to sustain a concussion (2.34 versus 0.21 injuries per 1000 A-Es, rate ratio = 11.1, 95% CI = 10.5, 11.8, $P < .01$).

Mechanism of Injury

The 3 primary injury mechanisms—player contact, other contact (eg, balls, blocking dummies, ground), and no contact—in fall games, fall practices, and spring practices with division and years combined are displayed in Figure 2. The majority of game (78%, $n = 23\,993$), fall practice (57%, $n = 223\,950$), and spring practice (69%, $n = 7578$) injuries resulted from player contact. Noncontact (ie, no direct contact to the injured body part) was the second highest injury mechanism in all 3 activities but was less than half as frequent as player contact.

Severe Injuries: 10+ Days of Activity Time Loss

The top injuries that resulted in at least 10 consecutive days of restricted or total loss of participation and their primary injury mechanisms, combined across divisions and years, are presented in Table 7. Time loss of 10+ days was, for this analysis, considered a measure of severe injury. A total of 27% of fall game injuries, 25% of fall practice injuries, and 34% of spring practice injuries resulted in a loss of participation for at least 10 days. In all 3 activities, knee and ankle problems, primarily resulting from player contact, accounted for most of the severe injuries. Concussions represented 4% of severe time-loss injuries in games and 3% in both fall and spring practices (data not shown). The time-loss data may be more variable for spring practices due to the limited “season” length (up to 15 practices) for that activity.

The top severe (10+ days of time loss) injuries in the fall

Table 6. Most Common Game and Practice Injuries, Men's Football, 1988–1989 Through 2003–2004*

Body Part	Injury Type	Frequency	Percentage of Injuries	Injury Rate per 1000 Athlete-Exposures	95% Confidence Interval
Fall games†					
Knee	Internal derangement	5493	17.8	6.17	6.01, 6.34
Ankle	Ligament sprain	4799	15.6	5.39	5.24, 5.54
Head	Concussion	2085	6.8	2.34	2.24, 2.44
Upper leg	Contusion	1129	3.7	1.27	1.19, 1.34
Upper leg	Muscle-tendon strain	1103	3.6	1.24	1.17, 1.31
Shoulder	Acromioclavicular joint injury	869	2.8	0.98	0.91, 1.04
Shoulder	Ligament sprain	808	2.6	0.91	0.85, 0.97
Shoulder	Subluxation	649	2.1	0.73	0.67, 0.79
Shoulder	Contusion	584	1.9	0.66	0.60, 0.71
Pelvis, hip	Muscle-tendon strain	581	1.9	0.65	0.60, 0.71
Shoulder	Muscle-tendon strain	567	1.8	0.64	0.58, 0.69
Pelvis, hip	Contusion	563	1.8	0.63	0.58, 0.68
Lower leg	Contusion	557	1.8	0.63	0.57, 0.68
Neck	Nerve injury	539	1.8	0.61	0.55, 0.66
Ribs	Contusion	445	1.4	0.50	0.45, 0.55
Neck	Muscle-tendon strain	407	1.3	0.46	0.41, 0.50
Knee	Contusion	398	1.3	0.45	0.40, 0.49
Lower back	Muscle-tendon strain	357	1.2	0.40	0.36, 0.44
Patella	Patella or patella tendon injury	341	1.1	0.38	0.34, 0.42
Foot	Ligament sprain	325	1.1	0.37	0.33, 0.40
Hand	Fracture	303	1.0	0.34	0.30, 0.38
Unspecified‡	Unspecified	298	1.0	0.33	0.30, 0.37
Lower leg	Fracture	294	1.0	0.33	0.29, 0.37
Fall practices§					
Knee	Internal derangement	5075	12.0	0.46	0.44, 0.47
Ankle	Ligament sprain	5011	11.8	0.45	0.44, 0.46
Upper leg	Muscle-tendon strain	4518	10.7	0.41	0.39, 0.42
Head	Concussion	2319	5.5	0.21	0.20, 0.22
Pelvis, hip	Muscle-tendon strain	2196	5.2	0.20	0.19, 0.21
General body	Heat illness	1632	3.9	0.15	0.14, 0.15
Lower back	Muscle-tendon strain	1143	2.7	0.10	0.10, 0.11
Shoulder	Subluxation	1006	2.4	0.09	0.08, 0.10
Shoulder	Muscle-tendon strain	878	2.1	0.08	0.07, 0.08
Shoulder	Ligament sprain	840	2.0	0.08	0.07, 0.08
Unspecified‡	Unspecified	811	1.9	0.07	0.07, 0.08
Upper leg	Contusion	798	1.9	0.07	0.07, 0.08
Shoulder	Acromioclavicular joint injury	769	1.8	0.07	0.06, 0.07
Patella	Patella or patella tendon injury	683	1.6	0.06	0.06, 0.07
Neck	Muscle-tendon strain	626	1.5	0.06	0.05, 0.06
Shoulder	Contusion	607	1.4	0.05	0.05, 0.06
Neck	Nerve injury	586	1.4	0.05	0.05, 0.06
Lower leg	Contusion	444	1.0	0.04	0.04, 0.04
Hand	Fracture	414	1.0	0.04	0.03, 0.04
Foot	Ligament sprain	404	1.0	0.04	0.03, 0.04
Spring practices					
Knee	Internal derangement	1793	16.4	1.58	1.50, 1.65
Ankle	Ligament sprain	1519	13.9	1.34	1.27, 1.40
Upper leg	Muscle-tendon strain	1179	10.8	1.04	0.98, 1.10
Head	Concussion	612	5.6	0.54	0.50, 0.58
Pelvis, hip	Muscle-tendon strain	377	3.4	0.33	0.30, 0.36
Shoulder	Subluxation	344	3.1	0.30	0.27, 0.33
Shoulder	Ligament sprain	218	2.0	0.19	0.17, 0.22
Shoulder	Acromioclavicular joint injury	215	2.0	0.19	0.16, 0.21
Lower back	Muscle-tendon strain	198	1.8	0.17	0.15, 0.20
Shoulder	Muscle-tendon strain	193	1.8	0.17	0.15, 0.19
Upper leg	Contusion	190	1.7	0.17	0.14, 0.19
Patella	Patella or patella tendon injury	176	1.6	0.15	0.13, 0.18
Neck	Nerve injury	152	1.4	0.13	0.11, 0.15
Unspecified‡	Unspecified	147	1.3	0.13	0.11, 0.15
Hand	Fracture	146	1.3	0.13	0.11, 0.15
Shoulder	Contusion	132	1.2	0.12	0.10, 0.14
Neck	Muscle-tendon strain	120	1.1	0.11	0.09, 0.12
Foot	Ligament sprain	115	1.1	0.10	0.08, 0.12
Lower leg	Contusion	108	1.0	0.09	0.08, 0.11
Shoulder	Dislocation	106	1.0	0.09	0.08, 0.11

*Only injuries that accounted for at least 1% of all injuries are included.

†“Fall games” includes any game that occurred between the start of preseason and the end of postseason.

‡“Unspecified” indicates injuries that could not be grouped into existing categories but that were believed to constitute reportable injuries.

§“Fall practices” includes any practice that occurred between the start of preseason and the end of postseason.




Figure 2. Game and practice injury mechanisms, all injuries, men's football, 1988–1989 through 2003–2004 (n = 30 797 game injuries, 42 355 fall practice injuries, and 10 943 spring practice injuries). "Other contact" refers to contact with items such as balls, blocking dummies, or the ground.

preseason and in season (games and practices combined) are presented in Table 8. The percentages of knee and ankle injuries were similar between the preseason and the regular season, but muscle-tendon strains of the upper leg were more common in preseason practices.

Heat Illness

Heat illness rates by year in preseason fall practices from 1988–1989 through 2003–2004 are shown in Table 9. The table indicates that the rate of reported heat illnesses in preseason fall practices was relatively stable over the years. (Note that, due to missing data, the practice injury rate for 1996–

Table 8. Summary of Preseason (Practice) Versus In Season (Game and Practice) 10+ Day Time-Loss Injuries, Men's Football, 1988–1989 Through 2003–2004

Body Part	Injury Type	Frequency	Percentage of All 10+ Day Time-Loss Injuries
Preseason practices			
Knee	Internal derangement	1602	26.0
Upper leg	Muscle-tendon strain	637	10.3
Ankle	Ligament sprain	536	8.7
In season (games and practices combined)			
Knee	Internal derangement	4050	32.8
Ankle	Ligament sprain	1459	11.8

1997 was imputed from adjacent years). We caution that these data contain only heat illnesses that resulted in time lost from the sport.

Game Injuries

The general game injury distribution by weighted position (over all years of analysis) is shown in Figure 3. The percentages of injury by position were calculated and then divided by the number of players at each position ("position weight") as follows: offensive line = 6, wide receiver = 2, quarterback = 1, running back = 2, defensive line = 4, linebacker = 3, and defensive back = 4. The offensive players with the highest number of injuries (by weighted position) were the quarterback (18%) and the running back (20%). All 3 general defensive positions accounted for approximately 15% of reported game injuries.

The game concussion injury distribution by weighted position (over all years of analysis) is shown in Figure 4. The offensive players with the highest number of concussions (by weighted position) were the running back (17%) and the quarterback (28%). The defensive player with the highest number

Table 7. Most Common Game and Practice Injuries Resulting in 10+ Days of Activity Time Loss, Men's Football, 1988–1989 Through 2003–2004

Body Part	Injury Type	Frequency	Percentage of Severe Injuries	Most Common Injury Mechanism
Fall games (27.0% of all injuries required 10+ days of time loss)				
Knee	Internal derangement	3017	36.2	Player contact
Ankle	Ligament sprain	1032	12.4	Player contact
Other		4275	51.4	
Total		8324		
Fall practices (24.9% of all injuries required 10+ days of time loss)				
Knee	Internal derangement	2787	26.4	Player contact
Ankle	Ligament sprain	1014	9.6	Player contact
Upper leg	Muscle-tendon strain	861	8.2	No contact
Other		5898	55.9	
Total		10 560		
Spring practices (33.5% of all injuries required 10+ days of time loss)				
Knee	Internal derangement	1109	30.2	Player contact
Ankle	Ligament sprain	410	11.2	Player contact
Upper leg	Muscle-tendon strain	290	7.9	No contact
Other		1863	50.7	
Total		3672		

Table 9. Fall Preseason Heat Illness by Year, Men's Football, 1988–1989 Through 2003–2004*

Year	No. of Heat Illnesses	Percentage of All Preseason Practice Injuries	Rate of Preseason Practice Heat Illnesses per 1000 Athlete-Exposures
1988–1989	77	5.35	0.43
1989–1990	38	2.81	0.22
1990–1991	68	3.90	0.29
1991–1992	78	4.22	0.31
1992–1993	41	2.82	0.18
1993–1994	123	6.12	0.47
1994–1995	94	4.98	0.37
1995–1996	246	12.49	0.98
1996–1997	123	6.80	0.52
1997–1998	67	4.54	0.33
1998–1999	86	5.20	0.41
1999–2000	145	5.63	0.48
2000–2001	124	6.75	0.56
2001–2002	163	7.59	0.63
2002–2003	103	5.45	0.41
2003–2004	111	6.02	0.53
Average	104.3	5.60	0.44

*Spring practice rates of heat illness could not be calculated because of small numbers of cases (14 from 1988–1989 through 2003–2004). Preseason practices are team practices before the first regular-season contest. Practice counts for fall 1996 were imputed from adjacent years.

of concussions was the defensive back (14%). Offensive players appeared to receive a higher number of concussions than defensive players.

Rule Changes

We analyzed the data for spring football, comparing 1988–1989 through 1996–1997 and 1997–1998 through 2003–2004 seasons, to assess the effect of a rule change that limited the amount of contact in spring practice (see “Commentary” section). The injury rate was slightly greater for the former time span than the latter, a finding that was statistically significant ($P < .01$) but perhaps not clinically significant given that the difference was only 0.38 injuries per 1000 A-Es. However, the spring injury rate declined gradually and steadily since the

highest injury rate (11.08 per 1000 A-Es) reported for the 1997–1998 season (Figure 1).

We also assessed the effect of the spring practice rule change on the rate of anterior cruciate ligament (ACL) injuries and concussions. The ACL injury rates between the 2 periods were similar with 187 injuries for 1988–1989 through 1996–1997 for an injury rate = 0.32 per 1000 A-Es, and 191 injuries for 1997–1998 through 2003–2004 for an injury rate = 0.34 per 1000 A-Es. The rate ratio comparing the 2 periods was 0.94. However, the number of reported concussions actually increased between the 2 periods: 225 concussions occurred in the 1988–1989 through 1996–1997 seasons (0.39 per 1000 A-Es) and 388 concussions in the 1997–1998 through 2003–2004 seasons (0.69 per 1000 A-Es), for a rate ratio of 0.57, 95% CI = 0.48, 0.67.

Knee Injuries

Approximately 85% of knee internal derangements were classified as new injuries. There were 3 major types of knee injuries ($n = 4841$) during games and practices: ACL, posterior cruciate ligament (PCL), and menisci. Athletes were more likely to injure the ACL, PCL, and menisci in fall games versus practices, and the contact injury mechanisms were more frequent than noncontact injury mechanisms for all 3 types of injury (Table 10).

Further description of the mechanism of knee injury is provided in Table 11. Potentially illegal plays (blocking below the waist, clipping in the legal zone, and clipping) accounted for 7% of the ACL injuries, 9% of the PCL injuries, and 6% of the meniscus injuries.

A total of 45% of all knee internal derangements ($n = 4767$) resulted in less than 10 days of time loss. However, for those knee injuries with more than 10 days of time loss, Table 12 presents a summary of treatments (operative versus nonoperative). For the reported ACL injuries ($n = 2159$), 78% of the cases were operative, but we have no data on the operative procedures performed and, therefore, cannot assume that all the patients received some type of ACL reconstruction. When the PCL was injured, 39% underwent a surgical procedure. A limitation of this study was the inability to identify isolated versus multiple ligament injuries in the treatment and surgical intervention.




Figure 3. Game injuries by weighted position, men's fall football, 1988–1989 through 2003–2004. Percentages of injury by position were calculated and then divided by the weight of position as follows: offensive line = 6, wide receiver = 2, quarterback = 1, running back = 2, defensive line = 4, linebacker = 3, and defensive back = 4.




Figure 4. Game concussion injuries by weighted position, men's fall football, 1988–1989 through 2003–2004. Percentages of injury by position were calculated and then divided by the weight of position as follows: offensive line = 6, wide receiver = 2, quarterback = 1, running back = 2, defensive line = 4, linebacker = 3, and defensive back = 4.

Table 10. Game and Practice Injury Rates and Mechanisms for Knee Structures per 1000 Athlete-Exposures, Men's Fall Football, 1988–1989 Through 2003–2004

Event and Knee Structure	Injury Mechanism		Rate Ratio*	95% Confidence Interval
	Contact	Non-contact		
Games				
Anterior cruciate ligament (n = 1059)	0.83	0.36	2.3	2.0, 2.6
Posterior cruciate ligament (n = 183)	0.18	0.02	7.3	4.7, 11.4
Menisci (n = 988)	0.82	0.29	2.9	2.5, 3.3
Practices				
Anterior cruciate ligament (n = 895)	0.05	0.03	1.5	1.3, 1.7
Posterior cruciate ligament (n = 122)	0.01	0.0029	2.8	1.9, 4.2
Menisci (n = 1098)	0.05	0.05	1.2	1.0, 1.3

*Contact versus noncontact. Cases involving other contact or unknown mechanism (n = 496, 10%) are not included.

Shoulder Injuries

The most frequent shoulder injuries were acromioclavicular joint injury, ligament sprains, dislocations/subluxations, contusions, muscle-tendon strains, and nerve injuries (Table 13). All these injuries occurred at a greater than 10-fold higher rate in games relative to practices.

COMMENTARY

Football is a high-impact collision sport, with injuries occurring in both contact and noncontact situations. It is also a complex sport that requires athletes to perform different skills and activities, depending on the position played. The sport has a major fall season and a minor spring season. Players' physical characteristics (eg, age, height, weight) vary widely, both within a team and among NCAA divisions.

The results from this 16-year study period showed little variation in the individual injury rates for games, fall practices, or spring practices over time (Figure 1). This is likely because the basic characteristics of the team and the typical exposure to football have not changed drastically over the years. Pow-

ell,³ using data from the late 1970s and early 1980s (the National Athletic Injury/Illness Reporting System, or NAIRS), reported that the average team consisted of 93 players exposed to 82 practices sessions, and the average game squad was 51 players who team averaged 11 games. The data from the ISS are virtually identical (Table 2) to the NAIRS data. Thus, exposure frequency in collegiate football has not changed much since NAIRS began in 1974. Differences among Divisions I, II, and III in their injury rates for preseason practices and in-season practices and games also appeared to be minimal.

The ISS injury rates were approximately 36 per 1000 A-Es for fall games, 4 per 1000 A-Es for fall practices, and 10 per 1000 A-Es for spring practices. Powell and Dompier⁴ reported that the time-loss football injury rate was 9.3 per 1000 A-Es in Division I, 10.2 per 1000 A-Es in Division II, and 10.4 per 1000 A-Es in Division III during 2 seasons (2000–2001 and 2001–2002 academic years). Although these injury rates are lower than those seen in the ISS data, Powell and Dompier⁴ did not distinguish between games and practices.

Other findings were that the game injury rate was more than 9 times higher than the in-season practice injury rate and that the spring injury rate was 2 times higher than the fall in-season practice injury rate. The game and practice injury rates are consistent with the data previously published by Powell.³ He reported the game injury rate was 8 to 9 times greater than the fall practice rate and stated that injury prevention programs should focus on game conditions rather than practices. However, when placed in the context for football, these differences in the injury rates between games and practices are logical. The intensity level and speed are generally considered higher in games than in practices, increasing the magnitude of collisions and, thus, increasing the risk of injury. Most player-to-player contact occurs during games, preseason practices, and spring practices. Practice times are usually longer and involve more contact work during the preseason and spring season, possibly accounting for the higher injury rates in these sessions.

In spring football, the concept of practice may be vastly different than in the fall season. The roster is typically smaller in the spring, with the seniors having reached the end of eligibility and the new signees not yet in school; therefore, the number of repetitions and injury exposures are increased for all participants. Also, the level of competition may be higher

Table 11. Knee Injury Mechanisms for Games and Practices in Detail, Men's Fall Football, 1988–1989 Through 2003–2004*

Injury Mechanism	Anterior Cruciate Ligament				Posterior Cruciate Ligament				Meniscus			
	Games		Practices		Games		Practices		Games		Practices	
	n	Percent-age	n	Percent-age	n	Percent-age	n	Percent-age	n	Percent-age	n	Percent-age
Blocked below waist	84	7.2	37	3.7	17	7.5	8	5.3	70	6.6	31	2.5
Tackling	92	7.9	67	6.8	12	5.3	6	4.0	102	9.6	85	6.9
Being tackled	195	16.8	106	10.7	52	22.9	23	15.3	167	15.7	93	7.6
Blocking	114	9.8	130	13.2	24	10.6	20	13.3	148	13.9	188	15.3
Being blocked	174	15.0	124	12.6	34	15.0	18	12.0	159	15.0	111	9.0
Impact with playing surface	7	0.6	23	2.3	24	10.6	21	14.0	13	1.2	32	2.6
Stepped on/fallen on/kicked	97	8.4	82	8.3	30	13.2	10	6.7	97	9.1	91	7.4
Sprints/running	8	0.7	16	1.6	1	0.4	3	2.0	6	0.6	38	3.1
Blocking a kick/punt	3	0.3	1	0.1	1	0.4	0	0.0	0	0.0	3	0.2
Noncontact (rotation about a planted foot)	332	28.6	343	34.8	21	9.3	25	16.7	252	23.7	442	35.9
Noncontact other	6	0.5	13	1.3	0	0.0	1	0.7	9	0.8	19	1.5
Overuse	1	0.1	9	0.9	0	0.0	3	2.0	4	0.4	37	3.0
Catching/blocking pass	11	0.9	17	1.7	2	0.9	7	4.7	11	1.0	25	2.0
Clipped by an offensive lineman in legal clip zone	6	0.5	3	0.3	1	0.4	1	0.7	2	0.2	2	0.2
Clipped	18	1.6	6	0.6	4	1.8	2	1.3	13	1.2	10	0.8
Impact with padded cast	13	1.1	10	1.0	4	1.8	2	1.3	9	0.8	24	1.9
Total	1161	100.0	987	100.0	227	100.0	150	100.0	1062	100.0	1231	100.0

*Caution is advised when interpreting individual categories with small sample sizes.

Table 12. Treatment of Anterior and Posterior Cruciate Ligament Injuries With 10+ Days of Time Loss, Men's Fall Football, 1988–1989 Through 2003–2004*

Treatment	Anterior Cruciate Ligament				Posterior Cruciate Ligament			
	Games		Practices		Games		Practices	
	n	Percentage	n	Percentage	n	Percentage	n	Percentage
Nonoperative	192	16.4	226	22.9	125	54.1	89	58.9
In-season operation	694	59.3	619	62.7	82	35.5	45	29.8
Postseason operation	258	22.0	105	10.6	17	7.4	4	2.6
Unknown	27	2.3	38	3.8	7	3.0	13	8.6
Total	1171	100.0	988	100.0	231	100.0	151	100.0

*Data with unknown injury treatment not included.

Table 13. Injury Rates and Rate Ratios for Selected Game and Practice Shoulder Injuries, Men's Fall Football, 1988–1989 Through 2003–2004

Shoulder Injury	Injury Frequency		Injury Rate per 1000 Athlete-Exposures		Rate Ratio (Games Versus Practices)	95% Confidence Interval
	Games	Practices	Games	Practices		
Acromioclavicular joint injury	869	769	0.98	0.07	14.14	12.7, 15.4
Ligament sprain	808	840	0.91	0.08	12.04	10.3, 12.5
Dislocation/subluxation	900	1262	1.01	0.11	8.93	8.4, 10.0
Contusion	607	584	0.68	0.05	13.01	12.1, 15.2
Muscle-tendon strain	567	878	0.64	0.08	8.08	7.2, 8.9
Nerve injury	229	257	0.26	0.02	11.15	10.9, 15.5

in the spring. In the fall, many teams scrimmage with their first teams versus their second teams or practice offense and defense with their first and second teams versus scout teams, simulating the opponents' plays. In the spring season, to prepare for the upcoming fall season and without needing to be ready for a game each week, many teams have significantly more contact work than in the fall season, often against a higher level of competition (eg, first-team offense scrimmaging first-team defense) than typically seen in the fall season. In

this instance, the football coaches may have the greatest influence on the injury rate, as they determine the amount of actual contact or hitting, the intensity level, who is contacted (eg, first string versus fourth string), how much contact is received during any given practice, and use of proper techniques and enforcement of the rules (eg, spearing). Games tend to reduce the influence of the coaches over the quantity and nature of body contact, as the game is played at high speed and high intensity and players expect to be involved in contacts.

Location of Injuries

It is important for the certified athletic trainer not only to examine the frequency of injury (Table 5) in games and practices but to consider how the injury rate varies among games and fall and spring practices. The game-to-practice rate ratio indicates that upper leg contusions, acromioclavicular joint injury, knee internal derangements, ankle ligament sprains, and concussions occur at greater than 10-fold higher rates in games compared with practices. The certified athletic trainer must be alert to these conditions and be prepared to evaluate the athlete and render emergency care during games. The frequency distribution for the top 5 injury types for games, fall practices, and spring practices was about the same, accounting for approximately 45% of the total injuries for that session type. In fact, the top 5 injury types were identical between fall practices and spring practices (knee internal derangement, ankle ligament sprain, upper leg muscle-tendon strain, concussion, and pelvis, hip muscle strain). Although the percentage of knee internal derangements was 4.4% greater during spring practices, few differences were seen in the percentages of injuries for upper leg muscle-tendon strain (0.1%), concussion (0.1%), and pelvis, hip muscle-tendon strain (1.8%) among the seasons. However, for games, upper leg contusions were the fourth most common injury, and the percentages of knee internal derangements, ankle ligament sprains, and concussions also increased over those for fall and spring practices. Buckley and Powell⁵ found a similar distribution pattern of injury frequency by body part using the NAIRS data, and Orchard and Powell⁶ reported that the knee accounted for 13% and the ankle for 12% of all injuries in National Football League games from 1989–1998. As previously discussed, more contact and higher-velocity collisions would result in a higher number of injuries during games.

Spring Practice Injuries

In 1997, in an attempt to decrease the injury rate disparity between fall and spring football, the NCAA Committee on Competitive Safeguards and Medical Aspects of Sports recommended changes in spring practices that limited the amount of full-team, full-contact practice.⁷ Of the 15 practices allowed for spring football, 3 are mandated to be noncontact, and headgear is the only piece of protective equipment allowed during noncontact practices. The first 2 practices must be noncontact, and 1 other noncontact practice must be among the 15 practices. Of the 12 permissible contact practices, 8 may involve tackling, and no more than 3 of the 8 tackling sessions may be devoted primarily (greater than 50% of practice time) to 11-on-11 scrimmages. This rule went into effect for the 1997–1998 spring practices. Our analyses detected a statistically significant decline in the rate of spring practice injuries ($P < .01$) after the rule change.

Albright et al.,⁸ in a controlled study using the Big Ten Conference Sports Injury Surveillance System, analyzed the spring football injury rate before and after the implementation of the new practice rule. From 1992 to 1997, they reported a spring injury rate of 19.8 per 1000 A-Es; the rate declined to 16.4 per 1000 A-Es for the 1998–2000 seasons (2 seasons of spring football data) for a decrease of 1.2 times. The authors concluded that the injury rate in spring practices was three-fold greater than for fall practices,⁸ and this rate is similar to the ISS data. Both the ISS data and the Albright et al.⁸ study demonstrate variations in the overall injury rate with the imple-

mentation of the spring practice rule. We believe this topic requires further study to determine the longitudinal effect of this rule change.

We also wanted to determine the effects of the spring practice rule change on knee injuries and concussions. The injury rate for ACL injuries between the 2 periods was approximately the same, as evidenced by a rate ratio close to 1. Even though the type of protective equipment worn during practice between the 2 periods differed, the activities fundamental to the sport of football remained the same. Athletes still participated in drills and activity sessions during spring football, which did not change the number of ACL injuries. The rate of reported concussions increased between the 2 periods. We attribute this increased number of concussions in the latter period in part to greater knowledge and recognition by certified athletic trainers and the medical staffs to the signs and symptoms of concussion and to better management of this injury.^{9,10} Thus, the increased rate of concussion during the second period may not necessarily be due to an actual increase in the number of concussions or the change in practice rules but to better and more accurate recognition by certified athletic trainers. It will be interesting to continue to monitor this trend of concussions during spring football practice in future investigations.

Preseason Acclimatization Period

The other major rule change implemented by the NCAA was the 5-day acclimatization period for preseason football that went into effect in the 2002–2003 season. The rule states that during the first 5 days of practice, student-athletes shall not engage in more than 1 on-field practice per day, not to exceed 3 hours in length. During the first 2 days of the acclimatization period, helmets are the only piece of protective equipment participants may wear. During the third and fourth days of the acclimatization period, helmets and shoulder pads are the only pieces of protective equipment student-athletes may wear. During the final day of the 5-day period and on any days thereafter, student-athletes may practice in full pads. However, after the 5-day acclimatization period, an institution may not conduct multiple on-field practice sessions on consecutive days. Student-athletes cannot engage in more than 3 hours of on-field practice activities on those days during which 1 practice is permitted. Student-athletes shall not engage in more than 5 hours of on-field activities on those days during which more than 1 practice is permitted. On those days when institutions conduct multiple practice sessions, student-athletes must be provided with at least 3 hours of continuous recovery time between the end of the first practice and the start of the last practice that day. Only 1 season of data with the new acclimatization rule change was available, which was insufficient for comparing injury rates or determining heat illness trends before and after the rule change. Future researchers should assess changes in these rates with the implementation of this new rule.

Knee Injuries

Knee injuries are common in football players. Athletes were more likely to injure the ACL, PCL, and menisci in games versus practices, and contact injury mechanisms were more frequent than noncontact injury mechanisms. Although we do not know the exact mechanisms of all injuries, most of the significant knee injuries (especially of the ACL) are believed

to occur with the foot planted and force applied. With contact mechanisms, the athlete who is injured may not be able to pick his foot up and eliminate the ground contact.

Injury prevention measures should continue to focus on the use of safe sport techniques and elimination of illegal plays. Neuromuscular training programs have shown promise in reducing ACL injury risk in females but have not been thoroughly studied in males.

Shoulder Injuries

The shoulder is a complex joint that receives substantial loads and axial forces during football. As with the other major joints, the greatest risk of injury was during games. The fall game versus practice rate ratios were all above 10 for acromioclavicular joint injury, ligament sprains, contusions, and nerve injuries (Table 13). Although the ISS cannot differentiate among the types of dislocations or the specific nerves involved, certified athletic trainers should be cognizant of these injury patterns. Anecdotally, in our clinical experience, we are seeing more shoulder injuries in linemen: in particular, labral conditions and posterior glenohumeral instability. In recent years, offensive linemen have started using blocking techniques in which they punch and block the opponent using extended arms. Although effective for blocking, this technique places greater stress on the glenoid labrum and posterior capsule structures, which could be a factor in increased labral injuries and posterior shoulder instability. This topic requires further analysis.

Exertional Heat Illnesses and Concussions

The other 2 injury trends that merit further attention are exertional heat illness and sport-related concussion, as both are high-profile issues with the potential for catastrophic outcomes. The National Athletic Trainers' Association has published position statements to guide practitioners in the recognition and management of exertional heat illness¹¹ and sport-related concussions.¹² The preseason period (defined as the first day of practice to the last practice before the first game) has been postulated to be the period when athletes are at the greatest risk of heat illness. We found an overall heat illness injury rate of 0.44 per 1000 A-Es for the preseason period, which accounted for 5.6% of all the reported injuries during this time (Table 9). Almost all heat illnesses occurred in preseason practices ($n = 1567/1632$), but it is important to remember that not all game heat illnesses may be captured because the ISS injury definition states that only injuries resulting in time loss are reportable. In a study of 5 NCAA football institutions in the southeastern United States from August through October 2003, Cooper et al¹³ noted an overall heat-related injury rate of 4.10 per 1000 A-Es and reported that exertional heat cramps accounted for 73% of the total exertional heat illnesses, followed by heat exhaustion at 21% and heat syncope at 6%. The first 3 weeks of August presented the greatest risk of exertional heat illnesses to collegiate football players. Certified athletic trainers must continually monitor athletes' responses during games and practices for signs and symptoms of exertional heat illnesses and the environment.

The percentage of concussion was 5.5% for fall practices (0.21 per 1000 A-Es), 6.8% for games (2.34 per 1000 A-Es), and 5.6% (0.05 per 1000 A-Es) for spring practices. When the

frequencies are standardized using A-Es, the fall game injury rate was 11 times higher than the fall practice rate (Table 6). Several authors have reported similar or lower concussion injury rates,^{14,15} but this percentage of concussions is similar to that in another report.¹⁶ Clearly, the greatest risk of concussion is in games, which have the greatest risk of high-speed collisions. Using an accelerometer system, Duma et al¹⁷ estimated that the average peak head linear acceleration was 32 ± 25 g in collegiate football players.

In the ISS data, the quarterback and running back had the greatest frequency of game concussions (Figure 4). It is not surprising that the offensive player who was struck in the open field was more susceptible to concussion than the other positions. In a previous report of concussion in collegiate football players,¹⁴ the concussion injury rate for the quarterback was 0.83 per 1000 A-Es; for the running back, 0.71 per 1000 A-Es; and for the wide receiver, 0.54 per 1000 A-Es. Higher rates were reported for the linebacker (0.99 per 1000 A-Es) and offensive lineman (0.95 per 1000 A-Es).¹⁵ About 15% of the athletes with concussions lost 10 or more days from participation (games = 15%, practices = 13%, spring practices = 15%).

The National Athletic Trainers' Association published a position statement on the management of sport-related concussion that provided 36 recommendations for certified athletic trainers to use in the recognition, management, assessment, and return to play after a concussion.¹² The general standard for management of a concussion includes a physical assessment and evaluation, including a neurologic examination, imaging studies, assessment of concussion-related self-report symptoms, measures of cognition and balance, and an exertional testing program before returning to play. Several authors^{12,18} have recommended that an athlete must be symptom-free for 7 days, both at rest and with exertion, before return to play can be considered. McCrea et al,¹⁹ in their NCAA study, tracked the recovery of concussion-related symptoms, neuropsychological performance, and balance in 94 athletes with concussion. Concussion related-symptoms gradually resolved within 7 days, cognitive functioning improved to baseline levels within 5 to 7 days, and balance deficits also returned to baseline within 3 to 5 days after injury. Recent advances by several manufacturers toward better-constructed helmets may aid in reducing the impact force and have a role in the short-term and long-term effects of the concussive injury.²⁰ Along these same lines, face mask technology has also improved, allowing more force to be absorbed and reducing the amount of force transmitted to the head.

CONCLUSIONS

This is the first published analysis of the ISS data over such an extended time. The ISS uses a time-loss definition for a reportable injury and relies on the volunteer efforts of certified athletic trainers to record and report the information. Frequency distributions, injury rates, and rate ratios were used to identify general injury patterns and trends. Games clearly presented the greater risk of injury to athletes than practices. Although injuries are an unfortunate but expected aspect of competitive football, the sports medicine team's goal must be injury prevention. This can be a difficult and complicated process, especially because the fundamentals of the game have not changed dramatically during the last 40 years. Certified athletic trainers, coaches, and administrators should focus on

improving equipment, rule changes, and modifications in sport-specific and coaching techniques. The development of new and improved equipment may aid in preventing and reducing injuries. Not only have the helmet and face mask changed during this study, but shoulder pads, flak jackets, and other padding garments have also seen technologic improvements. The enhanced force dissipation and the materials used in the equipment may aid in the reduction of forces transmitted to the body. Mouthguards are also better constructed with improved materials, and many institutions are using custom mouthguards to reduce dental injuries. Also, prophylactic knee and ankle taping and bracing has been studied with the aims of reducing injury rates and increasing the awareness of these injuries. Consistent data need to be collected so that we can determine the effectiveness of these protective devices and the resulting changes in injury rates.

It has been thought that strength and conditioning programs would have an effect on the injury rate. During the 16-year reporting period, although strength and conditioning programs have improved dramatically, the injury rates for games and practices have remained largely unchanged. Strength and conditioning programs may help to decrease injuries through improved conditioning. Yet we can also argue that stronger, faster athletes increase the speed and collision forces, causing more injuries.

In 2004, the NCAA Football Rules Committee changed the rules related to spearing and head-down contact in football, based on recommendations by a task force created specifically to reduce the incidence of head and cervical spine injuries in football players. As a part of this rule change, extensive education was initiated for athletes, coaches, and officials, including the creation of a videotape on proper tackling techniques and a locker-room poster to increase awareness. As injury patterns that may be rule based are identified, the rules must be re-examined with the goal of lowering the injury rate. In many instances, rule changes are closely related to technique changes and vice versa. The effects of rule and technique changes on the integrity of the game are an interesting topic for discussion.

Given the range of potential injuries, football creates unique challenges. The certified athletic trainer and the rest of the medical team must be prepared to evaluate, treat, and rehabilitate a myriad of injuries and illnesses. At 1 university, physician evaluations resulted in 79% musculoskeletal diagnoses and 21% general medical condition diagnoses, and football accounted for 22% of all physician evaluations in a comprehensive intercollegiate sports program.²¹ Other authors⁴ found that 57% of the therapeutic treatments by certified athletic trainers were associated with non-time-loss injuries and that more therapeutic treatments were required for these than for time-loss injuries during the course of a year. These data demonstrate the need for appropriate medical care for those schools that sponsor collegiate football. This is especially true for games, as the injury rate increased substantially over practices, but it is also important during practices as the volume of injuries increases. A disparity in medical coverage may occur whereby games may have multiple medical providers (certified athletic trainers, physicians, emergency medical technicians) and practices have significantly fewer. Although the injury rate was higher for games, these data also demonstrate that a greater number of injuries occurred during practices. This finding makes sense as practice sessions were far more numerous than games. The National Athletic Trainers' Association

published the recommendations and guidelines for appropriate medical coverage of collegiate athletes to help guide institutions in determining the number of certified athletic trainers needed for a comprehensive athletic health care program.²²

Football is a sport contested at many levels, with players having a variety of shapes and sizes, playing at different speeds and intensities. The reduction in catastrophic cervical spine injuries after spear tackling was prohibited provides incentive to pursue additional injury prevention endeavors in football.⁷ Future NCAA data collection and analysis will allow us to identify and study injury trends. Strategies to prevent injury will be developed, implemented, and evaluated during specific periods with the goal of improving the safety of the game for the participants. Evaluation of injury trends and risks in collegiate football by the sports medicine community through detailed epidemiologic investigations will allow us to improve the sport's safety for all participants.

ACKNOWLEDGMENTS

We thank John Powell, PhD, ATC, of Michigan State University, and Cathleen Brown, PhD, ATC, of the University of Georgia, for their insightful review of the manuscript.

DISCLAIMER

The conclusions in the Commentary section of this article are those of the Commentary authors and do not necessarily represent the views of the National Collegiate Athletic Association.

REFERENCES

1. 1981/82–2004/05 NCAA Sports Sponsorship and Participation Rates Report. Indianapolis, IN: National Collegiate Athletic Association; 2006.
2. Dick R, Agel J, Marshall SW. National Collegiate Athletic Association Injury Surveillance System commentaries: introduction and methods. *J Athl Train.* 2007;42:173–182.
3. Powell JW. Pattern of knee injuries associated with college football 1975–1982. *Athl Train J NATA.* 1985;20:104–109.
4. Powell JW, Dompier TP. Analysis of injury rates and treatment patterns for time-loss and non-time-loss injuries among collegiate student-athletes. *J Athl Train.* 2004;39:56–70.
5. Buckley WE, Powell J. NAIRS: an epidemiological overview of the severity of injury in college football 1975–1980 seasons. *Athl Train J NATA.* 1982;17:279–282.
6. Orchard JW, Powell JW. Risk of knee and ankle sprains under various weather conditions in American football. *Med Sci Sports Exerc.* 2003;35:1118–1123.
7. Reider B. An ounce of prevention. *Am J Sports Med.* 2004;32:1383–1384.
8. Albright, JP, Powell JW, Martindale A, et al. Injury patterns in Big Ten Conference football. *Am J Sports Med.* 2004;32:1394–1404.
9. Ferrara MS, McCreia M, Peterson CL, Guskiewicz KM. A survey of practice patterns in concussion assessment and management. *J Athl Train.* 2001;36:145–149.
10. Notebaert A., Guskiewicz KM. Current trends in athletic training practice for concussion assessment and management. *J Athl Train.* 2005;40:320–325.
11. Binkley HM, Beckett J, Casa DJ, Kleiner DM, Plummer PE. National Athletic Trainers' Association position statement: exertional heat injuries. *J Athl Train.* 2002;37:329–343.
12. Guskiewicz KM, Bruce SL, Cantu RC, et al. National Athletic Trainers' Association position statement: management of sport-related concussion. *J Athl Train.* 2004;39:280–297.
13. Cooper ER, Ferrara MS, Broglio SP. Exertional heat illness and environ-

- mental conditions during a single football season in the Southeast. *J Athl Train*. 2006;41:332–336.
14. Alles W, Powell JW, Buckley WE, Hunt EE. Three year summary of NAIRS football data. *Athl Train J NATA*. 1980;15:98–100.
 15. Buckley WE. Concussion in college football: a multivariate analysis. *Am J Sports Med*. 1988;16:51–56.
 16. Guskiewicz KM, McCrea M, Marshall SW, et al. Cumulative effects associated with recurrent concussion in collegiate football players: the NCAA Concussion Study. *JAMA*. 2003;290:2549–2555.
 17. Duma SM, Manoogian SJ, Bussone WR, et al. Analysis of real-time head accelerations in collegiate football players. *Clin J Sport Med*. 2005;15:3–8.
 18. Cantu RC. Posttraumatic retrograde and anterograde amnesia: pathology and implication in grading and safe return to play. *J Athl Train*. 2001;36:244–248.
 19. McCrea M, Guskiewicz KM, Marshall SW, et al. Acute effects and recovery time following concussion in collegiate football players: the NCAA Concussion Study. *JAMA*. 2003;290:2556–2563.
 20. Collins M, Lovell MR, Iverson GL, Ide T, Maroon J. Examining concussion rates and return to play in high school football players wearing newer helmet technology: a three-year prospective cohort study. *Neurosurgery*. 2006;58:275–286.
 21. Steiner ME, Quigley B, Wang F, Balint CR, Boland AL. Team physicians in college athletics. *Am J Sports Med*. 2005;33:1545–1551.
 22. NATA recommendations and guidelines for appropriate medical coverage of intercollegiate athletics, revised, May 2003. Available at: <http://www.nata.org/publicinformation/files/amciarecs%20andguidesrevised.pdf>. Accessed January 16, 2007.
-

Randall Dick, MS, FACS, contributed to conception and design; analysis and interpretation of the data; and drafting, critical revision, and final approval of the article. Michael S. Ferrara, PhD, ATC, contributed to analysis and interpretation of the data and drafting, critical revision, and final approval of the article. Julie Agel, MA, ATC, contributed to conception and design; analysis and interpretation of the data; and drafting, critical revision, and final approval of the article. Ron Courson, ATC, PT, NREMT-I, CSCS, contributed to analysis and interpretation of the data and drafting, critical revision, and final approval of the article. Stephen W. Marshall, PhD, contributed to conception and design; analysis and interpretation of the data; and drafting, critical revision, and final approval of the article. Michael J. Hanley, MS, ATC, and Fred Reifsteck, MD, contributed to analysis and interpretation of the data and drafting, critical revision, and final approval of the article.

Address correspondence to Michael S. Ferrara, PhD, ATC, University of Georgia, Department of Exercise Science, 300 River Road, Athens, GA 30602. Address e-mail to mferrara@uga.edu.