

CAFFEINE INTAKE AMONG LAW ENFORCEMENT OFFICERS PARTICIPATING IN SHIFT WORK

ABSTRACT

Objective: To investigate caffeine consumption among law enforcement officers during day shifts (DAY) and night shifts (NIGHT); to assess the types of caffeine-containing products and the frequency of product intake during DAY and NIGHT; to identify relationships between caffeine products and selected demographic characteristics; to identify relationships between caffeine intake and job-related characteristics.; and to identify associative relationships between caffeine intake and perceived concentration level, caffeine side effects, and tobacco use.

Participants: Police officers (PO) and Sheriff Department Deputies (SDD) in a rural region in Eastern North Carolina. Useable sample of 75 (N = 75).

Methods: Anonymous, self-administered caffeine food-frequency questionnaires (FFQ; three for DAY, three for NIGHT) and demographic questionnaire. Nonparametric tests were used. The Wilcoxon signed rank test was used to compare related samples. Spearman's correlation was used to determine relationships between two samples. The Mann-Whitney U test was used to compare two independent samples. The Kruskal-Wallis test was used to compare more than two independent samples.

Results: Caffeine consumption was similar from DAY to NIGHT (225 ± 227 mg DAY; 252 ± 232 mg NIGHT; $p=0.891$). Frequency of tea intake was greater during DAY ($p=0.032$). Greater caffeine consumption was found among (1) SDD versus PO; (2) expert officers (for night only); (3) lower concentration levels (for night only); (4) tobacco users; (5) those who experience caffeine side effects. In addition, younger officers consumed more energy drinks DAY and NIGHT.

Conclusions: It was found that certain job-related characteristics influence caffeine intake, but types of products used and frequency of intake tends to be the same regardless of shift or caffeine side effects. Energy drinks were found to be most popular among younger officers. Tobacco use and caffeine intake were found to have a correlate relationship. Caffeine may aid alertness, especially at night. Further

research is needed to assess other areas of shiftwork, more caffeine-containing products, especially energy drinks, and psychological or behavioral aspects of caffeine intake and product choice among shift workers. Limitations, gaps in the literature and implications are discussed.

CAFFEINE INTAKE AMONG LAW ENFORCEMENT OFFICERS PARTICIPATING IN
SHIFT WORK

A Thesis Submitted to
The Faculty of the Department of Nutrition and Dietetics
College of Human Ecology
East Carolina University

In Partial Fulfillment of the Requirements for the Degree
Master of Science in Nutrition

by

Maddison M. Greaves, November 2013

© Maddison M. Greaves, 2013

CAFFEINE INTAKE AMONG LAW ENFORCEMENT OFFICERS PARTICIPATING IN
SHIFT WORK

by

Maddison M. Greaves

APPROVED BY:

DIRECTOR OF THESIS: _____
Brenda M. Bertrand, PhD, RD

COMMITTEE MEMBER: _____
Christopher Duffrin, PhD

COMMITTEE MEMBER: _____
Xiangming Fang, PhD

COMMITTEE MEMBER: _____
Kimberly Myers, PhD, MHS, RD, LDN

CHAIR OF THE DEPARTMENT OF NUTRITION AND DIETETICS:

William A. Forsythe III, PhD

DEAN OF THE GRADUATE SCHOOL: _____
Paul J. Gemperline, PhD

I would like to give sincere gratitude to Dr. Bertrand and the members of my committee for their guidance and insight throughout the completion of this work.

TABLE OF CONTENTS

LIST OF TABLES.....	vii
LIST OF FIGURES.....	viii
LIST OF ABBREVIATIONS.....	ix
CHAPTER 1: REVIEW OF LITERATURE.....	1
CHAPTER 2: MANUSCRIPT: CAFFEINE INTAKE AMONG LAW ENFORCEMENT OFFICERS PARTICIPATING IN SHIFT WORK	7
Abstract.....	7
Introduction.....	9
Methods.....	11
Caffeine Food Frequency Questionnaire.....	11
Demographic and Caffeine Effects Questionnaire.....	12
Procedures.....	13
Statistical Analysis.....	13
Results.....	14
Assessment of Caffeine Consumption.....	14
Caffeine Product Use and Demographic Characteristics.....	15
Caffeine Intake and Job Related Characteristics.....	15
Associative Relationship between Caffeine Intake, Perceived Concentration Level, Caffeine Side Effects and Tobacco Use.....	16
Discussion.....	16
References.....	21
List of Tables.....	27
List of Figures.....	28

REFERENCES.....	41
APPENDIX A: IRB APPROVAL LETTER.....	48
APPENDIX B: PERMISSION LETTERS.....	49
APPENDIX C: FOOD FREQUENCY QUESTIONNAIRE.....	52
APPENDIX D: DEMOGRAPHIC AND CAFFEINE EFFECTS QUESTIONNAIRE.....	53
APPENDIX E: SERVING SIZE REFERENCE SHEET.....	54

LIST OF TABLES

TABLE 1. Mean caffeine content (mg) per serving of caffeine-containing products assessed on the food frequency questionnaire as reported by various sources.....	29
TABLE 2. Age and job-related characteristics of Police and Sheriff law enforcement officers.....	30
TABLE 3. Demographic characteristics of Police and Sheriff law enforcement officers.....	31
TABLE 4. Number of days or nights during a 3-day shift that caffeine-containing products were consumed among Police and Sheriff officers as reported by food frequency questionnaire.....	33
TABLE 5. Mean caffeine-containing product frequency during day and night shifts among Police and Sheriff officers as reported by food frequency questionnaire.....	34
TABLE 6. Mean distribution of ages for energy drink consumption during day and night shifts	35
TABLE 7. Caffeine intake between Police and Sheriff departments for 3-day period of day and night shifts as reported by food frequency questionnaire.....	36
TABLE 8. Associative relationships between caffeine intake and tobacco use, caffeine side effects, and perceived concentration level among Police and Sheriff officers.....	37

LIST OF FIGURES

Figure 1. Caffeine intake during a 3-night shift period among “rookie,” “seasoned,” and “expert” Police and Sheriff officers.....	38
Figure 2. Association of age and caffeine intake for those with caffeine side effects vs. those without side effects for day shifts among Police and Sheriff officers	39
Figure 3. Association of age and caffeine intake for those with caffeine side effects vs. those without side effects for night shifts among Police and Sheriff officers.....	40

LIST OF ABBREVIATIONS

DAY	Day shifts
NIGHT	Night shifts
PO	Police Officers
SDD	Sheriff Department Deputies
SD	Sheriff Department
FFQ	Food Frequency Questionnaire

CHAPTER 1

LITERATURE REVIEW

Participation in shift work has been found to interfere with normal behaviors, including eating, exercise, and sleep patterns, potentially leading to adverse health effects. Atkinson, Fullick, Grindey and Maclaren (2008), reviewed the behavioral and biological disturbances associated with shift work. Shift work has been linked with increased BMI, prevalence of obesity, and other health complications (Atkinson, et. al., 2008). Total energy intake during shift work does not appear to be affected; however, there is a tendency toward decreased meal frequency and increased snacking during night shift work (Atkinson, et. al., 2008). It was concluded that a multidisciplinary approach to investigating health and energy balance of shift workers is needed as the biological and behavioral factors influencing energy expenditure and intake are not independent of each other (Atkinson, et. al., 2008).

Holmback and colleagues (2002) studied the postprandial reaction related to macronutrient composition when food is consumed at different points during a 24-hour period. It was hypothesized in this study that subjects would respond metabolically different when given identical meals in the morning compared to the evening, glucose tolerance would decrease from morning to midnight, and night time eating would increase low density lipoprotein to high density lipoprotein ratio (LDL:HDL; Holmback, et. al., 2002). The nocturnal hormonal pattern was thought to influence these results, as well as contribute to the prevalence of obesity and cardiovascular disease seen in shift workers (Holmback, et. al., 2002).

Results showed that participants' basal metabolic rate (BMR) tended to be higher after day 6 of a high fat diet. Fasting blood glucose, non-esterified fatty acids (NEFA), and glycerol concentrations did not differ when given a high fat versus high carbohydrate diet, but the triglyceride concentration was greater for the high carbohydrate diet (Holmback, et. al., 2002). It was determined that a high fat diet impacted metabolic responses differently than a high carbohydrate diet, and these responses may put shift workers at an increased risk for metabolic disturbances, including obesity, cardiovascular disease, and gastrointestinal problems (Holmback, et. al., 2002).

De Assis, Kupek, Nahas and Bellisle (2003) studied the association between nutritional-and health-related problems associated with shift workers, specifically garbage collectors of Florianopolis, Brazil. Because of the increase in shift work to about 20% of the work force in industrialized countries, it is important to understand the cause and effect of health issues related to shift work (De Assis, et. al., 2003). There were no to date studies showing the influence of shift time on food intake in workers with very high energy expenditure, thus, this study investigated energy and nutrient intake of garbage collectors (a high energy job) based on meal recording during a 24-hour period (De Assis, et. al., 2003). Three different shifts were examined: morning, afternoon, and night. The work tasks were the same for the three shifts. The 66 total participants (n = 22 per shift) were asked to answer questionnaires regarding their eating habits. Age, body weight and Body Mass Index (BMI) were similar for all three groups as well as daily energy expenditure. Multiple regression analysis was used to compare factors influencing energy and nutrient intake (De Assis, et. al., 2003).

It was found that energy intake did not differ among the three shifts, even with differences in food choices and ingestion times (De Assis, et. al., 2003). Factors that were found to have a significant impact on ingestion included hour of day, time since last meal, age, and BMI (De Assis, et. al., 2003). When looking at macronutrients, only carbohydrate intake significantly differed among the three shifts. Soda consumption was high among all shifts, with night workers consuming the greatest amount of soda compared to morning and afternoon workers (De Assis, et. al., 2003). This may be due to the caffeine or high sugar content of sodas, which would help combat fatigue during the night shift. The results confirm the influence of factors other than shift time on health and nutrition, including time of day, time since last meal, age, and BMI (De Assis, et. al., 2003).

Similarly, Reeves, Newling-Ward, and Gissane (2004) investigated how shift work changes food intake and eating patterns and the impact of these changes on health. It was found that night workers did not consume more than day workers, but that they did eat smaller meals and snacks over more time. Furthermore, night workers showed a more significant difference in their food intake patterns on work

days and days off, while day workers did not (Reeves, et. al. 2004). Shift work impacts the timing of food consumption since there is typically less availability of food during the night shifts, which is thought to contribute to unhealthy eating patterns observed in shift workers (Reeves, et. al., 2004).

Fisher, Rutishauser and Read (1986) investigated how the specific type of job, work environment, and availability of food in the work place influence dietary habits among 25 oil refinery shift workers. Day, night and afternoon shifts were investigated using a 24-hour food intake record. Significant differences were found between the types and frequency of foods consumed during the different shifts (Fisher, et. al., 1986). These differences were thought to be related to change in type of meal consumed as well as the availability of food, with night shift workers being primarily influenced by the availability of food and time constraints, rather than by hunger (Fisher, et. al., 1986).

Lennernas, Hambraeus, and Akerstedt (1995) explored how work hours impact eating behaviors and dietary choices among 96 male industrial workers on day work (work 40h/week between 0654 and 1530 hours), two-shift work (work 38h/week between 0530 and 1400 hours one week and between 1400 and 2230 hours the next week), and three-shift work (work 36h/week between 0530 and 1400 hours, 1400 and 2230 hours, and 2230 and 0530 hours). Due to a lack of research connecting shift work, eating behaviors, and disease as well as a gap in knowledge regarding shift workers and their specific food intake, the researchers chose to evaluate how shift work influences eating habits and nutritional parameters (Lennernas, et. al. 1995). Not only did the researchers compare energy and nutrient intake, but they also considered coffee and tea consumption for 8-hour work shifts (day, morning, afternoon, night), and for the 24-hour periods that included all four shifts. It was observed that the intake of coffee was significantly lower among three-shift workers on days off compared to work days (Lennernas, et. al. 1995). Among all three groups of workers, a difference in food and beverage intake, including coffee and alcohol, was found between work days and days off; however, no difference was reported among the 24-hour work periods with varying work shifts (Lennernas, et. al. 1995). Overall, energy intake and quality of food were not significantly influenced by shift work. Two- and three- shift work appeared to change

circadian distribution of food and coffee intake, but not the overall 24-hour consumption (Lennernas, et. al. 1995).

Another aspect that has been shown to influence tolerance of shift work is aging, as studied by Smith and Mason (2001). The number of shift workers aged over 45 years is increasing. It was hypothesized that age is linked to decreased tolerance of shiftwork as it relates to aging, lifestyle, behavior, type of work and the shift worked (Smith & Mason, 2001). The participants in this study included 306 police officers working a new rotation for approximately six months. They were divided into three age groups for comparison (1=20-32.9, 2=33-39.9, 3=40+). A range of sleep-related variables, including fatigue, drowsiness, sleep quality, sleep duration, and caffeine consumption, were analyzed using multivariate analysis of covariance (Smith & Mason, 2001). Results showed that the three age groups differed significantly in terms of sleep needs (Smith & Mason, 2001). Younger officers tended to report better shift work attitudes, better adjustment to night shifts, higher job satisfaction and commitment to the work organization, less fatigue and longer sleep duration needs. Older officers reported significantly higher “morningness” (morning-oriented) and lower sleep needs (Smith & Mason, 2001). Concerning caffeine intake, older officers tended to consume more caffeine on all shifts worked (Smith & Mason, 2001). Older officers also tended to report less drowsiness on morning compared to night shifts. The results support the hypothesis that shiftwork tolerance is related to age; however, a link between the impacts of aging, lifestyle, behavior, type of work and the rotation worked has yet to be fully understood.

Caffeine has taken an important role in helping to maintain performance among shift workers as well as other sleep depriving situations. Snel and Lorist (2011) researched how caffeine can be used to alter mental state. While caffeine has been found to be useful for restoring low levels of wakefulness, it may result in detrimental effects on sleep, producing daytime fatigue (Snel & Lorist, 2011). During night shift work, where sleep deprivation is common, caffeine contributes to maintaining performance and wakefulness. However, the effects of caffeine may be due, in part, to caffeine expectancy and placebo effects (Snel & Lorist, 2011). Disturbances in the circadian rhythm, such as those seen with shift work,

disrupt normal sleeping patterns and cognition, which may result in less than optimal work performance. According to Snel and Lorist (2011), it is necessary to combat circadian disturbances and restore cognitive performance, especially during work.

A series of double-blind studies was conducted by Killgore, Kahn-Greene, Grugle, Killgore, and Balkin (2009) to determine whether or not caffeine is a viable intervention to restore cognition. Fifty-four participants (18-36 years old) who had been sleep deprived for 45-50 hours were given either a 600 mg dose of caffeine, 400 mg modafinil (approved by FDA to manage fatigue, sleep apnea, and shift work sleep disorder), 20 mg dextroamphetamine, or a placebo prior to completing various cognitive tests designed to measure formation of abstract concepts (Killgore, et. al., 2009). Results showed that the three different stimulants (caffeine, modafinil, and dextroamphetamine) affected the outcome of the cognitive tests differently, with caffeine affecting cognitive processing most significantly (Killgore, et. al., 2009). Thus, it is suggested that performance is influenced by sleep deprivation and that this deprivation can be counteracted, in part, by caffeine (Killgore, et. al., 2009).

In another study, Kilpelainen, Huttunen, Lohi, and Lyytinen (2010) focused on perceived effects of caffeine on sleepiness, motivation, mood, and task performance. Military pilot students (23-24 years old) underwent a series of flight simulator tests after a 37-hour sleep deprivation period. It was predicted that sleep deprivation would decrease amount of correct answers and increase reaction time during testing. The subjects received either a placebo or 200 mg caffeine twice a day. Vigilance and learning were assessed during the test (Kilpelainen, et. al., 2010). As expected, sleep deprivation decreased the number of correct decisions as well as increased the reaction time for both placebo and caffeine groups. Interestingly, the perceived feeling of success remained stable across sustained wakefulness in the caffeine group (Kilpelainen, et. al., 2010). Similarly, it was observed by Ker, Edwards, Felix, Blackhall, and Roberts (2010) that caffeine has a significant impact on reducing the number of errors compared to a placebo.

Pecotic, Valic, Kardum, Seva, and Dogas (2008) further investigated the impact of caffeine on the adaptability of sleep habits by shift workers. The researchers predicted that sleep habits would be negatively affected by caffeine consumption. Medical students, physicians, specialists, and nurses participated in this study. The variables found to influence the amount of sleep needed were age, gender, work demands, and work schedule (Pecotic, et. al., 2008). Surprisingly, those participants who consumed caffeine reported greater difficulty staying awake during lectures and while driving a car. Based on these results, the researcher's hypothesis that caffeine intake impairs sleep habits and quality of sleep was supported.

Even in well-rested individuals, such as those examined in a study conducted by Michael, Johns, Owen, and Patterson (2008), caffeine can reduce drowsiness that persists for 3-4 hours. Different results between caffeine and placebo were only observed 30 minutes post administration of treatment. While this supports the prediction that caffeine can beneficially impact alertness and work performance, the benefits may be short-lived (Michael, et. al., 2008). Furthermore, the findings from Snel and Lorist (2011) show that caffeine can alter mental state in terms of cognitive processing and alertness as caffeine has been shown to restore wakefulness and performance. This is an important conclusion and may help explain the use of caffeine during shift work as a means to counteract fatigue and cognitive dysfunction.

CHAPTER 2
MANUSCRIPT

CAFFEINE INTAKE AMONG LAW ENFORCEMENT OFFICERS PARTICIPATING IN SHIFT
WORK

Abstract

Objective: To investigate caffeine consumption among law enforcement officers during day shifts (DAY) and night shifts (NIGHT); to assess the types of caffeine-containing products and the frequency of product intake during DAY and NIGHT; to identify relationships between caffeine products and selected demographic characteristics; to identify relationships between caffeine intake and job-related characteristics.; and to identify associative relationships between caffeine intake and perceived concentration level, caffeine side effects, and tobacco use.

Participants: Police officers (PO) and Sheriff Department Deputies (SDD) in a rural region in Eastern North Carolina. Useable sample of 75 (N = 75).

Methods: Anonymous, self-administered caffeine food-frequency questionnaires (FFQ; three for DAY, three for NIGHT) and demographic questionnaire. Nonparametric tests were used. The Wilcoxon signed rank test was used to compare related samples (caffeine intake DAY to NIGHT, types of products used DAY to NIGHT, and frequency of product use DAY to NIGHT). Spearman's correlation was used to determine relationships between two samples. The Mann-Whitney U test was used to compare two independent samples. The Kruskal-Wallis test was used to compare more than two independent samples.

Results: Caffeine consumption was similar from DAY to NIGHT (225 ± 227 mg DAY; 252 ± 232 mg NIGHT; $p=0.891$). Frequency of tea intake was greater during DAY ($p=0.032$). Greater caffeine consumption was found among (1) SDD versus PO; (2) expert officers (for night only); (3) lower concentration levels (for night only); (4) tobacco users; (5) those who experience caffeine side effects. In addition, younger officers consumed more energy drinks DAY and NIGHT.

Conclusions: It was found that certain job-related characteristics influence caffeine intake, but types of products used and frequency of intake tends to be the same regardless of shift or caffeine side effects. Energy drinks were found to be most popular among younger officers. Tobacco use and caffeine intake were found to have a correlate relationship. Caffeine may aid alertness, especially at night. Further research is needed to assess other areas of shiftwork, more caffeine-containing products, especially energy drinks, and psychological or behavioral aspects of caffeine intake and product choice among shift workers. Limitations, gaps in the literature and implications are discussed.

CAFFEINE INTAKE AMONG LAW ENFORCEMENT OFFICERS PARTICIPATING IN SHIFT WORK

Introduction

Participation in shift work has been found to interfere with normal behaviors, including eating, exercise, and regular sleep patterns. Disruption in these normal behaviors can lead to adverse health effects. Shift work is associated with increased BMI and prevalence of obesity, cardiovascular disease, peptic ulcers, and gastrointestinal dysfunction (Atkinson, 2008; Holmback, 2002). De Assis and colleagues (2003) observed that night shift workers are more likely to consume energy-dense foods and beverages during shift work compared to morning and afternoon workers (2003). Fisher and colleagues (1986) reported that the types and frequency of foods consumed during the night shift are influenced by the availability of food and time constraints, rather than by hunger.

Disturbances in sleep patterns and normal circadian rhythms may contribute to food and beverage choices among shift workers (Atkinson, et al., 2008). According to Snel and Lorist (2011), caffeine is commonly used to offset fatigue and sleepiness and to assist with maintaining work performance. Lennernas and colleagues (1995) studied the impact of work hours on eating behaviors and dietary choices among male industrial workers on day work (work 40h/week between 0654 and 1530 hours), two-shift work (work 38h/week between 0530 and 1400 hours one week and between 1400 and 2230 hours the next week), and three-shift work (work 36h/week between 0530 and 1400 hours, 1400 and 2230 hours, and 2230 and 0530 hours). The findings indicated that intake of coffee was significantly higher among three-shift workers on work days compared to days off. Among all groups, differences in food and beverage intake, including coffee and alcohol, were found between work days and days off (Lennernas, Hambræus, & Akerstedt, 1995).

According to Schardt (2012), over 80 percent of American adults regularly consume caffeine. Burke (2008) states that 90 percent of adults worldwide consume caffeine daily. Some of the most common caffeine-containing beverages include coffee, tea, and soda, as assessed by Barone and Roberts

(1996). Both American adults and adolescents consume caffeine, but caffeine intake has been shown to be nearly three times greater among the adult population (Barone & Roberts, 1996). Energy drinks have been declared the second most popular supplement behind multi-vitamins, specifically among adolescents and young adults (Hoffman, 2010). In addition, caffeine-containing products are commonly used among athletes due to the perceived effects on performance and endurance (Hoffman, 2010; Desbrow & Leveritt 2006; Burke, 2008). Athletes that competed in the 2005 Ironman triathlon chose sodas, caffeinated gels, coffee, energy drinks, and NoDoz tablets most often (Desbrow & Leveritt, 2006).

Caffeine affects people differently, with reported side effects ranging from jittery, nervous, and restless to energetic or insomniac (Monroe, 1998). Use of caffeine can result in dependence, defined as craving caffeine regularly, and tolerance, defined as the body requiring more and more caffeine to obtain the same effect (Monroe, 1998). Along with dependence comes withdrawal, which often results in increased negative mood (Monroe, 1998; Smith, 2002). Caffeine side effects are more common among heavy daily caffeine users and include insomnia, restlessness, depression, rapid heartbeat, anxiety, stomach pain or heartburn, headaches, muscle gain, shaking, and ringing in the ears (Monroe, 1998). Some of the more positive side effects are enhanced wakefulness and alertness, which have attracted shift workers, truck drivers, military officers, athletes, and others who aim to combat fatigue (Burke, 2008).

While the majority of adults both worldwide and within the United States regularly use caffeine, it has been found to be most popular among certain populations, including athletes (Hoffman, 2010; Desbrow & Leveritt 2006; Burke, 2008) and night workers (Smith, 2002). Studies have shown caffeine intake to be common among various workers participating in night shifts, including industrial workers, electronic workers, steel and railway workers, and medical staff, including nurses (Glazner, 1991). Lieberman and colleagues (2012) assessed intake among US army soldiers, where coffee was the most popular product used overall, and energy drinks were most commonly used among young adult males. Literature is lacking on caffeine intake among law enforcement officers participating in shift work as well as among shift workers serving as their own control to compare day versus night shifts.

The purpose of this study was to evaluate caffeine intake from caffeinated beverages and energy boosting supplements among law enforcement officers while working day shifts (DAY) and night shifts (NIGHT). The specific research aims for this study were: 1) Assess the amount of caffeine consumed, the types of caffeine products used, and the frequency of caffeine products used between DAY and NIGHT; 2) Identify relationships between caffeine products used and demographic characteristics; 3) Identify relationships between caffeine intake and job-related characteristics; and 4) Identify associative relationships between caffeine intake and perceived concentration level, caffeine side effects, and tobacco use.

Methods

This was a quantitative prospective study design. Participants were Police Officers (PO) and Sheriff Department Deputies (SDD) of a rural city in the eastern region of North Carolina who worked rotating DAY and NIGHT. The county served was 652 square miles large with a population of 172,554 (United States Census Bureau, 2012). The city served accounted for 5% of the county (34.6 sq. mi.) and over half of the county population (population 87,242), making the target city the largest, most populated area of the county (United States Census Bureau, 2012). In 2012, the county index crime rate was 3,986.5 per 100,000 persons, which was slightly higher than the state index crime rate of 3,767.2 per 100,000 persons (State Bureau of Investigation, 2013).

Caffeine Food Frequency Questionnaire (FFQ)

A quantitative self-administered caffeine food frequency questionnaire (FFQ) was used to assess 24-hour caffeine intake from beverages and supplements for each of three DAY and NIGHT shifts worked. An initial FFQ with five drink categories was designed from the literature (Neuhours et al, 2009; Heckman et al, 2010; McCusker et al, 2003; Chin et al, 2008; USDA, 2011; McCusker et al, 2006; Chou et al, 2007; McCusker 2006; Consumer Reports 2011; (Starbucks Drinks., 2012; McDonald's Beverages, 2012). Additional drink categories and an energy-boosting supplement category were added following field research by a Registered Dietitian of area convenient stores, pharmacies, coffee shops, and grocery

stores. The final FFQ had seven drink and four energy boosting supplement categories. Drink categories included coffee, coffee drinks, espresso, caffeinated sodas, energy drinks, tea, and caffeinated water. The energy boosting supplements included energy shots, caffeine supplements, energy chewing gum, and energy strips. Participants indicated each drink and supplement that they consumed for the given shift, in addition to serving size and number of servings consumed for each item. A reference serving size was provided for each item. For example, eight ounces (8 oz.) for coffee (standard coffee mug size) and 12 oz. for soda (regular can of soda). Job-related information for that shift was also reported, including the shift worked (day or night), the number of hours worked, and how busy was the shift (not busy, slightly busy but less than normal, normal workload, slightly more busy than normal, or very busy).

The FFQ was content validated by two Registered Dietitians, the SD Captain, and a Health Educator. Revisions were made to drink categories and serving sizes from feedback provided. A pilot test was next conducted among five university fitness center employees, aged 22 to 40 years (two males and three females). Reference serving sizes for each drink were added and modifications to heading titles and instructions for completing the questionnaire were made from feedback provided. Table 1 presents the mean caffeine content (mg) of products assessed on the FFQ. A serving size reference sheet was developed from field research of area restaurants. The sheet listed standard serving sizes for each drink and supplement.

Demographic and Caffeine Effects Questionnaire

The Demographic and Caffeine Effects Questionnaire assessed demographic information and characteristics of shift work for the past two weeks. As part of this questionnaire, a 7-point Likert scale was used to assess perceived level of energy and concentration while participating in shift work (1 = extremely low, 7 = extremely high; Laerhoven, et al, 2004). The questionnaire also assessed side effects experienced from consuming caffeine. This questionnaire was reviewed for content validity by the SD Captain, a Health Educator, and the City Human Resources Safety and Risk Manager. Three additional job-related and two demographic questions were added from their feedback. Further, the phrasing of the

Likert scale question was modified. The study was approved by the university's Institutional Review Board, the SD Captain, the Chief of Police, and the City Human Resources Safety and Risk Manager. All participants provided written informed consent prior to study participation. Data collection took place from April to June 2012.

Procedures

Each participant received a 9-page packet. Page 1 included a description of each item in the packet, deadline for completion, and researcher contact information. Page 2 was the Demographic and Caffeine Effects Questionnaire. Pages 3-8 were color-coded FFQs (pages 3-5 were yellow for DAY; pages 6-8 were purple for NIGHT). Page 9 was the Serving Size Reference Sheet, which included a graphic outline of a 16-ounce cup for gauging drink size. To maintain participant anonymity, each page was ID-coded with the same randomly assigned three-digit number.

Initially, the researcher attended roll call for each patrol unit of the SD. The researcher used a script to describe the packet content and time frame for completing the study, instruct recruits that participation was voluntary and recorded anonymous, and to obtain informed consent. Participants returned the packets to an assigned deputy; these were then given to the Captain and returned to the researcher. The same protocol was used for recruiting PO, with the exception that an assigned PO distributed and collected the packets, which were then returned by the PO to the researcher. By telephone and email, the researcher was in contact with the SDD and PO bi-monthly throughout data collection. Neither questions nor concerns were reported throughout the data collection period.

Statistical Analysis

Because data were not normally distributed, nonparametric tests were used for analyses. The Wilcoxon Signed Rank Test for related samples was used to compare mean caffeine intake and frequency of product intake for DAY and NIGHT in order to assess mean population difference (Wilcoxon, 1945). Spearman's correlation was used to assess the significance of the relationship between two quantitative variables (Page, 1963). The Kruskal-Wallis test was used when comparing more than two independent

samples (Kruskal, Wallis, 1952), and The Mann-Whitney U Test for independent samples was used (Lund Research Limited, 2013) (Page, 1963) for comparisons between two non-related samples.

Results

There were 187 PO who serviced the city area. The PO operated 12-hour shifts and rotated from DAY to NIGHT (and vice versa) every three weeks. There were 199 SDD who serviced the county beyond police jurisdiction (City of Greenville, 2007; Pitt County Development Commission, 2007-2012). SDD also had 12-hour shifts, but rotated shifts every four weeks. SDD followed a Dupont schedule; officers worked two days during the week, then had two days off, and worked every other weekend (Friday, Saturday, and Sunday) for a total of seven 12-hour shifts during a 2-week period. There were 111 packets distributed between the PO and SDD. The response rate was 76% and 9 were excluded, resulting in a 68% useable response rate. Those who returned only one FFQ for DAY or NIGHT were excluded.

Participant demographic characteristics are reported in Tables 2 and 3. Mean participant age was 36 years, and ranged from 23 to 58 years (24% aged 20-29 years, 39% were 30-39 years, 30% were 40-49 years, and 7% were 50-58 years). The majority was male, married, non-Hispanic white, and did not use tobacco. On average, participants were in their current position for the past six years and in shift work for the past 10 years. The majority changed shift each three weeks (PO) and worked overtime 2 to 6 of the past 14 days.

Assessment of Caffeine Consumption

Overall, daily caffeine intake and types of caffeine products used were similar between DAY and NIGHT (255±227 milligrams per day (mg/d) ($M\pm SD$), Mdn 180 mg/d, 95% CI [202, 308] mg/d DAY; 252±232, 164, [198, 306] NIGHT; $p=0.891$), as assessed using the Wilcoxon Signed Rank Test. Frequency of product consumption was similar between DAY and NIGHT for all products except tea, which had greater frequency during DAY ($p=0.032$). The number of days and mean frequency of product intake for 3-day shift periods are reported in Tables 4 and 5. Of the four supplements assessed, energy

shots was the only one that was used by participants. The other products assessed included NoDoz, energy gum and energy strips.

Caffeine Product Use and Demographic Characteristics

Demographic characteristics tested in this study included tobacco use, age, sex, marital status, and race. Energy drinks was the only product that differed with age. Only eight of the 75 participants used energy drinks; their ages ranged from 24 to 40 years. Those with the highest intake of energy drinks (average one per shift) were younger for both DAY and NIGHT. Table 6 presents the mean distribution of ages for energy drink consumption during DAY and NIGHT. Mean caffeine intake and product consumption were similar across categories of sex, marital status, and race (data not shown).

Further looking at demographic characteristics between departments, SDD were, on average, five years older than PO (34 ± 8 years ($M \pm SD$), Mdn 33 years, 95% CI [32, 37] years for PO; 39 ± 7 years, 95% CI [36, 41] years for SDD; $p=0.016$) using the Kruskal-Wallis Test. Gender, marital status and race, were similar between departments (data not shown).

Caffeine Intake and Job Related Characteristics

Job-related characteristics assessed included shifts worked (8-10 hour night or day versus 12 hour night or day), length of time participating in current position and in shift work overall, how often the shift changed from day to night, perceived energy and concentration levels during shift work, caffeine side effects, overtime, and department. As illustrated in Table 7, mean caffeine intake was nearly twice as much for SDD versus PO for both DAY ($p=0.001$) and NIGHT ($p=0.001$). However, within each department, caffeine intake was similar between DAY and NIGHT ($p=0.206$ SDD; $p=0.258$ PO). Those who worked more years of shift work (>10 years = “expert”; $N=29$; Marsal, 2013) consumed more caffeine during NIGHT ($p=0.029$) regardless of department, as illustrated in Figure 1. Intake was similar based on years of shift work for DAY ($p=0.077$). In addition, caffeine intake was similar despite years worked at current job, overtime, how busy the current shift was, or the reported hours worked for that shift (data not shown).

Associative Relationship between Caffeine Intake, Perceived Concentration Level, Caffeine Side Effects and Tobacco Use

Associative relationships between caffeine intake and tobacco use, side effects from caffeine ingestion, and perceived level of concentration during shift work are reported in Table 8. Tobacco users consumed about 190 mg/d more caffeine for DAY and NIGHT as compared to those who did not use tobacco. Further, those who had caffeine side effects consumed about 130 mg/d more caffeine during NIGHT and 80 mg/d more during DAY as compared to those without side effects. As displayed in Figures 2 and 3, for both DAY and NIGHT the majority of participants who reported no caffeine side effects consumed less caffeine (between 0-200 mg) and a weak negative trend between caffeine intake and age was found; thus, as caffeine intake increased, age decreased. In contrast, among participants who did report caffeine side effects, caffeine intake was widely distributed and a weak positive correlation between caffeine intake and age was seen; thus, as caffeine intake increased, age also increased. Concentration level approached statistical significance for NIGHT only ($p=0.056$), with a greater mean caffeine intake observed for lower concentration. Intake was similar for DAY and NIGHT across reported energy levels (data not shown).

Discussion

Among a survey of law enforcement officers, the results of this study indicate that certain job-related factors influence caffeine consumption, including department (SDD versus PO), perceived concentration level and the number of years participating in shift work. Nearly double the intake of caffeine was found among SDD versus PO, regardless of shift. Whereas caffeine consumption and frequency of caffeine product use were similar between DAY and NIGHT, those who consumed more caffeine reported caffeine side effects, including headaches, stomach aches, heart palpitations, heart burn, and “jolt and crash” episodes.

These results suggest that habits play a large role in caffeine consumption regardless of shift. Regular caffeine use has been recognized as habit-forming and strongly addictive, as well as being

associated with withdrawal symptoms (Smith, 1987; Olekalns & Bardsley, 1996). Tobacco use is also habit-forming (Stolerman & Jarvis, 1995), which may explain why many who use tobacco also consume more caffeine. Since the majority of participants consumed the same amount of caffeine and caffeine-containing products for DAY and NIGHT, availability does not seem to be a factor. All of the products assessed are readily available at convenience stores, grocery stores and fast food restaurants; many of these establishments are open 24-hours per day or have extended hours of operation.

Higher caffeine intake for NIGHT was associated with reported lower concentration level. Working night shifts disrupts the normal circadian rhythm (Van Reeth, 1998). Even after years of shift work, it has been reported that circadian rhythm does not adjust to working nighttime hours (Van Reeth, 1998). The wide use of caffeine to combat sleepiness and circadian disturbances (Roehrs & Roth, 2008) may explain why a greater caffeine intake was reported at night among those with lower concentration levels and for those who participated in shift work for more years. There may be other contributing factors that impact caffeine intake and the choice of products that contain caffeine that were not assessed in the current study, such as co-worker influence, stress of the job, down time, location of patrol work (i.e., in city limits versus rural areas), and availability of products within the workplace.

Previous studies have found that food choices and frequency of intake are influenced more by habit and time availability rather than by hunger (Waterhouse, Buckley, Edwards, & Reilly, 2003). Waterhouse and colleagues (2003) found a higher intake of snacks and lower intake of large hot meals among night workers compared to day workers. Multiple researchers have reported that total energy intake for shift workers does not differ despite the shift worked (Debry, et al, 1967; Reinberg, et al, 1979; Pasqua & Moreno, 2004; Lennernas, et al, 1994), which is consistent with our findings that caffeine consumption was similar between DAY and NIGHT. Lowden and colleagues (2010), however, found that eating behaviors are influenced by night shift work. Psychological disruptions in eating habits, including meal times of shift workers compared to culturally normal meal times, have been associated with

increased snacking among night shift workers (Lowden, et al, 2010). Thus, further research should investigate timing of caffeine and caffeine-product intake among shift workers.

In contrast to the present study, Reeves and colleagues found that female night-shift workers consumed more cups of coffee and tea compared to female and male day-shift workers. These investigators also found that night workers smoked more cigarettes than day workers (Reeves, et al, 2004). Both caffeine and nicotine likely aid alertness and are habit-forming (Smith, 1987; Olekalns & Bardsley, 1996; Stolerman & Jarvis, 1995; Roehrs & Roth, 2008), which may explain why night shift workers consumed more of both. Similar to our findings, a strong relationship between tobacco use, namely cigarette smoking, and caffeine consumption has been reported (Friedman, et al, 1974; Dawber et al, 1974; Prineas, et al, 1980; Kannas, 1981; Thomas, 1973; Conway, et al, 1981). Roehrs and Roth (2008) reported that caffeine is a widely used substance among those with sleep deprivation, restriction, or circadian rhythm reversal to combat sleepiness because it enhances alertness and performance. The present study found that a lower reported concentration level was associated with greater caffeine intake at night, which may be explained by caffeine's ability to offset fatigue and restore performance. Weis and Laties (1962) concluded that caffeine restored numerous performance tasks while Smith (2002) further reported that caffeine improved attention and psychomotor functions as opposed to complex intellectual tasks.

Although age did not influence overall caffeine intake, it does seem to be a determinant toward chosen caffeine products. Our findings suggest that energy drinks are more popular among younger officers in their early-to-mid-twenties. Energy drinks have become a popular product in recent years, especially among teens and young adults (Torpy & Livingston, 2013; Rath, 2012; Hoffman, 2010). Energy drinks are considered the second most popular supplement, after multi-vitamins, among American adolescents and young adults. Caffeine is the primary active ingredient in energy drinks (Hoffman, 2010). In addition, many energy drinks combine caffeine with other stimulatory ingredients (Hoffman, 2010). Young adult males are the targeted market for energy drinks due to the reported ability of energy drinks to

enhance performance, increase attention, and prolong endurance (Reissig, et al, 2008). These performance effects might be considered beneficial among young male shift workers.

A limitation of the current study was that intake of caffeine-containing products was self-reported. The researchers did not control the products consumed, and were not able to determine whether or not reported sizes were consistent with the sizes provided on the FFQ. It has been commonly found that self-reported food intake underestimates actual intake (Gibson, 2005). Similarly, participants may have underestimated the size of a product consumed. Although the FFQ included many caffeine-containing beverages and supplements, it is possible that more products were available. In addition, most of the participants were male, limiting the researchers to distinguish between sex differences with regard to caffeine intake.

Future research should compare the present study's findings with other areas of shift work as well as other regions of the country and urban areas of residency. Future studies may better determine whether the length of time during a shift as well as how often one rotates from day to night shifts influences caffeine intake, since both participating departments in the present study worked 12-hour shifts and rotated from day to night in relatively the same amount of time (every three versus four weeks). Caffeine-containing products offered on the market today will change as new products become available and grow in popularity; therefore, it is important for future research to identify the variety of caffeine-containing products on the market. Energy drinks are one such product that continues to gain popularity for its perceived energy-boosting and performance-enhancing properties (Rath, 2012; Hoffman, 2010; Reissig, et al, 2008). Future research with a greater focus on energy drink consumption among shift workers is needed. Psychological aspects seem to play a significant role in determining food choices (Lowden, et al, 2010); however, there is a lack of research investigating the relationship between caffeine consumption and psychological factors among shift workers. Additional research aiming to explore this relationship is indicated to better understand caffeine intake among shift workers from a behavioral standpoint. Finally, there may be other factors that influence caffeine consumption among shift workers, such as number of

children in the participants' household and children's ages, health status of the participant, and how the participant perceives caffeine with relation to health or nutritional value.

REFERENCES FOR MANUSCRIPT

1. Anonymous (2011). Can 5-Hour Energy Kick your Afternoon Slump? *Consumer Reports*, 76(3), 16.
Retrieved from <http://www.consumerreports.org/cro/magazine-archive/2011/march/health/5-hour-energy/overview/index.htm>. Accessed September 2012.
2. Atkinson, G, Fullick, S, Grindey, C, & Maclaren, D. (2008). Exercise, Energy Balance and the Shift Worker. *Sports Medicine*, 38(8), 671-685.
3. Avitae Caffeinated Water. (2012) Retrieved from www.avitaeusa.com. Accessed September 2012.
4. Barone, JJ, & Roberts, HR. (1996). Caffeine Consumption. *Food and Chemical Toxicology*, 34(1), 119-129.
5. Burke, LM. (2008). Caffeine and Sports Performance. *Applied Physiology, Nutrition, and Metabolism*, 33(6), 1319-1334.
6. Center for Science in the Public Interest (2011). Caffeine Content of Food & Drugs. Retrieved from <http://www.cspinet.org/new/cafchart.htm>. Accessed September 2012.
7. Chin, JM, Merves, ML, Goldberger, BA, Sampson-Cone, A, & Cone, EJ. (2008). Caffeine Content of Brewed Teas. *Journal of Analytical Toxicology*, 32(8), 702-704.
8. Chou KH, & Bell, LN. (2007). Caffeine Content of Prepackaged National-Brand and Private-Label Carbonated Beverages. *Journal of Food Science*, 72(6), C337-C342.
9. Conway, TL, Vickers, RR, Ward, HW, & Rahe, RH. (1981). Occupational Stress and Variation in Cigarette, Coffee, and Alcohol Consumption. *Journal of Health and Social Behavior*, 22(2), 155-165.
10. Dawber, TR, Kannel, WB, & Gordon, T. (1974). Coffee and Cardiovascular Disease: Observations from the Framingham study. *New England Journal of Medicine*, 291(17), 871-874.
11. De Assis, MAA, Kupek, E, Nahas, MV, & Bellisle, F. (2003). Food Intake and Circadian Rhythms in Shift Workers with a High Workload. *Appetite*, 40(2), 175-183.

12. Debry G, Girault P, Lefort J, & Thiébaud J. (1967). Survey on the Eating Habits of Workers. *Bulletin Institute National de la Sante et de la Recherche Medicale.*, 22(6): 11 69-202.
13. Desbrow, B, & Leveritt, M. (2006). Awareness and Use of Caffeine by Athletes Competing at the 2005 Ironman Triathlon World Championships. *International Journal of Sport Nutrition and Exercise Metabolism*, 16(5), 545-558.
14. Fisher, M, Rutishauser, IHE, & Read, RSD. (1986). The Dietary Patterns of Shift Workers on Short Rotation Shifts. *Community Health Studies*, 10(1), 54-56.
15. Friedman, GD, Siegelau, AB, & Seltzer, CC. (1974). Drinking, Smoking, and Drug-Taking Patterns in a Predominantly Lower Socioeconomic Status Sample. *Medical Journal of Australia*, 2, 459-461.
16. Gibson, RS. (2005). Measurement Errors in Dietary Assessment. (2nd Ed.), *Principles of Nutritional Assessment* (pp. 105-128). New York, New York: Oxford University Press.
17. Glazner, LK. (1991). Shift work: Its Effect on Workers. *American Association of Occupational Health Nurses Journal*, 39(9), 416-421.
18. Heckman MA, Weil, J, & De Mejia, EG. (2010). Caffeine in Foods: A Comprehensive Review on Consumption, Functionality, Safety, and Regulatory Matters. *Journal of Food Science*, 75(3), R77-R87.
19. Hoffman, JR. (2010). Caffeine and Energy Drinks. *Strength and Conditioning Journal*, 32(1), 15-20.
20. Holmback, U, Forslund, A, Forslund, J, Hambraeus, L, Lennernas, M, Lowden, A, Stridsberg, M, & Akerstedt, T. (2002). Metabolic Responses to Nocturnal Eating in Men are affected by Sources of Dietary Energy. *The Journal of Nutrition*, 132(7), 1892-1899.
21. J. Marsal, personal communication, April 16, 2013.
22. Kannas, L. (1981). The Dimensions of Health Behavior among Young Men in Finland. *International Journal of Health Education*, 24, 146-155.

23. Kruskal, WH, & Wallis, WA. (1952). Use of Ranks in One-Criterion Variance Analysis. *Journal of the American Statistical Association*, 47(260), 583-621.
24. Lennernäs M, Abrahamsson L, Hambraeus L, & Akerstedt T. (1994). Nutrition and 3-Shift Work: The 24-Hour Intake of Energy and Nutrients. *Ecology of Food and Nutrition*, 32(3), 157-165.
25. Lennernäs, M, Hambraeus, L, & Akerstedt, T. (1994). Nutrient Intake in Day Workers and Shift Workers. *Work and Stress*, 8(4), 332-342.
26. Lieberman, HR, Stavinoha T, McGraw S, White, A, Hadden, L, & Marriott, BP. (2012). Caffeine Use among Active Duty US Army Soldiers. *Journal of the Academy of Nutrition and Dietetics*, 112(6), 902-912.
27. Lowden, A, Moreno, C, Homback, U, Lennernas, M, & Tucker, P. (2010). Eating and Shift Work – Effects on Habits, Metabolism, and Performance. *Scandinavian Journal of Work, Environment & Health*, 36(2), 150-162.
28. Lund Research Limited. (2013). Mann-Whitney U Test using SPSS. *Laerd Statistics*. Retrieved from <https://statistics.laerd.com/spss-tutorials/mann-whitney-u-test-using-spss-statistics.php>. Accessed August 9, 2013.
29. Malinauskas, BM, Aeby, V, Overton, R, Carpenter-Aeby, T, & Barber-Heidal, K. (2007). A Survey of Energy Drink Consumption Patterns among College Students. *Nutrition Journal*, 6(1), 35-41.
30. McCusker RR, Fuehrlein, B, Goldberger, BA, Gold, MS, & Cone, EJ. (2006). Caffeine Content of Decaffeinated Coffees. *Journal of Analytical Toxicology*, 30(8), 611-613.
31. McCusker RR, Goldberger, BA, & Cone, EJ. (2003). Caffeine Content of Specialty Coffees. *Journal of Analytical Toxicology*, 27(7), 520-522.
32. McCusker RR, Goldberger, BA, & Cone, EJ. (2006). Caffeine Content of Energy Drinks, Carbonated Sodas, and Other Beverages. *Journal of Analytical Toxicology*, 30(2), 112-114.
33. McDonald's Beverages (2012). Retrieved from <http://www.mcdonalds.com> . Accessed September 2012.

34. Monroe, J. (1998). Caffeine's Hook. *Current Health*, 24(5), 16.
35. Neuhouser, ML, Lilley, S, Lund, A, & Johnson, DB. (2009). Development and Validation of a Beverage and Snack Questionnaire for Use in Evaluation of School Nutrition Policies. *Journal of the American Dietetic Association*, 109(9), 1587-92.
36. Olekalns, N, & Bradsley, P. (1996). Rationale Addiction to Caffeine: An Analysis of Coffee Consumption. *Journal of Political Economy*, 104(5), 1100-1104.
37. Page, EB. (1963). Ordered Hypotheses for Multiple Treatments: A Significance Test for Linear Ranks. *Journal of the American Statistical Association*, 58(301), 216-230.
38. Pasqua IC, & Moreno CR. (2004). The Nutritional Status and Eating Habits of Shift Workers: A Chronobiological Approach. *Chronobiology International*, 21(6), 949-60.
39. Prineas, RJ, Jacobs, DR, Crow, RS, & Blackburn, H. (1980). Coffee, Tea and VPB. *Journal of Chronic Diseases*, 33(2), 67-72.
40. Rath, M. (2012). Energy Drinks: What is All the Hype? The Dangers of Energy Drink Consumption. *Journal of the American Academy of Nurse Practitioners*, 24, 70-76.
41. Reeves, SL, Newling-Ward, E, & Gissane, C. (2004). The Effect of Shift-Work on Food Intake and Eating Habits. *Nutrition and Food Science*, 35(5), 216-221.
42. Reinberg A, Migraine C, Apfelbaum M, Brigant L, Ghata J, Vieux N, Laporte, A, & Nicolai. (1979). Circadian and Ultradian Rhythms in the Feeding Behavior and Nutrient Intakes of Oil Refinery Operators with Shift-Work Every 3-4 Days. *Diabetes & Metabolism*, 5(1), 33-41.
43. Reissig, CJ, Strain, EC, & Griffiths, RR. (2008). Caffeinated Energy Drinks – A Growing Problem. *Drug and Alcohol Dependence*, 99(1-3), 1-10.
44. Roehrs, T, & Roth, T. (2008). Caffeine: Sleep and Daytime Sleepiness. *Sleep Medicine Reviews*, 12(2), 153-162.
45. Schardt, D. (2012). Caffeine! *Nutrition Action Health Letter*, 39(10), 7-8.

46. Smith, A. (2002). Effects of Caffeine on Human Behavior. *Food and Chemical Toxicology*, 40(9), 1243-1255.
47. Smith, R. (1987). Caffeine Withdrawal Headache. *Journal of Clinical Pharmacy and Therapeutics*, 12(1), 53-57.
48. Snel, J, & Lorist, M. (2011). Effects of Caffeine on Sleep and Cognition. *Progress in Brain Research*, 190, 105-117.
49. Starbucks Drinks. (2012). Retrieved from <http://www.starbucks.com/menu/drinks>. Accessed September, 2012.
50. State Bureau of Investigation. (2013). Crime in North Carolina – 2012. *Annual Summary Report of 2012 Uniform Crime Reporting Data*. Retrieved from <http://crimereporting.ncdoj.gov/Reports.aspx>. Accessed November 2013.
51. Stolerman, IP, & Jarvis, MJ. (1995). The Scientific Case that Nicotine is Addictive. *Psychopharmacology*, 117(1), 2-10.
52. Thomas, CB. (1973). The Relationship of Smoking and Habits of Nervous Tension. In W. Dunn, *Smoking Behavior: Motives and Incentives* (pp. 157-170). New York, Winston. The Halste Press Division.
53. Torpy, JM, & Livingston, EH. (2013). Energy Drinks. *Journal of the American Medical Association*, 309(3), 297.
54. United States Census Bureau. (2012). State and County Quick Facts. Retrieved from <http://quickfacts.census.gov>. Accessed July 2012.
55. USDA National Nutrient Database for Standard Reference, Release 23. U.S. Department of Agriculture. Retrieved from <http://www.nal.usda.gov/fnic/foodcomp/search>. Accessed July 6, 2011.

56. Van Laerhoven, H, van der Zaag-Loonen, HJ, & Derkx, BHF. (2004). A Comparison of Likert Scale and Visual Analogue Scales as Response Options in Children's Questionnaires. *Acta Paediatrica*, 93(6), 830-835.
57. Van Reeth, O. (1998). Sleep and Circadian Disturbances in Shift Work: Strategies for Their Management. *Hormone Research*, 49(3-4), 158-162.
58. Waterhouse, J, Buckley, P, Edwards, B, & Reilly, T. (2003). Measurement of, and Some Reasons for, Differences in Eating Habits between Night and Day Workers. *Cardiology International*, 20(6), 1075-1092.
59. Weiss, B, & Laties, VG. (1962). Enhancement of Human Performance by Caffeine and the Amphetamines. *Pharmacological Review*, 14(1), 1-36.
60. Wilcoxon, F. (1945). Individual Comparisons by Ranking Methods. *Biometrics Bulletin*, 1(6), 80-83.

LIST OF TABLES

TABLE 1. Mean caffeine content (mg) per serving of caffeine-containing products assessed on the food frequency questionnaire as reported by various sources

TABLE 2. Age and job-related characteristics of Police and Sheriff law enforcement officers

TABLE 3. Demographic characteristics of Police and Sheriff law enforcement officers

TABLE 4. Number of days or nights during a 3-day shift that caffeine-containing products were consumed among Police and Sheriff officers as reported by food frequency questionnaire

TABLE 5. Mean caffeine-containing product frequency during day and night shifts among Police and Sheriff officers as reported by food frequency questionnaire

TABLE 6. Mean distribution of ages for energy drink consumption during day and night shifts

TABLE 7. Caffeine intake between Police and Sheriff departments for 3-day period of day and night shifts as reported by food frequency questionnaire

TABLE 8. Associative relationships between caffeine intake and tobacco use, caffeine side effects, and perceived concentration level among Police and Sheriff officers

LIST OF FIGURES

Figure 1. Caffeine intake during a 3-night shift period among “rookie,” “seasoned,” and “expert” Police and Sheriff officers

Figure 2. Association of age and caffeine intake for those with caffeine side effects vs. those without side effects for day shifts among Police and Sheriff officers

Figure 3. Association of age and caffeine intake for those with caffeine side effects vs. those without side effects for night shifts among Police and Sheriff officers

Table 1. Mean caffeine content (mg) per serving of caffeine-containing products assessed on the food frequency questionnaire (FFQ) as reported by various sources.

Product	Mean caffeine content	Serving size
Coffee	148 mg	8 fl. oz.
Coffee drinks	138 mg	16 fl. oz.
Espresso	58 mg	1 fl. oz.
Caffeinated sodas	37 mg	12 fl. oz.
Energy drinks	64 mg	8 fl. oz.
Tea	38 mg	8 fl. oz.
Caffeinated water	45 mg	16.9 fl. oz. of Avitae regular
	90 mg	16.9 fl. oz. of Avitae 90
Energy shots	207 mg	2 fl. oz.

Note. Caffeine content not included for NoDoz, energy gum and energy strips; these were included on the FFQ, but were not used by participants. (Anonymous, 2011; Center for Science in the Public Interest, 2011; Chin, et al., 2008; Chou, et al., 2007; McCusker, et al., 2003; McCusker, et al., 2006; Avitae Caffeinated Water, 2012)

Table 2. Age and job-related characteristics of Police and Sheriff officers (N=75)

Characteristic	<i>M</i> (\pm <i>SD</i>)	<i>Mdn</i>	95% CI
Age (years)	36 (\pm 8.0)	36	[34.3, 38.0]
Years at current job	6 (\pm 4.8)	5	[5.0, 7.2]
Years shift work	10 (\pm 6.6)	9	[8.3, 11.3]
Overtime (days)	2 (\pm 1.7)	1	[1.1, 1.9]

Note. *M* = mean; *SD* = standard deviation; *Mdn* = median; CI = confidence interval.

Table 3. Demographic characteristics of Police and Sheriff officers (N=75)

Characteristic	Total (n)	Distribution (%)
Sex		
Male	66	88
Female	8	11
Missing	1	1
Marital status		
Single	13	17
Married	56	75
Divorced	4	5
Missing	2	3
Race		
Non-Hispanic White	59	79
Asian American	1	1
Hispanic/Latino	3	4
African American	9	12
Other	1	1
Missing	2	3
Shift change		
Each three weeks (Police)	44	59
Each four weeks (Sheriff)	29	38
Missing	2	3

Division		
Patrol	73	98
Sheriff	1	1
Missing	1	1

Tobacco use		
Yes	20	27
No	54	72
Missing	1	1

Table 4. Number of days or nights during a 3-day shift that caffeine-containing products were consumed among Police and Sheriff officers as reported by food frequency questionnaire (FFQ) (n = 64)

Product	3-day shift period						P-value
	Number of days			Number of nights			
	<i>M</i> (\pm <i>SD</i>)	<i>Mdn</i>	95% CI	<i>M</i> (\pm <i>SD</i>)	<i>Mdn</i>	95% CI	
Coffee	1.2 (\pm 1.4)	0.0	[0.8, 1.5]	1.2 (\pm 1.4)	0.0	[0.8, 1.5]	0.904
Coffee drinks	0.1 (\pm 0.3)	0.0	[0.0, 0.1]	0.1 (\pm 0.2)	0.0	[0.0, 0.1]	0.655
Espresso	0.1 (\pm 0.3)	0.0	[0.0, 0.1]	0.1 (\pm 0.4)	0.0	[0.0, 0.2]	0.414
Soda	1.4 (\pm 1.3)	1.0	[1.0, 1.7]	1.3 (\pm 1.4)	1.0	[1.0, 1.7]	0.623
Energy drinks	0.2 (\pm 0.6)	0.0	[0.0, 0.3]	0.2 (\pm 0.7)	0.0	[0.0, 0.4]	0.180
Tea	0.8 (\pm 1.2)	0.0	[0.5, 1.1]	0.7 (\pm 1.0)	0.0	[0.4, 0.9]	0.110
Water	0.1 (\pm 0.5)	0.0	[0.0, 0.2]	0.1 (\pm 0.3)	0.0	[0.0, 0.1]	0.655
Energy shots	0.1 (\pm 0.3)	0.0	[0.0, 0.1]	0.0 (\pm 0.2)	0.0	[0.0, 0.1]	0.414

Note. *M* = mean; *SD* = standard deviation; *Mdn* = median; CI = confidence interval. The *sample size* accounts for elimination of participants who completed less than three FFQs for DAY or NIGHT. The Wilcoxon Signed Rank Test was used to determine the p-values.

Table 5. Mean caffeine-containing product frequency during day and night shifts among Police and Sheriff officers as reported by food frequency questionnaire (FFQ) (n = 64)

Product	Frequency per 3-day shift period						P-value
	Day			Night			
	<i>M</i> (\pm <i>SD</i>)	<i>Mdn</i>	95% CI	<i>M</i> (\pm <i>SD</i>)	<i>Mdn</i>	95% CI	
Coffee	0.6 (\pm 0.9)	0.0	[0.4, 0.8]	0.6 (\pm 0.8)	0.0	[0.4, 0.8]	0.974
Coffee drinks	0.0 (\pm 0.2)	0.0	[0.0, 0.1]	0.0 (\pm 0.1)	0.0	[0.0, 0.1]	0.705
Espresso	0.0 (\pm 0.1)	0.0	[0.0, 0.1]	0.0 (\pm 0.1)	0.0	[0.0, 0.1]	0.414
Soda	0.9 (\pm 1.5)	0.3	[0.6, 1.3]	1.0 (\pm 1.7)	0.3	[0.6, 1.4]	0.585
Energy drinks	0.1 (\pm 0.2)	0.0	[0.0, 0.1]	0.1 (\pm 0.2)	0.0	[0.0, 0.1]	0.317
Tea	0.4 (\pm 0.7)	0.0	[0.2, 0.6]	0.3 (\pm 0.7)	0.0	[0.2, 0.5]	0.032
Water	0.0 (\pm 0.3)	0.0	[0.0, 0.1]	0.0 (\pm 0.1)	0.0	[0.0, 0.0]	0.655
Energy shots	0.0 (\pm 0.1)	0.0	[0.0, 0.0]	0.0 (\pm 0.1)	0.0	[0.0, 0.0]	0.414

Note. *M* = mean; *SD* = standard deviation; *Mdn* = median; CI = confidence interval. The sample size accounts for elimination of participants who completed less than three FFQs for DAY or NIGHT. The Wilcoxon Signed Rank Test was used to determine the p-values.

Table 6. Mean distribution of ages for energy drink consumption during day and night shifts (N=75)

		Age (years)							
		Day				Night			
	N	<i>M</i> (\pm <i>SD</i>)	<i>Mdn</i>	95% CI	P-value	<i>M</i> (\pm <i>SD</i>)	<i>Mdn</i>	95% CI	P-value
Mean number of energy drinks									
0.00	67	36.7 (\pm 7.8)	36.0	[34.8, 38.6]		37.0 (\pm 7.8)	37.0	[35.1, 38.9]	
0.33	4	39.0 (\pm 1.4)	39.0	[26.3, 51.7]	0.03	32.3 (\pm 6.9)	32.0	[21.2, 43.3]	0.003
1.00	4	24.3 (\pm 1.5)	24.0	[20.5, 28.1]		25.0 (\pm 1.8)	25.0	[22.1, 27.9]	

Note. *M* = mean; *SD* = standard deviation; *Mdn* = median; CI = confidence interval. Spearman's Correlation was used to determine the p-values.

Table 7. Caffeine intake between Police and Sheriff departments for 3-day period of day and night shifts as reported by food frequency questionnaire (FFQ) (N=75)

Department	Mean daily caffeine intake for 3-day shift (mg/d)							
	Day				Night			
	<i>M</i> (\pm <i>SD</i>)	<i>Mdn</i>	95% CI	P-value	<i>M</i> (\pm <i>SD</i>)	<i>Mdn</i>	95% CI	P-value
Police	176 (\pm 62)	135	126, 225	0.001	184 (\pm 184)	137	128, 240	0.001
Sheriff	381 (\pm 259)	319	282, 479		361 (\pm 260)	341	262, 460	

Note. *M* = mean; *SDI* = standard deviation; *Mdn* = median; CI = confidence interval. The Kruskal-Wallis Test was used to determine the p-values.


Table 8. Associative relationships between caffeine intake and tobacco use, caffeine side effects, and perceived concentration level among Police and Sheriff officers (N=75)

Characteristic	Caffeine intake (mg/day)									
	N	Day				P-value	Night			
		<i>M</i> (\pm <i>SD</i>)	<i>Mdn</i>	95% CI	<i>M</i> (\pm <i>SD</i>)		<i>Mdn</i>	95% CI	P-value	
Tobacco										
Use	20	397 (\pm 198)	338	304, 489	0.000	388 (\pm 227)	353	282, 494	0.001	
Do not use	54	208 (\pm 217)	139	149, 267		206 (\pm 214)	137	147, 264		
Caffeine side effects										
Yes	17	318 (\pm 161)	319	235, 401	0.024	352 (\pm 209)	305	245, 460	0.009	
No	58	239 (\pm 240)	139	176, 302		223 (\pm 230)	141	163, 284		
Perceived concentration level										
3	3	300 (\pm 260)	169	-346, 945	0.160	426 (\pm 467)	173	-733, 1586	0.055*	
4	35	293 (\pm 224)	261	216, 370		301 (\pm 239)	283	219, 383		
5	18	226 (\pm 233)	136	110, 342		202 (\pm 196)	126	105, 300		
6	12	275 (\pm 252)	218	115, 434		235 (\pm 198)	183	109, 361		
7	4	133 (\pm 143)	114	-94, 360		124 (\pm 148)	100	-122, 360		

Note. *M* = mean; *SD* = standard deviation; *Mdn* = median; CI = confidence interval. Side effects from caffeine ingestion included headaches, jolt and crash episodes, heart palpitations, stomach ache and heart burn (Malinauskas, et al, 2007); a 7-point Likert scale was used to evaluate perceived concentration level (1 = extremely low concentration, 4 = normal level of concentration, 7 = extremely high level concentration) (Laerhoven, et al, 2004). Spearman's correlation, the Kruskal-Wallis test and Mann-Whitney U Test were used to determine the p-values.

*For perceived concentration level and caffeine intake during night, using Spearman's Correlation, p=0.055, using Pearson Correlation, p=0.047, and using the Kruskal-Wallis Test, p=0.283. Spearman's Correlation was kept due to data was not normally distributed.

Figure 1. Caffeine intake during a 3-night shift period among “rookie,” “seasoned,” and “expert” Police and Sheriff officers (N=75)



Note. Rookie: < 1 year service (N=3); seasoned: 1-10 years (N=41); expert: > 10 years (N=29); (J. Marsal, personal communication, April 16, 2013). *p=0.029. The Kruskal-Wallis Test was used to determine the p-value.

Figure 2. Association of age and caffeine intake for those with caffeine side effects vs. those without side effects for day shifts among Police and Sheriff officers (n=75)





Figure 3. Association of age and caffeine intake for those with caffeine side effects vs. those without side effects for night shifts among Police and Sheriff officers (n=75)



REFERENCES

1. Anonymous (2011). Can 5-Hour Energy Kick your Afternoon Slump? *Consumer Reports*, 76(3), 16.
Retrieved from <http://www.consumerreports.org/cro/magazine-archive/2011/march/health/5-hour-energy/overview/index.htm>. Accessed September 2012.
2. Atkinson, G, Fullick, S, Grindey, C, & Maclaren, D. (2008). Exercise, Energy Balance and the Shift Worker. *Sports Medicine*, 38(8), 671-685.
3. Avitae Caffeinated Water. (2012) Retrieved from www.avitaeusa.com. Accessed September 2012.
4. Barone, JJ, & Roberts, HR. (1996). Caffeine Consumption. *Food and Chemical Toxicology*, 34(1), 119-129.
5. Burke, LM. (2008). Caffeine and Sports Performance. *Applied Physiology, Nutrition, and Metabolism*, 33(6), 1319-1334.
6. Center for Science in the Public Interest (2011). Caffeine Content of Food & Drugs. Retrieved from <http://www.cspinet.org/new/cafchart.htm>. Accessed September 2012.
7. Chin, JM, Merves, ML, Goldberger, BA, Sampson-Cone, A, & Cone, EJ. (2008). Caffeine Content of Brewed Teas. *Journal of Analytical Toxicology*, 32(8), 702-704.
8. Chou KH, & Bell, LN. (2007). Caffeine Content of Prepackaged National-Brand and Private-Label Carbonated Beverages. *Journal of Food Science*, 72(6), C337-C342.
9. Conway, TL, Vickers, RR, Ward, HW, & Rahe, RH. (1981). Occupational Stress and Variation in Cigarette, Coffee, and Alcohol Consumption. *Journal of Health and Social Behavior*, 22(2), 155-165.
10. Dawber, TR, Kannel, WB, & Gordon, T. (1974). Coffee and Cardiovascular Disease: Observations from the Framingham study. *New England Journal of Medicine*, 291(17), 871-874.
11. De Assis, MAA, Kupek, E, Nahas, MV, & Bellisle, F. (2003). Food Intake and Circadian Rhythms in Shift Workers with a High Workload. *Appetite*, 40(2), 175-183.

12. Debry G, Girault P, Lefort J, & Thiébaud J. (1967). Survey on the Eating Habits of Workers. *Bulletin Institute National de la Sante et de la Recherche Medicale.*, 22(6): 11 69-202.
13. Desbrow, B, & Leveritt, M. (2006). Awareness and Use of Caffeine by Athletes Competing at the 2005 Ironman Triathlon World Championships. *International Journal of Sport Nutrition and Exercise Metabolism*, 16(5), 545-558.
14. Fisher, M, Rutishauser, IHE, & Read, RSD. (1986). The Dietary Patterns of Shift Workers on Short Rotation Shifts. *Community Health Studies*, 10(1), 54-56.
15. Friedman, GD, Siegelauab, AB, & Seltzer, CC. (1974). Drinking, Smoking, and Drug-Taking Patterns in a Predominantly Lower Socioeconomic Status Sample. *Medical Journal of Australia*, 2, 459-461.
16. Gibson, RS. (2005). Measurement Errors in Dietary Assessment. (2nd Ed.), *Principles of Nutritional Assessment* (pp. 105-128). New York, New York: Oxford University Press.
17. Glazner, LK. (1991). Shift work: Its Effect on Workers. *American Association of Occupational Health Nurses Journal*, 39(9), 416-421.
18. Heckman MA, Weil, J, & De Mejia, EG. (2010). Caffeine in Foods: A Comprehensive Review on Consumption, Functionality, Safety, and Regulatory Matters. *Journal of Food Science*, 75(3), R77-R87.
19. Hoffman, JR. (2010). Caffeine and Energy Drinks. *Strength and Conditioning Journal*, 32(1), 15-20.
20. Holmback, U, Forslund, A, Forslund, J, Hambraeus, L, Lennernas, M, Lowden, A, Stridsberg, M, & Akerstedt, T. (2002). Metabolic Responses to Nocturnal Eating in Men are affected by Sources of Dietary Energy. *The Journal of Nutrition*, 132(7), 1892-1899.
21. J. Marsal, personal communication, April 16, 2013.
22. Kannas, L. (1981). The Dimensions of Health Behavior among Young Men in Finland. *International Journal of Health Education*, 24, 146-155.

23. Ker, K, Edwards, PJ, Felix, LM, Blackhall, K, & Roberts, I. (2010). Caffeine for the Prevention of Injuries and Errors in Shift Workers. *The Cochrane Database of Systematic Reviews*, 5, CD008508.
24. Killgore, WD, Kahn-Greene, ET, Grugle, NL, Killgore, DB, & Balkin, TJ. (2009) Sustaining executive functions during sleep deprivation: A comparison of caffeine, dextroamphetamine, modafinil, lab. *Sleep*, 32(2), 205-216.
25. Kilpeläinen, AA, Huttunen, KH, Lohi, JJ, & Lyytinen, H. (2010). Effect of caffeine on vigilance and cognitive performance during extended wakefulness. *The International Journal of Aviation Psychology*, 20(2), 144–159.
26. Kruskal, WH, & Wallis, WA. (1952). Use of Ranks in One-Criterion Variance Analysis. *Journal of the American Statistical Association*, 47(260), 583-621.
27. Lennernäs M, Abrahamsson L, Hambraeus L, & Akerstedt T. (1994). Nutrition and 3-Shift Work: The 24-Hour Intake of Energy and Nutrients. *Ecology of Food and Nutrition*, 32(3), 157-165.
28. Lennernäs, M, Hambraeus, L, & Akerstedt, T. (1994). Nutrient Intake in Day Workers and Shift Workers. *Work and Stress*, 8(4), 332-342.
29. Lennernas, M, Hambraeus, L, Akerstedt, T. (1995). Shift Related Dietary Intake in Day and Shift Workers. *Appetite*, 25(3), 253-265.
30. Lieberman, HR, Stavinoha T, McGraw S, White, A, Hadden, L, & Marriott, BP. (2012). Caffeine Use among Active Duty US Army Soldiers. *Journal of the Academy of Nutrition and Dietetics*, 112(6), 902-912.
31. Lowden, A, Moreno, C, Homback, U, Lennernas, M, & Tucker, P. (2010). Eating and Shift Work – Effects on Habits, Metabolism, and Performance. *Scandinavian Journal of Work, Environment & Health*, 36(2), 150-162.

32. Lund Research Limited. (2013). Mann-Whitney U Test using SPSS. *Laerd Statistics*. Retrieved from <https://statistics.laerd.com/spss-tutorials/mann-whitney-u-test-using-spss-statistics.php>. Accessed August 9, 2013.
33. Malinauskas, BM, Aeby, V, Overton, R, Carpenter-Aeby, T, & Barber-Heidal, K. (2007). A Survey of Energy Drink Consumption Patterns among College Students. *Nutrition Journal*, 6(1), 35-41.
34. McCusker RR, Fuehrlein, B, Goldberger, BA, Gold, MS, & Cone, EJ. (2006). Caffeine Content of Decaffeinated Coffees. *Journal of Analytical Toxicology*, 30(8), 611-613.
35. McCusker RR, Goldberger, BA, & Cone, EJ. (2003). Caffeine Content of Specialty Coffees. *Journal of Analytical Toxicology*, 27(7), 520-522.
36. McCusker RR, Goldberger, BA, & Cone, EJ. (2006). Caffeine Content of Energy Drinks, Carbonated Sodas, and Other Beverages. *Journal of Analytical Toxicology*, 30(2), 112-114.
37. McDonald's Beverages. Retrieved from <http://www.mcdonalds.com> . Accessed September 2012.
38. Michael, N, Johns, M, Owen, C, & Patterson, J. (2008). Effects of Caffeine on Alertness as Measured by Infrared Reflectance Oculography. *Psychopharmacology*, 200(2), 255–260.
39. Monroe, J. (1998). Caffeine's Hook. *Current Health*, 24(5), 16.
40. Olekalns, N, & Bradsley, P. (1996). Rationale Addiction to Caffeine: An Analysis of Coffee Consumption. *Journal of Political Economy*, 104(5), 1100-1104.
41. Page, EB. (1963). Ordered Hypotheses for Multiple Treatments: A Significance Test for Linear Ranks. *Journal of the American Statistical Association*, 58(301), 216-230.
42. Pasqua IC, & Moreno CR. (2004). The Nutritional Status and Eating Habits of Shift Workers: A Chronobiological Approach. *Chronobiology International*, 21(6), 949-60.
43. Pecotic, R, Valic, M, Kardum, G, Sevo, V, & Dogas, Z. (2008). Sleep Habits of Medical Students, Physicians and Nurses Regarding Age, Sex, Shift Work and Caffeine Consumption. *Lijecnicki Vjesnik*, 130(3–4), 87–91.

44. Prineas, RJ, Jacobs, DR, Crow, RS, & Blackburn, H. (1980). Coffee, Tea and VPB. *Journal of Chronic Diseases*, 33(2), 67-72.
45. Rath, M. (2012). Energy Drinks: What is All the Hype? The Dangers of Energy Drink Consumption. *Journal of the American Academy of Nurse Practitioners*, 24, 70-76.
46. Reeves, SL, Newling-Ward, E, & Gissane, C. (2004). The Effect of Shift-Work on Food Intake and Eating Habits. *Nutrition and Food Science*, 35(5), 216-221.
47. Reinberg A, Migraïne C, Apfelbaum M, Brigant L, Ghata J, Vieux N, Laporte, A, & Nicolai. (1979). Circadian and Ultradian Rhythms in the Feeding Behavior and Nutrient Intakes of Oil Refinery Operators with Shift-Work Every 3-4 Days. *Diabetes & Metabolism*, 5(1), 33-41.
48. Reissig, CJ, Strain, EC, & Griffiths, RR. (2008). Caffeinated Energy Drinks – A Growing Problem. *Drug and Alcohol Dependence*, 99(1-3), 1-10.
49. Roehrs, T, & Roth, T. (2008). Caffeine: Sleep and Daytime Sleepiness. *Sleep Medicine Reviews*, 12(2), 153-162.
50. Schardt, D. (2012). Caffeine! *Nutrition Action Health Letter*, 39(10), 7-8.
51. Shen, J, Botly, LCP, Chung, SA, Gibbs, AL, Sabanadzovic, S, & Shapiro, CM. (2006). Fatigue and Shift Work. *Journal of Sleep Research*, 15(1), 1-5.
52. Smith, A. (2002). Effects of Caffeine on Human Behavior. *Food and Chemical Toxicology*, 40(9), 1243-1255.
53. Smith, L, & Mason, C. (2001). Age and the Subjective Experience of Shiftwork. *Journal of Human Ergology*, 30(1-2), 307-313.
54. Smith, L, & Mason, C. (2001). Reducing Night Shift Exposure: A Pilot Study of Rota, Night Shift and Age Effects on Sleepiness and Fatigue. *Journal of Human Ecology*, 30(1-2), 83-87.
55. Smith, R. (1987). Caffeine Withdrawal Headache. *Journal of Clinical Pharmacy and Therapeutics*, 12(1), 53-57.

56. Snel, J, & Lorist, M. (2011). Effects of Caffeine on Sleep and Cognition. *Progress in Brain Research*, 190, 105-117.
57. Starbucks Drinks. (2012). Retrieved from <http://www.starbucks.com/menu/drinks>. Accessed September, 2012.
58. State Bureau of Investigation. (2013). Crime in North Carolina – 2012. *Annual Summary Report of 2012 Uniform Crime Reporting Data*. Retrieved from <http://crimereporting.ncdoj.gov/Reports.aspx>. Accessed November 2013.
59. Stolerman, IP, & Jarvis, MJ. (1995). The Scientific Case that Nicotine is Addictive. *Psychopharmacology*, 117(1), 2-10.
60. Takahashi, M, Nakata, A, Haratani, T, Otsuka, Y, Kaida, K, & Fukasawa, K. (2006). Psychosocial Work Characteristics Predicting Daytime Sleepiness in Day and Shift Workers. *Chronobiology International*, 23(6), 1409-1422.
61. Thomas, CB. (1973). The Relationship of Smoking and Habits of Nervous Tension. In W. Dunn, *Smoking Behavior: Motives and Incentives* (pp. 157-170). New York, Winston. The Halste Press Division.
62. Torpy, JM, & Livingston, EH. (2013). Energy Drinks. *Journal of the American Medical Association*, 309(3), 297.
63. United States Census Bureau. (2012). State and County Quick Facts. Retrieved from <http://http://quickfacts.census.gov>. Accessed July 2012.
64. USDA National Nutrient Database for Standard Reference, Release 23. U.S. Department of Agriculture. Retrieved from <http://www.nal.usda.gov/fnic/foodcomp/search>. Accessed July 6, 2011.
65. Van Laerhoven, H, van der Zaag-Loonen, HJ, & Derkx, BHF. (2004). A Comparison of Likert Scale and Visual Analogue Scales as Response Options in Children's Questionnaires. *Acta Paediatrica*, 93(6), 830-835.

66. Van Reeth, O. (1998). Sleep and Circadian Disturbances in Shift Work: Strategies for Their Management. *Hormone Research*, 49(3-4), 158-162.
67. Waterhouse, J, Buckley, P, Edwards, B, & Reilly, T. (2003). Measurement of, and Some Reasons for, Differences in Eating Habits between Night and Day Workers. *Cardiology International*, 20(6), 1075-1092.
68. Weiss, B, & Laties, VG. (1962). Enhancement of Human Performance by Caffeine and the Amphetamines. *Pharmacological Review*, 14(1), 1-36.
69. Wilcoxon, F. (1945). Individual Comparisons by Ranking Methods. *Biometrics Bulletin*, 1(6), 80-83.

APPENDIX A

IRB APPROVAL LETTER

Notification of Exempt Certification

From: Social/Behavioral IRB
To: Maddison Greaves
CC: Brenda Bertrand
Date: 2/16/2012
Re: UMCIRB 11-001291
Caffeine Intake among Shift Workers

I am pleased to inform you that your research submission has been certified as exempt on 2/15/2012. This study is eligible for Exempt Certification under category #2.

It is your responsibility to ensure that this research is conducted in the manner reported in your application and/or protocol, as well as being consistent with the ethical principles of the Belmont Report and your profession.

This research study does not require any additional interaction with the UMCIRB unless there are proposed changes to this study. Any change, prior to implementing that change, must be submitted to the UMCIRB for review and approval. The UMCIRB will determine if the change impacts the eligibility of the research for exempt status. If more substantive review is required, you will be notified within five business days.

The UMCIRB office will hold your exemption application for a period of five years from the date of this letter. If you wish to continue this protocol beyond this period, you will need to submit an Exemption Certification request at least 30 days before the end of the five year period.

The Chairperson (or designee) does not have a potential for conflict of interest on this study.

APPENDIX B
PERMISSION LETTERS

Statement of Informed Consent

Included in this packet:

- 6 Caffeine Surveys:
 - Complete 3 while on three separate day shifts (yellow paper).
 - Complete 3 while on three separate night shifts (blue paper).
- 1 Demographic Survey (pink paper)
- 1 Drink size reference sheet

Please complete all six caffeine surveys and demographic survey within **four weeks** of receiving this packet.

This survey is **anonymous**, so please do not write your name. Your participation in this research is **voluntary**. You may choose not to answer any or all questions, and you may stop at any time. There is **no penalty for not taking part in or withdrawing** from this research study at any point. If you have any questions please feel free to contact me, Maddie Greaves, at 619-980-3355, or the Office for Human Research Integrity (OHRI) at 252-744-2914 for questions about your rights as a research participant.

This is acknowledging that the Greenville Police Department has agreed to participate in the research study, Caffeine Consumption among Shift Workers, being conducted by Maddison Greaves, graduate student at East Carolina University.

J A Bartlett
Signature

2/7/12
Date

J. A. Bartlett
Print name

Interim Chief of Police
Title

This is acknowledging that the Pitt County Sheriff Department has agreed to participate in the research study, Caffeine Consumption among Shift Workers, being conducted by Maddison Greaves, graduate student at East Carolina University.

Neil Elks
Signature

2/13/12
Date

Neil Elks
Print name

Sheriff
Title

APPENDIX C

CAFFEINE FOOD FREQUENCY QUESTIONNAIRE

Caffeine Survey

Date: _____

Which shift did you just work? <input type="checkbox"/> 8-10 hour day <input type="checkbox"/> 12 hour day <input type="checkbox"/> 8-10 hour night <input type="checkbox"/> 12 hour night	How many regular work hours did you log this shift? _____ hours	Tell me how busy this shift was: <input type="checkbox"/> Not busy <input type="checkbox"/> Slightly busier than normal <input type="checkbox"/> Light workload <input type="checkbox"/> Very busy <input type="checkbox"/> Normal workload
---	---	---

Check all you had this shift.	What's your size and amount? (For each drink checked, indicate the size(s) and how many you had.)		
Drinks (all flavors, regular or diet, not decaffeinated)	Your Size	How many?	
<input type="checkbox"/> Coffee	Regular coffee mug = 8oz	X-small (8oz) Small (12oz) Medium (16oz) Large (20oz) X-large (24oz)	0 _____ 0 _____ 0 _____ 0 _____ 0 _____
<input type="checkbox"/> Coffee drinks (Frappuccinos, etc.)	Starbucks "Tall" = 12oz	X-small (9.5oz) Small (12oz) Medium (16oz) Large (20oz) X-large (24oz)	0 _____ 0 _____ 0 _____ 0 _____ 0 _____
<input type="checkbox"/> Espresso (Latte, mocha, espresso shot, etc.)	Small mocha = 1 shot	Small (1 shot) Medium (2 shots) Large (3 shots)	0 _____ 0 _____ 0 _____
<input type="checkbox"/> Caffeinated sodas (colas, root beer, Mountain dew)	Regular can soda = 12oz	X-small (12oz) Small (16oz) Medium (20oz) Large (24oz) X-large (32oz)	0 _____ 0 _____ 0 _____ 0 _____ 0 _____
<input type="checkbox"/> Energy drinks (Red Bull, Rockstar, Monster Starbucks Double Shot, etc.)	Regular can Red Bull = 8oz	Small (8oz) Medium (12oz) Large (16oz) X-large (24oz)	0 _____ 0 _____ 0 _____ 0 _____
<input type="checkbox"/> Tea (Green, black, sweet tea)	Small McDonald's sweet tea = 16oz	X-small (12oz) Small (16oz) Medium (21oz) Large (32oz)	0 _____ 0 _____ 0 _____ 0 _____
<input type="checkbox"/> Caffeinated water (Avitae)	1 bottle = 16.9oz	Avitae (16.9oz) Avitae 90 (16.9oz)	0 _____ 0 _____
Energy Boosting Supplements (all flavors, regular or diet, not decaffeinated)	Standard Serving Size	How many?	
<input type="checkbox"/> Energy shots (5-hour Energy, 6 Hour Power, etc.)	1 bottle	_____	
<input type="checkbox"/> Caffeine supplements (NoDoz)	1 capsule	_____	
<input type="checkbox"/> Energy chewing gum (Jolt Caffeine Energy Gum)	2 pieces	_____	
<input type="checkbox"/> Sheets Energy Strips	2 strips	_____	

APPENDIX D

DEMOGRAPHIC AND CAFFEINE EFFECTS QUESTIONNAIRE

Demographic Survey																	
1. In the past two weeks, which shifts did you work (mark all that apply), & how many of each shift did you work? <table border="0"> <tr> <td style="text-align: center;"><u>Shift</u></td> <td style="text-align: center;"><u>How many</u></td> </tr> <tr> <td><input type="checkbox"/> 8-10 hour night</td> <td>_____</td> </tr> <tr> <td><input type="checkbox"/> 8-10 hour day</td> <td>_____</td> </tr> <tr> <td><input type="checkbox"/> 12 hour night</td> <td>_____</td> </tr> <tr> <td><input type="checkbox"/> 12 hour day</td> <td>_____</td> </tr> </table>				<u>Shift</u>	<u>How many</u>	<input type="checkbox"/> 8-10 hour night	_____	<input type="checkbox"/> 8-10 hour day	_____	<input type="checkbox"/> 12 hour night	_____	<input type="checkbox"/> 12 hour day	_____	2. How long have you been at your current position? _____			
<u>Shift</u>	<u>How many</u>																
<input type="checkbox"/> 8-10 hour night	_____																
<input type="checkbox"/> 8-10 hour day	_____																
<input type="checkbox"/> 12 hour night	_____																
<input type="checkbox"/> 12 hour day	_____																
3. How many years have you participated in shift work? _____				4. How often do you switch from one shift to another? (Example: every 4 weeks.) _____													
5. For each statement listed below, circle the number that best represents how you feel.																	
Extremely low → Normal → Extremely high																	
What is your energy level when participating in shift work?		1	2	3	4	5	6	7									
What is your concentration level while participating in shift work?		1	2	3	4	5	6	7									
6. In the past two weeks, did you experience any of the following side effects related to caffeine intake? (Mark all that apply.)																	
<input type="checkbox"/> Headaches				<input type="checkbox"/> Stomach ache													
<input type="checkbox"/> "Jolt and crash" episodes				<input type="checkbox"/> Heart burn													
<input type="checkbox"/> Heart palpitations				<input type="checkbox"/> Don't have any of the side effects listed.													
7. In the past two weeks, how many days did you work overtime? _____ days				8. What is your department / division?													
				<input type="checkbox"/> Patrol <input type="checkbox"/> Investigation <input type="checkbox"/> Sheriff's <input type="checkbox"/> Dispatch													
9. Do you currently use tobacco? <input type="checkbox"/> Yes <input type="checkbox"/> No				10. Age: _____ years		11. Gender: <input type="checkbox"/> Male <input type="checkbox"/> Female											
12. Marital Status: <input type="checkbox"/> Single <input type="checkbox"/> Married <input type="checkbox"/> Divorced <input type="checkbox"/> Widowed				13. Race: <input type="checkbox"/> Non-Hispanic White <input type="checkbox"/> Asian American <input type="checkbox"/> Hispanic or Latino <input type="checkbox"/> Black or African American <input type="checkbox"/> American Indian or Alaskan Native <input type="checkbox"/> Hawaiian or other Pacific Islander <input type="checkbox"/> Other. Specify: _____													

APPENDIX E

SERVING SIZE REFERENCE SHEET

Drink Sizes

Drink	Sizes & Examples
Coffee	X-small (8oz).....Small coffee mug; Starbucks <i>short</i> Small (12oz).....Starbucks <i>tall</i> ; Sheetz small Medium (16oz).....Starbucks <i>grande</i> ; Sheetz medium; regular size travel mug Large (20oz).....Starbucks <i>venti</i> ; Sheetz large X-large (24oz).....Sheetz X-large
Coffee drinks	X-small (9.5oz).....Bottled Starbucks Frappuccino Small (12oz).....Starbucks <i>tall</i> ; Sheetz small Medium (16oz).....Starbucks <i>grande</i> ; Sheetz medium Large (20oz).....Starbucks <i>venti</i> ; Sheetz large X-large (24oz).....Sheetz X-large
Espresso / Espresso drinks	Small (1oz).....1 shot espresso; small/medium latte Medium (2oz).....2 shots espresso; large latte Large (3oz).....3 shots espresso; large latte with added shot espresso
Sodas	X-small (12oz).....Regular soda can Small (16oz).....16oz soda bottle; McDonald's/ Bojangles small Medium (20oz).....McDonald's medium; Bojangles regular Large (24oz).....24oz bottle X-large (32oz).....McDonald's/ Bojangles large
Energy drinks	Small (8oz).....Smallest can Red Bull, Rockstar, & Monster Energy Medium (12oz).....Medium size can Red Bull Large (16oz).....Largest can Red Bull; medium can Rockstar & Monster Energy X-large (24oz).....Largest can Rockstar & Monster Energy
Tea	X-small (12oz).....McDonald's <i>child</i> size tea Small (16oz).....McDonald's & Bojangles small tea Medium (21oz).....McDonald's medium tea; Bojangles regular tea Large (32oz).....McDonald's & Bojangles large tea

Miscellaneous examples:

Small coffee mug = 8-10 oz

Larger coffee mugs = 16-20 oz

Standard travel mug/tumbler = 16oz

Sheetz and other gas station cup sizes = 12oz, 16oz, 20oz, 24oz

Big gulp or large soda/tea from McDonald's or Bojangles= 32oz

Starbucks cup sizes = *short* 8oz, *tall* 12oz, *grande* 16oz,
venti 20oz, *venti iced* 24oz

Use the image below to gauge the size drink you had. This is

