

Elsevier Editorial System(tm) for Clinical Biomechanics
Manuscript Draft

Manuscript Number: CLBI-D-11-00638R3

Title: MIRROR GAIT RETRAINING FOR THE TREATMENT OF PATELLOFEMORAL PAIN IN FEMALE RUNNERS

Article Type: Research Paper

Keywords: patellofemoral pain; running; rehabilitation; hip; patella

Corresponding Author: Dr. Richard Willy, PhD

Corresponding Author's Institution: Ohio University

First Author: Richard W Willy, PT, PhD

Order of Authors: Richard W Willy, PT, PhD; John P Scholz, PT, PhD; Irene S Davis, PT, PhD

Abstract: **Background:** Abnormal hip mechanics are often implicated in female runners with patellofemoral pain. We sought to evaluate a simple gait retraining technique, using a full-length mirror, in female runners with patellofemoral pain and abnormal hip mechanics. Transfer of the new motor skill to the untrained tasks of single leg squat and step descent was also evaluated. **Methods:** Ten female runners with patellofemoral pain completed 8 sessions of mirror and verbal feedback on their lower extremity alignment during treadmill running. During the last 4 sessions, mirror and verbal feedback were progressively removed. Hip mechanics were assessed during running gait, a single leg squat and a step descent, both pre- and post-retraining. Subjects returned to their normal running routines and analyses were repeated at 1-month and 3-month post-retraining. Data were analyzed via repeated measures analysis of variance. **Findings:** Subjects reduced peaks of hip adduction, contralateral pelvic drop, and hip abduction moment during running ($P < 0.05$, effect size=0.69-2.91). Skill transfer to single leg squatting and step descent was noted ($P < 0.05$, effect size=0.91-1.35). At 1 and 3 months post retraining, most mechanics were maintained in the absence of continued feedback. Subjects reported improvements in pain and function ($P < 0.05$, effect size=3.81-7.61) and maintained through 3 months post retraining. **Interpretation:** Mirror gait retraining was effective in improving mechanics and measures of pain and function. Skill transfer to the untrained tasks of squatting and step descent indicated that a higher level of motor learning had occurred. Extended follow-up is needed to determine the long term efficacy of this treatment.

Division of Physical Therapy
Ohio University
Athens, Ohio 45701

6 December 2011

Dear Sir/Madam,

For the attached manuscript titled, "Mirror gait retraining for the treatment of patellofemoral pain syndrome in female runners," There are no commercial relationships which may lead to a conflict of interest for any author. Additionally, there are no conflicts of interests on the part of the authors.

Yours sincerely,

*Richard Willy

* Corresponding author

Richard Willy
W273 Grover Center
Division of Physical Therapy
Ohio University
Athens, Ohio 45701

Tel: 740 593 1209
Fax: 740 593 0293
Email: willyr@ohio.edu

Division of Physical Therapy
Ohio University
Athens, Ohio 45701

26 July 2012

Dr. Burton,

Please consider the resubmission of the manuscript titled, "Mirror gait retraining for the treatment of patellofemoral pain syndrome in female runners" for publication in *Clinical Biomechanics* as an original research article. Within the revised manuscript, our revisions are highlighted in yellow. Within this document, the reviewers' comments are in bold and our responses are in standard text.

This work has not been published nor submitted elsewhere. Each author was fully involved in the study and preparation of the manuscript and concurs with the content in the final manuscript. The material within the article has not been and will not be submitted for publication elsewhere except as an abstract. Finally, there are no commercial relationships which may lead to a conflict of interest for any author.

Yours sincerely,

*Richard Willy

* Corresponding author

Richard Willy
W273 Grover Center
Division of Physical Therapy
Ohio University
Athens, Ohio 45701

Tel: 740 593 1209
Fax: 740 593 0293
Email: willyr@ohio.edu

Response to review of Manuscript CLBI-D-11-00638: MIRROR GAIT RETRAINING FOR THE TREATMENT OF PATELLOFEMORAL PAIN IN FEMALE RUNNERS

We have addressed each of your comments in the manuscript body and/or following each comment below. Your comments are listed in **bold** and our responses are in standard text. Within the body of the manuscript, any subsequent edits are highlighted in **yellow**.

Main concerns:

You were previously asked to avoid abbreviations in the abstract: kindly make the corrections now.

All abbreviations have been removed from the abstract.

1 **TITLE:** MIRROR GAIT RETRAINING FOR THE TREATMENT OF PATELLOFEMORAL
2 PAIN IN FEMALE RUNNERS

3 **Authors:** Richard W. Willy, PT, PhD^{1*}

4 John P. Scholz, PT, PhD²

5 Irene S. Davis, PT, PhD³

6 ¹Division of Physical Therapy, Ohio University, Athens, OH, USA

7 ²Department of Physical Therapy, University of Delaware, Newark, Delaware, USA

8 ³Spaulding National Running Center, Harvard Medical School, Cambridge, MA, USA

9

10 ***Corresponding Author:**

11 Richard Willy, PT, PhD

12 W273 Grover Center, Division of Physical Therapy

13 Ohio University, Athens, Ohio 45701 USA

14 email: willyr@ohio.edu

15 Telephone: (740) 593-1209

16 Fax: (740) 593-0293

17 Word Counts: Abstract= 250

18 Text= 3987

19 Number of Figures= 3

20 Number of Tables= 2

1 **MIRROR GAIT RETRAINING FOR THE TREATMENT OF PATELLOFEMORAL**
2 **PAIN IN FEMALE RUNNERS**

3 **Abstract**

4 *Background:* Abnormal hip mechanics are often implicated in female runners with
5 patellofemoral pain. We sought to evaluate a simple gait retraining technique, using a full-length
6 mirror, in female runners with patellofemoral pain and abnormal hip mechanics. Transfer of the
7 new motor skill to the untrained tasks of single leg squat and step descent was also evaluated.

8 *Methods:* Ten female runners with patellofemoral pain completed 8 sessions of mirror
9 and verbal feedback on their lower extremity alignment during treadmill running. During the last
10 4 sessions, mirror and verbal feedback were progressively removed. Hip mechanics were
11 assessed during running gait, a single leg squat and a step descent, both pre- and post-retraining.
12 Subjects returned to their normal running routines and analyses were repeated at 1-month and 3-
13 month post-retraining. Data were analyzed via repeated measures analysis of variance.

14 *Findings:* Subjects reduced peaks of hip adduction, contralateral pelvic drop, and hip
15 abduction moment during running ($P < 0.05$, effect size=0.69-2.91). Skill transfer to single leg
16 squatting and step descent was noted ($P < 0.05$, effect size=0.91-1.35). At 1 and 3 months post
17 retraining, most mechanics were maintained in the absence of continued feedback. Subjects
18 reported improvements in pain and function ($P < 0.05$, effect size=3.81-7.61) and maintained
19 through 3 months post retraining.

1 *Interpretation:* Mirror gait retraining was effective in improving mechanics and measures
2 of pain and function. Skill transfer to the untrained tasks of squatting and step descent indicated
3 that a higher level of motor learning had occurred. Extended follow-up is needed to determine
4 the long term efficacy of this treatment.

5 Key Words: patellofemoral pain, running, rehabilitation, hip, patella

6

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21

Introduction

Running is one of the most popular forms of exercise with upwards of 16 million Americans participating in the sport today (National Sporting Goods Association, 2009). However, runners report an alarmingly high annual injury rate between 19.4 and 79.3% (van Gent et al. 2007). Patellofemoral pain syndrome (PFP) is the most prevalent type of knee pain among runners (Taunton et al. 2002). As with most knee injuries, there is a sex bias associated with PFP with 68% of sufferers of PFP being female (Taunton et al. 2002).

Abnormal hip mechanics in females with PFP have been reported during running, squatting, and during a step down maneuver (Willson and Davis, 2008, Dierks et al. 2008, Souza and Powers, 2009a, b, Noehren et al. 2012). These motions include excessive contralateral pelvic drop (Dierks et al., 2008, Willson et al., 2008), hip adduction (Willson et al. 2008, Dierks et al, 2008), and hip internal rotation (Dierks et al. 2008, Souza et al. 2009 a,b). Together, these motions likely increase the dynamic Q-angle of the knee (Powers 2003, 2010) resulting in an increase in joint stress on the lateral aspect of the patellofemoral joint (Besier et al. 2008, Huberti and Hayes, 1984, Lee et al. 2001).

The gluteus medius and maximus are the primary stabilizers of the hip in the frontal and transverse planes. Because weakness of this musculature has been identified in individuals with active PFP, hip strengthening is often advocated for the treatment of PFP. Hip strengthening has clearly been shown to result in short term pain reduction in PFP cohorts (Dolak et al. 2011, Mascal et al. 2003, Ferber et al. 2011, Earl et al. 2011, Nakagawa et al. 2007, Boling et al. 2006). However, the ability of hip strengthening to change lower extremity mechanics is largely

1 unproven. After a hip strengthening program in healthy females with normal running mechanics,
2 Snyder and colleagues (Snyder et al. 2007) actually reported an *increase* of hip adduction
3 excursion, the opposite of the desired direction. More recently, Willy and Davis (2011)
4 suggested that a 6-week program that significantly increased hip strength had no effect on
5 abnormal hip mechanics during running. In females with PFP, Earl et al. (Earl et al. 2011) failed
6 to find any change in hip and knee kinematics after a hip strengthening and flexibility program.
7 Therefore, it appears that hip strengthening alone is insufficient to change the proximal
8 mechanics that have been associated with PFP in females. If the underlying mechanics of PFP
9 are not addressed by a treatment modality, then symptoms will likely eventually return.

10 Treatments for PFP that directly address faulty proximal mechanics have shown promise.
11 Gait retraining, using real-time kinematic feedback, has been suggested as an effective treatment
12 of PFP in female runners (Noehren et al. 2010). Using a real time motion analysis system, the
13 hip adduction angle during each stance phase of treadmill running was provided to the runner in
14 real time. Subjects were encouraged match their hip angle to a normative target range provided
15 on the monitor. Significant improvements in hip mechanics, knee pain and overall function were
16 found at post-gait retraining. Interestingly, the reductions in abnormal hip mechanics during
17 running transferred to the untrained task of single leg squatting. Importantly, these subjects were
18 able to maintain their new movement patterns at the 1-month follow up. Improved pain and
19 function scores also persisted, thus suggesting the potential for long-term changes in mechanics,
20 pain, and function.

21 There are limitations to the work by Noehren and colleagues (Noehren et al. 2010). First,
22 most rehabilitation settings do not have the resources to purchase and/or operate a real-time

1 motion capture system. Thus, the clinical utility of real-time kinematic gait retraining is limited
2 to research oriented treatment centers at best. In order to implement this gait retraining technique
3 in clinical settings, it is possible that visual feedback could be provided by a mirror rather than a
4 motion capture system. If successful, mirror feedback would provide a simple, cost-effective
5 method of retraining. In addition, it might be more informative to assess the transfer of the new
6 movement pattern during running to a more functional activity than the single leg squat, such as
7 step descent. Finally, subjects were only followed for 1 month post-gait retraining. Most studies
8 define chronic PFP as having a duration of at least 3 months (Crossley et al. 2006, Souza and
9 Powers 2009a,b, Nakagawa et al., 2009). Thus, investigations of potential treatments of PFP
10 should likely follow subjects for at least 3 months post-intervention to demonstrate stronger
11 evidence of symptom resolution.

12 Therefore, the purpose of this study was to examine the effect of a clinically applicable,
13 gait retraining program on hip mechanics, pain and function in runners with PFP. We
14 hypothesized that mirror gait retraining would improve measures of abnormal hip mechanics
15 during running. We further hypothesized that these changes would persist through 3 months
16 post-intervention. We also hypothesized that subjects would successfully transfer the new
17 movement pattern to the untrained tasks of SLS and step descent with similar persistence through
18 3 months. Finally, it was hypothesized that subjects would report a decrease in reported pain and
19 an increase in function through 3 months post-intervention.

Methods

The data collection procedures and informed consent document were approved by the University of Delaware Human Subjects Research Board. An a priori power analysis was conducted using data from pilot work examining the difference between females with PFP and healthy controls. Using the variable peak hip adduction (HADD), it was revealed that 9 subjects (effect size=1.43 $\alpha=0.05$, $\beta=0.20$) were required to adequately power this study. Therefore, 10 qualified subjects were recruited for this study.

Subjects were female, between 18 and 40 years of age, running at least 10 km/week, comfortable with treadmill running at 3.35 m/sec, and free of any cardiac risk factors. All subjects were required to have retropatellar or peripatellar pain that was insidious in nature and self-rated at least at a “3” (moderate) on a visual analog scale of “0” to “10” during running. These symptoms were required to be present during running and at least one of the activities of jumping, squatting, kneeling, prolonged sitting, or stair descent. All subjects with patellofemoral instability or other knee diagnoses, history of any lower extremity surgery, or who were otherwise unhealthy were excluded.

All qualified subjects were invited for the kinematic screening session. In the presence of bilateral knee pain, the knee with the highest self-rated pain was analyzed. When pain was equal bilaterally, the most dominant limb was analyzed, defined as the limb used to kick a soccer ball. To analyze overall function, subjects completed the Lower Extremity Functional Scale (LEFS). The LEFS is a 20-question clinical measure. Subjects ranked the amount a lower extremity injury affects various tasks and activities on a scale of 0-4 with a “0” signifying “extreme

1 difficulty” and a “4” signifying “no difficulty” with a score of 80/80 corresponding to no
2 limitations. The LEFS has previously been validated in PFP populations and a minimal
3 clinically important difference of 9 points has been established (Brinkley et al.1999).

4 Thirty retro-reflective markers were attached to the involved lower extremity to analyze running,
5 SLS and step descent mechanics. To control for the effect of footwear on mechanics, subjects
6 wore standardized neutral running shoes (Nike Pegasus, Beaverton, OR). The use of
7 standardized running shoes was particularly important as qualified runners were enrolled in this
8 study for 3 months. During this time period, the shoes utilized by the subjects during their
9 everyday running likely experienced considerable wear or were even replaced. Thus, our use of
10 standardized running shoes controlled for this likelihood and allowed us to control for this
11 potential influence on running mechanics. Placement of anatomical markers was recorded with a
12 marker placement device (MPD). This device was used to improve marker placement reliability
13 when gait data are being measured over time. Intraclass correlation coefficient values of at least
14 0.9 were reported for all hip and knee variables when using the MPD (Noehren et al. 2010). The
15 marker placement measurements were used for all subsequent data collections for each subject.

16 Three-dimensional marker coordinates were captured with an 8-camera, MX Vicon
17 motion analysis system (VICON, Oxford, UK). A standing calibration trial was collected while
18 the subject stood on a force plate (Bertec, Worthington, OH) mounted in the center of the capture
19 volume. To assure standardized lower extremity position for standing calibration trials at follow-
20 up data sessions, each subject’s foot position was recorded with a foot tracing for the baseline
21 standing calibration trial. Next, a spherical hip trial was collected to calculate the functional hip
22 joint center (Hicks and Richards, 2005).

1 All kinematic and kinetic data were sampled at 200 Hz and 1000 Hz, respectively. After
2 completing approximately 10 warm-up trials, running data were collected as subjects traversed a
3 25-meter runway at 3.35 m/sec (8 min/mile). Next, single leg squat data were collected as
4 subjects performed a squat to approximately 60 degrees knee flexion. Movement data were then
5 collected as subjects descended an 8-inch instrumented step. Both single leg squat and step
6 descent speed was standardized to a 1 Hz count. Approximately 8 trials of both the SLS and step
7 descent were collected to obtain 5 trials with the correct speed of movement. Finally, we
8 collected baseline video during treadmill running for future education purposes.

9 Kinematic and kinetic data from 5 trials were filtered with an 8-Hz and 50-Hz, low-pass,
10 fourth order, zero-lab Butterworth filter, respectively. Internal joint moments were calculated
11 utilizing segment inertial properties (Dempster et al. 1959) and normalized to body mass and
12 height. Single leg squat mechanics were analyzed at 45 degrees knee flexion as this index
13 represents typical peak knee flexion seen during running in our lab. Step descent mechanics
14 were indexed at the point of peak knee extensor moment, which has been reported to correspond
15 to peak quadriceps muscle force (Andriacchi et al. 1984). Customized software (LabVIEW 8.0,
16 National Instruments, Austin, TX) was used to extract the discrete variables of interest from five
17 individual curves for the motion files. Means and standard deviations of these values were
18 calculated.

19 Those subjects who demonstrated abnormal hip alignment during running were invited to
20 participate in the gait retraining phase. Abnormal hip alignment during running was
21 operationally defined as peak HADD greater than 1 standard deviation above the mean of our
22 lab's normative database of running (peak HADD qualifying criterion= 20°). For qualified

1 subjects, data collected during the kinematic screening served as their baseline data for the gait
2 retraining protocol. Runners who did not meet this kinematic inclusion criterion were dismissed
3 from further participation in the study.

4 Participants who met the kinematic qualification criteria attended a total of eight gait
5 retraining sessions over the course of two weeks. During the first training session, subjects were
6 first shown their baseline video so they could visualize the abnormal hip and knee alignment that
7 they exhibited during running. During gait retraining, visual feedback was provided by a full
8 length mirror that was placed directly in front of the treadmill. Participants received scripted
9 verbal cueing at the beginning of each session, consisting of “run with your knees apart with
10 your kneecaps pointing straight ahead” and “squeeze your buttocks.” Subjects received
11 additional verbal feedback during each training session if they were not maintaining the desired
12 gait modifications. During all training sessions, each subject’s response to the cueing was
13 analyzed subjectively using a standard video camera and compared with their baseline video.
14 Pilot work suggested that subjects may attempt to run with a widened stance or an increased toe
15 out in a maladapted attempt to reduce HADD and HIR, respectively. If either of these
16 maladaptations was noted on the video feed, subjects were immediately cued to correct them.

17 Feedback exposure and treadmill runtime were tightly controlled. Runners attended 8
18 retraining visits over the course of the 2 weeks, during which treadmill runtime was gradually
19 increased from 15 minutes to 30 minutes (Figure 1). This schedule is based on previous studies
20 (Noehren et al. 2010, Barrios et al. 2010, Crowell and Davis, 2010). Feedback was gradually
21 removed during the final 4 training sessions, in accordance with the feedback schedule, to shift
22 dependence from external to internal cues and reinforce learning (Winstein and Schmidt, 1990).

1 This was accomplished by decremental reductions in verbal cueing in addition to reductions in
2 visual feedback by turning the mirror around so that subjects could not see themselves while
3 running. During each period of feedback removal, running mechanics were monitored via a
4 standard video stream that was only visible to the investigator. Once feedback was resumed,
5 runners received retrospective verbal cueing on their running mechanics during the preceding
6 feedback removal period. To ensure that feedback was consistent across subjects, runners were
7 not permitted to run outside of the lab while participating in the gait retraining phase. Subjects
8 were asked to verbally attest to the restrictions on running activity. Subjects were required to
9 complete all 8 sessions.

10 An instrumented gait analysis was repeated at the conclusion of the 2-week gait
11 retraining program (POST). During this session, anatomical markers were placed according to
12 the measurements obtained at the baseline visit using the marker placement device. Running,
13 single leg squat and step descent data were collected in the same manner as during the baseline
14 visit. Pain rating and LEFS data were recorded. After the post- retraining data collection,
15 subjects were asked to return to their normal running routine. Further follow-up instrumented
16 motion analyses, pain ratings, and LEFS were collected at 1MO and 3MO post-retraining. The
17 variables of interest during running, squatting, and step descent were contralateral pelvic drop
18 referenced to the lab (CPD), HADD, thigh adduction referenced to the lab (thigh ADD), hip
19 abduction moment (HABDM), and hip internal rotation (HIR). Peak values were utilized for the
20 analysis of running and the appropriately indexed values were utilized for analysis of the SLS
21 and step descent.

1 Repeated measure, one way ANOVA's were used for statistical analysis of running, SLS,
2 and step descent. Sphericity of the data was assessed with Mauchly's test with $\alpha < 0.05$. In the
3 case of a positive Mauchly's test, a Huhn-Feldt correction was conducted during the analysis to
4 generate accurate α scores (Field, 2008). Post hoc 2-tailed comparisons were conducted using a
5 criterion $\alpha < 0.05$, while a trend was defined as between $\alpha \leq 0.10$ and $\alpha > 0.05$. Post hoc
6 comparisons were conducted from PRE to POST. Comparisons were then made between POST
7 and 1MO and again between POST and 3MO. Effect sizes were also calculated. A large effect
8 size was defined as ≥ 0.80 , moderate ≥ 0.40 , and small < 0.40 (Cohen, 1992).

9 **Results**

10 A total of 10 subjects completed the study (Table 1). An additional 3 subjects were lost to
11 drop out. Two dropped out as they were unable to comply with the restriction of no running
12 outside of the training sessions. The third subject dropped out due to health issues unrelated to
13 the study. None of the 3 dropouts completed the 8-session training phase. Thus, an intention to
14 treat analysis was not conducted.

15 Subjects demonstrated a visible reduction in their peak HADD and CPD during running
16 at POST (Figure 2). In addition, the ANOVA was significant for the variables of peak CPD,
17 HADD, thigh ADD, and HABDM, leading to the examination of pair-wise comparisons (Table
18 2). At POST, subjects significantly reduced peak CPD (Figure 3a) and maintained these changes
19 for peak CPD through 1MO and 3MO. At POST, subjects successfully reduced peak HADD by a
20 mean of 5.9° ($sd=1.5$), (Figure 3b). However, at 1MO peak HADD increased by 1.1° , $sd=1.2$,
21 and again by 0.6° , $sd=2.1$ at 3MO, moving closer to PRE levels. While significant, these changes

1 were small and were associated with relatively small effect sizes ($d=0.37$ and $d=0.23$ at 1MO
2 and 3MO, respectively). Peak thigh ADD was reduced at POST ($d=1.32$), suggesting that
3 changes in both the thigh and the pelvis segment contributed to the overall reduction in peak
4 HADD. Reductions in peak thigh ADD were maintained at 1MO and 3MO. At POST, peak
5 HABDM was successfully reduced (Figure 3a). This reduction in peak HABDM was maintained
6 through 1MO. However, peak HABDM increased significantly towards baseline levels at 3MO.
7 The ANOVA was nonsignificant for peak HIR baseline values (Figure 3b).

8 The improved hip mechanics noted during running transferred to the untrained task of
9 SLS, with the exception of CPD (Table 2). The ANOVA was significant for HADD ($P=0.000$),
10 thigh ADD ($P=0.01$) and HABDM ($P=0.008$), but not for CPD ($P=0.08$). Analysis of pairwise
11 comparisons revealed that HADD and thigh ADD were significantly reduced at POST with
12 reductions maintained at 1MO, However, a trend of drifting (by 1.1 deg) of HADD towards
13 baseline levels ($P=0.10$, $d=0.69$) was noted at 3MO. Interesting, thigh adduction also drifted
14 towards baseline levels at 3MO during the SLS ($d=0.92$). Similarly, HABDM was reduced at
15 POST ($d=0.91$) and with a trend towards baseline levels at POST ($P=0.058$). However, HABDM
16 drifted towards baseline at 3MO ($P=0.017$) and this drift was associated with a large effect size
17 ($d=1.19$). As with running, no changes were noted for HIR during the SLS.

18 Similar to SLS, the improved hip mechanics during running transferred to the untrained
19 task of step descent. During step descent, the ANOVA was significant for CPD ($P=0.001$),
20 HADD ($P=0.009$) and thigh ADD ($P=0.01$), but not HIR ($P=0.81$) or HABDM ($P=0.14$).
21 Examination of pairwise comparisons revealed that CPD was not reduced at POST, but
22 interestingly was reduced from POST values at 1MO ($d=1.09$) and reduced even more at 3MO

1 ($d=2.98$). HADD was reduced at POST, and maintained at 1MO and 3MO. There was a trend
2 ($P=0.10$) of reduction in thigh ADD at POST with no significant changes between POST-1MO
3 and POST-3MO, indicating retention of the kinematic change.

4 Both pain and LEFS scores were improved from PRE to POST (Figure 4a,b). These
5 changes were associated with large effect sizes ($d=7.61$ and $d= 3.81$, respectively) and the score
6 change of 9 points required to be a minimal clinically important difference (Brinkley et al.,
7 1999). Subjects maintained these improvements through 1MO and 3MO. The 1.8 point ($sd=2.2$)
8 increase in LEFS score between POST and 3MO was significant, yet did not exceed the
9 threshold for a minimal clinical important difference.

10

Discussion

11 Our primary goal was to determine if mirror gait retraining would reduce abnormal
12 running mechanics in females with PFP and if these changes would persist through 3MO. We
13 also investigated if the new movement skill during running would transfer and persist in the
14 untrained tasks of single leg squat and step descent. Finally, we sought to determine if these
15 changes in mechanics would be reflected in improvements in pain and function. Based on these
16 data, it appears that mirror gait retraining is an effective treatment to reduce abnormal mechanics
17 during running. These changes generally persisted through at least 3MO. Interestingly, this new
18 movement skill transferred to the untrained tasks of single leg squat and step descent. As with
19 running, alterations in mechanics during these untrained tasks were either maintained or
20 continued to improve through 3MO. Reflecting these changes in dynamic alignment, measures of
21 pain and function improved and remained at these levels through 3MO.

1 During running, the reductions in peak HADD were greater than any other variable
2 tested. Interestingly, post hoc examination of subject characteristics (age, baseline running
3 volume, baseline pain levels, and years of running experience) failed to yield any discernible
4 factor that predicted a runner's ability to decrease HADD after the retraining intervention.
5 HADD was likely the easiest variable for subjects to visualize by focusing on the space between
6 their knees. As the gluteus medius musculature is the primary stabilizer of the hip in the frontal
7 plane, runners may have altered the neuromuscular control of this muscle to achieve reductions
8 in HADD and CPD. Indeed, increased latency and decreased duration of activation of the gluteal
9 musculature has been implicated in PFPS in females during running and stair negotiation
10 (Willson et al. 2012, Cowan et al. 2009, Brindle et al. 2003). However, electromyography is
11 necessary to investigate changes in neuromuscular control of the gluteal musculature. It is
12 unknown if changes in gluteal strength accompanied changes in hip mechanics. However, the
13 retraining period was brief (2 weeks) and thus, was likely insufficient to stimulate true
14 strengthening. It is also possible that subjects may have shifted their trunk over the stance limb
15 to decrease CPD, resulting in less HADD. Our marker set did not include the trunk and thus, we
16 are unable to comment on the potential contribution of altered trunk mechanics. However, peak
17 thigh adduction during running was also reduced by 2.6° indicating that alterations in CPD did
18 not account for all of the reduction in peak HADD. The large reduction in HADD likely had a
19 marked effect on the dynamic Q-angle (Powers 2003, 2010). Decreasing the dynamic Q-angle
20 has been suggested to decrease lateral tracking of the patella, thus decreasing lateral joint stress
21 of the patellofemoral joint (Huberti and Hayes 1984, Lee et al. 2003, Besier et al. 2008). The
22 decrease in lateral patellofemoral joint stress would decrease forces on the subchondral bone,
23 leading to a decrease in pain (Besier et al. 2008).

1 Unlike the frontal plane measures of the hip during all tasks, we did not find any changes
2 in the transverse plane at POST. This was somewhat surprising as HADD and HIR are coupled
3 motions and participants also received specific cueing to reduce HIR during retraining sessions.
4 The reason for this discrepancy may be attributed to the fact that our kinematic inclusion
5 criterion was solely excessive peak HADD during running and not excessive peak HIR. Indeed,
6 most subjects did not exhibit excessive peak HIR during running at baseline. As with excessive
7 peak HADD, we operationally defined “excessive” as peak HIR greater than 1 standard deviation
8 above our lab’s normative database (peak HIR during running: present study= 8.6° , $sd=5.4$
9 versus our lab’s normative database= 5.0 , $sd=6.7$). Therefore, our participants may have had a
10 high capacity to change HADD but a limited capacity to make changes to HIR.

11 Following the overall significant improvement in hip mechanics during running at POST, the
12 majority of these changes persisted through 3 months. There was significant, albeit slight,
13 drifting of HADD in the direction of baseline levels by 8/10 runners. Suggesting the importance
14 of sufficient continued practice, the two runners with the greatest drifting of peak HADD during
15 running from PRE to 3MO had the lowest reported running volume. However, the drift from
16 POST to 1MO and to 3MO was only 1.1° and 1.6° , respectively. This drifting pattern may have
17 been in response to an initial overcorrection by the subjects. Subjects, in fact, exhibited a very
18 low value of $14.8^\circ \pm 3.1$ for HADD following the retraining. This value was well below the peak
19 HADD of our reference, normative database of runners ($16.5^\circ \pm 3.5$). At 3MO, peak HADD
20 ($16.4^\circ \pm 2.4$) was nearly identical to that of our database. This suggests that the mechanics may
21 have settled towards a more normal value at 3MO. Further evidence that a normalization of
22 running occurred, peak CPD at 3MO was nearly identical to peak CPD of our normative
23 database ($-8.0^\circ \pm 2.8$). Regardless, the general retention of changes in peak CPD, HADD, thigh

1 ADD, and HABDM at 1MO and 3MO suggest that subjects learned a new motor skill (Sherwood
2 and Lee, 2003, Salmoni et al. 1984).

3 Further evidence of skill acquisition was noted by the transfer of the new movement
4 pattern learned during running to the untrained tasks of SLS and step descent (Salmoni et al.
5 1984, Schmidt, 1972, and Sherwood and Lee, 2003). HADD and thigh adduction (a trend) were
6 reduced during SLS and step descent immediately following the retraining. Interestingly, CPD
7 was not reduced at POST during step descent. Nevertheless, it was reduced at 1MO and then
8 even further at 3MO. We speculate that subjects may have later reduced CPD at 1MO and 3MO
9 due to continued practice of this new skill in the community setting. In contrast to the drifting of
10 thigh ADD and the trend towards drifting noted in HADD during the SLS, these measures
11 remained consistent during step descent at 3MO. While SLS is often used as a functional test, it
12 is a movement that is rarely utilized during activities of daily living. Therefore, the SLS may not
13 have been as well-reinforced during normal activities as the step descent. Future studies should
14 examine changes in neuromuscular control of the gluteal musculature utilizing surface
15 electromyography.

16 The focus of the retraining was to alter excessive HADD during running. However, the
17 overarching goal of this intervention is to reduce pain and improve function in runners with PFP.
18 The reductions in pain visual analog scores noted in this study were of greater magnitude (-
19 90.5%, $sd=7.8$ greater than that reported in studies of other interventions of PFP (-43.1--87.5%)
20 (Nakagawa et al. 2009, Crossley et al. 2002, Ferber et al. 2011, Earl et al. 2011, Boling et al.
21 2006). Furthermore, the duration of this intervention (2 weeks) was considerably shorter than
22 that of previous intervention studies (3-8 weeks) of PFP that were focused on hip and knee

1 strengthening. (Nakagawa et al. 2009, Crossley et al. 2002, Ferber et al. 2011, Earl et al. 2011,
2 Boling et al. 2006). The greater reduction in pain and shorter duration of treatment makes gait
3 retraining an appealing intervention to minimize financial and time investment for clinical
4 settings. The 12.1 ± 2.7 point increase in LEFS scores at POST exceeded the 9-point minimally
5 clinically important difference associated with this measure (Brinkley et al. 1999). The increase
6 in this score indicates improved function and suggests that the new movement pattern translated
7 to functional movements that were previously painful. It is possible that participants experienced
8 these improvements in pain and function due to the decrease in running volume while enrolled in
9 the 2-week gait retraining phase. However, all runners reported returning to their pre-enrollment
10 running volume at 1MO and 3MO. In fact, 6/10 of the runners reported higher running volume at
11 3MO, attributed to a decrease in their knee pain. Importantly, participants maintained POST
12 levels of pain and LEFS through 3MO, suggesting potential for benefits lasting beyond the time
13 interval examined in this study.

14 Overall, the changes in running and squatting mechanics, as well as pain and function in
15 the current study compare favorably with real-time kinematic gait retraining (Noehren et al.
16 2010). The reductions seen in peak HADD and CPD during running are of the same magnitude,
17 or greater, than Noehren et al. However, those authors also reported a trend in reduction in peak
18 HIR, which was not seen in this study. The discrepancy is likely due to the fact that the subjects
19 in Noehren's study exhibited excessive HIR at baseline (Noehren: $11.0^\circ \pm 4.1$ versus norm:
20 $5.0^\circ \pm 7.1$). Thus, the subjects in Noehren et al. had greater capacity for change for HIR than
21 participants in the current study. Despite this difference, mirror retraining resulted in nearly
22 identical improvements in pain and function compared with real-time kinematic feedback.
23 Additionally, the reduction in HADD during the SLS was also similar to the changes in HADD

1 changes. The results of this study are promising, as this technique requires only a treadmill and a
2 full-length mirror. Further study is necessary to determine the long term efficacy of this
3 treatment technique.

4 **Acknowledgements**

5 This work was supported by a Promotional of Doctoral Studies scholarship from the Foundation
6 for Physical Therapy (Doctoral studies support for co-author RW), Drayer Physical Therapy
7 Institute (Doctoral studies support for co-author RW and subject reimbursement), and NIH 1 S10
8 RR022396 (Shared instrumentation grant).

9

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19

References

- Andriacchi TP, Andersson GB, Ortengren, R, Mikosz, RP, 1984. A study of factors influencing muscle activity about the knee joint. *J Orthop Res.* 12, 266-75.
- Barrios J, Crossley K, Davis I., 2010. Gait retraining to reduce the knee adduction moment through real-time visual feedback of dynamic knee alignment. *Journal of Biomech.* 91, 2208-13.
- Besier TF, Gold GE, Delp SL, Fredericson M, Beaupre GS., 2008. The influence of femoral internal and external rotation on cartilage stresses within the patellofemoral joint. *J Orthop Res.* 26, 1627-35.
- Boling M, Bolgla L, Mattacola C, Uhl T, Hose R., 2006. Outcomes of weightbearing rehabilitation program for patients diagnosed with patellofemoral pain syndrome. *Arch. Of Phys. Med. And Rehab.* 87, 1428-35.
- Brindle TJ, Mattacola C, McCrory J. 2003. Electromyographic changes in the gluteus medius during stair ascent and descent in subjects with anterior knee pain. *Knee Surg Sports Traumatol Arthrosc.* 11, 244-251.
- Brinkley J, Stratford P, Lott S, Riddle D., 1999. The lower extremity functional scale (LEFS): scale development, measurement properties and clinical application. *Phys Ther.* 79:371-83.

- 1 Cohen J. 1992. A power primer. *Psych. Bull.* 112, 155-159.
- 2 Cowan SM, Crossley KM, Bennell KL. 2009. Altered hip and trunk muscle function in
3 individuals with patellofemoral pain. *Br J Sports Med.* 43, 584-588.
- 4 Crossley K, Bennell K, Breen S, Cowan S, McConnell J., 2002. Physical therapy for
5 patellofemoral pain: A randomized, double-blinded, placebo-controlled trial. *Am. J. of Sp.*
6 *Med.* 30, 857-67.
- 7 Crowell HP, Davis IS., 2011. Gait retraining to reduce lower extremity loading in runners. *Clin*
8 *Biomech.* 26, 78-83.
- 9 Dempster, W.T., Gabel, W.C., Felts, W.J., 1959. The anthropometry of the manual work space
10 for the seated subject. *Am. J. Phys. Anthropol.* 17. 289–317.
- 11 Dierks TA, Manal KT, Hamill J, Davis IS., 2008. Proximal and distal influences on hip and knee
12 kinematics in runners with patellofemoral pain during and prolonged run. *J. Orthop*
13 *Sports Phys Ther.* 38, 448-456.
- 14 Dolak KL, Silkman C, Medina McKeon J, Hosey RG, Latterman C, Uhl TL., 2011. Hip
15 strengthening prior to functional exercises reduces pain sooner than quadriceps
16 strengthening in females with patellofemoral pain syndrome: a randomized clinical trial.
17 *J. Orthop Sports Phys Ther.* 41, 560-570.

18

- 1 Earl J, Hoch A., 2010. A proximal strengthening program improves pain, function, and
2 biomechanics in women with patellofemoral pain syndrome. *Am J Sports Med.* 59, 110-2.
- 3 Field A. Discovering Statistics using SPSS. 2nd Edition. Sage Publishing, London, Eng.
4 (2005).Pg. 451.
- 5 Ferber R., Kendall KD, Farr L., 2011. Changes in knee biomechanics after a hip-abductor
6 strengthening protocol for runners with patellofemoral pain syndrome *.J. Athl Train.* 46,
7 142-9.
- 8 Hicks J, Richards J., 2005. Clinical applicability of using spherical fitting to find hip joint
9 centers. *Gait & Posture.* 22, 138-45.
- 10 Huberti HH, Hayes WC., 1984. Patellofemoral Contact Pressures- The influence of Q-angle and
11 tendofemoralcontact. *J. of Bone and Joint Surg. Am.* 66A, 715-24.
- 12 Lee TQ, Yang BY, Sandusky MD, McMahon PJ., 2001. The effects of tibial rotation on the
13 patellofemoral joint: assessment of the changes in in situ strain in the peripatellar
14 retinaculum and the patellofemoral contact pressures and areas. *J. Rehabil. Res. Dev.* 38,
15 463-69.
- 16 Mascal CL, Landel R, Powers C., 2003. Management of patellofemoral pain targeting hip,
17 pelvis, and trunk muscle function: 2 case reports. *J Orthop Phys Ther.* 33, 647-60.
- 18 Nakagawa T, Muniz T, Baldon R, Maciel C, Reiff B, Serrao V., 2008. The effect of additional
19 strengthening of hip abductor and lateral rotator muscles in patellofemoral pain
20 syndrome: a randomized controlled pilot study. *Clin. Rehabil.* 22, 1051-60.

- 1 Noehren B, Manal K, Davis I., 2010. Improving day to day reliability using a marker placement
2 device. *J. of Biomech.* 28, 1405-10.
- 3 Noehren B, Pohl MD, Sanchez Z, Cunningham T, Latterman C. 2012. Proximal and distal
4 kinematics in female runners with patellofemoral pain. *Clin. Biomech.* 27, 366-371.
- 5 Noehren B, Scholz J, Davis I., 2010. The effect of real-time gait retraining on hip kinematics,
6 pain and function in subjects with patellofemoral pain syndrome. *Br J Sports Med.* 34,
7 691-696.
- 8 National Sporting Goods Association Web site [internet]. Mount Prospect, (Ill): [cited 2009 Nov.
9 30 11/30/09]. Available from :[http://www.nsga.org](http://www.nsga.org/files/public/2008RankedByTotal_4Web_080423.pdf)
10 [/files/public/2008RankedByTotal_4Web_080423.pdf](http://www.nsga.org/files/public/2008RankedByTotal_4Web_080423.pdf).
- 11 Powers C., 2003. The influence of altered lower extremity kinematics on patellofemoral joint
12 dysfunction: a theoretical perspective. *J. Orthop Sports Phys Ther.* 33, 639-46.
- 13 Powers C., 2010. The influence of abnormal hip mechanics on knee injury: a biomechanical
14 perspective. *J Orthop Sports Phys Ther.* 40, 42-1.
- 15 Salmoni, A.W., Schmidt, R.A., Walter, C.B., 1984. Knowledge of results and motor learning: a
16 review and critical reappraisal. *Psychological Bulletin:* 95, 355–86.
- 17 Schmidt RA., 1972. The case against learning and forgetting scores. *Journal of Motor Behavior:*
18 4, 79-8.

- 1 Selkowitz DM, Souza RB, Powers CM., 2009. Effect of femoral strapping on lower extremity
2 kinematics and pain response in females with patellofemoral pain. Proceedings of the
3 International Retreat: Patellofemoral pain syndrome: proximal, distal, and local factors.
4 April 30- May2, 2009. Baltimore, MD. *J. Orthop Sports Phys. Ther.* 3, A39-40.
- 5 Sherwood D, Lee T., 2003. Schema theory: critical review and implications for the role of
6 cognition in a new theory of motor learning. *Res Quart Exer Sport.* 74, 376-391.
- 7 Souza, RB, Powers CM., 2009. Differences in hip kinematics, muscle strength, and muscle
8 activation between subjects with and without patellofemoral pain. *J. Orthop. Sp. Phys.*
9 *Ther.* 39, 12-9.
- 10 Souza RB, Powers CM., 2009. Predictors of hip internal rotation during running: an evaluation
11 of hip strength and femoral structure in women with and without patellofemoral pain.
12 *Am. J. Sport. Med.* 37, 579-587.
- 13 Stathopulu E, Baildam E., 2003. Anterior knee pain: a long term follow-up. *Rheumatology.* 42,
14 380-2.
- 15 Taunton J, Ryan M, Clement D, McKenzie C, Lloyd-Smith R, Zumbo B., 2002. A retrospective
16 case control analysis of 2002 running injuries. *Br J Sports Med.* 36, 95-101.
- 17 van Gent B, Siem GD, Middelkoop M, van Os T, Bierma-Zeinstra S, Koes B., 2007 Incidence
18 and determinants of lower extremity running injuries in long distance runners: A
19 systematic review. *Br J Sports Med.* 41, 469-80.

- 1 Willson JD, Davis IS., 2008. Lower extremity mechanics of females with and without
2 patellofemoral pain across activities with progressively greater task demands. *Clin*
3 *Biomech.* 23, 203-11.
- 4 Willson J, Kernozek,T, Arndt, Reznichak D, Straker S. Gluteal muscle activation during running
5 in females with and without patellofemoral pain syndrome. *Clin. Biomech.* 2011; 26, 735-
6 740.
- 7 Willy RW, Davis IS., 2011. The Effect of a Hip Strengthening Program on Mechanics during
8 Running and during a Single Leg Squatting. *J Orthop Phys Ther.* 41, 625-32.
- 9 Winstein C, Schmidt R., 1990. Reduced frequency of knowledge of results enhances motor skill
10 learning. *J. Exp. Psyc.* 16, 677-91.

11

12

13

14

Table 1: Subjects demographics. Mean (SD).

	<u>Group mean (sd)</u>
Age (years)	22.4(5.0)
Running volume (km/week)	23.7(11.3)
Body Mass Index (kg/m ²)	22.0 (2.5)
Years running	5.8 (2.9)
Years with knee pain	4.3 (2.5)

Table 2: Group mean (SD) variables during for the variables of interest at the 4 time points. Variables for running were indexed to their peak values and were indexed to the discreet time points for the squat and step descent. Pairwise comparisons are indicated for PRE-POST, POST-1MO, and POST-3MO with respective effect sizes (*d*). * indicates $P < 0.05$, n/s indicates repeated measures ANOVA was non-significant. PRE= baseline measures, POST= post-gait retraining measures, 1MO= measures at 1 month post-gait retraining, 3MO= measures at 3 months post-gait retraining, CPD=contralateral pelvic drop, HADD= hip adduction, Thigh ADD= Thigh adduction, HABDM=Internal hip abduction moment, HIR= hip internal rotation.

<u>Variable</u>	<u>PRE</u>	<u>P</u>	<u>d</u>	<u>POST</u>	<u>P</u>	<u>d</u>	<u>1MO</u>	<u>P</u>	<u>d</u>	<u>3MO</u>
<u>Run</u>										
CPD	-9.0°(2.5)	0.02*	0.82	-7.1° (2.2)	0.171	0.19	-7.5° (2.3)	0.782	0.05	-7.0° (2.2)
HADD	20.7°(1.0)	0.000*	2.91	14.8° (3.1)	0.02*	0.37	15.9° (2.7)	0.05*	0.60	16.4° (2.5)
Thigh ADD	9.8°(1.2)	0.17*	1.32	7.2°(2.7)	0.92	0.04	7.3°(1.8)	0.29	0.47	8.1°(1.4)
HABDM (N*m/kg*m)	-1.180 (0.185)	0.042*	0.69	-1.054 (0.184)	0.605	0.11	-1.074 (0.173)	0.05*	0.61	-1.153 (0.145)
HIR	8.6° (5.4)	n/s	0.21	7.1° (8.7)	n/s	0.11	6.2° (7.9)	n/s	0.19	5.7° (6.3)
<u>Squat</u>										
CPD	0.6° (2.0)	n/s	0.11	2.3° (2.5)	n/s	0.10	2.6° (2.6)	n/s	0.04	2.2° (2.4)
HADD	11.6° (3.4)	0.002*	1.35	7.6° (2.6)	0.83	0.06	7.7° (2.6)	0.10	0.69	9.2° (2.1)
Thigh ADD	11.5°(2.0)	0.04*	0.68	10.1°(2.2)	0.48	0.15	9.8°(2.2)	0.92	0.03	10.1°(1.9)
HABDM (N*m/kg*m)	-0.470 (0.064)	0.004*	0.91	-0.412 (0.070)	0.06	0.28	-0.431 (0.071)	0.02*	0.18	-0.477 (0.039)
HIR	3.0° (6.4)	n/s	0.38	5.9° (8.5)	n/s	0.24	3.9° (7.6)	n/s	0.25	4.0° (6.4)
<u>Step</u>										
CPD	-5.0° (3.5)	0.72	0.07	-5.2° (2.7)	0.05*	1.09	-4.0° (3.0)	0.001*	1.28	-1.8° (2.6)
HADD	15.1° (6.8)	0.03*	0.69	11.6° (3.2)	0.51	0.07	11.1° (4.2)	0.88	0.04	11.8° (4.2)
Thigh ADD	7.8°(2.7)	0.10	0.42	6.8°(2.2)	0.13	0.12	6.2°(2.2)	0.55	0.16	6.4°(2.0)
HABDM (N*m/kg*m)	-0.556 (0.122)	n/s	0.34	-0.520 (0.085)	n/s	0.10	-0.498 (0.0813)	n/s	0.18	-0.506 (0.083)
HIR	7.0° (5.7)	n/s	0.03	7.3° (8.9)	n/s	0.16	6.0° (8.8)	n/s	0.10	8.0° (5.1)




Figure 1: The gait retraining schedule: Over the first four visits, runtime and feedback time is increased from 15 minutes to 24 minutes. Over the last four visits, runtime is increased to 30 minutes while feedback is faded to 3 minutes by the eighth visit.

Figure 2

[Click here to download high resolution image](#)



Figure 2: Representative subject during treadmill running a) PRE and b) POST. Note the reduction of contralateral pelvic drop and hip adduction at POST. This subject demonstrated a 3.8° , 8.1° , 4.5° reduction in contralateral pelvic drop, hip adduction, and thigh adduction respectively.

Figure_3

[Click here to download high resolution image](#)




Figure 3: A) Subjects reduced CPD from PRE to POST and maintained these changes through 3MO. B) Subjects reduced HADD from PRE to POST with a small amount of drifting towards baseline at 1MO and 3MO. C) Subjects reduced HABDM from PRE to POST and maintained these levels at 1MO. However, values returned to baseline levels at 3MO. D) No reductions were noted for HIR.




Figure 4: A) Subjects reported a significant reduction in visual analog score of pain from PRE to POST and maintained this reduction through 3MO. B) Lower extremity functional scale scores increased PRE to POST and maintained this reduction from POST to 1MO. A significant increase in this score resulted from POST to 3MO, albeit less than the minimally clinically important difference for this measure.