

Selected reactive oxygen species and antioxidant enzymes in common bean after *Pseudomonas syringae* pv. *phaseolicola* and *Botrytis cinerea* infection

Agata Nowogórska · Jacek Patykowski

Received: 30 April 2014 / Revised: 19 September 2014 / Accepted: 14 November 2014 / Published online: 30 November 2014
© The Author(s) 2014. This article is published with open access at Springerlink.com

Abstract *Phaseolus vulgaris* cv. Korona plants were inoculated with the bacteria *Pseudomonas syringae* pv. *phaseolicola* (*Psp*), necrotrophic fungus *Botrytis cinerea* (*Bc*) or with both pathogens sequentially. The aim of the experiment was to determine how plants cope with multiple infection with pathogens having different attack strategy. Possible suppression of the non-specific infection with the necrotrophic fungus *Bc* by earlier *Psp* inoculation was examined. Concentration of reactive oxygen species (ROS), such as superoxide anion (O_2^-) and H_2O_2 and activities of antioxidant enzymes such as superoxide dismutase (SOD), catalase (CAT) and peroxidase (POD) were determined 6, 12, 24 and 48 h after inoculation. The measurements were done for ROS cytosolic fraction and enzymatic cytosolic or apoplastic fraction. Infection with *Psp* caused significant increase in ROS levels since the beginning of experiment. Activity of the apoplastic enzymes also increased remarkably at the beginning of experiment in contrast to the cytosolic ones. Cytosolic SOD and guaiacol peroxidase (GPOD) activities achieved the maximum values 48 h after treatment. Additional forms of the examined enzymes after specific *Psp* infection were identified; however, they were not present after single *Bc* inoculation. Subsequent *Bc* infection resulted only in changes of H_2O_2 and SOD that occurred to be especially important during plant–pathogen interaction. Cultivar Korona of common bean is considered to be resistant to *Psp*

and mobilises its system upon infection with these bacteria. We put forward a hypothesis that the extent of defence reaction was so great that subsequent infection did not trigger significant additional response.

Keywords Reactive oxygen species · Antioxidant enzymes · *Pseudomonas syringae* pv. *phaseolicola* · *Botrytis cinerea* · Common bean

Introduction

During ontogenesis, a plant is exposed to many environmental factors including biotic and abiotic stresses. Generally it is accepted that the reactions of plants to necrotrophic pathogens are associated with jasmonic acid (JA) and/or ethylene (ET) signal transduction pathways, while biotrophic pathogens induce salicylic acid (SA) pathway (Glazebrook 2005). In addition, the recent studies have demonstrated that abscisic acid (ABA) is a regulator of plant defence against necrotrophs (Laluk et al. 2011). Adie et al. (2007) observed that ABA-impaired mutants simultaneously showed high susceptibility to *Pythium irregulare* and *Alternaria brassicicola* and increased resistance to *Botrytis cinerea*. Moreover, the authors presented that ABA functioning affected JA-dependent genes and their biosynthesis. Interactions between the mentioned hormones can be either synergistic or antagonistic (Koronneef and Pieterse 2008) bringing about either a positive effect, i.e. optimization of defence response to a pathogen, or producing a negative outcome during multiple infection with pathogens with different lifestyles (Beckers and Spoel 2005). Plant response to various environmental factors is also regulated by ROS and other molecules (Bowler and Fluhr 2000).

Communicated by R. Baczek-Kwinta.

A. Nowogórska (✉) · J. Patykowski
Department of Plant Physiology and Biochemistry, Faculty of
Biology and Environmental Protection, University of Łódź,
Banacha 12/16, 90-237, Łódź, Poland
e-mail: nowogorska.agata@gmail.com

Each plant–pathogen interaction may be characterised by specific features. It can be different depending on plant species and pathogen strains. Moreover, the stage of this interaction influences shifts in functioning of hormone signalling pathways (Derksen et al. 2013). Sometimes a plant exhibits resistance to a specific race of pathogen and it is related to gene-for-gene interactions (Flor 1971). Therefore, comprehensive defence against pathogen attack in plants is a result of a complex interaction between hormones, molecules and their signalling pathways, genetic mechanisms as well as their effects.

Common bean occurs in many varieties both resistant and susceptible to certain pathogens. Most of these plants are susceptible to the necrotrophic fungus *Botrytis cinerea* (*Bc*). This is mainly due to the attack strategy of this pathogen which consists in direct penetration of a cell wall or lesions using appressorium-like structures developed at the tip of conidial germ tubes (Garcia-Arenal and Sagasta 1980; Williamson et al. 2007). Halo blight caused by *P. syringae* pv. *phaseolicola* (*Psp*) is another disease affecting common bean. At the early stages of infection, this pathogen penetrates tissues through lesions and stomata and colonises apoplastic space (Glazebrook 2005). However, in later phases, *Psp* is necrogenic thus considered as a hemibiotroph (Thaler et al. 2004; Alfano and Collmer 1996).

In response to a biotic stress, a plant triggers a number of defence reactions which may be related either to direct immune responses or to a signal transduction pathway. Oxidative burst resulting in increased activity of the enzymes involved in ROS scavenging is a direct reaction after a pathogen infection. More intensive generation of ROS may increase resistance to some biotrophic pathogens (Lamb and Dixon 1997), however, cell death can promote growth of necrotrophic pathogens. Such reactions were confirmed in both cytosol and apoplast. The latter one is considered to be the first line of defence during pathogen attack and is involved in coordination of various defence reactions (Bolwell et al. 2001).

Peroxidases take part in many physiological processes throughout the whole life of a plant. Due to a large number of their isoforms and heterogeneous regulation of their expression (Passardi et al. 2005) they are good indicators of both biotic and abiotic stresses. Isoenzymes of class III peroxidases are mostly soluble apoplastic and cell wall-bound forms (Mika et al. 2010). In addition, they can be generated by post-transcriptional and post-translational modifications (Welinder et al. 2002). Peroxidases play a key role in plant–pathogen interactions, as they strengthen a cell wall by the formation of lignin, dityrosine bonds and of extension cross-links (Passardi et al. 2005). This kind of enzymes may use phenolic compounds especially syringaldazine (Goldberg et al. 1983) and ferulic acid

(Oudgenoeg et al. 2002) as substrates for hydrogen peroxide (H_2O_2) oxidation.

Superoxide dismutase (SOD) is an enzyme also involved in the metabolism of reactive oxygen species (ROS) by dismutation of superoxide anion to molecular oxygen and H_2O_2 . Moreover, particular isoforms of SOD interact with other enzymes protecting cells from the oxidative damage (Sakamoto et al. 1995). SOD through the production of H_2O_2 cooperates with catalase to control the appropriate level of this reactive oxygen form during infection.

Up to now, there is lack of information how plants cope with sequential infection with incompatible and non-specific pathogens. The aim of this study was to determine whether local reactions of *P. vulgaris* to the incompatible pathogen *Psp* induce a local response, which would prevent or slow down the non-specific infection with the necrotrophic fungus *Bc*. Thus, the levels of ROS and enzymes associated with their metabolism, especially those related to strengthening the structure of the cell wall during infection (including the creation of diferulic bonds), were studied. Therefore, especial attention was paid to reactions in the apoplastic spaces of infected bean leaves.

Materials and methods

Plant material and pathogen treatment

4-day-old seedlings of *P. vulgaris* cv. Korona were planted in pots with soil and grown in a growth chamber at temperature of 24 °C with 16/8 h day/night photoperiod at 70 % humidity. Light was supplied by white fluorescent lamps (36 W Philips TLD 36/84) at $350 \mu\text{E m}^{-2}\text{s}^{-1}$ intensity. Then, 14-day-old plants were divided into two groups: control and treated one. Those treated were inoculated with a single pathogen: *Psp* or *Bc* or sequentially with both pathogens: first the bacterial pathogen and after 2 days the necrotrophic fungus (*Psp+Bc*). 48 h was considered to be the optimum time inducing relatively permanent effects after plant response to first pathogen infection. The culture of *Psp* (strain No IOR 551 obtained from the Institute of Plant Protection in Poznań, Poland) was grown for 48 h on liquid King B medium at 24 °C with vigorous shaking, subsequently harvested, centrifuged at 10,000 rpm, rinsed twice, resuspended in sterile distilled water and adjusted to 10^7 cfu/cm³. The leaves were infiltrated with the bacterial suspension or water (control) using a needleless syringe. *B. cinerea* Pers. (obtained from the Institute of Plant Protection in Poznań, Poland) inoculation was performed by spraying plants with conidia suspension (1.5×10^6 (cm³)⁻¹) supplemented with 0.3 mM H_2KPO_4 and 2.2 mM glucose.

Areas of the infected leaves were calculated using the leaf scans imposed on a millimeter scale.

Determination of enzyme activities

Preparation of apoplastic extract

Apoplastic extracts were obtained from 1 g fresh weight of common bean leaves. Fluids were obtained according to the method of Polle et al. (1994) modified by Patykowski (2008) as amended, to adapt it to the bean leaves. About 4 cm² fragments of leaves were washed in deionized water for 5 min and subsequently vacuum-infiltrated with 10 mM NaCl in 50 mM phosphate buffer pH 6.5 at -0.9 kPa pressure at 4 °C for 10 min. Then the pieces were gently surface dried and centrifuged at 900g for 15 min at 4 °C in a syringe barrel placed in a centrifuge tube. Apoplastic fluid was rapidly collected and immediately used for enzymatic assays. Purity of the apoplastic fraction was confirmed by the activity of glucose-6-phosphate dehydrogenase (Creissen et al. 1999). Contamination with cytoplasm constituents was always less than 1.5 %.

Preparation of cytosol extract

1 g of the leaf tissue was homogenised in 5 cm³ of 1 M NaCl in 50 mM phosphate buffer pH 7.0 containing 1 mM EDTA and 1 % PVP (polyvinylpyrrolidone). Subsequently, the homogenate was centrifuged at 15,000 rpm for 15 min at 4 °C and the supernatant was used to analyse enzyme activities.

Guaiacol peroxidase (GPOD, EC 1.11.1.7) activity was measured spectrophotometrically by the modified method of Maehly and Chance (1954) in absorbance at 470 nm ($\epsilon = 26.6 \text{ mM}^{-1} \text{ cm}^{-1}$). The final reaction mixture contained 25 mM acetate buffer pH 5.6, 5 mM guaiacol, 15 mM H₂O₂ and the enzyme extract.

Peroxidase with syringaldazine (SPOD) capacity was measured at 530 nm ($\epsilon = 27 \text{ mM}^{-1} \text{ cm}^{-1}$) according to the method of Imberty et al. (1985). The total reaction mixture of 2 cm³ contained 50 mM potassium phosphate buffer pH 6.0, 41.6 μM syringaldazine, 0.11 mM H₂O₂ and the enzyme extract.

Peroxidase with ferulic acid (FPOD) activity was measured spectrophotometrically in absorbance at 310 nm ($\epsilon = 11.3 \text{ mM}^{-1} \text{ cm}^{-1}$) by the method of Takahama (1995). The total reaction mixture of 2 cm³ contained 50 mM potassium phosphate buffer pH 7.0, 0.14 mM ferulic acid, 0.1 mM H₂O₂ and the enzyme extract.

Analysis of superoxide dismutase (SOD, EC 1.15.1.1) activity was based on monitoring its ability to inhibit the

photochemical reduction of NBT following the method of Beauchamp and Fridovich (1971). The incubation mixture contained 50 mM potassium phosphate buffer pH 7.8, 13 mM methionine, 75 μM NBT, 2 μM riboflavin, 0.1 mM EDTA and the enzyme extract. The reaction was started by turning on the UV lamp. The absorbance was measured after 10 min at wavelength of 560 nm. SOD ability to inhibit 50 % of the photochemical reduction of NBT was used as one unit of its activity.

Catalase (CAT, EC 1.11.1.6) activity was determined using the assay of Dhindsa et al. (1981), which is based on H₂O₂ degradation ($\epsilon = 36.1 \text{ mM}^{-1} \text{ cm}^{-1}$). The total reaction mixture of 2 cm³ contained 50 mM potassium phosphate buffer pH 7.0, 15 mM H₂O₂ and the enzyme extract.

Assays of ROS

The hydrogen peroxide content was determined using the method described by Capaldi and Taylor (1983). 0.5 g of the leaf tissue was homogenised in 2.5 cm³ of 5 % trichloroacetic acid (TCA) with addition of 50 mg of activated charcoal in chilled mortars. Subsequently, the homogenates were centrifuged for 10 min at 15,000 rpm. Supernatants were adjusted to pH 3.6 using 4 M KOH and were treated as extracts. The reaction mixture contained: 0.2 cm³ of extract and 0.1 cm³ of 3.4 mM 3-methyl-2-benzothiazolinone hydrazone (MBTH) as well as 0.5 cm³ of horseradish peroxidase (90 U per 100 cm⁻³) solution in 0.2 M sodium acetate buffer (pH 3.6), which started the reaction. After 2 min, addition of 1.4 cm³ of 1 M HCl inhibited the reaction. The absorbance at 630 nm was measured after 15 min.

Histochemical visualisation of hydrogen peroxide was performed according to Thordal-Christensen et al. (1997) DAB staining. The leaves were incubated in an acetate buffer pH 3.8 containing 0.1 % DAB overnight and subsequently rinsed and discoloured with 96 % ethanol.

Superoxide anion concentration was measured according to the method of Doke (1983). Bean leaves were cut into discs 9 mm in diameter. Then, the discs were incubated for 1 h in the mixture containing: 50 mM phosphate buffer (pH 7.8), 0.05 % NBT, 0.1 mM EDTA and 0.065 % NaN₃. The mixture was then heated at 85 °C for 15 min and subsequently cooled. The NBT-reducing activity was presented as an increase in absorbance at 580 nm g⁻¹ of fresh weight.

For detection of superoxide anion, the leaves were incubated in 50 mM phosphate buffer (pH 7.8) containing 0.05 % NBT for 2 h according to the method of Romero-Puertas et al. (2004) modified by Shi et al. (2010). Then, the leaves were immersed in 96 % ethanol to wash out the chlorophyll.

Native PAGE

Native protein electrophoresis was performed at 100 V, on 7.5 % polyacrylamide gel, prepared using modified buffer system of Laemmli (1970). SOD isoenzyme activities were visualised according to the procedure of Beauchamp and Fridovich (1971). The reaction mixture contained: 50 mM phosphate buffer pH 7.8, 0.245 mM blue nitrotriazolium (NBT), 1 mM EDTA, 2.8 mM TEMED and 22 μ M riboflavine.

The gels were incubated for 15–20 min under UV light till light bands appeared. To identify different isoforms of SOD, 5 mM KCN (Cu/Zn-SOD inhibitor) or 5 mM H₂O₂ (Cu/Zn-SOD and Fe-SOD inhibitor) were used before staining. To visualise POD isoforms, the gels were incubated in 50 mM acetate buffer (pH 5.6) with addition of 30 mM H₂O₂ and 0.1 % 4-chloro-1-naphthol (dissolved in a minimal volume of 96 % ethanol) for 15 min and subsequently washed with distilled water. To detect isoenzymes of catalase, the gels were incubated in 0.005 % H₂O₂ for 10 min, rinsed twice with distilled water and stained with the reaction mixture containing 2 % ferricyanide [K₃Fe(CN)₆] and 2 % ferric chloride until light bands appeared.

Preparation of protein extracts

Bean leaves of each variant were homogenised in 1:5 ratio (w/v) in 50 mM TRIS–HCl buffer (pH 8.0) containing 3 mM EDTA, 1 mM DTT, 1 mM MgCl₂ and 2 % PVP. Then the homogenates were centrifuged at 20,000 rpm for 20 min. For each lane, 10–15 μ l of supernatants enriched with sucrose and 0.025 % bromophenol blue were applied.

Analysis of protein content

Protein content was measured by the method of Bradford (1976) using bovine serum albumin as a standard.

Statistical analysis

Sample variability for three or four replicates was given as standard deviation of mean. The statistical analyses were performed using Mann–Whitney rank sum test. Statistically significant differences were accepted at $P \leq 0.05$.

Results

Infected leaf area and ROS generation

Plant–pathogen interaction is often associated with an increase in the concentrations of reactive oxygen species

and enzymes involved in their scavenging/generation. Inoculation of the plant with pathogens triggered a range of morphological symptoms. Chlorotic and necrotic lesions after *Psp* infiltration and necrosis caused by *Bc* development were observed. However, the infected area after subsequent *Bc* infection was not much larger compared to the single *Psp* variant (Table 1). These changes mainly resulted from the accumulation of reactive oxygen species in plant tissues, especially hydrogen peroxide and superoxide anion, which was confirmed by DAB and NBT staining (Fig. 1). Increase in ROS generation was observed especially in the areas nearest to those infected with *Bc* and *Psp*. Superoxide anion is produced as a result of one-electron reduction of molecular oxygen and can be quite toxic to cell compartments. After *Psp* treatment, the level of O₂^{•−} increased starting from 12th hour to the end of the experiment (Fig. 2). Significantly higher superoxide anion concentration after *Bc* inoculation was only noticeable 6 h after plant treatment. Infection of leaves with both pathogens resulted in significantly increased O₂^{•−} level as early as 6 h after *Bc* inoculation and reached the maximum value (212 % of the control) after 12 h. Significantly higher concentrations of hydrogen peroxide were observed as early as 6 h after *Psp* inoculation, with the largest value 48 h after treatment (281 % of the control) which persisted until 96th h of the experiment (Fig. 3c). *Bc* treatment caused less intensive production of H₂O₂ compared to the *Psp* variant. The results of the combined *Psp*+*Bc* treatment were similar to those for *Psp* alone.

Assessment of enzyme activities

Superoxide dismutase is an enzyme responsible for O₂^{•−} dismutation to H₂O₂. This reactive oxygen form was measured in the whole leaf tissue, although it can be generated both by superoxide dismutase derived from cytosolic and apoplastic fraction. After bacterial treatment, the only significant changes in the apoplastic SOD activity were observed 6, 12 and 24 h after inoculation and in the cytosolic pool of SOD after 60, 72 and 96 h (Fig. 3a, b).

Table 1 Area of infected leaves of *P. vulgaris* 48 h after inoculation with *Bc*, *Psp* or *Psp*+*Bc* analysed 96 h after *Psp* inoculation and 48 h after *Bc* treatment

Variant	Control	<i>Psp</i>	<i>Bc</i>	<i>Psp</i> + <i>Bc</i>
% of infected leaf area	0	7.24 ± 0.61*	6.73 ± 0.96*	8.17 ± 0.85*

Values are means with SD

* Significant differences from the control at $P < 0.05$. No differences between particular treatments were found




Fig. 1 Visualisation of H₂O₂ (b) and O₂⁻ (c) in the leaves of *P. vulgaris* 48 h after inoculation with *Bc* and *Psp*; *Psp+Bc* analysed 96 h after *Psp* inoculation and 48 h after *Bc* treatment; a leaves without staining




Fig. 2 Superoxide anion concentration in the primary leaves of *P. vulgaris* inoculated with *P. syringae* pv. *phaseolicola* (*Psp*) or *B. cinerea* (*Bc*) or with both pathogens sequentially (*Psp+Bc*) compared to the control (C). Hours in parentheses refer to the time after *Bc* inoculation. Values are means with SD; asterisk indicates significant differences from the control at *P* < 0.05

Treating plants with the fungal pathogen caused the maximum increase (twofold) in cytosolic pool of SOD amount 48 h after inoculation. Sequential *Psp+Bc* infection resulted in the significant SOD enhancement after 48 h. Catalase is an enzyme responsible for the removal of high concentrations of H₂O₂. *Psp* inoculation resulted in a slight decrease (6–24 h after inoculation) in the cytosolic activity of this enzyme, compared to the control (Fig. 4a). Statistically significant change in the catalase amount was observed only 6 h after *Bc* treatment. In other variants, no significant changes were observed. Peroxidase is another enzyme considered to be involved in H₂O₂ scavenging. It is supposed that phenolic compounds, such as syringaldazine




Fig. 3 Time course of changes of superoxide dismutase activity in the apoplasmic (a) and cytosolic (b) fraction and hydrogen peroxide concentration (c) in common bean leaves inoculated with *P. syringae* pv. *phaseolicola* (*Psp*) or *B. cinerea* (*Bc*) or with both pathogens sequentially (*Psp+Bc*) compared to the control (C). Hours in parentheses refer to the time after *Bc* inoculation. Values are means with SD; asterisk indicates significant differences from the control at *P* < 0.05

and ferulic acid, take part in cell wall strengthening, and increased activity of peroxidases assayed with these substrates is of particular importance during infection. Increased levels of GPOD in the apoplasmic fraction started 6 h after *Psp* treatment and remained quite high to the end of the experiment, which indicate involvement of the total pool of peroxidase during infection (Fig. 5b). Fungal inoculation caused systematic increase in guaiacol peroxidase with the maximum value after 48 h. Its activities in the *Psp+Bc* variant were very similar to *Psp*. The cytosolic fraction showed a slight increase in GPOD activities from 54th up to 96th h after *Psp* treatment, while *Bc* inoculation caused significant changes starting from 6th h until 48th h




Fig. 4 **a** Time course of changes of catalase activity in the cytosolic fraction of common bean leaves inoculated with *P. syringae* pv. *phaseolicola* (*Psp*) or *B. cinerea* (*Bc*) or with both pathogens sequentially (*Psp+Bc*) compared to the control (C). Hours in parentheses refer to the time after *Bc* inoculation. Values are means with SD; asterisk indicates significant differences from the control at $P < 0.05$. **b** Identification of CAT and POD isoforms from *Phaseolus vulgaris* leaves inoculated with *Psp* and *Bc* on native PAGE and the corresponding densitograms. Electrophoretic patterns for control (C), *Psp* (P) and *Bc* (B) single variants were obtained 48 h after inoculation, while *Psp+Bc* (PB) variant was analysed 96 h after *Psp* inoculation and 48 h after *Bc* treatment

(Fig. 5a). The most marked enhancement of this enzyme was noticed for *Psp+Bc* 48 h after fungal inoculation. High activities of FPOD and SPOD in apoplast were observed 6 h after *Psp* inoculation, and over time decreased to the values comparable to the control in 96th h of experiment (Fig. 6a, b). On the contrary, *Bc* treatment caused gradual increase in these enzymes with the maximum value after 48 h, which was better noticeable in FPOD activity. Peroxidase activity measured with ferulic acid after sequential infection did not differ significantly from the *Psp* variant.




Fig. 5 Time course of changes of guaiacol peroxidase activity in the cytosolic (**a**) and apoplastic (**b**) fraction of common bean leaves inoculated with *P. syringae* pv. *phaseolicola* (*Psp*) or *B. cinerea* (*Bc*) or with both pathogens sequentially (*Psp+Bc*) compared to the control (C). Hours in parentheses refer to the time after *Bc* inoculation. Values are means with SD; asterisk indicates significant differences from the control at $P < 0.05$

Native PAGE

Analysis of SOD isoenzymes on native PAGE showed the presence of 2 Mn-SODs isoforms, 2 Cu/Zn-SODs isoforms and 2 Fe-SODs isoenzymes, of whose one was specific for *Psp* inoculation and persisted also after *Psp+Bc* infection (Fig. 7). CAT patterns on native PAGE showed 1 isoform characteristic of the control plants and *Bc* infection and 1 additional form of this enzyme in both *Psp* and *Psp+Bc* variants (Fig. 4b).

Native PAGE revealed the presence of 3 POD isoforms, of whose an extra POD form appeared after *Psp* inoculation and persisted also in the case of sequential infection with *Bc* (Fig. 4b).

Discussion

Botrytis cinerea is a necrotrophic fungus secreting a number of specific toxins leading to cell death. It was shown that fluorescently labelled *Bc*NEP1 and *Bc*NEP2 toxins derived from *B. cinerea* were associated with plasma membranes and the nuclear envelope, as well as they were present in the nuclei of responding plant cells. These toxins




Fig. 6 Time course of changes of peroxidase with syringaldazine (a) and ferulic acid (b) activity in the apoplasmic fraction of common bean leaves inoculated with *P. syringae* pv. *phaseolicola* (*Psp*) or *B. cinerea* (*Bc*) or with both pathogens sequentially (*Psp+Bc*) compared to the control (C). Hours in parentheses refer to the time after *Bc* inoculation. Values are means with SD; asterisk indicates significant differences from the control at $P < 0.05$

may induce a strong hydrogen peroxide accumulation that was observed in chloroplasts. The cell death may occur as apoptosis, necrosis or intermediate forms (Schouten et al. 2007). A cell wall is a complicated structure that serves both as a constitutive defence and as an active and dynamic barrier in response to signals from pathogens (Huckelhoven 2007). Moreover, hydrogen peroxide as well as peroxidases can participate in strengthening a cell wall (Mayer et al. 2001; Burr and Fry 2009). The changes of peroxidase activity in cell wall surroundings during creation of cross-linking differulate in the wall were probably governed by fluctuations in H_2O_2 and/or presence of its inhibitors (Encina and Fry 2005). The same authors found that cross-linking of [¹⁴C]feruloyl-arabinoxylans also occurred in vitro, in the presence of endogenous peroxidases plus exogenous H_2O_2 . Van Baarlen et al. (2007) confirmed that cell-wall-located ROS production during *Botrytis* sp. colonisation was often accompanied by production of phenolic compounds. *Psp* colonising mainly apoplasmic spaces affects cell walls and specific enzymes bound with them which results in increase in their activity.

Our studies showed that high activity of peroxidase within a cell wall as early as 6 h after *Psp* infection was correlated with great concentration of hydrogen peroxide,




Fig. 7 Identification of different SOD isoforms from *P. vulgaris* leaves inoculated with *Psp* and *Bc* on native PAGE with KCN and H_2O_2 inhibition and the corresponding densitograms. Electrophoretic patterns for control (C), *Psp* (P) and *Bc* (B) single variants were obtained 48 h after inoculation, while *Psp+Bc* (PB) variant was analysed 96 h after *Psp* inoculation and 48 h after *Bc* treatment

which is the major form of ROS taking part in oxidative burst (Wojtaszek 1997). It can be assumed that overproduction of H_2O_2 is used to form diferulic bridges responsible for cell wall strengthening and participates in defence reactions against attack of incompatible pathogens. No significant additional enzymatic activity or ROS generation after sequential infection may suggest that the plant response to the bacterial pathogen was strong enough, mainly due to the fact that it is considered to be resistant to *Psp*. However, sequential infection by *Bc* induced slight increase in SOD activity in the cytosolic fraction 6 h after inoculation which was correlated with increased ROS generation. This may suggest that some mobilisation of resistance enzymes after second infection is possible. It can be assumed that the additional form of Fe-SOD appearing after *Psp* infection and persisting after sequential *Psp+Bc* infection might inhibit *Bc* development through the production of hydrogen peroxide, providing that CAT activity was low. Thus, the suppression of CAT activity contributed to higher generation of toxic H_2O_2 whose greater concentration may inhibit not only incompatible pathogen infection. On the other hand, high CAT activity 6 h after *Bc* infection resulted in reduced amount of hydrogen peroxide

which could increase *Bc* expansion. Such results confirm a combined action of SOD and CAT in response to contact with microorganisms observed by Baptista et al. (2007) and Schwacke and Hager (1992). CAT was also found to be down-regulated at the transcription level during HR (Dorey et al. 1998). It is interesting that the induction of the additional Fe-SOD form, sensitive to high concentrations of H₂O₂, coincided with increased plant resistance (even after sequential infection). This fact may suggest that the produced H₂O₂ participated in the plant–pathogen interaction, therefore, its level was not high enough to inhibit Fe-SOD.

The increased GPOD activity after bacterial infection may suggest more intensive use of overgenerated H₂O₂ in the oxidation of phenolic compounds which can be the inhibitors of both bacteria and fungi growth. Their ability to inhibit the activity of enzymes, i.e. those derived from pathogens, and thereby to suppress growth of pathogens has been repeatedly confirmed (Mellersh et al. 2002; Lebeda and Sedlarova 2003; Sedlářová et al. 2004). In our work, the GPOD activity measured in the apoplastic fraction increased after both *Psp* and sequential infection with both pathogens. It proves that the most important processes during plant–pathogen interaction occur around a cell wall, which is the first barrier of cell defence. *Psp+Bc* infection resulted in increased GPOD activity, comparable to that after single *Psp* infection. Such results showed that after sequential infection with *Bc*, there was no additional increase in the activity above that obtained after *Psp* application.

The results obtained for the SPOD and FPOD activities seem to be of special interest. Peroxidase activity measured with syringaldazine as a substrate is assumed to be involved in cell wall lignification. Our results showed its increased activity after *Psp* treatment but subsequent *Bc* inoculation (*Psp+Bc*) did not change SPOD level in comparison to the *Psp* variant. Only *Bc* infection caused an increased SPOD activity at the end of the experiment.

Since pathogenic bacteria and fungi exhibit various toxicity towards plants they may be expected to induce different kinds of response, e.g. necrotrophs produce extracellular enzymes damaging plant cell walls and inducing apoptotic-like cell death (Shlezinger et al. 2011). Therefore, it was important to determine FPOD activity in the apoplastic fraction. It was found out that its activity associated with the formation of additional diferulic bonds by apoplastic peroxidase increased after *Psp* infection since the beginning of the experiment and even stronger than that of SPOD after *Bc* inoculation. Sequential *Psp+Bc* infection did not cause increase in the FPOD activity above the levels obtained after single *Psp* or *Bc* inoculation.

Plant response to bacterial and fungal infection can differ remarkably and depends on a broad spectrum of reactions such as interactions between different plant

hormones regulating defence gene expression and pathogen resistance (Thomma et al. 1998). Enhanced generation of reactive oxygen species is characteristic of reactions to both biotrophic and necrotrophic pathogens. These molecules not only participate in intercellular signalling, but locally at excessive concentration they can be also toxic to pathogens (Mehdy 1994, Mayer et al. 2001). Stress related to infection induces alterations in plant gene expression patterns mainly involved in redeployment of cell functions from metabolic processes to defence reactions (Takahashi et al. 2004). Finiti et al. (2013) demonstrated that transgenic tomato plants, having greater susceptibility to *P. syringae*, exhibited increased expression of the marker gene of jasmonic acid pathway after infection. They showed that endo-1,4- β -glucanases were differentially expressed upon many different plant–pathogen challenges, hormone treatments and many abiotic stresses. They demonstrated that the reduced activity of this enzyme could alter the impact of plant pathogens and could contribute to the respective signalling and resulted in a variety of plant responses to biotic and abiotic stresses.

Recent studies on plant resistance mechanisms have confirmed the presence of genes responsible for the response to both biotic and abiotic factors (AbuQamar et al. 2013). Mengiste et al. (2003) in the research on *A. thaliana* demonstrated the presence of *BOS1* gene activated by *B. cinerea* infection. *BOS1* gene expression resulted in inhibition of infection not only by *B. cinerea*, but also by another necrotroph *A. brassicicola*, suggesting a similar defensive strategy against these pathogens. *bos1* mutants with greater susceptibility to *B. cinerea* exhibited also increased sensitivity to oxidative stress caused by abiotic factors. Furthermore, *BOS1* expression was blocked in plants with *coil* mutation which causes insensitivity to jasmonic acid and higher sensitivity to *Botrytis*. It has been suggested that the protein encoded by *BOS1* (transcript R2R3MYB protein) together with JA is responsible for the activation of a signal cascade of reactive oxygen intermediates (ROI).

Studies on tomato plants (*Solanum lycopersicum*) proved the presence of *SIAMI1* gene whose expression is regulated by abscisic acid (ABA) (AbuQamar et al. 2009). Plants with reduced expression of this gene were characterised by increased susceptibility to both *B. cinerea* infection and salt stress. The authors suggested a correlation between ABA, the ability to maintain Na⁺ homeostasis, oxidative stress and response to pathogens.

AbuQamar et al. (2013) also demonstrated that similarly to the results obtained for *A. thaliana*, *EXLA2* gene encoding expansin, involved in cell wall modification, played an important role during *B. cinerea* infection. The authors showed that in *exla2* mutants with decreased susceptibility to *B. cinerea* sensitivity to infection with *P. syringae* pv. *tomato* was not changed. Moreover, *EXLA2*

gene activation was induced by abiotic factors such as salinity, cold or exogenous ABA application. It has been suggested that lack or deregulation of *EXLA2* gene expression may be of importance for increasing resistance to *B. cinerea*. As in the previous study, they demonstrated that *exla2* mutants were hypersensitive to cold and salinity for which the response is regulated by a metabolic pathway of ABA.

It seems that in the case of our study on bean, the presence of a specific gene activated upon the infection with *Psp*, whose expression might down-regulate the early development of the second infection with *Bc*, cannot be excluded at least at the beginning of this process. The area of lesions 48 h after subsequent *Bc* inoculation was smaller than in the case of a single primary infection with this necrotroph. Certainly, ROS generation initiated by *Psp* inoculation may be involved in further resistance to subsequent pathogen attack.

Conclusion

Our results suggest, that *P. vulgaris* cv. Korona, resistant to *Psp*, after inoculation with this pathogen mobilised the immune system to such an extent that after contact with the necrotrophic pathogen only H₂O₂ and SOD levels, crucial for further resistance, changed. This may indicate that the induction of enzymatic response after *Psp* allowed to delay growth of the necrotrophic pathogen. Increased FPOD and SPOD activities were observed both after *Psp* and *Bc* infection which may indicate their important role in strengthening plant cell walls during different kinds of infections. However, increased generation of H₂O₂ and higher SOD activity as well as additional Fe-SOD form after *Psp* inoculation with slight increase in these parameters after subsequent *Bc* infection may confirm the specificity of the response to *Psp* which confers increased resistance to *Bc* at the early stages of infection.

Author contribution Agata Nowogórska was responsible for carrying out the experiment and working out the results and Jacek Patykowski was responsible for the idea and coordination of the experiment.

Open Access This article is distributed under the terms of the Creative Commons Attribution License which permits any use, distribution, and reproduction in any medium, provided the original author(s) and the source are credited.

References

AbuQamar S, Luo H, Laluk K, Mickelbart MV, Mengiste T (2009) Crosstalk between biotic and abiotic stress responses in tomato is mediated by the AIM1 transcription factor. *Plant J* 58:347–360

- AbuQamar S, Ajeb S, Sham A, Enan MR, Iratni R (2013) A mutation in the *expansin-like A2* gene enhances resistance to necrotrophic fungi and hypersensitivity to abiotic stress in *Arabidopsis thaliana*. *Mol Plant Pathol* 14(8):813–827
- Adie BA, Pérez-Pérez J, Mm Pérez-Pérez, Godoy M, Sánchez-Serrano JJ, Schmelz EA, Solano R (2007) ABA is essential signal for plant resistance to pathogens affecting JA biosynthesis and the activation of defences in *Arabidopsis*. *Plant Cell* 19:1665–1681
- Alfano JR, Collmer A (1996) Bacterial pathogens in plants: life up against the wall. *Plant Cell* 8:1683–1698
- Baptista P, Martins A, Pais MS, Tavares RM, Lino-Neto T (2007) Involvement of reactive oxygen species during early stages of ectomycorrhiza establishment between *Castanea sativa* and *Pisolithus tinctorius*. *Mycorrhiza* 17:185–193
- Beauchamp C, Fridovich I (1971) Superoxide dismutase: improved assays and an assay applicable to acrylamide gels. *Anal Biochem* 44:276–287
- Beckers GJM, Spoel SH (2005) Fine-tuning plant defence signalling: salicylate versus jasmonate. *Plant Biol* 8:1–10
- Bolwell PP, Page A, Piślewska M, Wojtaszek P (2001) Pathogenic infection and the oxidative defences in plant apoplast. *Protoplasma* 217:20–32
- Bowler C, Flur R (2000) The role of calcium and activated oxygens as signals for controlling cross-tolerance. *Trends Plant Sci* 5:241–246
- Bradford MM (1976) A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal Biochem* 72:248–254
- Burr SJ, Fry SF (2009) Extracellular cross-linking of maize arabinoxylans by oxidation of feruloyl esters to form oligoferuloyl esters and ether-like bonds. *Plant J* 58:554–567
- Capaldi DJ, Taylor KE (1983) A new peroxidase colour reaction: oxidative coupling of 3-methyl-2-benzothiazolinone hydrazone (MBTH) with its formaldehyde azine application to glucose and choline oxidases. *Anal Biochem* 129:329–336
- Creissen G, Firmin J, Freyer M, Kular B, Leyland N, Reynold H, Pastori G, Wellburn F, Baker N, Wellburn A, Mullineaux P (1999) Elevated glutathione biosynthetic capacity in the chloroplasts of transgenic tobacco paradoxically caused increased oxidative stress. *Plant Cell* 11:1277–1291
- Derksen H, Rampitsch C, Daayf F (2013) Signaling cross-talk in plant disease resistance. *Plant Sci* 207:79–87
- Dhindsa RS, Plumb-Dhindsa P, Thorpe TA (1981) Leaf senescence correlated with increased levels of membrane permeability and lipid peroxidation, and decreased levels of superoxide dismutase and catalase. *J Exp Bot* 32:93–101
- Doke N (1983) Involvement of superoxide anion generation in the hypersensitive response of potato tuber tissues to infection with an incompatible race of *Phytophthora infestans* and to the hyphal wall components. *Physiol Plant Pathol* 23:345–357
- Dorey S, Bailleul F, Saindrenan P, Fritig B, Kauffmann S (1998) Tobacco class I and II catalases are differentially expressed during elicitor induced hypersensitive cell death and localized acquired resistance. *Mol Plant Microb Int* 11:1102–1109
- Encina A, Fry SC (2005) Oxidative coupling of a feruloylarabinoxylan trisaccharide (FAXX) in the walls of living maize cells requires endogenous hydrogen peroxide and is controlled by a low-Mr apoplastic inhibitor. *Planta* 223:77–89
- Finiti I, Leyva MO, López-Cruz J, Calderan Rodrigues B, Vicedo B, Angulo C, Bennett AB, Grant M, García-Agustín P, González-Bosch C (2013) Functional analysis of endo-1,4- β -glucanases in response to *Botrytis cinerea* and *Pseudomonas syringae* reveals their involvement in plant–pathogen interactions. *Plant Biol* 15:819–831

- Flor HH (1971) Current status of the gene-for-gene concept. *Annu Rev Phytopathol* 9:275–296
- García-Arenal F, Sagasta EM (1980) Scanning electron microscopy of *Botrytis cinerea* penetration of bean (*Phaseolus vulgaris*) hypocotyls. *Phytopathol Z* 99:37–42
- Glazebrook J (2005) Contrasting mechanisms of defense against biotrophic and necrotrophic pathogens. *Annu Rev Phytopathol* 43:205–227
- Goldberg R, Cateson AM, Czaninski Y (1983) Some properties of syringaldazine oxidase, a peroxidase specifically involved in the lignification processes. *Z Pflanzenphysiol* 110:267–279
- Huckelhoven R (2007) Cell wall-associated mechanisms of disease resistance and susceptibility. *Annual Rev Phytopathol* 45:101–127
- Imberty A, Goldberg R, Cateson AM (1985) Isolation and characterization of *Populus* isoperoxidases involved in the last step of lignification. *Planta* 164:221–226
- Koornneef A, Pieterse CMJ (2008) Cross talk in defense signalling. *Plant Physiol* 146:839–844
- Laemmli UK (1970) Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* 227:680–685
- Laluk K, AbuQamar S, Mengiste T (2011) The Arabidopsis mitochondrial localized pentatricopeptide repeat protein PGN functions in defense against necrotrophic fungi and abiotic stress tolerance. *Plant Physiol* 156:2053–2068
- Lamb C, Dixon RA (1997) The oxidative burst in plant disease resistance. *Annu Rev Plant Physiol Plant Mol Biol* 48:251–275
- Lebeda A, Sedlarova M (2003) Cellular mechanisms involved in the expression of specificity in *Lactuca* spp.–*Bremia lactucae* interactions. In: Hintum ThJL, Van-Lebeda A, Pink DAC, Schut JW (eds) *Eucarpia leafy vegetables conference 2003*. CGN, Wageningen, The Netherlands, pp 55–60
- Maehly AC, Chance B (1954) The assay of catalases and peroxidases. In: Glick D (ed) *Methods of biochemical analysis*. Wiley, New York, pp 357–425
- Mayer AM, Staples RC, Gilad NL (2001) Mechanisms of survival of necrotrophic fungal plant pathogens in hosts expressing the hypersensitive response. *Phytochemistry* 58(1):33–41
- Mehdy MC (1994) Active oxygen species in plant defense against pathogens. *Plant Physiol* 105(2):467–472
- Mellersh DG, Foulds IV, Higgins VJ, Heath MC (2002) H₂O₂ plays different roles in determining penetration failure in three diverse plant–fungal interactions. *Plant J* 29:257–268
- Mengiste T, Chen X, Salmeron J, Dietrich R (2003) The *BOTRYTIS SUSCEPTIBLE1* gene encodes an R2R3MYB transcription factor protein that is required for biotic and abiotic stress responses in Arabidopsis. *Plant Cell* 15:2551–2565
- Mika A, Boenisch MJ, Hopff D, Lüthje S (2010) Membrane-bound guaiacol peroxidases from maize (*Zea mays* L.) roots are regulated by methyl jasmonate, salicylic acid, and pathogen elicitors. *J Exp Bot* 61(3):831–841
- Oudgenoeg G, Dirksen E, Ingemann S, Hilhorst R, Gruppen H, Boeriu CG, Piersma SR, van Berkel WJH, Laane C, Voragen AGJ (2002) Horseradish peroxidase-catalyzed oligomerization of ferulic acid on a template of a tyrosine-containing tripeptide. *J Biol Chem* 24(277):21332–21340
- Passardi F, Cosio C, Penel C, Dunand C (2005) Peroxidases have more functions than a Swiss army knife. *Plant Cell Rep* 24:255–265
- Patkowski J (2008) Rola apoplastu w interakcji rośliny—patogen po infekcji pomidora przez *Botrytis cinerea*. Wydawnictwo Uniwersytetu Łódzkiego, Łódź
- Polle A, Otter T, Seifert F (1994) Apoplastic peroxidases and lignification in needles of Norway Spruce (*Picea abies*). *Plant Physiol* 106:53–60
- Romero-Puertas MC, Rodríguez-Serrano M, Corpas FJ, Gómez M, Del Río LA, Sandalio LM (2004) Cadmium-induced subcellular accumulation of O₂^{•−} and H₂O₂ in pea leaves. *Plant Cell Environ* 27:1122–1134
- Sakamoto A, Okumura T, Kaminaka H, Sumi K, Tanaka K (1995) Structure and differential response to abscisic acid of two promoters for the cytosolic copper/zinc-superoxide dismutase genes, *SodCc1* and *SodCc2*, in rice protoplasts. *FEBS Lett* 358:62–66
- Schouten A, van Baarlen P, van Ka JA (2007) Phytotoxic Nep1-like proteins from the necrotrophic fungus *Botrytis cinerea* associate with membranes and the nucleus of plant cell. *New Phytol* 177:493–505
- Schwacke R, Hager A (1992) Fungal elicitors induce a transient release of active oxygen species from cultured spruce cells that is dependent on Ca²⁺ and protein-kinase activity. *Planta* 187:136–141
- Sedlářová M, Lebeda A, Luhová L (2004) The role of active oxygen species in resistance of *Lactuca* spp. to *Bremia lactucae*. *Acta Fytotech Zootech* 7:272–274
- Shi J, Fu XZ, Peng T, Huang XS, Fan QJ, Liu JH (2010) Spermine pretreatment confers dehydratase tolerance of citrus in vitro plants via modulation of antioxidative capacity and stomatal response. *Tree Physiol* 30:914–922
- Shlezniger N, Minz A, Gur Y, Hatam I, Dagdas YF, Talbot NJ, Sharo A (2011) Anti-apoptotic machinery protects the necrotrophic fungus *Botrytis cinerea* from host-induced apoptotic like cell death during plant infection. *PLoS Pathog* 7(8):e100218
- Takahama U (1995) Oxidation of hydroxycinnamic acid and hydroxycinnamyl alcohol derivatives by laccase and peroxidase. Interactions among *p*-hydroxyphenyl, guaiacyl and syringyl groups during the oxidation reactions. *Physiol Plant* 93:61–68
- Takahashi S, Seki M, Ishida J, Satou M, Sakurai T, Narusaka M, Kamiya A, Nakajima M, Enju A, Akimaya K, Yamaguchi-Shinozaki K, Shinozaki K (2004) Monitoring the expression profiles of genes induced by hyperosmotic, high salinity, and oxidative stress and abscisic acid treatment in Arabidopsis cell culture using a full length cDNA microarray. *Plant Mol Biol* 56:29–55
- Thaler JS, Owen B, Hoggins VJ (2004) The role of the jasmonate response in plant susceptibility to diverse pathogens with a range of lifestyles. *Plant Physiol* 135:1–9
- Thomma BPHJ, Eggermont K, Penninckx IAMA, Mauch-Mani B, Vogelsang R, Cammue BPA, Broekaert WF (1998) Separate jasmonate-dependent and salicylate-dependent defense-response pathways in Arabidopsis are essential for resistance to distinct microbial pathogens. *Proc Natl Acad Sci USA* 95:15107–15111
- Thordal-Christensen H, Zhang Z, Wei Y, Collinge DB (1997) Subcellular localization of H₂O₂ in plants. H₂O₂ accumulation in papillae and hypersensitive response during the barley–powdery mildew interaction. *Plant J* 11:1187–1194
- Van Baarlen P, Woltering EJ, Staats M, van Kan JAL (2007) Histochemical and genetic analysis of host and non-host interactions of Arabidopsis with three *Botrytis* species: an important role for cell death control. *Mol Plant Pathol* 8(1):41–54
- Welinder KG, Justesen AF, Kjaersgard IV, Jensen RB, Rasmussen SK, Jespersen HM, Duroux L (2002) Structural diversity and transcription of class III peroxidases from Arabidopsis thaliana. *Eur J Biochem* 269:6063–6081
- Williamson B, Tudzynski B, Tudzynski P, van Kan JAL (2007) *Botrytis cinerea*: the cause of grey mould disease. *Mol Plant Pathol* 8:561–580
- Wojtaszek P (1997) Oxidative burst: an early plant response to pathogen infection. *Biochem J* 322:681–692