

Collection: 3rd International Elm Conference, Florence (Italy - 2013)
 “The elms after 100 years of Dutch Elm disease”

Guest Editors: A. Santini, L. Ghelardini, E. Collin, A. Solla, J. Brunet, M. Faccoli, A. Scala, S. De Vries, J. Buiteveld

Seven *Ulmus minor* clones tolerant to *Ophiostoma novo-ulmi* registered as forest reproductive material in Spain

Juan Antonio Martín ⁽¹⁾, Alejandro Solla ⁽²⁾, Martín Venturas ⁽¹⁾, Carmen Collada ⁽¹⁾, Jorge Domínguez ⁽¹⁾, Eva Miranda ⁽¹⁾, Pablo Fuentes ⁽³⁾, Margarita Burón ⁽¹⁾, Salustiano Iglesias ⁽⁴⁾, Luis Gil ⁽¹⁾

□ (1) ETSI Montes, Universidad Politécnica de Madrid, Ciudad Universitaria s/n, E-28040 Madrid (Spain); (2) Ingeniería Forestal y del Medio Natural, Universidad de Extremadura, Avenida Virgen del Puerto 2, E-10600 Plasencia (Spain); (3) Institute of Evolutionary Biology, The University of Edinburgh, West Mains Rd., Edinburgh EH9 9JT (United Kingdom); (4) Dirección General de Desarrollo Rural y Política Forestal, Ministerio de Medio Ambiente y Medio Rural y Marino, c/ Ríos Rosas 24, E-28003 Madrid (Spain)

@ Luis Gil (luis.gil@upm.es)

Received: Dec 30, 2013 - Accepted: May 21, 2014

Citation: Martín JA, Solla A, Venturas M, Collada C, Domínguez J, Miranda E, Fuentes P, Burón M, Iglesias S, Gil L, 2015. Seven *Ulmus minor* clones tolerant to *Ophiostoma novo-ulmi* registered as forest reproductive material in Spain. *iForest* 8: 172-180 [online 2014-08-13] URL: <http://www.sisef.it/forest/contents/?id=ifor1224-008>

Communicated by: Alberto Santini

Asian elms, including *Ulmus pumila*, *U. chenmoui*, *U. davidiana* var. *japonica* and *U. wallichiana*, have been the main sources of resistance in the Dutch, American and Italian elm breeding programmes (Heybroek 1993, Smalley & Guries 2000, Santini et al. 2011, Buiteveld et al. 2014). As a result of crossing these species with native elms, a wide range of hybrid clones of varying tolerance levels and genetic backgrounds is now available on the market. The Spanish programme took advantage of the knowledge, methodologies and plant materials previously developed by the Dutch and Italian programmes. In the first 14 years, *U. pumila* was used as the main source of resistance, giving rise to 10 crossings tolerant to *O. novo-ulmi* (Solla et al. 2000). The tolerance of these crossings was tested in clone replicates ($N > 16$) over several years at various locations, and clone adaptation to different environments in Spain was evaluated. Five crossings with Asian background were recently selected to be released onto the market for ornamental use.

In the 1990s the Spanish programme included some native elms, mainly *U. minor*, in the *O. novo-ulmi* susceptibility trials. In the following decade the programme focused mainly on selecting native elms. This new strategy complied with European and Spanish legislation governing the quality and genetic background of forest reproductive materials for production and marketing. In the European Union, forest reproductive materials are governed by Council Directive

The Spanish elm programme began in 1986 in response to the devastating impact of Dutch elm disease on natural elm stands and urban trees. Its main objectives were to conserve remaining genetic resources and select and breed tolerant native elm genotypes. After 27 years of work conducting susceptibility trials on thousands of elm genotypes, the first seven tolerant *Ulmus minor* trees are now being registered by the Spanish Environmental Administration. This paper presents the results of the susceptibility tests on these clones and their distinctive genetic, morphological and phenological features. In all susceptibility trials the commercial “Sapporo Autumn Gold” clone, which is highly tolerant to *O. novo-ulmi*, was used as a control. The registered clones were named “Ademuz”, “Dehesa de la Villa”, “Majadahonda”, “Toledo”, “Dehesa de Amaniel”, “Retiro” and “Fuente Umbria”. The most tolerant clone was “Dehesa de Amaniel”, as its wilting values were below 5% during the two consecutive inoculation trials performed in Madrid. “Fuente Umbria”, tested over four consecutive years in Guadalajara and Palencia, was the Spanish clone with the most reliable tolerance level to *O. novo-ulmi*. The “Ademuz” and “Majadahonda” clones had the highest ornamental scores and are promising trees for use in urban environments and tree breeding for ornamental quality. These two genotypes showed a later bud burst phenology than the other *U. minor* clones, demonstrating suitability to areas with late frost events. The Spanish programme aims to substantially increase the range of tolerant native elms through new selections and crossings to gain a better understanding of the genetic basis of resistance.

Keywords: Dutch Elm Disease, Breeding, Plant Release, Resistance, Invasive Species

Introduction

In the first half of the 20th century, the first Dutch elm disease (DED) pandemic caused a massive loss of elms in Europe and North America. The much more aggressive *O. novo-ulmi* Brasier took the place of the causal agent, *Ophiostoma ulmi* (Buisman) Nannf., in the second half of the century. The second pathogen has caused the disappearance of adult elms in many European and North American locations (Brasier & Kirk 2010). *O. novo-ulmi* is almost impossible to control through chemical, biological or silvicultural methods due to its high virulence and highly effective transmission via small beetles of the *Scolytus* and *Hylorgopinus* genera (Weber 2000). Tolerant elm genotype selection and breeding has been the most successful strategy for elm recovery, particularly in ur-

ban environments (Santini et al. 2004, 2011, Solla et al. 2005a, 2014). The Spanish elm breeding and conservation programme began in 1986 as the result of an agreement between the Spanish Environmental Administration and the Technical University of Madrid School of Forestry Engineering. Its two main objectives were to conserve remaining elm genetic resources and to transmit their variability to future generations of tolerant elms obtained through breeding; i.e., hybridisation of selected progenitors (native or tolerant Asian elms) to obtain tolerant trees with the appearance of native elm species.

The first elm breeding programme began in the Netherlands in 1928 (Heybroek 1993) and was followed by several programmes in the United States and various European countries (Mittempergher & Santini 2004).

1999/105/CE, and Annex I of the directive lists the permitted forest species. Although *Ulmus* species are not included in Annex I, Article 3.2 of the directive allows particular Member States to add to the list. In Spain, several regions expressed the need to certify the plant origin and genetic quality of some forest species not included in Annex I that are traditionally used in reforestation programmes. As a result, Annex XII of Spanish Royal Decree-Law 289/2003 listed 24 additional forest species, including *U. glabra* and *U. minor* (Iglesias 2005). The major spread of *U. pumila* in Spain and its extensive hybridisation with the native *U. minor* (Cogoludo-Agustín et al. 2000) led to conservation concerns for the native species. To preserve the genetic integrity of the Spanish elms, artificial hybrids in the genus *Ulmus* were not included in Annex XII of Royal Decree-Law 289/2003. This means that hybrids with Asian background cannot be marketed for forest use in Spain, although they can be used for urban planting.

Progress in selecting elms was slow due to the long periods required to propagate trees and evaluate their tolerance with a scientifically sound base, as plant material needs to be at least four years old (Solla et al. 2005c). In the case of selecting pure *U. minor* material, a further difficulty was the very limited number of native elms exhibiting some degree of tolerance to *O. novo-ulmi*, which was around 0.5% in comparison to 2-5% for hybrids with Asian background (unpublished results). Fortunately, susceptibility trials performed in the last 10 years provided some native individuals with low leaf wilting values. After 27 years of activity, the Spanish programme continues to breed and conserve Spanish elms with the ultimate goal of recovering their forest and ornamental uses.

Native elms can be registered by the Environmental Administration as “qualified forest reproductive material” when they show

low (0-30%) crown wilting or symptoms similar to the tolerant “Sapporo Autumn Gold” clone after two consecutive years of artificial inoculation with *O. novo-ulmi*. At least six replicates of the tested clone must be inoculated. Replicates are grown in a plot in which a susceptible control clone has to exhibit more than 70% wilting symptoms. When a tested clone is registered, it can be propagated, marketed and used for forest purposes. The “qualified” category is provisional and after 10 years it becomes “controlled material” and acquires a permanent category (Iglesias 2005). Before clones are registered as “controlled material”, they must meet the same requirements as “qualified material” when tested at a second location. This paper reports the selection and the features of the first *U. minor* clones registered for forest use in Spain. The potential use of these clones and the future of the Spanish breeding programme are discussed.

Material and methods

Plant material

From 1990 to 2002, plant material was propagated from trees selected during surveys of adult elms in natural forests, rural areas, parks, and other urban environments in Spain (Tab. 1). The main selection criterion was good sanitary status, i.e., putative tolerance if trees had survived in a DED affected area. Trees were propagated using seeds, root cuttings and grafts (Tab. 1). In the case of seed propagation, seedlings selected for their tolerance to *O. novo-ulmi* were propagated by hardwood cuttings and at least six ramets per seedling were obtained. This procedure was used for the “Dehesa de Amaniel”, “Retiro”, “Toledo”, and “Fuente Umbría” clones.

Ramets of the seven clones were planted with 157 other elm clones in five different inoculation plots in Spain (Tab. 1), under a

Mediterranean phytoclimate (Allué-Andrade 1990). All clones except “Fuente Umbría” were planted in Puerta de Hierro Forest Breeding Center, Madrid (40° 27' 24" N; 3° 45' 0" W; 600 m a.s.l.). This location has an average annual rainfall of 397 mm and an average annual temperature of 14°C. “Fuente Umbría” ramets were planted in El Serranillo Forest Breeding Center (Guadalajara - 40° 40' 13" N; 3° 9' 39" W; 685 m a.s.l.; 457 mm, 13.5 °C) and Calabazanos Forest Health Center (Palencia - 41° 57' 7" N; 4° 30' 60" W; 739 m a.s.l.; 412 mm, 11.7 °C).

Plots were designed in two blocks, with random experimental units of three to four ramets per block. Spacing in Puerta de Hierro and El Serranillo was 0.5 to 1 m between plants and 1 to 1.5 m between rows. In Calabazanos spacing was 5 × 5 m. To avoid side effects, a tree border line surrounded all plots. Plants were watered in spring and summer to ensure growth. Their main stems were fastened to supports to avoid wind shake. “Sapporo Autumn Gold”, highly tolerant to *O. novo-ulmi* (Smalley & Lester 1973), and UPM089, a Spanish *U. minor* clone classified by the Spanish elm breeding programme as very susceptible to *O. novo-ulmi*, were used as control clones. At least six replicates of each control were included in each inoculation plot.

Inoculations

Local strains of *O. novo-ulmi* were used to evaluate the tolerance level of the clones (Tab. 1). Strains NA-PE, CU-HU and Z-BU1 were isolated from DED-infected trees in Peralta (Navarra), Huelves (Cuenca) and Bubierca (Zaragoza), respectively. Strains NA-PE and CU-HU were isolated in 2002 and Z-BU1 was isolated in 2009. Inoculations were performed at the end of April or beginning of May, depending on the plant phenological stage, about 15-30 days after full leaf development. This is the time *U. mi-*

Tab. 1 - Plant material specifications. (a) R: root cutting; G: graft; S: seed; (b) numbers in brackets indicate the year of inoculation.

Clone	Origin in Spain	Initial propagation				Inoculation test			
		Type ^a	Year	Plot	N	Years	Location in Spain	<i>O. novo-ulmi</i> ssp. <i>americana</i> strain ^b	
Ademuz	Valencia 40° 4' 52" N, 1° 16' 55" W	R	1996	XXIV	10	2008, 2009	Puerta de Hierro	NA-PE (2008), CU-HU (2009)	
Dehesa de la Villa	Madrid 40° 27' 29" N, 3° 44' 00" W	R	1990	XXV	10	2009, 2010	Puerta de Hierro	CU-HU	
Majadahonda	Madrid 40° 28' 90" N, 3° 52' 19" W	G	1993	XXIV	6	2008, 2009	Puerta de Hierro	NA-PE (2008), CU-HU (2009)	
Toledo	Toledo 39° 51' 51" N, 4° 1' 30" W	S	1999	XXX	7	2011, 2012	Puerta de Hierro	Z-BU1	
Dehesa de Amaniel	Madrid 40° 27' 37" N, 3° 43' 17" W	S	1999	XXX	12	2011, 2012	Puerta de Hierro	Z-BU1	
Retiro	Madrid 40° 24' 56" N, 3° 41' 10" W	S	2002	XXX	7	2011, 2012	Puerta de Hierro	Z-BU1	
Fuente Umbría	Valencia 39° 25' 23" N, 0° 56' 46" W	S	1995	V and A	>10	2010-2013	El Serranillo and Calabazanos	CU-HU (2010), Z-BU1 (2011-2013)	

nor takes to reach its susceptibility peak to *O. novo-ulmi* in Madrid (Solla et al. 2005b).

A bud-cell suspension of the pathogen was prepared by adding 2 × 2 mm plugs from the edge of 7-day-old cultures on malt extract agar to 50 ml Tchernoff's liquid medium (Tchernoff 1965) in sterile Erlenmeyer flasks, followed by shaking in the dark for four days at room temperature. Spore suspensions were centrifuged at 50 × g for 20 min to eliminate the medium and then suspended in sterile distilled water. Pathogen inoculation was performed by inserting 0.1 ml of the spore suspension at 10⁶ spores ml⁻¹ into an incision made in the trunk base with a razor blade, allowing the suspension to be absorbed by the sap flow. The elms were at least four years old and 1.5 m in height, to obtain maximum disease symptoms (Solla et al. 2005c). Disease development was evaluated by three independent observers who recorded the percentage of wilting leaves in the crown at 30, 60 and 120 days post inoculation (dpi).

Molecular characterization

In the marketing of forest reproductive material, characterization and traceability of trees are of pivotal importance. Various techniques using molecular markers are efficient tools for this purpose. Genetic characterization of the seven *U. minor* clones was performed at two levels. Trees were analyzed firstly with chloroplast DNA markers to determine the lineages of the individuals selected, and secondly with nuclear DNA markers to quantify genotypic diversity (Gil et al. 2004).

For the lineage study, two chloroplast markers were used. One corresponds to the chloroplast fragment SFm and was developed from the sequence of the SFm fragment to differentiate the *U. minor* lineages in Spain (Collada et al., unpublished data). The other marker corresponds to microsatellite *cmp2*, which was developed for tobacco (Weising & Gardner 1999) and transferred to *U. minor* to enable differentiation of variants within lineages.

For the genetic description, 12 nuclear microsatellites were selected. Four of these were described in *U. minor* (Ulmi1-98, Ulmi1-165, Ulmi2-16 and Ulmi2-20 - Collada et al. 2004), three were transferred from *U. laevis* (Ulm2, Ulm3 and Ulm8 - Whiteley et al. 2003) and five were transferred from *U. rubra* (UR 123, UR 141, UR 153, UR 158 and UR 159 - Zalapa et al. 2008).

Leaves from at least two individuals for each selected clone were collected, labeled and stored in silica gel. After DNA extraction, 2.5 ng µl⁻¹ dilutions were used in amplification reactions conducted following the literature (Weising & Gardner 1999, Collada et al. 2004, Whiteley et al. 2003, Zalapa et al. 2008).

Tree morphology and phenology assessments

Clones were morphologically described following specific literature (Richens 1955, Jeffers & Richens 1970, Ipinza 1990). Quantitative data on subdistal leaves of new shoots were measured on four leaves per tree. The parameters measured are shown in Fig. 1. The number of nerve pairs, total number of main teeth per leaf, and type of leaf margin serration (simple, double or triple) were also determined.

Height growth of each clone was assessed in Puerta de Hierro Forest Breeding Center (Madrid), as well as ornamental qualities of trees such as growth habit and branching (erect, spreading, or pendulous), leaf density (abundance of leaves per crown volume, estimated as high, medium or low), crown shape (conical, spindle, globular or irregular) and leaf size. The ornamental value of each clone was quantified on a scale from 1 to 5, where 5 corresponded to the most frequent features of Spanish *U. minor* according to the clone collection (N = 363) held at Puerta de Hierro Forest Breeding Center (*i.e.*, erect branching, globular-spindle crown, medium-high leaf density, leaf size of about 50 mm length and 30 mm width) and 1 corresponded to unusual *U. minor* features. The presence of corky tissue was also recorded but not considered for ornamental evaluation. After four independent observers

had assessed ornamental quality, the average value was calculated.

The unfolding of elm leaf buds was characterized in 2011 using the methodology described by Santini et al. (2005). Leaf phenology is divided into five stages from bud formation to complete leaf expansion: phase 1: dormant buds; phase 2: swelling buds but with closed flakes; phase 3: flakes open and the first leaf ends are visible in the apex of the buds; phase 4: the ends of all the leaves are visible but the leaves are not expanded; Phase 5: two or more leaves are fully expanded. To compare leaf phenology between genotypes, these stages were grouped into three classes: dormancy (phases 1 and 2), bud break (phases 3 and 4), and leaf expanded (phase 5).

Statistical analysis

For each inoculation plot and year, wilting percentages at 30, 60 and 120 dpi were analyzed using repeated measures ANOVA, considering time since inoculation, block, and genotype as the main factors and tree height as a covariate. Fisher's least significant difference (LSD) test was applied to compare average wilting values (least square means of wilting percentages at 30, 60 and 120 dpi) between clones ($P < 0.05$). Analyses were performed using the STATISTICA v. 7.0 package (StatSoft Inc., Tulsa, OK, USA).




Fig. 1 - Foliar parameters measured to describe the *Ulmus minor* clones. (MW): maximum foliar width; (ML): maximum foliar length; (BA): basal asymmetry; (PL): petiole length; (TD): tooth depth; (TL): tooth length; (TB): tooth breadth.

Tab. 2 - Results (p-values) of repeated measures ANOVA of the wilting values shown by elm trees at 30, 60 and 120 dpi (repeated variable) considering time since inoculation, genotype, block, and genotype x block interaction as factors, and plant height as a covariate. (a): The plot had one block.

Inoculation year	Plot	Source of variation				Plant height
		Time dpi	Genotype (G)	Block (B)	G × B	
2008	XXIV	< 0.001	< 0.001	< 0.001	0.001	0.708
2009	XXIV	0.15	< 0.001	0.029	0.65	0.928
2009	XXV	0.002	< 0.001	0.032	< 0.001	0.078
2010	XXV	< 0.001	< 0.001	0.064	< 0.001	0.513
2010	V	< 0.001	< 0.001	0.163	0.751	0.951
2011	V	< 0.001	< 0.001	0.185	0.572	0.839
2011	XXX	< 0.001	< 0.001	0.001	< 0.001	< 0.001
2012	XXX	< 0.001	< 0.001	0.068	0.01	0.91
2012	A ^a	< 0.001	< 0.001	-	-	0.623
2013	A ^a	0.001	< 0.001	-	-	0.086

Results and Discussion

Seven *U. minor* clones were selected for their tolerance to *O. novo-ulmi* in various susceptibility tests conducted in Spain. The results of the repeated measures ANOVA of leaf wilting at 30, 60 and 120 dpi (Tab. 2) showed that genotype was a highly significant factor, block was an important source of variation in some plots (XXIV, XXV and XXX), and plant height was not a significant effect, except in 2011 in plot XXX. The ef-

fect of time since inoculation on the wilting values observed was significant in all susceptibility tests, except in 2009 in plot XXIV (Tab. 2). The highest wilting values were recorded at 60 dpi (data not shown). After pathogen inoculation, the seven clones showed leaf wilting values similar to or lower than “Sapporo Autumn Gold” (Fig. 2). In all tests, the susceptible control clone UPM089 showed wilting values above 70%, confirming the virulence of the isolates used and

the correct inoculation methodology. The most tolerant clone was “Dehesa de Amanuel”, with wilting values below 5% during the two consecutive inoculation trials performed in Madrid.

Six clones were tested at one location (Madrid), but the “Fuente Umbría” clone was tested over four consecutive years in Guadalajara and Palencia (Fig. 2d). These two locations are 280 km apart and have different climate conditions. “Fuente Umbría” should therefore be regarded as the Spanish clone with the most reliable tolerance level to *O. novo-ulmi*. The six other clones will need to pass a second inoculation test under a different environment from Madrid before they can be registered as controlled material. Given the availability of ramets from this material, the second test will be performed in 2016 for the “Ademuz”, “Dehesa de la Villa”, “Dehesa de Amanuel”, “Retiro” and “Toledo” clones, and in 2018 for the “Majadahonda” clone.

Environmental conditions can strongly influence elm susceptibility to *O. novo-ulmi* (Smalley 1963, Sutherland et al. 1997, Solla & Gil 2002, Martín et al. 2010a). Use of registered material in areas with a similar environment to the area of the susceptibility test is therefore highly recommended. The seven clones performed well during the inoculation years under the environmental conditions described, but their long-term tolerance to pos-




Fig. 2 - Susceptibility of the seven *Ulmus minor* clones (least squares means of wilting percentages at 30, 60 and 120 dpi) after tests performed at various experimental plots in Spain. (a) “Ademuz” and “Majadahonda” clones, tested in Puerta de Hierro Forest Breeding Center, Madrid; (b) “Dehesa de Amanuel”, “Retiro” and “Toledo” clones, tested in Puerta de Hierro; (c) “Dehesa de la Villa” clone, tested in Puerta de Hierro; (d) “Fuente Umbría” clone, tested in 2010 and 2011 in El Serranillo Forest Breeding Center, Guadalajara, and in 2012 and 2013 in Calabazanos Forest Health Centre, Palencia. (T): tolerant; (S): susceptible.

Tab. 3 - Morphological description of the seven *Ulmus minor* clones. Numbers in brackets indicate range values. (a): on a scale from 1 to 5, where 5 = most attractive (typical *U. minor* traits), 1 = least attractive (unusual *U. minor* traits).

Feature	Clone						
	Ademuz	Dehesa de la Villa	Majadahonda	Toledo	Dehesa de Amanuel	Retiro	Fuente Umbría
Height growth in Puerta de Hierro, Madrid (cm year ⁻¹)	100.0	63.0	60.8	89.3	90.0	70.5	51.7
Petiole length (mm)	5.2 (4.2-6.2)	6.3 (3.5-10.0)	11.0 (10-12.7)	5.8 (4.6-7.4)	2.6 (1.9-3.4)	7.3 (6.1-8.0)	10.2 (8.3-12.2)
Leaf basal asymmetry (mm)	1.6 (1.1-1.9)	2.9 (2.0-3.8)	3.8 (2.0-4.8)	1.3 (0.7-2.4)	1.3 (0.8-1.8)	1.2 (0.3-1.7)	3.1 (2.2-4.2)
Maximum foliar length (mm)	53.7 (43.5-65.1)	55.4 (44.0-70.0)	50.4 (46.7-53.6)	47.0 (35.6-71.5)	38.5 (36.2-39.6)	71.4 (63.9-79.4)	75.9 (69.8-85.9)
Maximum foliar width (mm)	33.8 (30.0-38.6)	35.6 (28.0-45.0)	28.8 (25.5-30.6)	26.6 (19.6-35.2)	29.7 (27.3-33.0)	42.2 (36.4-48.9)	44.9 (39.3-48.7)
Tooth breadth (mm)	2.2 (1.9-2.6)	3.5 (3.0-4.9)	1.1 (0.8-1.3)	4.1 (3.6-4.7)	2.2 (1.2-2.6)	2.8 (2.5-3.7)	2.8 (1.9-3.8)
Tooth depth (mm)	3.4 (2.8-4.0)	3.7 (2.4-5.0)	1.8 (1.4-2.3)	3.2 (2.6-4.4)	2.7 (2.1-3.5)	2.2 (1.4-2.8)	2.3 (2.1-2.6)
Tooth length (mm)	4.2 (3.8-4.4)	4.6 (4.0-5.1)	2.3 (1.9-2.9)	5.4 (4.8-5.9)	3.5 (2.8-4.2)	2.9 (2.1-3.8)	3.0 (2.2-4.1)
Teeth per leaf (N)	38 (35-42)	44 (30-64)	54 (52-57)	30.0 (28-33)	33 (31-35)	49 (45-53)	32 (28-34)
Pairs of secondary nerves (N)	10 (9-11)	10 (9-12)	12 (11-12)	8.8 (8-10)	9.3 (8-10)	13 (12-15)	12 (11-13)
Leaf serration	Double	Double	Simple	Double	Triple	Double	Double
Presence of corky tissue	No	No	No	No	Yes	No	Yes
Foliar density	Medium	High	High	Medium	High	High	Medium
Branching	Erect	Erect	Erect	Erect	Spreading	Erect	Erect
Crown shape	Spindle	Spindle	Globular	Irregular	Irregular	Globular	Irregular
Ornamental value ^a	4.5	4.1	4.3	2.9	3.0	4.0	3

sible emerging races of the pathogen under different climate conditions need to be assessed. Before the clones are used in other areas, it would be advisable to establish adaptation trials to allow quantification of their susceptibility to drought, frosts, flooding and pests such as bark beetles, Hemiptera and the elm leaf-beetle, *Galerucella luteola*.

DED fungi have caused mortality not only of large elm groves of *U. minor* in Spain, but also of many centuries-old elms that had decorated parks, gardens and town and city squares. Therefore, although registration of the seven clones was focused on forest use, recovery of *U. minor* for urban use was also a key objective of the programme. To this end, distinct morphological features and appreciation of the ornamental value of the seven clones are shown in Tab. 3, Fig. 3 and Fig. 4. Three of the clones (“Ademuz”, “Toledo” and “Dehesa de Amanuel”) showed growth rates very similar to “Sapporo Autumn Gold”, which grows 94.5 cm in height per year in Puerta de Hierro. Foliar density of the seven clones was medium or high compared to “Sapporo Autumn Gold”, which shows low or medium-low density in Madrid. The “Ademuz” and “Majadahonda” clones have the highest ornamental scores and are promising trees for use in urban environments and tree breeding for ornamental

quality. These two genotypes showed a later bud burst phenology than the other *U. minor* clones (Fig. 5), which demonstrates probable suitability to areas with late frost events. The earlier phenology in “Sapporo Autumn Gold” (*U. pumila* × *U. davidiana* var. *japonica* hybrid) than in the Spanish elms is due to its Asian background. Asian elms exhibit earlier bud burst phenology than European elms (Ghelardini et al. 2010).

The results of the genetic characterization of the seven clones are shown in Tab. 4. This description is intended to guarantee the traceability of the plant material during use, especially if clones are commercialized in the near future. From the experience of the Spanish elm breeding programme, the high level of polymorphism obtained with the microsatellites selected allows rigorous identification of each clone.

Tab. 4 - Genetic characterisation of the seven *Ulmus minor* clones, showing alleles for the two chloroplast and 12 nuclear microsatellites used. (a): not amplified.

Organelle	DNA marker	Clone														
		Ademuz		Dehesa de la Villa		Majadahonda		Toledo		Dehesa de Amanuel		Retiro		Fuente Umbría		
Chloroplast	ccmp2	237	236	236	236	215	215	236	236	216	216	215	215	215		
	SFm	278	278	278	278	297	297	278	278	297	297	297	297	297		
Nuclear	Ulm 2	102	108	102	102	106	108	102	102	106	108	108	108	102	102	
	Ulm 3	161	176	176	176	176	180	161	180	176	176	176	180	161	161	
	Ulm 8	196	196	196	196	194	196	194	198	196	196	196	196	194	194	
	UR 123	250	250	252	254	250	254	242	250	250	254	252	254	255	259	
	UR 141	150	152	152	160	152	158	152	152	158	158	158	158	152	158	
	UR 153	178	190	184	188	188	188	188	188	188	178	188	186	188	178	190
	UR 158	195	195	179	195	195	195	179	179	179	199	179	179	179	179	
	UR 159	258	260	260	278	278	278	260	278	278	280	278	278	258	258	
	Ulm 1-98	151	151	151	151	151	151	151	151	151	151	151	151	151	151	154
	Ulm 1-165	204	204	146	146	164	164	130	148	204	204	156	156	160	160	
Ulm 2-16	90	90	90	90	90	90	-	-	90	90	82	94	90	90		
Ulm 2-20	- ^a	-	184	202	172	206	206	220	186	202	180	184	220	220		




Fig. 3 - Registered *Ulmus minor* clones grown in the clonal bank of Puerta de Hierro Forest Breeding Center, Madrid. (a) “Ademuz”; (b) “Dehesa de la Villa”; (c) “Majadahonda”; (d) “Toledo”; (e) “Dehesa de Amanuel”; and (f) “Retiro” clones met the requirements to be registered as “qualified forest reproductive material” in Spain. Scale bars in leaf close-ups = 1 cm.

Fig. 4 - Registered “Fuente Umbría” clone grown in the clonal bank of Puerta de Hierro Forest Breeding Center (Madrid). This *Ulmus minor* clone met the requirements to be registered as “controlled forest reproductive material” in Spain. Scale bars in leaf close-up = 1 cm.




The Spanish programme aimed to directly control DED (Solla & Gil 2003, Martín et al. 2010b, 2012, Vivas et al. 2012) and gain a better understanding of tolerance through new techniques (Martín et al. 2005, 2007, 2008a). One of its medium-term priorities is to increase the genetic diversity of tolerant

native elms. Elm tolerance to *O. novo-ulmi* has been shown to be inheritable (Townsend 2000, Guries & Smalley 2000, Solla et al. 2014, Venturas et al. 2014) and polygenic (quantitative) in nature (Aoun et al. 2010). It also depends on constitutive and inducible mechanisms of defence (e.g., Martín et al.

2007, 2008b, 2013). In addition, the defense mechanisms of elms to *O. novo-ulmi* seem to differ between genotypes. This is the case of some anatomical features of the xylem associated with pathogen dispersal, such as pit and vessel size (Martín et al. 2009, 2013). When more resistance mechanisms are ga-

Fig. 5 - Leaf phenology in 2011 of the seven registered *Ulmus minor* clones and the “Sapporo Autumn Gold” control clone in Puerta de Hierro Forest Breeding Centre, Madrid.



thered in the same genotype, the chances of overcoming an infection are likely to increase. Thus it would be desirable to perform controlled crossings between genotypes that express different, and preferably complementary, defense mechanisms. If multiple resistance layers act jointly and in a complementary fashion, the influence of environmental factors in tree tolerance to *O. novo-ulmi* would probably be lower. The possibility of any emerging variant or pathogen mutation overcoming the resistance mechanisms would also decrease. Understanding the genetic basis of elm tolerance to *O. novo-ulmi* is our second main research challenge.

To broaden the genetic base of tolerant native elms, the Spanish programme has grown 1400 seedlings from controlled F₁ crossings between the seven *U. minor* clones. In 2016, when seedlings are four years old, they will be inoculated with *O. novo-ulmi*. The tolerance of new genotypes from different provenances in Spain to *O. novo-ulmi* will be assessed through clonal replicates ($N \geq 6$) in the near future. Most of these genotypes have shown high tolerance levels when individually tested as seedlings. With this material, the Spanish programme expects to substantially increase the number of *U. minor* clones tolerant to *O. novo-ulmi*.

Conclusions

Although tree selection, trial establishment and breeding cycles require a major investment in time and effort, breeding programmes are the most reliable option for recovery of native elm populations. Results reported here show that selection of tolerant native *U. minor* genotypes is possible. The seven clones registered as forest reproductive material have shown high tolerance to DED in Spain. Their form and foliage are attractive and they are fast-growing trees. Longitudinal monitoring of the performance of the selected clones under different environments will make it possible to determine the suitable environmental range of each clone. New elm varieties likely to show low wilting values after *O. novo-ulmi* inoculation will be obtained in the next few years.

Acknowledgements

This paper is dedicated to Margarita Burón, who spent the last ten years of her life working on the conservation and breeding of Spanish elms.

The authors are grateful to everyone who has participated in the Spanish elm breeding programme during its 27 years of existence. We would also like to express our gratitude to the Dutch and Italian elm breeding programmes and the participants in the EU Project RESGEN CT96-78 for sharing their knowledge and materials. The programme is funded by the Spanish Directorate-General of Rural Development and Forestry Policy

(Dirección General de Desarrollo Rural y Política Forestal) and the Spanish Ministry of Agriculture, Food and Environmental Affairs (Ministerio de Agricultura, Alimentación y Medio Ambiente).

References

Allué-Andrade JL (1990). Atlas fitoclimático de España. Taxonomías [Phytoclimatic atlas of Spain. Taxonomy]. Instituto Nacional de Investigaciones Agrarias, Ministerio de Agricultura Pesca y Alimentación, Madrid, Spain. [in Spanish]

Aoun M, Jacobi V, Boyle B, Bernier L (2010). Identification and monitoring of *Ulmus americana* transcripts during in vitro interactions with the Dutch elm disease pathogen *Ophiostoma novo-ulmi*. *Physiological and Molecular Plant Pathology* 74: 254-266. - doi: [10.1016/j.pmpp.2010.04.003](https://doi.org/10.1016/j.pmpp.2010.04.003)

Brasier CM, Kirk SA (2010). Rapid emergence of hybrids between two subspecies of *Ophiostoma novo-ulmi* with a high level of pathogenic fitness. *Plant Pathology* 59: 186-199. - doi: [10.1111/1/j.1365-3059.2009.02157.x](https://doi.org/10.1111/1/j.1365-3059.2009.02157.x)

Buiteveld J, van der Werf B, Hiemstra JA (2014). Comparison of commercial elm cultivars and promising unreleased Dutch clones for resistance to *Ophiostoma novo-ulmi*. iForest (early view). - doi: [10.3832/ifer1209-008](https://doi.org/10.3832/ifer1209-008)

Cogolludo-Agustín MA, Agúndez D, Gil L (2000). Identification of native and hybrid elms in Spain using isozyme gene markers. *Heredity* 85: 157-166. - doi: [10.1046/j.1365-2540.2000.00740.x](https://doi.org/10.1046/j.1365-2540.2000.00740.x)

Collada C, Fuentes-Utrilla P, Gil L, Cervera MT (2004). Characterization of microsatellite loci in *Ulmus minor* Miller and cross-amplification in *U. glabra* Hudson and *U. laevis* Pall. *Molecular Ecology Notes* 4: 731-732. - doi: [10.1111/j.1471-8286.2004.00798.x](https://doi.org/10.1111/j.1471-8286.2004.00798.x)

Ghelardini L, Santini A, Black-Samuelsson S, Myking T, Falusi M (2010). Bud dormancy release in elm (*Ulmus* spp.) clones - a case study of photoperiod and temperature responses. *Tree Physiology* 30: 264-74. - doi: [10.1093/treephys/tpp110](https://doi.org/10.1093/treephys/tpp110)

Gil L, Fuentes-Utrilla P, Soto A, Cervera MT, Collada C (2004). English elm (*Ulmus procera*) is a 2000-year-old Roman clone. *Nature* 431: 1053. - doi: [10.1038/4311053a](https://doi.org/10.1038/4311053a)

Guries RP, Smalley EB (2000). Once and future elms: classical and molecular approaches to Dutch elm disease resistance. In: "The elms: breeding, conservation and disease management" (Dunn CP ed), Kluwer Academic Publishers, Boston, USA, pp. 231-248.

Heybroek HM (1993). The Dutch elm breeding program. In: "Dutch Elm Disease Research: Cellular and Molecular Approaches" (Sticklen MB, Sherald JL eds). Springer Verlag, New York, USA, pp. 16-25.

Iglesias S (2005). Normativa de comercialización de material forestal de reproducción [Marketing regulations for forest reproductive material]. In: "Manual para la Comercialización y Producción de Semillas y Plantas Forestales. Materiales de

Base y Reproducción" (Alía A, Alba N, Agúndez D, Iglesias S eds), Organismo Autónomo Parques Nacionales, Ministerio de Medio Ambiente, Madrid, Spain, pp. 39-58. [in Spanish]

Ipinza R (1990). Algunos aspectos relevantes sobre la taxonomía de los olmos ibéricos [Some relevant aspects of the taxonomy of Iberian elms]. In: "Los olmos y la grafiosis en España" (Gil L ed). ICONA, Madrid, pp. 69-98. [in Spanish]

Jeffers J, Richens R (1970). Multivariate analysis of the English elm population. *Silvae Genetica* 19: 31-38. [online] URL: http://sauerlaender-verlag.com/fileadmin/content/dokument/archiv/silvae-genetica/19_1970/19-1-31.pdf

Martín JA, Solla A, Woodward S, Gil L (2005). Fourier transform-infrared spectroscopy as a new method for evaluating host resistance in the Dutch elm disease complex. *Tree Physiology* 25: 1331-1338. - doi: [10.1093/treephys/25.10.1331](https://doi.org/10.1093/treephys/25.10.1331)

Martín JA, Solla A, Woodward S, Gil L (2007). Detection of differential changes in lignin composition of elm xylem tissues inoculated with *Ophiostoma novo-ulmi* using Fourier transform infrared spectroscopy. *Forest Pathology* 37: 187-191. - doi: [10.1111/j.1439-0329.2007.00495.x](https://doi.org/10.1111/j.1439-0329.2007.00495.x)

Martín JA, Solla A, Coimbra MA, Gil L (2008a). Metabolic fingerprinting allows discrimination between *Ulmus pumila* and *U. minor* and between *U. minor* clones of different susceptibility to Dutch elm disease. *Forest Pathology* 38: 244-256. - doi: [10.1111/j.1439-0329.2007.00542.x](https://doi.org/10.1111/j.1439-0329.2007.00542.x)

Martín JA, Solla A, Domingues MR, Coimbra MA, Gil L (2008b). Exogenous phenol increase resistance of *Ulmus minor* to Dutch elm disease through formation of suberin-like compounds on xylem tissues. *Environmental and Experimental Botany* 64: 97-104. - doi: [10.1016/j.envexpbot.2008.05.004](https://doi.org/10.1016/j.envexpbot.2008.05.004)

Martín JA, Solla A, Esteban LG, de Palacios P, Gil L (2009). Bordered pit and ray morphology involvement in elm resistance to *Ophiostoma novo-ulmi*. *Canadian Journal of Forest Research* 39: 420-429. - doi: [10.1139/X08-183](https://doi.org/10.1139/X08-183)

Martín JA, Solla A, Gil L, García-Vallejo MC (2010a). Phenological and histochemical changes of *Ulmus minor* due to root absorption of phenol: implications for resistance to DED. *Environmental and Experimental Botany* 69: 175-182. - doi: [10.1016/j.envexpbot.2010.04.001](https://doi.org/10.1016/j.envexpbot.2010.04.001)

Martín JA, Solla A, Witzell J, Gil L, García-Vallejo MC (2010b). Antifungal effect and reduction of *Ulmus minor* symptoms to *Ophiostoma novo-ulmi* by carvacrol and salicylic acid. *European Journal of Plant Pathology* 127: 21-32. - doi: [10.1007/s10658-009-9567-3](https://doi.org/10.1007/s10658-009-9567-3)

Martín JA, Solla A, García-Vallejo MC, Gil L (2012). Chemical changes in *Ulmus minor* xylem tissue after salicylic acid or carvacrol treatments are associated with enhanced resistance to *Ophiostoma novo-ulmi*. *Phytochemistry* 83: 104-109. - doi: [10.1016/j.phytochem.2012.07.017](https://doi.org/10.1016/j.phytochem.2012.07.017)

Martín JA, Solla A, Ruiz-Villar M, Gil L (2013). Vessel length and conductivity of *Ulmus* branches: ontogenetic changes and relation to resistance to Dutch elm disease. *Trees* 27: 1239-

1248. - doi: [10.1007/s00468-013-0872-2](https://doi.org/10.1007/s00468-013-0872-2)
- Mitterpergher L, Santini A (2004). The history of elm breeding. *Forest Systems* 13: 161-177. [online] URL: <http://revistas.inia.es/index.php/fs/article/view/821>
- Richens RH (1955). Studies on *Ulmus*. I. The range of variation of East Anglian elms. *Watsonia* 3: 138-153. [online] URL: <http://www.archive.bsbi.org.uk/Wats3p138.pdf>
- Santini A, Fagnani A, Ferrini F, Mitterpergher L, Brunetti M, Crivellaro A, Macchioni N (2004). Elm breeding for DED resistance: the Italian clones and their wood properties. *Forest Systems* 13: 179-184. [online] URL: <http://revistas.inia.es/index.php/fs/article/view/822>
- Santini A, Fagnani A, Ferrini F, Ghelardini L, Mitterpergher L (2005). Variation among Italian and French elm clones in their response to *Ophiostoma novo-ulmi* inoculation. *Forest Pathology* 35: 183-193. - doi: [10.1111/j.1439-0329.2005.00401.x](https://doi.org/10.1111/j.1439-0329.2005.00401.x)
- Santini A, Pecori F, Pepori A, Brookes A (2011). "Morfeo" elm: a new variety resistant to Dutch elm disease. *Forest Pathology* 42:171-176. - doi: [10.1111/j.1439-0329.2011.00737.x](https://doi.org/10.1111/j.1439-0329.2011.00737.x)
- Smalley EB (1963). Seasonal fluctuations in susceptibility of young elm seedlings to Dutch elm disease. *Phytopathology* 53: 846-853.
- Smalley EB, Guries RP (2000). Asian elms: sources of disease and insect resistance. In: "The elms: breeding, conservation and disease management" (Dunn CP ed). Kluwer Academic Publishers, Boston, USA, pp. 215-230.
- Smalley EB, Lester DT (1973). "Sapporo autumn gold" elm. *Horticultural Science* 8: 514-515.
- Solla A, Gil L (2002). Influence of water stress on Dutch elm disease symptoms in *Ulmus minor* Miller. *Canadian Journal of Botany* 80: 810-817. - doi: [10.1139/b02-067](https://doi.org/10.1139/b02-067)
- Solla A, Gil L (2003). Evaluating *Verticillium dahliae* for biological control of *Ophiostoma novo-ulmi* in *Ulmus minor*. *Plant Pathology* 52: 579-585. - doi: [10.1046/j.1365-3059.2003.00921.x](https://doi.org/10.1046/j.1365-3059.2003.00921.x)
- Solla A, Burón M, Iglesias S, Gil L (2000). Spanish program for the conservation and breeding of elms against DED. In: "The Elms: Breeding, Conservation and Disease Management" (Dunn CP ed), Kluwer Academic Publishers, Boston, pp. 295-303.
- Solla A, Bohnens J, Collin E, Diamandis S, Franke A, Gil L, Burón M, Santini A, Mitterpergher L, Pinon J, van den Broeck A (2005a). Screening European elms for resistance to *Ophiostoma novo-ulmi*. *Forest Science* 51: 134-141.
- Solla A, Martín JA, Corral P, Gil L (2005b). Seasonal changes in wood formation of *Ulmus pumila* and *U. minor* and its relation with Dutch elm disease. *New Phytologist* 166:1025-1034. - doi: [10.1111/j.1469-8137.2005.01384.x](https://doi.org/10.1111/j.1469-8137.2005.01384.x)
- Solla A, Martín JA, Ouellette G, Gil L (2005c). Influence of plant age on symptom development in *Ulmus minor* following inoculation by *Ophiostoma novo-ulmi*. *Plant Disease* 89:1035-1040. - doi: [10.1094/PD-89-1035](https://doi.org/10.1094/PD-89-1035)
- Solla A, López-Almansa JC, Martín JA, Gil L (2014). Genetic variation and heritability estimates of *Ulmus minor* and *U. pumila* hybrids for budburst, growth and tolerance to *Ophiostoma novo-ulmi*. *iForest* [submitted].
- Sutherland ML, Pearson S, Brasier CM (1997). The influence of temperature and light on defoliation levels of elm by Dutch elm disease. *Phytopathology* 87: 576 -581. - doi: [10.1094/PHYTO.1997.87.6.576](https://doi.org/10.1094/PHYTO.1997.87.6.576)
- Tchernoff V (1965). Methods for screening and for the rapid selection of elms for resistance to Dutch elm disease. *Acta Botanica Neerlandica* 14: 409-452. - doi: [10.1111/j.1438-8677.1965.tb00204.x](https://doi.org/10.1111/j.1438-8677.1965.tb00204.x)
- Townsend AM (2000). USDA genetic research on elms. In: "The Elms: Breeding, Conservation and Disease Management" (Dunn CP ed). Kluwer Academic Publishers, Boston, USA, pp. 271-278.
- Venturas M, López R, Martín JA, Gascó A, Gil L (2014). Heritability of *Ulmus minor* resistance to Dutch elm disease and its relationship to vessel size, but not to xylem vulnerability to drought. *Plant Pathology* - doi: [10.1111/ppa.12115](https://doi.org/10.1111/ppa.12115).
- Vivas M, Martín JA, Gil L, Solla A (2012). Evaluating methyl jasmonate for induction of resistance to *Fusarium oxysporum*, *F. circinatum* and *Ophiostoma novo-ulmi*. *Forest Systems* 21: 289-299. - doi: [10.5424/fs/2012212-02172](https://doi.org/10.5424/fs/2012212-02172)
- Webber JF (2000). Insect vector behavior and the evolution of Dutch elm disease. In: "The Elms: Breeding, Conservation and Disease Management" (Dunn CP ed). Kluwer Academic Publishers, Boston, MS, USA, pp. 47-60.
- Weising K, Gardner RC (1999). A set of conserved PCR primers for the analysis of simple sequence repeat polymorphisms in chloroplast genomes of dicotyledonous angiosperms. *Genome* 42: 9-19. - doi: [10.1139/g98-104](https://doi.org/10.1139/g98-104)
- Whiteley RE, Black-Samuelsson S, Clapham D (2003). Development of microsatellite markers for the European white elm (*Ulmus laevis* Pall.) and cross-species amplification within the genus *Ulmus*. *Molecular Ecology Notes* 3: 598-600. - doi: [10.1046/j.1471-8286.2003.00525.x](https://doi.org/10.1046/j.1471-8286.2003.00525.x)
- Zalapa JE, Brunet J, Guries RP (2008). Isolation and characterization of microsatellite markers for red elm (*Ulmus rubra* Muhl.) and cross-species amplification with Siberian elm (*Ulmus pumila* L.). *Molecular Ecology Resources* 8: 109-112. - doi: [10.1111/j.1471-8286.2007.01805.x](https://doi.org/10.1111/j.1471-8286.2007.01805.x)