

University of New Hampshire
University of New Hampshire Scholars' Repository

Earth Sciences Scholarship

Earth Sciences

12-13-1999

Application of Spaceborne Synthetic Aperture Radar to Monitoring Seasonal Ecological and Hydrologic Processes in Boreal Forest

Kyle McDonald
California Institute of Technology

John Kimball
University of Montana, Missoula

A R. Keyser
University of Montana, Missoula

S W. Running
University of Montana, Missoula

Steve Frolking
University of New Hampshire - Main Campus, steve.frolking@unh.edu

See next page for additional authors

Follow this and additional works at: https://scholars.unh.edu/earthsci_facpub

Recommended Citation

McDonald, KC, Kimball J, Keyser A, Running S, Frolking S, Zimmermann R, & Way JB. 1999. Application of Spaceborne Synthetic Aperture Radar to Monitoring Seasonal Ecological and Hydrologic Processes in Boreal Forest. EOS Trans. Suppl. 80:F96. Fall 1999 AGU.

This Conference Proceeding is brought to you for free and open access by the Earth Sciences at University of New Hampshire Scholars' Repository. It has been accepted for inclusion in Earth Sciences Scholarship by an authorized administrator of University of New Hampshire Scholars' Repository. For more information, please contact nicole.hentz@unh.edu.

Authors

Kyle McDonald, John Kimball, A R. Keyser, S W. Running, Steve Frolking, Reiner Zimmermann, and JoBea Way

Session B: Biogeosciences

for Poster Presentation

Application of Spaceborne Synthetic Aperture Radar to Monitoring Seasonal Ecological and Hydrologic Processes in Boreal Forest

Kyle C. MCDONALD(1), John KIMBALL(2), Alisa KEYSER(2), Steve RUNNING(2), Steve FROLKING(3), Reiner ZIMMERMANN(3), and JoBea WAY(1)

(1)Jet Propulsion Laboratory, California Institute of Technology, Mail Stop 300-233
4800 Oak Grove Drive, Pasadena, CA 91109
818-354-3263, FAX: 818-354-9476, kyle.mcdonald@jpl.nasa.gov

(2)NTSG School of Forestry, University of Montana
Missoula, MT 59802

(3)Complex Systems Research Institute,
University of New Hampshire, Durham, NH

(4)Bayreuth Institute for Terrestrial Ecosystem Research, University of Bayreuth
Plant Ecology II, Dr. Hans Frisch Str. 1, D-95448, Bayreuth, GERMANY
(49)-921-555624, FAX: (49)-921-555799, reiner.zimmermann@bitoek.uni-bayreuth.de

Freeze/thaw transitions in boreal landscapes drive critical dynamics in ecosystem and hydrologic activity. A capability for accurate, repeated, and reliable monitoring of landscape freeze/thaw dynamics would improve our ability to quantify the interannual variability of boreal hydrology and river runoff/flood dynamics and to assess the period of photosynthetic activity in boreal and arctic ecosystems, thus improving estimates of annual carbon budgets and of the interannual variability of regional carbon fluxes. Results from BOREAS experiments have indicated that the boreal forest has a net annual carbon flux near zero. A first step in assessing and monitoring year-to-year changes in the boreal carbon flux is to determine the annual variation in growing season length. We apply imagery from the ERS spaceborne Synthetic Aperture Radars (SARs) to estimate landscape freeze/thaw dynamics over selected areas of the BOREAS region of Canada. A temporal series of freeze/thaw maps are derived that provide fractional estimates of frozen and thawed landscape. The inferred landscape freeze/thaw state is validated against temperature measurements obtained from a distributed temperature monitoring network and from meteorological observations. We examine the relationships of radar-estimated thaw patterns with topography and landcover. SAR-derived timing of spring thaw is compared with initiation of streamflow. Ecological process models are used to estimate Net Ecosystem Exchange (NEE) and Net Primary Productivity (NPP) on the landscape scale. Model results are compared with timing and spatial distribution of freeze and thaw events. As the timing of spring thaw is a major factor influencing the net annual carbon flux, we seek to incorporate the radar-based measure of landscape freeze/thaw dynamics as direct input to ecological process models to provide a capability for improved ecosystem carbon flux estimates at regional scales using spaceborne radar.

This work was carried out at the Jet Propulsion Laboratory, California Institute of Technology, under contract to the National Aeronautics and Space Administration.