#### University of New Hampshire University of New Hampshire Scholars' Repository

Center for Coastal and Ocean Mapping

Center for Coastal and Ocean Mapping

1-22-2001

### Geographic Information Systems Tools for Collecting and Accessing Árctic Bathymetry: International Bathymetric Chart of the Arctic Ocean

Martin Jakobsson University of New Hampshire, Durham

Follow this and additional works at: https://scholars.unh.edu/ccom



Part of the Oceanography and Atmospheric Sciences and Meteorology Commons

#### Recommended Citation

Jakobsson, Martin, "Geographic Information Systems Tools for Collecting and Accessing Arctic Bathymetry: International Bathymetric Chart of the Arctic Ocean" (2001). Arctic Geographic Information Systems (AGIS). 135. https://scholars.unh.edu/ccom/135

This Conference Proceeding is brought to you for free and open access by the Center for Coastal and Ocean Mapping at University of New Hampshire Scholars' Repository. It has been accepted for inclusion in Center for Coastal and Ocean Mapping by an authorized administrator of University of New Hampshire Scholars' Repository. For more information, please contact nicole.hentz@unh.edu.

















# GIS Tools for Accessing Arctic Bathymetry: International Bathymetric Chart of the Arctic Ocean (IBCAO)



### **Martin Jakobsson**

Center for Coastal and Ocean Mapping
Joint Hydrographic Center
University of New Hampshire
















- 1. The IBCAO bathymetry model
- 2. Users of IBCAO
- 3. GIS tools for accessing the IBCAO grid model
- 4. Future IBCAO GIS products (Error estimation of the IBCAO bathymetry, vector contours)



### **IBCAO**

Grid model representing
Arctic Ocean bathymetry
and topography

**Grid cell spacing:** 

2.5 x 2.5 km

**Projection:** 

Polar stereographic

True scale: 75° N

Datum: WGS 84














### TRACKLINE SOURCES

#### ....

US and British Royal Navy nuclear submarine cruises (1958 - 1988).

#### 00000


Soundings collected by nuclear submarines during the SCICEX program (1993-1999

#### 00000

Soundings collected by surface vessels primarely obtained from four archives:

- (1) the US National Geophysical Data Center (NGDC)
- (2) the US Naval Research Laboratory (NRL)
- (3) the Canadian Hydrographic Service (CHS)
- (4) the Royal Danish Administration of Navigation and Hydrography (RDANH)

Recent acquisitions provided by agencies that mobilized missions aboard the Swedish icebreaker Oden (1991, 1996) and the German research vessel Polarstern (1990, 1994, 1995, 1997) are also marked in blue.















### MAP SOURCES



Contours drawn during this present work in order to faciliate the computer gridding.



Navigational charts published by the Russian Federation's Head Department of Navigation and Oceanography (HDNO).



Newly-published Russian map (Head Dep. of Navigation and Oceanography et al., 1999).



Bathymetry in the Gulf of Bothnia was derived from a digital grid compiled by Seifert and Kayser (1995).



Charts published by the US Naval Research Laboratory (Perry et al., 1985; Cherkis et al., 1991; Matishov et al., 1995).




Contours retrieved from GEBCO Digital Atlas.



Greenland DTM developed by KMS, the Danish Cadaster and Mapping Agency (Ekholm, 1996).



GTOPO30 topographic model (U.S. Geological Survey, 1997).


















## SOURCES
















### **GRID COMPILATION**


**Data gathering** 

(MGE)

Data
cleaning & \_
mining
(MGE)











Gridding using continuous curvature splines in tension (Smith & Wessel, 1990)







































### **IBCAO** web site:

#### http://www.ngdc.noaa.gov/mgg/bathymetry/arctic/arctic.html
















### **USERS OF THE IBCAO BATHYMETRY GRID**

Oceanography modeling (e.g. Arctic ice-ocean model for the Polar Ice prediction System (PIPS), US Naval Postgraduate School; ARCICE, Southampton Oceanography Centre, UK)

Geophysical modeling
lce sheet modeling
Cartographic applications

#### **IBCAO** web site news:

February, 26 652 accesses and 99297 hits

















## GIS Tools for accessing the IBCAO model

**ESRI:** ArcView, ArcInfo (Spatial Analyst, Arc Grid)

**Intergraph:** MGE (Terrain Analyst)

Geomedia (MFworks from ThinkSpace Inc.)

Clarks Labs: Idrisi32

HIPS Caris:























## Geophysical and Mapping softwares

**IVS:** Fledermaus



**Public domain:** Generic Mapping Tools



Web GIS tools

Intergraph: Geomedia Web Map

ESRI: ArcIMS3















### THE NESSECITY FOR AN ERROR ESTIMATION

Regional grid models representing bathymetry/topography are often used as a base for oceanographic/climate/geophysical/ice sheet modeling

The modeling results are often interpreted without knowledge of the spatial differences in reliability of the underlying bathymetry/topography model

The idea is to create a grid with the same structure as the bathymetry/topography grid containing an estimate of the standard deviation of the errors of the bathymetry/topography
















# USING THE MONTE CARLO METHOD FOR ERROR ESTIMATION

- 1. Randomly vary the source data within constraints determined by meta data (navigation, echo sounder etc)
- 2. Go through the process with which the bathymetry grid is compiled (block median filtering + continuous curvature splines in tension)



This processes are continued until......

















## **STATISTICS**



## Standard deviation of estimated error

(% of block median depth)



### **IBCAO** contours in Geomedia

