

## RESEARCH REPORT

**Influence of sericin in alleviating the hydrogen peroxide induced oxidative stress in silkworm *Bombyx mori*: role of the amino acids****AS Micheal, M Subramanyam***Department of Life Science, Bangalore University, Bangalore 560 056, India**Accepted September 15, 2014***Abstract**

Sericin is an important peptide derived from silk fibre spun by the silkworm *Bombyx mori* and has various biological activities. The aim of the present study was to characterize the major constituents of sericin that are providing cytoprotective effect against hydrogen peroxide-induced cell damage in midgut epithelial cells and hemocytes of silkworm. Extracted sericin was subjected to LCMS analysis for amino acid composition. Isolated cells of midgut and hemocytes were incubated with sericin or with mixture of serine and aspartic acid prior to suboptimal concentration of hydrogen peroxide treatment. Sericin as well as amino acid mixture reduced the activity of antioxidant enzymes triggered by hydrogen peroxide, inhibited oxidative derivatives such as protein carbonyl and malondialdehyde and increased antioxidant capacity in both the cells studied. Furthermore, sericin and amino acid mixture significantly decreased intracellular reactive oxygen species as assessed by fluorescent detection. These results suggest that major constituent amino acids of sericin defend midgut epithelial cells and hemocytes against oxidative damage by scavenging reactive oxygen species rather than activating antioxidant enzyme system thereby inhibiting cell damage.


**Key Words:** amino acids; antioxidant enzymes; *Bombyx mori*; oxidative stress; reactive oxygen species; sericin**Introduction**

Reactive oxygen species (ROS) are produced during normal cellular metabolism or are derived from exogenous sources, and play an important role in cellular homeostasis. Exogenous sources, including prooxidant allelochemicals, severely affect herbivorous insects during host interactions and stress such as starvation may contribute to endogenous sources of ROS accumulation. Unchecked or increased levels of ROS can cause severe damage to various cellular compartments, including DNA, protein and lipids, thereby causing oxidative injury. Oxidative injury to the cell membrane results in structural changes and increased permeability to ions and fluids (Nagasaka *et al.*, 2004). Nonetheless, insects have evolved a complex antioxidant mechanism to overcome the toxic effects of ROS (Krisnhnan and Kodrik, 2006). Antioxidant defence is primarily contributed by antioxidant enzymes (AOEs) such as peroxidase (POX, E.C. 1.11.1.7), superoxide dismutase (SOD, E.C. 1.15.1.1) and catalase (CAT, E.C. 1. 11. 1.6)

(Felton and Summer, 1995). Glutathione peroxidase (GPx) reduces H<sub>2</sub>O<sub>2</sub> and hydroperoxides, thereby scavenging oxidative radicals in cell membranes, whereas SOD converts O<sub>2</sub><sup>-</sup> to molecular O<sub>2</sub> and H<sub>2</sub>O<sub>2</sub> (Maiorino *et al.*, 2003; Imlay, 2008). H<sub>2</sub>O<sub>2</sub> is subsequently scavenged by CAT, resulting in the production of water and molecular oxygen (Kashiwagi *et al.*, 1997).

Several naturally occurring macromolecules (Balsano and Alisi, 2009) and amino acids have been found to be effective as antioxidants (Wu *et al.*, 2003a; Atmaca, 2004; Liu *et al.*, 2004; Movahedian *et al.*, 2006; Selvaraju and Subbashinidevi, 2011). Among the macromolecular proteins that combat oxidative stress (Elisa *et al.*, 2008; Medina-Navarro *et al.*, 2010), the silk protein sericin role as an antioxidant is commendable because of the presence of polyphenols and flavanoids (Devi *et al.*, 2011; Prasong, 2011). Sericin is an active biomolecule with several implications as a therapeutic agent (Khudaiderdier, 1997; Aramwit and Sangcakul, 2007; Li *et al.*, 2008) and component of cosmetics (Padamwar and Pawar 2004; Patel and Modasiya, 2011). Silk protein filament has two protein fractions, fibroin, a fibrous component and glue-like sericin that holds the fibroin components together and often discarded as waste product in silk industry. Sericin exhibits

*Corresponding author:*  
Muthangi Subramanyam  
Department of Life Science  
Bangalore University  
Bangalore 560 056, India  
E-mail: [asuba@vsnl.net](mailto:asuba@vsnl.net)




**Fig. 1** (A) LCMS spectra of sericin showing prominent amino acids. (B) Spectra of the major amino acids serine (a), aspartic acid (b), glycine (c), threonine (d) and glutamic acid (e). The X-axis is retention time and the Y-axis is intensity in cps. (C) Prominent amino acid composition of sericin protein in *B. mori*.

several pharmacological effects, such as enhancing digestion and cryoprotection. Dietary sericin reduces lipids and ameliorates glucose tolerance in rats fed a high fat diet and also acts as an anticoagulant upon sulphonation (Kato *et al.*, 1998; Sasaki *et al.*, 2000; Tsujimoto *et al.*, 2001; Okazaki *et al.*, 2010). Sericin is specifically synthesized in the middle silk gland of the silkworm *B. mori*. It is a polypeptide with 18 amino acids, most of which have strong polar side groups such as hydroxyl, carbonyl and amino groups (Wu *et al.*, 2007) and is especially rich in serine (~32 %) (Kwang *et al.*, 2003). In general, amino acids found in proteins have the potential to interact with free radicals if the energy of the radicals insult is high (Elias *et al.*, 2008) and antioxidant activity of proteins in radical

mediated oxidation reactions may be due to their ability to act as radical trapping devices (Ostdal *et al.*, 2002).

The silkworm *Bombyx mori* is a monophagous insect known worldwide because of the lustrous silk it produces in the final instar of the larval stage. One of the reasons for the decline in silk production and the survivability of highly domesticated mulberry silkworm is due to food contaminated with pesticides (Vyjayanthi and Subramanyam, 2002a, b). In an earlier study, we demonstrated a transient increase in antioxidant defence mechanisms in silkworms stressed with short-term exposure to low temperature, hypoxia and viral infection (Micheal and Subramanyam, 2013). The purpose of the present study was to assess the effect of sericin on




**Fig. 2** Malondialdehyde levels in silkworm *B. mori* midgut epithelial cells (a) and hemocytes (b) treated with hydrogen peroxide and pre-incubated with sericin or amino acid mixture prior to hydrogen peroxide treatment. Data are shown as mean  $\pm$  SE (n = 6);  $p < 0.05$  was considered significant. Values between the treatments are represented by lower case letters (a, b) and between the instars are represented by upper case letters (A, B). Those not sharing the same letters are significant.

hydrogen peroxide-induced oxidative stress in the midgut epithelial cells and hemocytes of silkworm and to explore the possible components of antioxidant activity that alleviate oxidative injury caused by transient exposure to hydrogen peroxide. We have demonstrated that the major amino acids of sericin acts as scavenger of radical oxygen species in hydrogen peroxide induced oxidative stress.

## Materials and Methods

### Chemicals

Thiobarbituric acid (TBA), glutathione reductase, horseradish peroxidase and dinitrophenylhydrazine (DNPH) were purchased from Sigma-Aldrich, (St. Louis, MO, USA). H<sub>2</sub>DCFDA was purchased from Molecular Probes (Eugene, USA). Reduced glutathione (GSH), NADPH, t-butylhydroperoxide,




**Fig. 3** Protein carbonyl levels in silkworm *B. mori* midgut epithelial cells (a) and hemocytes (b) treated with hydrogen peroxide and pre-incubated with sericin or amino acid mixture prior to hydrogen peroxide treatment. Data are shown as mean  $\pm$  SE (n = 6);  $p < 0.05$  was considered significant. Values between the treatments are represented by lower case letters (a, b) between the instars are represented by upper case letters (A, B). Those not sharing the same letters are significant.

hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>), Triton X-100, epinephrine, sodium dodecyl sulphate (SDS), acetic acid, butanol, pyridine, tetra methoxy propane (TMP) and 2,4,6-tripyridyl-S-triazine (TPTZ) were purchased from Sisco Research Laboratory (Mumbai, India). L-Aspartic acid and L- Serine from Spectochem (Mumbai, India).

#### *Insects and experimental design*


The present study was approved by the Institutional Animal Ethics Committee (IAEC) of Bangalore University, Bangalore, India. Second instar larvae were procured from the Kunigal seed area, Karnataka, India and were maintained in the laboratory throughout the larval stages and were fed



**Fig. 4** SOD activity in midgut epithelial cells (a) and hemocytes (b) of IV and V instar of the silkworm *B. mori* treated with hydrogen peroxide and pre-incubated with sericin or amino acid mixture prior to hydrogen peroxide treatment. Data are shown as mean  $\pm$  SE (n = 6);  $p < 0.05$  was considered significant. Values between the treatments are represented by lower case letters (a, b) between the instars are represented by upper case letters (A, B). Those not sharing the same letters are significant.

*ad libitum* with M5 variety mulberry leaves (Vyjayanthi and Subramanyam, 2002a, b). The uniformly grown healthy larvae of IV and V instars were used in all experiments and were maintained at 24 - 25 °C with a relative humidity of 70 - 75 %. Midgut epithelial cells were isolated by micro dissection and 1 % collagenase treatment. Dissociated cells were washed repeatedly with cold phosphate buffer and were allowed to stand in cold phosphate buffer of pH 7.4 for further usage. Hemolymph was collected in pre-cooled 2 ml vials

containing 5 mg thiourea by gentle incision on the caudal horn of the larvae. Hemocytes were separated by centrifuging the diluted hemolymph at 3,000 rpm for 10 min in cold. Cold phosphate buffer of pH 7.4 was used for the tissue homogenate preparation and for the separation of hemocytes. Isolated cells were exposed to H<sub>2</sub>O<sub>2</sub> (20  $\mu$ M) for 10 min to induce oxidative stress. To evaluate the antioxidant nature of sericin, isolated cells were incubated with 28 mg/ml sericin for 10 min at RT prior to treatment with H<sub>2</sub>O<sub>2</sub>. The isolated cells were




**Fig. 5** CAT activity in midgut epithelial cells (a) and hemocytes (b) of IV and V instar of the silkworm *B. mori* treated with hydrogen peroxide and pre-incubated with sericin or amino acid mixture prior to hydrogen peroxide treatment. Data are shown as mean  $\pm$  SE ( $n = 6$ );  $p < 0.05$  was considered significant. Values between the treatments are represented by lower case letters (a, b) between the instars are represented by upper case letters (A, B). Those not sharing the same letters are significant.

also incubated with 0.5 mM each of L-serine and L-aspartic acid for 10 min at room temperature prior to treatment with H<sub>2</sub>O<sub>2</sub> to access the role of amino acids as antioxidants.

#### Extraction of sericin

Sericin was extracted from silk cocoon according to (Wu *et al.*, 2007) with slight modification. Briefly, multi-voltine cocoons were boiled for several hours

in distilled water to extract sericin. The water extract was condensed further by evaporating the water at 50 °C, and the concentrate was spray dried and collected as a powder. The powder was dissolved in distilled water in a 10:1 ratio (w/v), and was chilled overnight at 4 °C. Pure chilled ethanol was added to the sericin solution with constant stirring to obtain a final ethanol concentration of 75 % (w/v). The obtained mixture was then kept at -20 °C overnight,



**Fig. 6** GPx-like activity in midgut epithelial cells (a) and hemocytes (b) of IV and V instar of the silkworm *B. mori* treated with hydrogen peroxide and pre-incubated with sericin or amino acid mixture prior to hydrogen peroxide treatment. Data are shown as mean  $\pm$  SE (n = 6);  $p < 0.05$  was considered significant. Values between the treatments are represented by lower case letters (a, b) between the instars are represented by upper case letters (A, B). Those not sharing the same letters are significant.

followed by centrifugation at 3500 rpm for 20 min. Alcohol evaporation was performed at 40 °C and the samples were lyophilized and stored until use.

#### LC-MS analysis of sericin protein

LC-MS analysis was performed using an API 3000 LC-MS system fitted with a turbo ion spray source and a quadrupole mass spectrometer (Perkin Elmer Sciex, Thornton, Canada). The instrument was operated in positive ion mode with a spray voltage of 5500 V and a source temperature

of 475 °C using a Phenomenex column (2540x6.6 mm) with methanol:water (3:1) as the mobile phase. Data were analysed with Analyst software version 1.4.2.

#### Lipid peroxidation level

Malondialdehyde (MDA), a product of LPO, was determined as described by Ohkawa *et al.* (1979). In brief, 200 µl of cell homogenates were added to 8.1 % SDS, vortexed and incubated for 10 min. 375 µl of 20 % acetic acid and 0.6 % thiobarbituric acid

were added to the reaction mixture and placed in a boiling water bath for 60 min. The samples were allowed to cool and 1.25 ml of a butanol:pyridine mixture (15:1, v/v) was added and centrifuged at 640 g for 5 min. Absorbance was measured at 532 nm using 1,3,3-tetramethoxy propane (TMP) as the standard. The MDA concentration was expressed as nmol/mg protein.

#### *Protein carbonyl level*

Protein carbonyl (PrC) was measured according to the method of Uchida and Stadtman (1993). 0.1 % DNPH in 2 N HCl was added to 800  $\mu$ l of cell homogenate. Samples were kept in the dark for 1 h. The protein was precipitated with 20 % trichloroacetic acid and centrifuged. The pellets were washed thrice with ethanol and ethyl acetate (1:1, v/v) and were dissolved in 2 ml of 8 M guanidine hydrochloride, and centrifuged. The supernatant was used to measure the absorbance at 365 nm and the PrC level was calculated using a molar absorption coefficient of 22,000 M<sup>-1</sup> cm<sup>-1</sup>. The results were expressed as  $\mu$ M/mg protein.

#### *Superoxide dismutase (SOD, E.C. 1.15.1.1) activity*

SOD activity was measured according to Misra and Fridovich (1972) with slight modifications. Briefly, 100  $\mu$ l of a 5 % cell homogenate was added to 880  $\mu$ l of carbonate buffer (0.5 M, pH 10.2). 20  $\mu$ l of epinephrine (30 mM in 0.05 % acetic acid) were added to the mixture and measured spectrophotometrically (model: Genova MK3, Jenway, UK) at 480 nm for 4 min. SOD activity was measured as the amount of enzyme that inhibits oxidation of epinephrine by 50 %, which is equal to 1 unit.

#### *Catalase (CAT, E.C. 1.11.1.6) activity*

Catalase was determined by the method of Aebi (1984). Briefly, 100  $\mu$ l enzyme samples with 10  $\mu$ l of absolute alcohol were incubated for 30 min at 0 °C followed by the addition of 10  $\mu$ l Triton X-100. An aliquot of 50  $\mu$ l was placed in 1.25 ml of 0.066 M H<sub>2</sub>O<sub>2</sub> in phosphate buffer and the decrease in absorbance was measured at 240 nm for 60 s in a spectrophotometer. An extinction coefficient of 43.6 M cm<sup>-1</sup> was used to determine enzyme activity and was expressed as one  $\mu$ mole of H<sub>2</sub>O<sub>2</sub> degraded/min/mg protein.

#### *Glutathione peroxidase-like enzyme (GPx, E. C. 1.11.1.9) activity*

GPx-like enzyme activity was analysed by the method of Flohe and Gunzler (1984). 50  $\mu$ l of 0.1 M phosphate buffer (pH 7.0), 100  $\mu$ l of the enzyme sample, 100  $\mu$ l glutathione reductase (0.24 units) and 100  $\mu$ l of 10 mM GSH were mixed. The mixture was pre incubated for 10 min at 37 °C followed by the addition of 100  $\mu$ l of 1.5 mM NADPH in 0.1 % NaHCO<sub>3</sub>. 50  $\mu$ l of 12 mM t-butylhydroperoxide was added to monitor the hydrogen peroxide independent concentration of NADPH for 3 min. The overall reaction was started by adding 100  $\mu$ l of pre-warmed H<sub>2</sub>O<sub>2</sub>, and the decrease in absorption at 340 nm was monitored for 5 min. GPx-like activity was expressed as  $\mu$ M NADPH oxidized/min/mg protein.

#### *Antioxidant capacity*

Antioxidant capacity was analysed by the modified method of Benzie and Strain (1996) using ascorbic acid as the standard. FRAP reagent was prepared fresh from acetate buffer (pH 3.6). 10 mM TPTZ diluted with 40 mM HCl and 20 mM ferric chloride solution at a ratio of 10:1:1 (v/v), respectively, were warmed to 37 °C prior to use. 100  $\mu$ l of cell homogenate and 3 ml of the FRAP reagent were vortexed and absorbance was measured at 593 nm at 0 min. The samples were incubated at 37 °C and absorbance was recorded after 30 min. The antioxidant capacity of the cell homogenate was expressed in  $\mu$ M Fe<sup>2+</sup> /100 mg cell mass.

#### *Fluorescent microscopy studies*

To assess the intracellular ROS levels in isolated midgut epithelial cells and hemocytes, the cells were loaded with an ROS-sensitive indicator, CM-H<sub>2</sub>DCFDA of 20  $\mu$ M for 10 min at RT. The excess stain was washed off with insect Ringer's solution. In cells, esterase cleaves CM-H<sub>2</sub>DCFDA to release CM-H<sub>2</sub>DCFH, which is converted to the fluorescent product CM-H<sub>2</sub>DCF when exposed to ROS (Xie *et al.*, 1999). CM-H<sub>2</sub>DCFDA was excited at 520 nm and the emitted light was collected at 570 nm using a fluorescent microscope (Olympus IX 71, Japan). ROS was quantified using Image Pro Express version 6.3 software from stored images.

#### *Statistical analysis*

Data are shown as the mean  $\pm$  SD of six observations. Changes between the groups were analysed by MANOVA and further tested by the Bonferroni post-hoc test using Statistical Package for Social Sciences (SPSS) software (Huberty and Olejnik, 2006) and *p* < 0.05 was considered significant. Statistically significant data are presented in the text.

## **Results**

#### *Composition of silk protein sericin*

A flow rate of 0.3 ml/min through the Phenomenex column was used in the LC-MS studies to provide information on the retention times and the endogenous concentrations of various amino acids in the silk protein sericin. The retention times from the samples were compared with synthetic standards, as shown in Figures 1A and B. Among the amino acids, serine was the most abundant, followed by glycine, aspartic acid, threonine and glutamic acid (Fig. 1C).

#### *Oxidative stress markers*

Incubation with 20  $\mu$ M hydrogen peroxide significantly increased MDA, a product of lipid peroxidation, in both midgut epithelial cells and hemocytes irrespective of the instars studied. However, the MDA level was significantly higher in IV instar larvae when compared to V instar larvae. Incubation with sericin or a combination of serine and aspartic acid for 10 min prior to hydrogen peroxide treatment inhibited hydrogen peroxide-induced lipid peroxidation in both midgut epithelial cells and hemocytes (Figs 2a, b). A significant increase in protein carbonyl content (a product of



**Table 1** Antioxidant capacity of sericin and the effect of sericin and amino acid mixture to induce antioxidant capacity in midgut epithelial cells and hemocytes of IV and V instar silkworm *B. mori* treated with hydrogen peroxide

Nature of the treatment	Midgut epithelial cells ( $\mu\text{M Fe}^{2+}$ / 100mg cell mass)		Hemocytes ( $\mu\text{M Fe}^{2+}$ / 100mg cell mass)	
	IV	V	IV	V
Control	13.6 <sup>a</sup> ±0.4	11.8 <sup>a</sup> ±0.9	15.2 <sup>a</sup> ±0.3	14.9 <sup>a</sup> ±0.3
Incubation with H <sub>2</sub> O <sub>2</sub>	8.6 <sup>a</sup> ±0.6	7.2 <sup>a</sup> ±0.5	14.5 <sup>a</sup> ±0.4	12.7 <sup>a</sup> ±0.5
Incubation with sericin prior to H <sub>2</sub> O <sub>2</sub>	23.8 <sup>b</sup> ±0.4	17.2 <sup>b</sup> ±0.7	25.3 <sup>b</sup> ±0.4	28.8 <sup>b</sup> ±0.4
Incubation with amino acids prior to H <sub>2</sub> O <sub>2</sub>	36.5 <sup>b</sup> ±0.9	17.4 <sup>b</sup> ±0.8	24.8 <sup>b</sup> ±0.8	23.3 <sup>b</sup> ±0.9
FRAP value of sericin	35.8 <sup>c</sup> ( $\mu\text{M Fe}^{2+}$ / 100mg sample) ±0.2			

Data are means  $\pm$  SE (n = 6);  $p < 0.05$  was considered significant. Values between the treatments are represented in lower cases (a, b, c). Those not sharing the same letters are significant.

protein oxidation), as a result oxidative stress, was evident following treatment with hydrogen peroxide in both in midgut epithelial cells and hemocytes (Figs 3a, b). Prior incubation with sericin and amino acid mixture significantly inhibited protein oxidation in both types of cells.

#### *Antioxidant enzymes*

Incubation of isolated midgut epithelial cells and hemocytes with 20  $\mu\text{M}$  hydrogen peroxide significantly increased SOD activity in both instars, but more so in IV instar over V instar silkworms irrespective of the treatment. Incubation with sericin and serine, aspartic acid mixture prior to hydrogen peroxide treatment inhibited the hydrogen peroxide-induced increase in enzymatic activity. SOD activity in midgut epithelial cells was significantly higher than in hemocytes per assay (Figs 4a, b).

Catalase hydrolyses hydrogen peroxide into H<sub>2</sub>O and O<sub>2</sub>. To assess the role of sericin as a probable antioxidant, isolated midgut epithelial cells and hemocytes were incubated with 20  $\mu\text{M}$  hydrogen peroxide with or without prior treatment with sericin. CAT activity was found to be significantly higher in IV instar larvae than in V instar larvae of both tissues studied. Incubation with hydrogen peroxide significantly increased CAT activity in midgut epithelial cells as well as in hemocytes. Cells treated with sericin or mixture of aspartic acid and serine before the induction of oxidative stress did not elicit a rise in CAT activity under stress (Figs 5a, b).

A significant increase in GPx-like activity was noted in the context of oxidative stress in both midgut epithelial cells and hemocytes. Prior treatment with sericin or the mixture of aspartic acid and serine prevented the oxidative stress-induced increase in enzymatic activity in both tissues (Figs


6a, b). The instar-dependent decrease in antioxidant enzymes such as SOD and CAT was not observed for the GPx-like enzyme.

#### *Antioxidant capacity of silk protein and a mixture of serine and aspartic acid*

The antioxidant capacity of the silk protein sericin was determined by a FRAP assay using ascorbic acid as the standard. The residual antioxidant capacity was found to be 13.6 and 11.8  $\mu\text{M Fe}^{2+}$ /100 mg cell mass in midgut epithelial cells, while in hemocytes it was found to be 14.5 and 14.9  $\mu\text{M Fe}^{2+}$ /100 mg cell mass in IV and V instar silkworms, respectively. Cells incubated with sericin and also with combination of serine/aspartic acid prior to hydrogen peroxide treatment showed significantly increased antioxidant capacity by 0.72 fold and 0.45 fold in midgut epithelial cells, while in hemocytes the increase was 0.74 fold and 1.6 fold in IV and V instar silkworms, respectively (Table 1).

#### *Hydrogen peroxide-induced increase in reactive oxygen species*

H<sub>2</sub>DCFDA staining was used to study the H<sub>2</sub>O<sub>2</sub>-induced change in reactive oxygen species in midgut epithelial cells and hemocytes under a fluorescent microscope. Under phase contrast, blebbing of the plasma membrane and swelling of the cells were observed in cells treated with H<sub>2</sub>O<sub>2</sub> at concentrations above 20  $\mu\text{M}$ . In contrast, pre incubation with 28 ng/ml sericin or with mixture of 0.5 mM serine and aspartic acid inhibited H<sub>2</sub>O<sub>2</sub>-induced morphological changes in both types of cells studied (Figs 7, 8). H<sub>2</sub>DCFDA stained cells revealed a higher intensity of fluorescence, indicating an increase in ROS upon exposure to 20  $\mu\text{M}$  H<sub>2</sub>O<sub>2</sub> (Supplementary Figs S1, S2).



**Fig. 7** Effect of sericin and amino acid mixture on the hydrogen peroxide-induced increase in reactive oxygen species in the midgut epithelial cells of the silkworm *B. mori*. (a, b) Normal midgut epithelial cells, (c, d) hydrogen peroxide treated cells, (e, f) cells incubated with sericin prior to hydrogen peroxide treatment and (g, h) cells incubated with amino acid mixture prior to hydrogen peroxide treatment.


## Discussion

Hydrogen peroxide is membrane permeable, diffusible, less reactive and longer-lived than  $\text{OH}^-$  or  $\text{O}_2^-$  (Stone and Yang, 2006). The physiological range of the intracellular hydrogen peroxide concentration appears to be remarkably conserved in different forms of life (Meller, 2000). Notably, among the ROS, hydrogen peroxide is the only species that is generated by several specific enzymes, which suggests that the intracellular concentration of hydrogen peroxide is tightly regulated and may serve several specific cellular functions. It potentiates antioxidant mechanisms; however, if produced in excess, it has deleterious consequences (Veal *et al.*, 2007). It has been reported to be less effective as an intracellular signalling molecule when added exogenously than endogenously produced hydrogen peroxide (Choi *et al.*, 2005). In the present study, 20  $\mu\text{M}$  hydrogen peroxide, when applied exogenously, caused close to 50 % mortality in midgut epithelial cells and hemocytes with a concomitant increase in ROS, as reflected in the fluorescent microscopy study.

Hydrogen peroxide can cause oxidative modifications in proteins (Stadtman, 1992; Dalle-Donne *et al.*, 2002) and peroxidise unsaturated lipids in the cell membrane (Fridovich, 1978). In our experiments, exposure of midgut epithelial cells and hemocytes to 20  $\mu\text{M}$  hydrogen peroxide significantly increased the level of lipid peroxidation products, *i.e.*, MDA; prior treatment with sericin and with a mixture of amino acids was found to inhibit hydrogen peroxide-mediated effects in the cell

membrane. Sericin was found to suppress lipid peroxidation in rat brain homogenates (Kato *et al.*, 1998) and several amino acids have the property of reducing the lipid peroxidation (Movahedian *et al.*, 2006; Selvaraju and Subbashinidevi, 2011). In our present study a mixture of serine and aspartic acid have similar effect of reducing the lipid peroxidation in midgut epithelial cells and hemocytes of silkworm. However, there has been no evidence of a role of sericin as an inhibitor of protein oxidation, as protein thiols were found to inhibit protein hydroperoxide formation (Platt and Gieseg, 2003). The current study clearly indicates that sericin functions as an inhibitor of protein oxidation.

Up-regulation of antioxidant enzymes (AOEs) affords protection against ROS (Fulda *et al.*, 2010). In current study, exogenous treatment with  $\text{H}_2\text{O}_2$  resulted in a significant increase in the antioxidant enzymes SOD, CAT and GPx in the midgut epithelial cells and hemocytes of the silkworm prior to the treatment with sericin or with combination of serine and aspartic acid. Such an increase in SOD (Tsai *et al.*, 2011), CAT (Dash *et al.*, 2008a) and GPx (Caldinin *et al.*, 1998) upon exposure to hydrogen peroxide was observed in fibroblasts. An increase in SOD, CAT and GPx were reported in silkworms infected with Bm NPV and reared on an artificial diet containing  $\text{CeCl}_3$  (Li *et al.*, 2011). The increased AOEs were explained by the fact that  $\text{CeCl}_3$  acts as scavenger of hydroxyl radicals in pathogenic states. Also, treatment with 4-methylumbelliferone, a model drug, up-regulated the activities of CAT and GPx against oxidative stress in fat body cells of the silkworm (Fang *et al.*, 2014). Our



**Fig. 8** Effect of sericin and amino acid mixture on the hydrogen peroxide-induced increase in reactive oxygen species in hemocytes of the silkworm *B. mori*. (a - f) Normal hemocytes, (g - l) hydrogen peroxide treated hemocytes, (m - r) hemocytes incubated with sericin prior to hydrogen peroxide treatment and (s - x) hemocytes incubated with amino acid mixture prior to hydrogen peroxide treatment.

results show a significant increase in AOE activities upon hydrogen peroxide treatment. A possible explanation for increased AOE activities seen in  $H_2O_2$  treated cells, but not in cells pre-treated with sericin and amino acids, is that the increased antioxidant defence reduces more of the generated ROS, such that the ROS signal is not generated or ROS-mediated damage may not occur. In our study, the

instar-dependent increase in antioxidant enzymes such as SOD and CAT was not observed for GPx. Similar variations in the antioxidant system as an ontogenic effect has been reported in the beetle *Tenebrio molitor* (Gulevsky *et al.*, 2006a, b). In the present study prior treatment with sericin and a mixture of amino acids reduced the AOE activities that were evoked by hydrogen peroxide exposure

alone. Contradictory findings have been reported regarding the effect of sericin and amino acids on AOE's. Sericin elevated SOD, CAT and GPx activity in rat livers treated with alcohol (Li *et al.*, 2008, Selvaraju and Subhashinidevi, 2011), while sericin had no effect on SOD in fibroblasts (Dash *et al.*, 2008a). It is possible that sericin and amino acids may participate in detoxifying harmful radicals rather than potentiating the activity of AOE's, thereby ameliorating oxidative stress-induced cell damage.

The ability of a compound to reduce  $Fe^{3+}$  and  $Fe^{2+}$  serves as an important indicator of its potential antioxidant power (Benzine and Strain, 1996; Yen *et al.*, 1999). In the present study, treatment with 20  $\mu M$   $H_2O_2$  failed to evoke a change in antioxidant capacity in midgut epithelial cells and hemocytes. Perhaps, the elevated antioxidant status upon oxidative stress might have been utilized by the cells to overcome the radicals in the initial period. Nonetheless, prior incubation with sericin and also with combination of serine and aspartic acid significantly increased the FRAP values in both tissues subjected to oxidative stress.

Exposure to hydrogen peroxide in excess of 20  $\mu M$  induced changes such as swelling and blebbing in both tissues studied. Similar changes in morphology were also observed in fibroblasts (Dash *et al.*, 2008a) upon hydrogen peroxide treatment. Furthermore, an increased hydrogen peroxide concentration was reported in keratinocytes irradiated with UVB (Dash *et al.*, 2008b). Changes in the intracellular concentration of peroxy radicals were evident from our fluorescent microscopy studies. The cells labelled with  $H_2DCFDA$  showed a higher intensity upon treatment with hydrogen peroxide, and prior incubation with sericin or with amino acids obliterated the accumulation of peroxy radicals induced by hydrogen peroxide. Our observations are in concurrence with earlier work of Fan *et al.* (2009). It was evident from our study that hydrogen peroxide induces peroxide radical generation and sericin and mixture of serine and aspartic acid may act as a scavenger of free radicals.

The ability of proteins to scavenge free radicals has been reported in several systems (Kong and Xiong, 2006; Sakanaka and Tachibana, 2006; Elias *et al.*, 2008). The results of our study substantiate earlier studies and provide evidence for free radical scavenging by sericin and LCMS studies clearly indicate that sericin is rich in serine, followed by aspartic acid, glycine, threonine and glutamine. Evidence for protective role of amino acids against oxidative stress is well documented in several experimental models (Patterson *et al.*, 2003; Movahedian *et al.*, 2006; Selvaraju and Subhashinidevi, 2011). In the present study a mixture of serine and aspartic acid 0.5 mM each significantly scavenged ROS thereby prevented protein oxidation and lipid peroxidation without altering the antioxidant enzymes. Serine acts as an antioxidant (Maralani *et al.*, 2012), efficient nucleophilic reagent and particularly play a vital role in enzyme catalyst (Anderson *et al.*, 1961). Whereas, aspartic acid is known as an antioxidant (Liu *et al.*, 2004) and also undergo transamination to function directly or indirectly as an antioxidant.

Although L-serine or L aspartic acid alone scavenged ROS to certain extent, a mixture of the amino acids significantly decreased the ROS. Perhaps synergistic effect of serine and aspartic acid may be responsible for the scavenging effects however, such synergistic effects were not found on antioxidation among the compounds used in combination (Wu *et al.*, 2003b). Our present result further support antioxidant properties of L-serine (Kitazawa *et al.*, 2005) and L- aspartic acid (Chen and Nawar, 1991; Wang and Xiong, 2005). Earlier studies on antioxidant property of sericin were appertained to phenol compound (Prasong, 2011) and flavonoids (Devi *et al.*, 2011). However, our results on a mixture of serine, aspartic acid have clearly indicated the scavenging of ROS by these amino acids without altering the antioxidant enzymes. Here we report for the first time that the antioxidant property of sericin as scavenger of free radicals rather than enhancing the activities of AOE's. We suggest that fortification of silkworm diet with a mixture of serine and aspartic acid can alleviate the effects of ROS thereby larvae can withstand environmental born stressors to certain extent. Although, the mechanism of protection offered is through scavenging of free radicals by major constituent hydroxyl amino acids, suboptimal hydrogen peroxide also helped to enhance the endogenous antioxidant machinery as defense against initial ROS signaling or ROS damage.

#### Acknowledgments


This work was supported in part by the Promotion of University Research and Scientific Excellence (PURSE), Department of Science and Technology, New Delhi (No.SR/S9/Z-23/2010138). Fellowship offered by Council for Scientific and Industrial Research (CSIR), New Delhi, to one of the author (ASM) is greatly acknowledged.

#### References


- Aebi H. Catalase *in vitro*. Methods Enzymol.. 105: 121-126, 1984.
- Anderson BM, Cordes EH, Jencks WP. Reactivity and reactions of the serine hydroxyl group and of o-acyl serines. J. Biol. Chem. 236: 455-463, 1961.
- Aramwit P, Sangcakul A. The effects of sericin cream on wound healing in rats. Biosci. Biotechnol. Biochem. 71: 2473-2477, 2007.
- Atmaca G. Antioxidant effects of sulfur- containing amino acids. Yonsei Med. J. 45: 776-788, 2004.
- Balsano C, Alisi A. Antioxidant effect of natural bioactive compounds. Curr. Pharm. Des. 15: 3063-3073, 2009.
- Benzie IF, Strain JJ. The Ferric reducing ability of plasma (FRAP) as a measure of "Antioxidant power": The FRAP assay. Anal. Biochem. 239: 70-76, 1996.
- Caldini R, Chevanne M, Mocali A, Tombaccini D, Paoletti F. Premature induction of ageing in sub lethally  $H_2O_2$ - treated young MRC5 fibroblasts correlates with increased glutathione peroxidase levels and resistance to DNA breakage. Mech. Ageing Develop. 105: 137-150, 1998.
- Chen ZY, Nawar WW. The role of amino acids in the autoxidation of milk fat. J. Amer. Oil Chem. Soc. 68: 47-50, 1991.

- Choi MH, Lee IK, Kim GW. Regulation of PDGF signaling and vascular remodeling by peroxiredoxin II. *Nature* 435: 347-353, 2005.
- Dalle Donne I, Rossi R, Giustarini D, Milazani A. Protein carbonyl groups as biomarkers of oxidative stress. *Clin. Chim. Acta* 329: 23-138, 2002.
- Dash R, Acharya C, Bindu PC, Kundu SC. Antioxidant potential of silk protein sericin against hydrogen peroxide-induced oxidative stress in skin fibroblasts. *BMB Rep.* 41: 236-241, 2008a.
- Dash R, Mandal M, Ghosh SK, Kundu SC. Silk sericin protein of tropical tasar silkworm inhibits UVB induced apoptosis in human skin keratinocytes. *Mol. Cell Biochem.* 311: 111-119, 2008b.
- Davies KJA. The broad spectrum of responses to oxidants in proliferating cells: A new paradigm for oxidative stress. *IUBMB Life* 48: 41-47, 1999.
- Devi R, Deori M, Devi D. Evaluation of antioxidant activities of silk protein sericin secreted by silkworm *Antheraea assamensis* (Lepidoptera: Saturniidae). *J. Pharm. Res.* 4: 4688-4691, 2011.
- Elias RJ, Kellerby S, Decker A. Antioxidant activity of proteins and peptides. *Crit. Rev. Food Sci. Nutr.* 48: 430-441, 2008.
- Fan JB, Wu LP, Mao XY, Ren FZ. Antioxidant activities of silk sericin from silkworm *Bombyx mori*. *J. Food Chem.* 33: 74-88, 2009.
- Fang Y, Wang H, Zhu W, Wang L, Liu H, He Y, Xu X, Yin W, Sima Y, Xu S. Antioxidative capacity in the fat body of *Bombyx mori* increased following oral administration of 4-mrthylumbelliferone. *Comp. Biochem. Physiol.* 159C: 31-37, 2014.
- Felton GW, Summers CB. Antioxidant systems in insects. *Arch. Insect Biochem. Physiol.* 29: 187-197, 1995.
- Flohe L, Gunzler WA. Assays of glutathione peroxidase. *Methods Enzymol.* 105: 115-121, 1984.
- Fridovich I. 1978. The biology of oxygen radicals. *Science* 201: 875-879, 1984.
- Fulda S, Gorman AM, Hori O, Samali A. Cellular stress responses: cell survival and cell death. *Int. J. Cell Biol.* 2010:214074, 2010, doi: 10.1155/2010/214074.
- Gulevsky AK, Grishchenkova YA, Relina LI. Antioxidant system in the darkling beetle (*Tenebrio molitor*) in ontogenesis. *Ukr Biokhim Zhm.* 78: 86-90, 2006a.
- Gulevsky AK, Relina LI, Grishchenkova YA. Variation of the antioxidant system during development of the cold-tolerant beetle, *Tenebrio molitor*. *Cryo Lett.* 27: 283-290, 2006b.
- Huberty CJ, Olejnik S. Applied MANOVA and discriminate analysis. John Wiley and Sons, New Jersey, 2006.
- Imlay JA. Cellular Defenses against Superoxide and Hydrogen Peroxide. *Annu. Rev. Biochem.* 77: 755-776, 2008.
- Kashiwagi A, Kashiwagi K, Takase M, Hanada H, Nakamura M. Comparison of catalase in diploids and haploid *Ranarugosa* using heat and chemical inactivation techniques. *Comp. Biochem. Physiol.* 118B: 499-503, 1997.
- Kato N, Sato S, Yamanaka A, Yamada H, Fuwa N, Nomura M. Silk protein, sericin, inhibits lipid peroxidation and tyrosine activity. *Biosci. Biotechnol. Biochem.* 62: 145-147, 1998.
- Khudaiderdier HA. Synthesis of copolymer processing an anticoagulant action. *Chem. Nat. Comp.* 33: 603-604, 1997.
- Kitazawa M, Ishitsuka Y, Kobayashi M, Nakano T, Iwasaki K, Sakamoto K, *et al.* Protective effects of an antioxidant derived from serine and vitamin B6 on skin photoaging in hairless mice. *Photochem. Photobiol.* 81: 970-974, 2005.
- Kong B, Xiong XL. Antioxidant activity of zein hydrolysates in a liposome system and the possible mode of action. *J. Agric. Food Chem.* 54: 6059-6068, 2006.
- Krishnan N, Kodrik D. Antioxidant enzymes in *Spodoptera littoralis* (Biosduval): are they enhanced to protect gut tissue during oxidative stress. *J. Insect Physiol.* 52: 11-20, 2006.
- Kwang YC, Jae YM, Yong WL, Kwang GL, Joo HY, Hae YK. Preparation of self-assembled silk sericin nanoparticles. *Int. J. Biol. Macromol.* 32: 36-42, 2003.
- Li B, Xie Y, Cheng Z, Cheng J, Hu R, Cui Y, *et al.* Effects of added CeCl<sub>3</sub> on resistance of fifth-instar larvae of silkworm to *Bombyx mori* Nucleopolyhedrovirus Infection. *Biol. Trace Elem. Res.* 146: 318-24, 2012.
- Li YG, Ji DF, Chen S, Hu GY. Protective effects of sericin protein on alcohol-mediated liver damage. *Alcohol Alcoholism* 43: 246-253, 2008.
- Liu DZ, Lin YS, Hou WC. Monohydrates of aspartic acid and glutamic acid exhibit antioxidant and angiotensin converting enzyme inhibitory activities. *J. Agric. Food Chem.* 52: 2386- 2390, 2004.
- Maiorino M, Scapin M, Ursini F, Biasolo M, Bosello V, Flohe L. Distinct promoters determine alternative transcription of *gpx-4* into phospholipid-hydroperoxide glutathione peroxidase variants. *J. Biol. Chem.* 278: 34286-34290, 2003.
- Maralani MN, Movahedian A, Javanmard ShH. Antioxidant and cytoprotective effects of L-serine on human endothelial cells. *Res. Pharm. Sci.* 7: 209-215, 2012.
- Medina-Navarro R, Duran-Reye G, Diaz-Flores M, Vilar-Rojas C. Protein antioxidant response to the stress and the relationship between molecular structure and antioxidant function. *PLOS One* 5: e8971, 2010.
- Meller S. Sensitive and non-enzymatic measurement of hydrogen peroxide in biological system. *Free Radic. Biol. Med.* 29: 410-415, 2000.
- Micheal AS, Subramanyam MVV. Antioxidant enzymes as defense mechanism against oxidative stress in midgut tissue and haemocytes of silkworm *Bombyx mori* larvae subjected to various stressors. *Arch. Insect Biochem. Physiol.* 84: 222-234, 2013.
- Misra HP, Fridovich I. The role of superoxide anion in the autoxidation of epinephrine and a simple assay of superoxide dismutase. *J. Biol. Chem.* 247: 3170-3175, 1972.

- Movahedian A, Naderi GA, Dashti GR, Asgary S, Zadoosh F. Antioxidant effects of L-serine against fatty streak formation in hypercholesterolemic animals. *ARYA Atherosclerosis J.* 2: 126-129, 2006.
- Nagasaka R, Okamoto N, Ushio H. Partial oxidative stress perturbs membrane permeability and fluidity of fish nucleated red blood cells. *Comp. Biochem. Physiol.* 139C: 259-266, 2004.
- Ohkawa H, Ohishi M, Yagi K. Assay for lipid peroxidation in animal tissues by thiobarbituric acid reaction. *Anal. Biochem.* 95: 351-358, 1979.
- Okazaki Y, Kakehi S, Xu Y, Tsujimoto K, Sasaki M, Ogawa H, *et al.* Consumption of sericin reduces serum lipids, ameliorates glucose tolerance and elevates serum adiponectin in rats fed on a high fat diet. *Biosci. Biotechnol. Biochem.* 74: 1534-1538, 2010.
- Padamwar MN, Pawar AP. Silk sericin and its application: a review. *J. Sci. Industrial Res.* 63: 323-329, 2004.
- Patel RJ, Modasiya MK. Sericin: Pharmaceutical applications. *Int. J. Res. Pharm. Biomed. Sci.* 2: 913-917, 2011.
- Patterson RA, Lamb DJ, Leake DS. Mechanisms by which cysteine can inhibit or promote the oxidation of low density lipoprotein by copper. *Atherosclerosis J.* 169: 87-94, 2003.
- Platt AA, Gieseg SP. Inhibition of protein hydroperoxide formation by protein thiols. *Redox Rep.* 8: 81-86, 2003.
- Prasong S. Screening of antioxidant activity of some *Samia ricini* (Eri) silks: comparison with *Bombyx mori*. *J. Biol. Sci.* 11: 336-339, 2011.
- Sakanaka S, Tachibana Y. Active oxygen scavenging activity of egg yolk protein hydrolysates and their effects on lipid oxidation in beef and tuna homogenates. *Food Chem.* 95: 243-249, 2006.
- Sasaki M, Kato N, Watanabe H, Yamada H. Silk protein, sericin, suppresses colon carcinogenesis induced by 1, 2-dimethylhydrazine in mice. *Oncol. Rep.* 7: 1049-1101, 2000.
- Selvaraju R, Subbashinidevi K. Impact of glycine on antioxidant defence system in rats with alcohol induced liver injury. *Int. J. Res. Pharm. Biomed. Sci.* 2: 1314-1320, 2011.
- Stadtman ER. Protein oxidation and ageing. *Science* 257: 1220-1224, 1992.
- Stone JR, Yang S. Hydrogen peroxide: a signaling messenger. *Antioxidant Redox signaling* 8: 243-270, 2006.
- Tsai CC, Wu SB, Cheng CY, Kao SC, Kau HC, Lee SM, *et al.* Increased response to oxidative stress challenge in Grave's ophthalmopathy orbital fibroblasts. *Mol. Vision* 17: 2782-2788, 2011.
- Tsujimoto K, Takagi H, Takahashi M, Yamada H, Nakamori S. Cryoprotective effects of the serine rich repetitive sequence in silk protein sericin. *J. Biol. Chem.* 276: 979-986, 2001.
- Uchida K, Stadtman ER. Covalent modification of 4-hydroxynonenal to glyceraldehyde-3-phosphate. *J. Biol. Chem.* 268: 6388-6393, 1993.
- Veal EA, Day AM, Morgan BA. Hydrogen peroxide sensing and signaling. *Mol. Cell* 26: 1-14, 2007.
- Vyjayanthi N, Subramanyam M. Effect of Fenvalerate - 20EC on sericigenous insect: I. Food utilization in the late age larva of silkworm *B. mori*. *Ecotoxicol. Environ. Saf.* 53: 206-211, 2002a.
- Vyjayanthi N, Subramanyam M. Effect of Fenvalerate - 20EC on sericigenous insect: II. Digestive enzymes in the nutritive physiology of silkworm *B. mori*. *Ecotoxicol. Environ. Saf.* 53: 212-220, 2002b.
- Wang LL, Xiong YL. Inhibition of lipid oxidation in cooked beef patties by hydrolyzed potato protein is related to its reducing and radical scavenging ability. *J. Agric. Food Chem.* 53: 9186-9192, 2005.
- Wu HC, Chen HM, Shiau CY. Free amino acids and peptides as related to antioxidant properties in protein hydrolysates of mackerel (*Scomber austriasicus*). *Food Res. Int.* 36: 949-957, 2003a.
- Wu HC, Shiau CY, Chen HM, Chiou TK. Antioxidant activities of carnosine, anserine, some free amino acids and their combination. *J. Food Drug Anal.* 11: 148-153, 2003b.
- Wu JH, Wang Z, Xu SY. Preparation and characterization of sericin powder extracted from silk industry waste water. *Food Chem.* 103: 1255-1262, 2007.
- Xie Z, Kometiani P, Liu J, Li J, Shapiro JI, Askari A. Intracellular reactive oxygen species mediates the linkage of Na<sup>+</sup>/K<sup>+</sup>-ATPase to hypertrophy and its marker genes in cardiomyocytes. *J. Biol. Chem.* 274: 19323-19328, 1999.
- Yen GC, Chen HY, Lee CE. Measurement of antioxidant activity in metal ion-reduced lipid peroxidation systems. *J. Sci. Food Agric.* 79: 1213-1217, 1999.



**Fig. S1** Effect of sericin and amino acid mixture on the hydrogen peroxide induced increase in reactive oxygen species in hemocytes (a) prohemocytes, (b) plasmocytes and (c) granular hemocytes of the silkworm *B. mori*. Data are shown as mean  $\pm$  SE (n = 3);  $p < 0.05$  was considered significant. Values between the stressors are represented by lower case letters (a, b). Those not sharing the same letters are significant.



**Fig. S2** Effect of sericin and amino acid mixture on the hydrogen peroxide induced increase in reactive oxygen species in midgut epithelial cells of the silkworm *B. mori*. Data are shown as mean  $\pm$  SE ( $n = 3$ );  $p < 0.05$  was considered significant. Values between the stressors are represented by lower case letters (a, b). Those not sharing the same letters are significant.