

Catalysis and mechanistic studies of ruthenium and osmium on synthesis of anthranilic acids

P. Karthikeyan^a, Rajenahally V. Jagadeesh^{a*}, Y. Sree Sandhya^a, Puttaswamy^b, P. Nithya^a, S. Senthil Kumar^a and P.R. Bhagat^a



Ruthenium, osmium and ruthenium + osmium catalyzed synthetic methodology was developed for the synthesis of anthranilic acids from indoles in good to excellent yields using bromamine-B in alkaline acetonitrile–water (1 : 1) at 313 K. Detailed catalysis studies of ruthenium, osmium and the mixture of both were carried out for the synthetic reactions. The positive synergistic catalytic activity of Ru(III) + Os(VIII) was observed to a large extent with the activity greater than the sum of their separate catalytic activities. Detailed kinetic and mechanistic investigations for each catalyzed reactions were carried out. The kinetic pattern and mechanistic picture of each catalyzed reaction were found to be different for each catalyst and to obey the underlying rate laws:

$$\text{rate} = k[\text{BAB}]_t[\text{Indole}][\text{Ru(III)}]^x[\text{OH}^-]^y$$

$$\text{rate} = k[\text{BAB}]_t[\text{Indole}][\text{Os(VIII)}][\text{OH}^-]^y$$

$$\text{rate} = k[\text{BAB}]_t[\text{Indole}]^0[\text{Ru(III)} + \text{Os(VIII)}][\text{OH}^-]^y$$

where, $x, y < 1$. The reactions were studied at different temperatures and the activation parameters were evaluated for each catalyzed reaction. Under the identical set of experimental conditions, the kinetics of all the three catalyzed reactions were compared with uncatalyzed reactions, revealing that the catalyzed reactions were 6- to 42-fold faster. The catalytic efficiency of aforementioned catalysts followed the order: Ru(III) + Os(VIII) > Os(VIII) > Ru(III). This trend may be attributed to the different d-electronic configuration of the catalysts. The proposed mechanisms and the rigorous kinetic models derived give results that fit well with the experimental data in each catalyzed reaction. Copyright © 2010 John Wiley & Sons, Ltd.

Supporting information may be found in the online version of this article.

Keywords: Ru(III)/Os(VIII)/Ru(III) + Os(VIII) catalysis; anthranilic acids; indoles; reaction mechanism; bromamine-B

Introduction

The importance of anthranilic acids resides in their well established anti-inflammatory activity and their potential uses as intermediates for the production of dyes, pigments and saccharin. Anthranilic acid and its ester derivatives are used in preparing perfumes, pharmaceuticals and UV-absorber as well as corrosion inhibitors for metals and mould inhibitors in soy sauce. Thus, many derivatives of anthranilic acids have been synthesized with the objective of discovering new pharmacological agents.^[1–4] Because of the versatile properties and pharmaceutical applications of anthranilic acids, synthesis of anthranilic acids is an important task for exploring convenient and efficient methodologies. There are several reports available in the literature about the synthesis of anthranilic acids with different approaches.^[4,5] However, there is still a need to introduce safer and more efficient reagents including favorable methods for the synthesis of anthranilic acids.

Synthetic methodology, as the building block of organic synthesis, continuously seeks new reagents, better reaction conditions and more efficient and selective methods. In this regard, a large group of compounds entitled sodium *N*-haloarenesulfonamidates (organic haloamines) are widely used in fine organic synthesis.^[6,7] *N*-haloamines act as good oxidants and reagents both in alkaline and acidic media, and have been widely used for the oxidation and synthesis of variety of organic and bio molecules.^[7–9]


Bromamine-B (C₆H₅SO₂NBrNa.1.5H₂O or PhSO₂NBrNa or BAB), the bromine derivative of sulfonamide, is gaining importance as an oxidant and reagent for the synthesis of a variety of organic molecules. Although the mechanistic aspects of many of haloamine reactions have been well documented,^[9–12] similar studies on bromine analogs are sparse. In view of these facts, there is a considerable scope for the study with BAB to achieve a better insight into the speciation of BAB reaction models and to understand its redox chemistry in solutions.

In recent times, studies on the use of transition metal ions as catalysts in many synthetic and redox reactions have been gaining interest.^[13–15] The catalytic activities are due to the existence of variable oxidation states, as a consequence of partially filled d or f orbitals. Their ability to form both σ and π bonds with other moieties or ligands is one of the chief facts for imparting catalytic

* Correspondence to: Rajenahally V. Jagadeesh, Division of Organic Chemistry, School of Advanced Sciences, VIT University, Vellore-632014, India. E-mail: rvjdeesh@yahoo.com

a Division of Organic Chemistry, School of Advanced Sciences, VIT University, Vellore-632014, India

b Department of Chemistry, Bangalore University, Central College Campus, Bangalore-560001, India



Scheme 1. Catalyzed synthesis of anthranilic acids.

properties to transition metals as well as their complexes. Most of the d-block elements show characteristic interligand migration reactions and such a process forms one of the most important types of reactions in homogeneous catalysis. Ruthenium (III) chloride [Ru(III)] and Os(VIII) oxide [Os(VIII)] have been widely used as homogeneous catalysts in various redox reactions.^[15–19] The catalytic effect of Ru(III) and Os(VIII) on various reactions is well known, but sparse information is available about Ru(III) + Os(VIII) mixed catalyst. Hence in the present work, we used Ru(III) and Os(VIII), both alone and in mixtures as a catalyst. The main objective of the present study is the development of catalytic methodology for the synthesis of anthranilic acids and the exploration of catalytic efficiency and mechanism of Ru(III), Os(VIII) and Ru(III) + Os(VIII) mixture and also synergism of Ru(III) + Os(VIII) catalyst on the title reaction.

The wide range of applications of anthranilic acids and the usefulness of transition metals catalysis in organic reactions instigated us to carry out the title reaction to develop a protocol for the synthesis of anthranilic acids from indoles with an interest in studying the reaction mechanism and kinetics. By keeping above points in mind, we report herein a new and simple method for the preparation of anthranilic acids from indoles using bromamine-B and ruthenium/osmium catalyst (Scheme 1).

Experimental

Materials

Melting points were determined on X-4 apparatus and are uncorrected. IR spectra were obtained using a Shimadzu FTIR-8900 spectrometer. Mass spectrometry data were obtained on a 17A Shimadzu gas chromatograph with a QP-5050A Shimadzu mass spectrometer. The mass spectrum was obtained using the electron impact ionization technique. Indole and substituted indoles (Lancaster and Sigma Aldrich), were of acceptable grade of purity and used as received. Aqueous solutions of the indoles were prepared by dissolution in aqueous acetonitrile (1 : 1 water–acetonitrile) and were employed for the kinetic study. A 0.1

M stock solutions of RuCl_3 (Merck) was prepared in 0.01 mol dm^{-3} HCl in order to avoid the hydrolysis of RuCl_3 in water. The 0.1 M OsO_4 (Merck) stock solution was prepared in 0.01 mol dm^{-3} NaOH to retain the stable oxidation state (+8) of Os without any formation of aquo complex with water. Allowances were made for the amount of NaOH and HCl present in the catalyst solutions while preparing solutions for kinetic runs. Solvent isotope studies were made with D_2O (99.4%) supplied by Bhabha Atomic Research Center, Mumbai, India. Reagent-grade chemicals and double-distilled water were used throughout. Bromamine-B was prepared by the known procedure.^[16]

Preparation of Bromamine-B


Bromamine-B was prepared^[16] by the partial debromination of dibromamine-B (DBB), which was obtained as follows. Pure chlorine was bubbled through an aqueous solution of chloramine-B (30 g in 560 ml of water) and liquid bromine (6 ml) was added drop-wise with constant stirring. The resulting yellow precipitate of dibromamine-B (DBB) was thoroughly washed with water, filtered under suction and dried in a desiccator. Dibromamine-B (31.5 g) was digested in batches with constant stirring in 50 ml of 4 mol dm^{-3} NaOH. The mass obtained was cooled in ice, filtered under suction and the product (BAB) was dried over anhydrous calcium chloride. The purity of bromamine-B was assayed iodometrically to determine the active halogen content. Aqueous solutions of the oxidant were standardized by iodometric procedure and preserved in brown bottles to prevent photochemical deterioration. Bromamine-B was confirmed by mass spectral analysis.

Reaction Stoichiometry

Reaction mixtures containing varying ratios of BAB to indoles in the presence of $2.0 \times 10^{-3} \text{ mol dm}^{-3}$ NaOH and 2.0×10^{-6} catalyst were equilibrated at 313 K for 24 h. Determination of unreacted BAB in reaction mixture showed that 1 mole of indole consumed 3 moles of BAB in all the three catalyzed reactions, confirming the stoichiometry in Scheme 2.

















Synthesis of Anthranilic Acids

To a stirred solution of indoles (10 mmol), bromamine-B (30 mmol) in alkaline (20 mmol NaOH) acetonitrile–water (1 : 1) mixture (20 ml), catalyst (2 mmol) was added and the mixture was heated at 313 K for 3–6 h. The reaction progress was monitored by TLC. After completion of the reaction, the reduction product of bromamine-B, benzenesulfonamide (PhSO_2NH_2) was extracted



Scheme 2. Stoichiometric equation.

Table 1. Synthesis of anthranilic acids from indoles

Entry	Substrate	Product	Reaction time (hr. minutes)	Yield (%)	mp (°C)
1			5.00	95	145 (146–148)
2			4.30	96	177 (178–179)
3			4.20	96	237 (235–236)
4			5.30	94	213 (211–216)
5			4.0	96	171 (172)
6			5.30	95	210 (210)
7			5.35	92	335 (336)
8			5.30	92	180 (183–184)

mp given in paranthesis refer to authentic samples

with ethyl acetate, identified by TLC and confirmed by mass spectral analysis. The aqueous part of the reaction mixture was neutralized with acid, the solvent was evaporated under reduced pressure and the residue was dissolved in dichloromethane. The dichloromethane layer was washed twice with water and dried over sodium sulfate. The solvent was evaporated under reduced pressure and the residue thus obtained was purified by passing through a short silica gel column using dichloromethane as eluent. Evaporation of the solvent yielded anthranilic acids around 95% yield. Alternatively, after extracting out benzenesulfonamide, the aqueous layer was neutralized with acid and the anthranilic acids are estimated as their zinc anthranilates. The procedure for the estimation is as follows: anthranilate was precipitated as its zinc salt.^[20] To the reaction mixture, a calculated volume of 1 mol dm⁻³ HCl was added, followed by 10 ml of pH 5.0 buffer and 10 ml of 1% zinc sulfate. The precipitate formed was filtered, dried at 105–110 °C and weighed, and the recovery of (C₇H₆O₂N)₂ Zn was found to be around 95%. The reaction times and yields are given in Table 1. The products were identified by TLC and melting point

by comparison with authentic samples. Further, the compounds were confirmed by mass spectral analysis.

Kinetic Measurements

The detailed kinetic experiments were made with respect to indole as model compounds. The reactions were carried out under pseudo first-order conditions with a known excess of [Indole]₀ over [BAB]₀ at 313 K. The reaction was carried out in stoppered Pyrex boiling tubes whose outer surfaces were coated black to eliminate photochemical effects. For each run, requisite amounts of solutions of indole, NaOH, catalyst and aqueous acetonitrile (1 : 1 water–acetonitrile; to keep the total volume constant for all runs) were introduced into the tube and thermostated at 313 K until thermal equilibrium was attained. A measured amount of BAB solution, also thermostated at the same temperature, was rapidly added with stirring to the mixture in the tube. The progress of the reaction was monitored by the iodometric determination of unreacted BAB in aliquots (5 ml each) of the reaction mixture

withdrawn at different intervals of time. The course of the reaction was studied for at least two half-lives. The pseudo first-order rate constants (k') calculated from the linear plots of $\log [\text{BAB}]$ vs time were reproducible within $\pm 5\%$.

Results

Synthesis of anthranilic acid and substituted anthranilic acids was achieved using catalytic amounts of ruthenium, osmium and a mixture of both in acetonitrile–water (1 : 1) at 40°C by bromamine-B with 1 : 3 indole–bromamine-B ratio in the presence of alkali. The products and the yields were summarized in Table 1. In general substrates containing electron-donating moieties were found to be slightly more reactive and required shorter reaction times compared to substrates containing electron-withdrawing groups.

The synthesis of anthranilic acids from indoles proceeds with the formation of isatins as the intermediates. Indoles first utilize 2 moles of bromamine-B to form isatins. Thus formed isatins consume another mole of bromamine-B to yield the ultimate and desired compounds, anthranilic acids. The detailed mechanisms for reactions are shown in Schemes 4, 6 and 8.

The reactions were studied in various solvents (acetonitrile, 1, 2-dichloromethane, ethanol and acetonitrile–water (1 : 1) mixture). The mixture of acetonitrile–water (1 : 1) was found to be the best solvent system, perhaps due to the dielectric constant and the solubility of the reactants. Indoles are not very soluble in water, but markedly soluble in acetonitrile–water mixture. Moreover the organic haloamines furnish different species better in aqueous (water) medium rather than in organic medium. For all these reasons acetonitrile–water with a 1 : 1 ratio system was the better choice of solvent system in the present synthetic study. The reactions were found to be highly dependent upon the pH of the system. To evaluate the effect of pH, the reactions were carried out under similar experimental conditions at different pHs using NaOH. At neutral pH, the oxidation reactions were found to be very slow. Reaction rates increase with increasing pH (addition of NaOH). This behavior of the reaction is attributed to the dissociation of bromamine-B in aqueous medium by furnishing different oxidizing species. This behavior of dissociation of bromamine-B is well explained in the Discussion. Because of the increase rate in the presence of NaOH, the reactions were carried out at $2.0 \times 10^{-3} \text{ mol dm}^{-3}$ NaOH. The kinetics of the oxidative conversion of indole by BAB was investigated at several initial concentrations of the reactants in NaOH medium in presence of Ru(III), Os(VIII) and Ru(III) + Os(VIII) mixture catalysts at 313 K under identical experimental conditions.

Effect of Varying Reactant Concentrations on the Rate

With the indole in excess at constant $[\text{Indole}]_0$, $[\text{NaOH}]$, $[\text{catalyst}]$ and temperature, plots of $\log [\text{BAB}]$ vs time were linear, indicating a first-order dependence of the rate on $[\text{BAB}]_0$ in all the three catalyzed reactions. The pseudo first-order rate constants (k') are recorded in Table 2. Further these k' values were unaffected by the variation of $[\text{BAB}]_0$, confirming the first-order dependence of the rate on $[\text{BAB}]_0$. Under the same experimental conditions, the rate of the reaction increased with increase in $[\text{Indole}]_0$ for Ru(III) and Os(VIII)-catalyzed reactions and plots of $\log k'$ vs $\log [\text{Indole}]$ (Fig. 1) were found to be linear with unit slopes, indicating first-order dependence on $[\text{Ru(III)}]$ and $[\text{Os(VIII)}]$ catalysts. However, the order with respect to Ru(III) + Os(VIII) catalyst was found to be zero order. These results are given in Table 2.

Table 2. Effect of varying reactant concentrations on the reaction rate at 313 K

$10^4 [\text{BAB}]_0$ (mol dm^{-3})	$10^3 [\text{Indole}]_0$ (mol dm^{-3})	$10^4 k' \text{ (s}^{-1}\text{)}$		
		Ru(III)	Os(VIII)	Ru(III) + Os(VIII)
0.2	1.0	3.30	6.75	21.8
0.5	1.0	3.19	6.67	21.9
1.0	1.0	3.21	6.70	21.7
2.0	1.0	3.28	6.69	22.0
4.0	1.0	3.16	6.77	21.5
1.0	0.2	0.81	1.42	21.2
1.0	0.5	1.60	3.32	21.9
1.0	1.0	3.21	6.70	21.7
1.0	2.0	6.50	14.0	21.4
1.0	4.0	13.0	27.5	22.0

$[\text{NaOH}] = 2.0 \times 10^{-3} \text{ mol dm}^{-3}$; $[\text{catalyst}] = 2.0 \times 10^{-6} \text{ mol dm}^{-3}$;
 $l = 0.3 \text{ mol dm}^{-3}$ [in the case of Os(VIII) and Ru(III) + Os(VIII) catalysis].




Figure 1. Plots of $\log k'$ versus $\log [\text{Indole}]$.

Effect of Varying NaOH and Catalyst Concentrations on the Rate

The kinetic experiments showed that the rate is unaffected with aqueous acetonitrile markedly. The reactions were carried out with different amounts of acetonitrile and it was found that the rate is not affected significantly by acetonitrile. However, rate is significantly affected by alkali. The rate of the reaction increased with increase in $[\text{NaOH}]$ in all cases (Table 3). The log–log plots of rate vs $[\text{NaOH}]$ (Fig. 2) showed that the orders in $[\text{NaOH}]$ were less than unity, suggesting a fractional-order dependence on $[\text{OH}^-]$ in all three catalyzed reactions. The reaction rate increased with increase in $[\text{catalyst}]$ in all the cases (Table 3). The order with respect to $[\text{Ru(III)}]$ was found to be fractional but it was found to be unity for $[\text{Os(VIII)}]$ and $[\text{Ru(III)} + \text{Os(VIII)}]$ catalysts. This clearly indicates a fractional-order dependence on $[\text{Ru(III)}]$, and first-order dependence on $[\text{Os(VIII)}]$ and $[\text{Ru(III)} + \text{Os(VIII)}]$ catalysts. The plots of $\log k'$ vs $\log [\text{catalyst}]$ are presented in Fig. 3.

Table 3. Effect of varying NaOH and catalyst concentrations on the reaction rate at 313 K

10 ³ [NaOH] (mol dm ⁻³)	10 ⁶ [catalyst] (mol dm ⁻³)	10 ⁴ <i>k'</i> (s ⁻¹)		
		Ru(III)	Os(VIII)	Ru(III) + Os(VIII)
0.5	2.0	1.5	3.32	10
1.0	2.0	2.02	4.80	13.5
2.0	2.0	3.21	6.70	21.7
4.0	2.0	5.00	9.00	31.6
6.0	2.0	6.96	11.4	40.0
2.0	0.5	1.1	1.72	5.00
2.0	1.0	1.71	3.26	10.0
2.0	2.0	3.21	6.70	21.7
2.0	4.0	6.38	14.3	43.0
2.0	8.0	10.0	28.5	88.0

[BAB]₀ = 1.0 × 10⁻⁴ mol dm⁻³; [Indole]₀ = 1.0 × 10⁻³ mol dm⁻³; *I* = 0.3 mol dm⁻³ [in the case of Os(VIII) and Ru(III) + Os(VIII) catalysis].

case of Ru(III). However, the rate of the reaction increased with increasing ionic strength in the case of Os(VIII) catalysis and the rate decreased in the case of Ru(III) + Os(VIII) catalysis. Plots of log *k'* vs *I*^{1/2} were found to be linear with slopes +0.50 and -1.60 in Os(VIII) and Ru(III) + Os(VIII) catalysis respectively (Table 4). Hence, the ionic strength of the medium was maintained at a constant concentration of 0.30 mol dm⁻³ of NaClO₄ in the case of Os(VIII) and Ru(III) + Os(VIII) catalysis for kinetic runs in order to swamp the reaction.

Effect of Solvent Isotope on the Rate

Studies of the reaction rate in D₂O medium for Ru(III), Os(VIII) and Ru(III) + Os(VIII) catalyzed reactions revealed that *k'* (H₂O) was equal to 3.20 × 10⁻⁴, 6.70 × 10⁻⁴ and 21.7 × 10⁻⁴ and *k'* (D₂O) = 4.50 × 10⁻⁴, 8.45 × 10⁻⁴ and 29.1 × 10⁻⁴, respectively. The solvent isotope effect, *k'* (H₂O)/*k'* (D₂O) was found to be 0.71, 0.79 and 0.75 for the three catalyzed reactions. These results are given Table 5.

Effect of Varying Ionic Strength of the Medium on the Rate

An increase in ionic strength of a reaction system by addition of NaClO₄ showed a negligible effect on the reaction rate in the

Effect of Varying Temperature on the Rate

The reactions were studied at different temperatures (303–323 K), keeping other experimental conditions constant. From the linear Arrhenius plots of log *k'* vs 1/*T*, values of activation parameters



**Figure 2.** Plots of log *k'* versus log[NaOH].**Figure 3.** Plots of log *k'* versus log[catalyst].

Table 4. Effect of varying ionic strength of the medium on the reaction rate at 313 K

Ionic strength (<i>I</i>) (mol dm ⁻³)	10 ⁴ <i>k'</i> (s ⁻¹)		
	Ru(III)	Os(VIII)	Ru(III) + Os(VIII)
0.1	3.28	5.00	48.9
0.2	3.19	6.00	28.8
0.3	3.21	6.7	21.7
0.4	3.30	7.60	14.8
0.5	3.17	8.70	11.2

[BAB]₀ = 1.0 × 10⁻⁴ mol dm⁻³; [Indole]₀ = 1.0 × 10⁻³ mol dm⁻³;
[NaOH] = 2.0 × 10⁻³ mol dm⁻³; [catalyst] = 2.0 × 10⁻⁶ mol dm⁻³.

Table 5. Effect of solvent isotope on the rate at 313 K

	<i>k'</i> (Water)	<i>k'</i> (D ₂ O)	<i>k'</i> (H ₂ O)/ <i>k'</i> (D ₂ O)
Ru(III)	3.20 × 10 ⁻⁴ s ⁻¹	4.50 × 10 ⁻⁴ s ⁻¹	0.71
Os(VIII)	6.70 × 10 ⁻⁴ s ⁻¹	8.45 × 10 ⁻⁴ s ⁻¹	0.79
Ru(III) + Os(VIII)	21.7 × 10 ⁻⁴ s ⁻¹	29.1 × 10 ⁻⁴ s ⁻¹	0.75

[BAB]₀ = 1.0 × 10⁻⁴ mol dm⁻³; [Indole]₀ = 1.0 × 10⁻³ mol dm⁻³;
[NaOH] = 2.0 × 10⁻³ mol dm⁻³; [catalyst] = 2.0 × 10⁻⁶ mol dm⁻³;
I = 0.3 mol dm⁻³ [in the case of Os(VIII) and Ru(III) + Os(VIII) catalysis].

Table 6. Temperature dependence and values of composite activation parameters for the synthesis of anthranilic acid by BAB in the presence and absence of catalyst

Temperature (K)	10 ⁴ <i>k'</i> (s ⁻¹)			
	Ru(III)	Os(VIII)	Ru(III) + Os(VIII)	Uncatalyzed
303	1.31	3.34	14.0	0.18
308	2.01	5.01	17.8	0.32
313	3.21	6.70	21.7	0.52
318	5.26	10.0	26.4	0.90
323	7.01	13.4	33.5	1.58
<i>E</i> _a (kJ mol ⁻¹)	70.3	52.3	29.7	85.4
Δ <i>H</i> [‡] (kJ mol ⁻¹)	67.4	49.8	26.7	82.8
Δ <i>G</i> [‡] (kJ mol ⁻¹)	94.6	92.6	90.0	98.4
Δ <i>S</i> [‡] (J K ⁻¹ mol ⁻¹)	-80.0	-141	-206	-52.7
log <i>A</i>	8.62	5.85	2.45	10.4

[BAB]₀ = 1.0 × 10⁻⁴ mol dm⁻³; [Indole]₀ = 1.0 × 10⁻³ mol dm⁻³;
[NaOH] = 2.0 × 10⁻³ mol dm⁻³; [catalyst] = 2.0 × 10⁻⁶ mol dm⁻³;
I = 0.3 mol dm⁻³ [in the case of Os(VIII) and Ru(III) + Os(VIII) catalysis].

(*E*_a, Δ*H*[‡], Δ*G*[‡], Δ*S*[‡] and log *A*) for the composite reaction were deduced. All these data are summarized in Table 6.

Effect of Varying Benzenesulfonamide Concentrations on the Rate

Addition of benzenesulfonamide (BSA; 1.0 × 10⁻³ to 8.0 × 10⁻³ mol dm⁻³), a reduction product of BAB, to the reaction mixture did not affect the rate significantly. This indicates that BSA

is not involved in any step prior to the rate-limiting step in the schemes proposed.

Effect of Varying Halide Ion Concentrations on the Rate

The rate remained unchanged with the addition of Cl⁻ or Br⁻ ions in the form of NaCl and NaBr (1.0 × 10⁻² to 8.0 × 10⁻² mol dm⁻³). These results showed that halide ions play no role in the reaction.

Test for Free Radicals

The addition of the reaction mixtures to the aqueous acrylamide monomer solution did not initiate polymerization indicating the absence of *in situ* formation of free radical species in the reaction sequence. The control experiments were also performed under the same reaction conditions but without the oxidant, BAB.


Discussion

Reactive Species of Bromamine-B

Investigations of Morris *et al.*,^[18] Bishop and Jennings,^[19] Pryde and Soper^[20] and Hardy and Johnston^[21] have shown the existence of similar equilibria of *N*-metallo-*N*-aryhalosulfonamides in aqueous media. Bromamine-B, like chloramine-T, acts as an oxidizing agent in both acidic and alkaline solutions. Depending on the pH, BAB exhibits different equilibria in aqueous solutions.^[18–21] In alkaline solutions of BAB, PhSO₂NBr₂ does not exist,^[8] and the possible oxidizing species are PhSO₂NBr⁻ and OBr⁻ anions, which could be transformed into the more reactive species PhSO₂NHBr and HOBr during the course of the reaction in alkali-retarding steps. Several workers have observed the retarding effect of OH⁻ ions on the rate of haloamines with a number of substrates^[22–27] and have suggested that the reactivity of weakly alkaline solutions of haloamines is due to the formation of the conjugate acid PhSO₂NHBr from PhSO₂NBr⁻ in the OH⁻ retarding step. However, in the present investigations, the OH⁻ ions increase the rate of the reaction, clearly indicating PhSO₂NBr⁻ as the reactive oxidizing species. In our earlier work^[28,29] the positive influence of OH⁻ ion on the rate of haloamine reactions with a number of substrates has been observed and RNX⁻ (R = Ts or PhSO₂; X = Cl or Br) suggested as the reactive oxidizing species. In the present investigations, the rate of the reaction accelerated by OH⁻ ions clearly indicates that the anion PhSO₂NBr⁻ is the most likely reactive species involved in the synthesis of anthranilic acids from indoles by BAB in all three catalyzed reactions.

Mechanism and Rate Law of Ru(III) Catalysis

Ru(III) chloride acts as catalyst in many organic reactions, particularly in an alkaline medium.^[30–33] Under the experimental conditions [OH⁻] ≫ [Ru(III)] and owing to the fact that [OH⁻] increases the rate, ruthenium (III) is predominantly present as the hydroxylated species [Ru(H₂O)₅OH]²⁺ and its formation is shown in the following equilibrium:




Similar equilibria have been reported between Ru(III) catalyzed reactions in alkaline medium.^[31,33]

The existence of a complex between the catalyst and BAB was evidenced from the UV–vis spectra of both Ru(III) and Ru(III)–BAB

mixture. Absorption maxima of Ru(III) in HCl medium, BAB and Ru(III)–BAB in aqueous alkaline acetonitrile medium appear at 352, 223 and 338 nm respectively. A hypsochromic shift of 14 nm from 352 to 338 nm of Ru(III) indicates the formation of complex between Ru(III) and BAB.

Further, for a general equilibrium (3)




between two metal species, M and MS_n , having different extinction coefficients, Ardon^[34] has derived the following equation (4):

$$\frac{1}{\Delta A} = \left(\frac{1}{[S]^n}\right) \frac{1}{(\Delta E)K[\text{Catalyst}]_{\text{Total}}} + \frac{1}{(\Delta E)[\text{Catalyst}]_{\text{Total}}} \quad (4)$$

where K is the formation constant of the complex, $[S]$ is the concentration of BAB, ΔE is the difference in extinction coefficient between two metal species, $[\text{Catalyst}]_{\text{Total}}$ is the total concentration of catalyst species and ΔA is the absorbance difference of solution containing with and without BAB. Equation (4) is valid provided that $[\text{BAB}]$ is much greater than $[\text{Catalyst}]_{\text{Total}}$. According to equation (4), a plot of $1/\Delta A$ vs $1/[\text{BAB}]$ or $1/[\text{BAB}]^2$ should be linear with an intercept in the case of 1:1 or 1:2 type of complex formation between M and S. The ratio of intercept to slope of this linear plot gives the value of K .

Ruthenium (III) in aqueous alkaline acetonitrile medium containing BAB showed an absorption peak at 338 nm (λ_{max} for the complex). The complex formation studies were made at this λ_{max} of 338 nm. In a set of experiments, the solutions were prepared by taking different amounts of BAB (2.0×10^{-4} – 4.0×10^{-3} mol dm⁻³) at constant amounts of RuCl₃ (2.0×10^{-6} mol dm⁻³) and NaOH (2.0×10^{-2} mol dm⁻³) in aqueous acetonitrile at 313 K. The absorbance of these solutions was measured at 338 nm. The absorbance of the solution in the absence of BAB was also measured at the same wavelength. The difference of these absorbances (with and without BAB) gives the differential absorbance, ΔA . A plot of $1/\Delta A$ vs $1/[\text{BAB}]$ was linear with an intercept suggesting the formation of a 1:1 complex between BAB and Ru(III) catalyst. Similar behavior for the formation of complex has been reported in earlier works.^[34,35] Further, the plot of $\log(1/\Delta A)$ vs $\log(1/[\text{BAB}])$ was also linear ($r = 0.9801$). From the slope and intercept of the plot $1/\Delta A$ vs $1/[\text{BAB}]$, the value of the formation constant, K , of the complex was deduced and found to be 6.06×10^2 .

In view of the above observations, a general mechanism (Scheme 3) is proposed for the Ru(III)-catalyzed synthesis of anthranilic acids from indoles by BAB to account for the observed experimental results.



Scheme 3. Ru(III)-catalyzed mechanistic scheme.

Here, X and X' are the intermediate complexes whose structures are illustrated in Scheme 4, where a detailed mechanistic interpretation of indole-BAB reaction in presence Ru(III) has been proposed.

If $[\text{BAB}]_t$ represents the total effective concentration of BAB, then

$$[\text{BAB}]_t = [\text{PhSO}_2\text{NHBr}] + [\text{PhSO}_2\text{NBr}^-] + [\text{X}] \quad (5)$$

By substituting for $[\text{PhSO}_2\text{NHBr}]$ and $[\text{PhSO}_2\text{NBr}^-]$ into equation (5) and solving for $[\text{X}]$ we get:

$$[\text{X}] = \frac{K_1 K_2 [\text{BAB}]_t [\text{OH}^-] [\text{Ru(III)}]}{[\text{H}_2\text{O}] + K_1 [\text{OH}^-] + K_1 K_2 [\text{OH}^-] [\text{Ru(III)}]} \quad (6)$$

From the slow and rate-determining step of Scheme 3, the rate of reaction is given by

$$\text{Rate} = -d[\text{BAB}]/dt = k_3 [\text{Indole}][\text{X}] \quad (7)$$

On substituting the value of $[\text{X}]$ from equation (6) into equation (7), one obtains the rate law (8):

$$\text{Rate} = \frac{-d[\text{BAB}]}{dt} = \frac{K_1 K_2 k_3 [\text{BAB}]_t [\text{Indole}][\text{OH}^-] [\text{Ru(III)}]}{[\text{H}_2\text{O}] + K_1 [\text{OH}^-] + K_1 K_2 [\text{OH}^-] [\text{Ru(III)}]} \quad (8)$$

Rate law (8) is in complete agreement with experimental data. Hence the proposed mechanism is valid.


Since $\text{rate} = k'[\text{BAB}]_t$, rate law (8) can be transformed into the following equations (9)–(12):

$$k' = \frac{K_1 K_2 k_3 [\text{Indole}][\text{OH}^-] [\text{Ru(III)}]}{[\text{H}_2\text{O}] + K_1 [\text{OH}^-] + K_1 K_2 [\text{OH}^-] [\text{Ru(III)}]} \quad (9)$$

$$\frac{1}{k'} = \frac{[\text{H}_2\text{O}]}{K_1 K_2 k_3 [\text{Indole}][\text{OH}^-] [\text{Ru(III)}]} + \frac{1}{K_2 k_3 [\text{Indole}][\text{Ru(III)}]} + \frac{1}{k_3 [\text{Indole}]} \quad (10)$$

$$\frac{1}{k'} = \frac{1}{[\text{OH}^-]} \left\{ \frac{[\text{H}_2\text{O}]}{K_1 K_2 k_3 [\text{Indole}][\text{Ru(III)}]} \right\} + \left\{ \frac{1}{K_2 k_3 [\text{Indole}][\text{Ru(III)}]} + \frac{1}{k_3 [\text{Indole}]} \right\} \quad (11)$$

$$\frac{1}{k'} = \frac{1}{[\text{Ru(III)}]} \left\{ \frac{[\text{H}_2\text{O}]}{K_1 K_2 k_3 [\text{Indole}][\text{OH}^-]} + \frac{1}{K_2 k_3 [\text{Indole}]} \right\} + \frac{1}{k_3 [\text{Indole}]} \quad (12)$$



Scheme 4. Ru(III)-catalyzed mechanism for the synthesis of anthranilic acids.

A plot of $1/k'$ vs $1/[\text{OH}^-]$ from equation (11) yields

$$\text{Slope} = \frac{[\text{H}_2\text{O}]}{K_1 K_2 k_3 [\text{Indole}] [\text{Ru(III)}]} \quad \text{and Intercept} \\ = \frac{1}{K_2 k_3 [\text{Indole}] [\text{Ru(III)}]} + \frac{1}{k_3 [\text{Indole}]}$$

Similarly, a plot of $1/k'$ vs $1/[\text{Os(VIII)}]$ from equation (12) yields

$$\text{Slope} = \frac{[\text{H}_2\text{O}]}{K_1 K_2 k_3 [\text{Indole}] [\text{OH}^-]} + \frac{1}{K_2 k_3 [\text{Indole}]} \\ \text{and Intercept} = \frac{1}{k_3 [\text{Indole}]}$$

Therefore from the slopes and intercepts of equations (11) and (12), the values of equilibrium constants K_1 and K_2 and decomposition constant k_3 were calculated for the standard run with $[\text{BAB}]_0 = 1.0 \times 10^{-4} \text{ mol dm}^{-3}$; $[\text{Indole}] = 1.0 \times 10^{-3} \text{ mol dm}^{-3}$; $[\text{NaOH}] = 2.0 \times 10^{-3} \text{ mol dm}^{-3}$ $[\text{Ru(III)}] = 2.0 \times 10^{-6} \text{ mol dm}^{-3}$ at 313 K. The values obtained were $k_3 = 6.14 \times 10^{-2} \text{ s}^{-1}$; $K_2 = 8.00 \times 10^5 \text{ dm}^3 \text{ mol}^{-1}$; $K_1 = 7.89 \times 10^{-3} \text{ dm}^3 \text{ mol}^{-1}$.

The proposed Scheme 3 and the derived rate law (8) are substantiated by the experimental observations discussed below.

For a reaction involving a fast pre-equilibrium H^+ or OH^- ion transfer, the rate increases in D_2O since D_3O^+ and OD^- are 2–3 times stronger acids and stronger bases,^[36,37] respectively, than H_3O^+ and OH^- ions. In the present studies the observed solvent isotope effect of $k'(\text{H}_2\text{O})/k'(\text{D}_2\text{O}) < 1$ is due to the greater basicity of OD^- compared with OH^- . This isotope effect observed in the present case confirms to the above fact. However, the magnitude of increase of rate in D_2O is small [$k'(\text{H}_2\text{O})/k'(\text{D}_2\text{O}) = 0.71$] compared with the expected value of 2–3 times greater, which can be attributed to the fractional-order dependence of rate on $[\text{OH}^-]$.

It was felt reasonable to compare the reactivity of BAB towards indole in the absence of Ru(III) catalyst under identical experimental conditions in order to evaluate the catalytic efficiency of Ru(III). The reactions were carried out at different temperatures (303–323 K) and from the plot of $\log k'$ vs $1/T$ (Fig. 4), activation parameters were also evaluated for the uncatalyzed reactions (Table 6). However, the Ru(III)-catalyzed reactions were found to be about 6 times faster than uncatalyzed reactions. The activation parameters evaluated for the catalyzed and uncatalyzed reactions explain the catalytic effect on the reaction. The catalyst Ru(III) forms a complex (X) with the BAB, which increases the reactivity of BAB.

The general equation relating for uncatalyzed and catalyzed reactions has been derived by Moelwyn and Hughes^[38] as:

$$k_1 = k_0 + K_C [\text{catalyst}]^x \quad (13)$$

Here k_1 is the observed pseudo first-order rate constant obtained in the presence of Ru(III) catalyst, k_0 is the pseudo first-order constant for the uncatalyzed reaction, K_C is the catalytic constant and x is the order of the reaction with respect to $[\text{Ru(III)}]$. In the present investigations, x was found to be 0.8. Then the value of K_C was calculated using the equation (13). The values of K_C were evaluated at different temperatures (303–323 K) and K_C was found to vary with temperature. Further, a plot of $\log K_C$ vs $1/T$ (Fig. 5) was linear and the values of energy of activation and other activation parameters for the Ru(III) catalyst were computed and are summarized in Table 7.

The proposed mechanism is supported by the observed moderate values of energy of activation and other thermodynamic parameters. The high positive values of the free energy of activation and of the enthalpy of activation suggest that the transition state is highly solvated, while fairly high negative entropy of activation (Table 6) indicates the formation of a rigid associated transition state. Variation of ionic strength of the




Figure 4. Plots of $\log k'$ versus $1/T$.





Figure 5. Plots of $\log K_c$ vs. $1/T$.

medium did not alter the reaction rate, which indicates that the noninvolvement of ionic species in the rate-determining step. Addition of the reduction product of BAB, BSA, does not influence the rate, showing that it is not involved in pre-equilibrium. Addition of halide ions has no effect on the rate, indicating that there is no role for halide ions in the reaction. All these observations also confirm the proposed mechanism and derived rate law.

Mechanism and Rate Law of Os (VIII) Catalysis

It has been shown that osmium has a stable +8 oxidation state^[14,39,40] and exists in the following equilibria in alkaline solutions:



The complexes $[\text{OsO}_4(\text{OH})(\text{H}_2\text{O})]^-$ and $[\text{OsO}_4(\text{OH})_2]^{2-}$, which can be reduced to $[\text{OsO}_2(\text{OH})_4]^{2-}$, with octahedral geometries are less likely to form species of higher coordination. It is more realistic to postulate that OsO_4 , which has tetrahedral geometry, as the active catalyst species can effectively form a complex.

The existence of a complex between indole and Os(VIII) was evidenced from the UV-vis spectra of indole, Os(VIII) and indole-Os(VIII) mixture. Absorption maxima in aqueous alkaline acetonitrile medium appeared at 286 nm for indole, 319 nm for Os(VIII) and 300 nm for a mixture of both. A bathochromic shift of 14 nm from 286 to 300 nm of indole suggests that complexation occurred between indole and Os(VIII). According to equation (4) a plot of $1/\Delta A$ vs $1/[\text{Indole}]$ with an intercept suggests the formation of 1:1 complex between Os(VIII) and indole. Further, the plot of $\log 1/\Delta A$ vs $\log 1/[\text{Indole}]$ was also linear ($r = 9898$). From the slope and intercept of the plot $1/\Delta A$ vs $1/[\text{Indole}]$, the value of formation constant, K , of the complex was found to be 8.23×10^2 .

Based on the experimental results, it is likely that $\text{PhSO}_2\text{NBr}^-$ itself also acts as the reactive oxidant species in the present case.

Table 7. Values of catalytic constant (K_C) at different temperatures and activation parameters calculated using K_C values

Temperature (K)	$K_C \times 10$		
	Ru(III)	Os(VIII)	Ru(III) + Os(VIII)
303	5.65	31.6	138
308	8.45	46.9	172
313	13.4	62.8	211
318	21.8	91	255
323	54.3	118	319
E_a (kJ mol ⁻¹)	88	50.6	41.8
ΔH^\ddagger (kJ mol ⁻¹)	85.1	47.3	38.1
ΔG^\ddagger (kJ mol ⁻¹)	93.4	70.2	62.2
ΔS^\ddagger (J K ⁻¹ mol ⁻¹)	-70.8	-52.3	-45.7
log A	13.1	8.98	5.86

[BAB]₀ = 1.0 × 10⁻⁴ mol dm⁻³; [Indole]₀ = 1.0 × 10⁻³ mol dm⁻³;
[NaOH] = 2.0 × 10⁻³ mol dm⁻³; [catalyst] = 2.0 × 10⁻⁶ mol dm⁻³;
 I = 0.3 mol dm⁻³ [in the case of Os(VIII) and Ru(III) + Os(VIII) catalysis].

Considering the above facts and all the observed experimental data, the following reaction, Scheme 5, can be suggested for Os(VIII) catalyzed synthesis of anthranilic acids from indoles by BAB in alkaline medium:

Here, X'' and X''' are the intermediate complexes whose structures are given in Scheme 6, wherein a detailed mechanistic pathway for Os(VIII) catalysis reaction is illustrated. The total effective concentration of BAB is given by:

$$[\text{BAB}]_t = [\text{PhSO}_2\text{NHBr}] + [\text{PhSO}_2\text{NBr}^-] \quad (16)$$

From step (ii) of Scheme 5,

$$[\text{X}'''] = K_6[\text{Indole}][\text{Os(VIII)}] \quad (17)$$

From the slow and rate determining step (iii) of Scheme 5, the rate of reaction is given by:

$$\text{Rate} = -d[\text{BAB}]_t/dt = k_7[\text{X}'''][\text{PhSO}_2\text{NBr}^-] \quad (18)$$

On substituting for [PhSO₂NBr⁻] and [X'''] into equation (18), the rate law (19) was obtained:

$$\text{Rate} = \frac{K_5 K_6 k_7 [\text{BAB}]_t [\text{Indole}] [\text{OH}^-] [\text{Os(VIII)}]}{[\text{H}_2\text{O}] + K_5 [\text{OH}^-]} \quad (19)$$

The rate law (19) is in agreement with the experimentally observed results. Since rate = $k'[\text{BAB}]_t$, equation (19) can be transformed

into equations (20) and (21):

$$k' = \frac{K_5 K_6 k_7 [\text{Indone}] [\text{OH}^-] [\text{Os(VIII)}]}{[\text{H}_2\text{O}] + K_5 [\text{OH}^-]} \quad (20)$$

$$\frac{1}{k'} = \frac{[\text{H}_2\text{O}]}{K_5 K_6 k_7 [\text{Indole}] [\text{Os(VIII)}] [\text{OH}^-]} + \frac{1}{K_6 k_7 [\text{Indole}] [\text{Os(VIII)}]} \quad (21)$$


A plot of $1/k'$ vs $1/[\text{OH}^-]$ according to equation (21), with other experimental conditions held constant, was found to be linear ($r > 0.9899$). K_5 and $K_6 k_7$ values were calculated from the slope and intercept of such a plot and was found to be 2.67×10^3 and $2.0 \times 10^{-5} \text{ dm}^3 \text{ mol}^{-1} \text{ s}^{-1}$ for the standard run, $[\text{BAB}]_0 = 1.0 \times 10^{-4} \text{ mol dm}^{-3}$; $[\text{Indole}]_0 = 1.0 \times 10^{-3} \text{ mol dm}^{-3}$; $[\text{Os(VIII)}] = 2.0 \times 10^{-6} \text{ mol dm}^{-3}$; and $I = 0.3 \text{ mol dm}^{-3}$ at 313 K. Scheme 5 and rate law (19) can explain the following observed experimental results.


The ionic strength (I) effect on the reaction rates was described according to the theory of Bronsted and Bjerrum,^[41] which postulates the reaction through the formation of an activated complex. According to this theory, the effect of ionic strength on the rate for a reaction involving two ions is given by the relationship

$$\log k' = \log k_0 + 1.02 Z_A Z_B I^{1/2} \quad (22)$$

Here Z_A and Z_B are the valency of the ions A and B, and k_0 are the rate constants in the presence and absence of the added electrolyte, respectively. A plot of $\log k'$ against $I^{1/2}$ should be linear with a slope of $1.02 Z_A Z_B$. If Z_A and Z_B have similar signs, the quantity $Z_A Z_B$ is positive and the rate increases with the ionic strength, having a positive slope, while if the ions have dissimilar charges, the quantity $Z_A Z_B$ is negative and the rate will decrease with the increase in ionic strength, having a negative slope. In the present case, a primary salt effect was observed as the rate increased with the increase in ionic strength of the medium,^[44] supporting the involvement of ions of same sign in the rate-limiting step (Scheme 6). The Debye-Huckel plot ($\log k'$ against $I^{1/2}$) gave a straight line with a slope of 0.6. In the present system, a two negative ions ($Z_A = -1$ and $Z_B = -1$) were involved in the rate-determining step and the expected slope of +1 was not found. This may be due to the fact that the ionic strength employed was beyond the formal Debye-Huckel limiting range. Alternatively, there could be formation of ion pairs in concentrated solutions, as suggested by Bjerrum.^[44]

The reactivity of BAB towards indole in the absence of Os(VIII) catalyst was compared with the Os(VIII) catalyzed reaction under identical experimental conditions. Rate constants revealed that the Os(VIII) catalyzed reactions were about 13-fold faster than

**Scheme 5.** Os(VIII)-catalyzed mechanistic scheme.



Scheme 6. Os(VIII)-catalyzed mechanism for the synthesis of anthranilic acids.

uncatalyzed reactions (Table 6). The values of K_C were determined at different temperatures and, from a plot of $\log K_C$ vs $1/T$, the values of activation parameters for Os(VIII) catalyst were computed (Table 7). The negligible effect of the addition of BSA and halide ions on the rate of the reaction and also activation parameters was in good agreement with the mechanism proposed and the rate law derived.

Mechanism and Rate Law of Ru(III) + Os(VIII) Catalysis

The kinetic data suggested that $\text{PhSO}_2\text{NBr}^-$ is also the reactive species of BAB in Ru(III) + Os(VIII) catalyzed reaction. UV-visible spectral studies revealed that there is transient existence of a complex between Ru(III) and Os(VIII) in aqueous alkaline acetonitrile medium. Absorption maxima in aqueous alkaline acetonitrile medium appeared at 352 nm for Ru(III), 319 nm for Os(VIII) and 336 nm for a mixture of both. This clearly indicates the formation of a transient complex between Ru(III) and Os(VIII) and this transient complex $[\text{Ru(III)} + \text{Os(VIII)}]$ acts as reactive catalyst species in this case.

The existence of other complex between Ru(III) + Os(VIII) and BAB was also evidenced from the UV-vis spectra of both BAB, Ru(III) + Os(VIII) and a mixture of both in alkaline medium. Absorption maxima in aqueous alkaline acetonitrile medium appeared at 223 nm for BAB, 336 nm for Os(VIII) and 245 nm for a mixture of both.

From the above preceding discussions, Scheme 7 has been proposed to account for the observed kinetic results. In this scheme, X^{IV} and X^{V} represent the complex intermediate species. Scheme 8 depicts the mode of indole-BAB reaction in the presence of Ru(III) + Os(VIII) mixture. The total effective concentration of BAB is

$$[\text{BAB}]_t = [\text{PhSO}_2\text{NHBr}] + [\text{PhSO}_2\text{NBr}^-] \quad (23)$$

From the slow step of Scheme 7,

$$\text{rate} = -d[\text{BAB}]_t = k_{10}[\text{PhSO}_2\text{NBr}^-][\text{Ru(III)} + \text{Os(VIII)}] \quad (24)$$

By substituting for $[\text{PhSO}_2\text{NBr}^-]$ into equation (24) the following rate law is obtained:



$$\text{rate} = \frac{-d[\text{BAB}]_t}{dt} = \frac{K_9 k_{10} [\text{BAB}]_t [\text{Ru(III)} + \text{Os(VIII)}] [\text{OH}^-]}{[\text{H}_2\text{O}] + K_9 [\text{OH}^-]} \quad (25)$$

Rate law (25) is in good agreement with the experimental data.

Since $\text{rate} = k'[\text{BAB}]_t$, equation (25) can be transformed into equations (26) and (27):

$$k' = \frac{K_9 k_{10} [\text{Ru(III)} + \text{Os(VIII)}] [\text{OH}^-]}{[\text{H}_2\text{O}] + K_9 [\text{OH}^-]} \quad (26)$$

$$\frac{1}{k'} = \frac{[\text{H}_2\text{O}]}{K_9 k_{10} [\text{Ru(III)} + \text{Os(VIII)}] [\text{OH}^-]} + \frac{1}{k_9 [\text{Ru(III)} + \text{Os(VIII)}]} \quad (27)$$

**Scheme 7.** Ru(III)+Os(VIII)-catalyzed mechanistic scheme.**Scheme 8.** Ru(III) + Os(VIII)-catalyzed mechanism for the synthesis of anthranilic acids.

Based on equation (27), a plot of $1/k'$ vs $1/[\text{OH}^-]$ at constant $[\text{BAB}]_0$, $[\text{Indole}]_0$, $[\text{Ru(III)} + \text{Os(VIII)}]$ ionic strength and temperature was found to be linear ($r = 0.9899$). From the values of slope and intercept of such a plot, the equilibrium constant (K_9) and decomposition constant (k_{10}) were calculated and found to be $2.35 \times 10^4 \text{ dm}^3 \text{ mol}^{-1}$ and $1.26 \times 10^{-2} \text{ s}^{-1}$ respectively.

The proposed mechanism and the derived rate law were substantiated by the following experimental facts. The ionic strength (I) effect on the reaction rates was studied and a primary salt effect was observed as the rate decreases with increase in ionic strength of the medium,^[41] supporting the involvement of ions of opposite sign in the rate-determining step (Scheme 8). The Debye–Huckel plot ($\log k'$ against $I^{1/2}$) gave straight line with slope of -1.60 . In the present system, a negatively charged ion and two positively charged ions were involved in the rate-determining step (Scheme 8) and the expected slope of -2 was not found. This may be due to the fact that the ionic strength employed was beyond the formal Debye–Huckel limiting range. Alternatively, there could be formation of ion pairs in concentrated solutions, as suggested by Bjerrum.^[44]

The reactions were also studied in absence of Ru(III) + Os(VIII) catalyst and rates were compared with that of Ru(III) + Os(VIII)

catalyzed reaction under identical experimental conditions. Rate constants revealed that the Ru(III) + Os(VIII) catalyzed reactions are about 42-fold faster than uncatalyzed reactions (Table 6). The values of K_C were determined at different temperatures and from a plot of $\log K_C$ vs $1/T$; values of activation parameters for Ru(III) + Os(VIII) catalyst were computed (Table 7). The negligible effect on the addition of BSA and halide ions on the rate of the reaction and also activation parameters was in good agreement with the mechanism proposed and the rate law derived.

The kinetics of all the three catalyzed reactions was compared with uncatalyzed reactions, under similar experimental conditions, and it was found that the catalyzed reactions were 6- to 42-fold faster. For the catalyzed reactions, it is seen from the Table 6 that the activation energy was highest for the slowest reaction and vice-versa. From the inspection of rate constants and the values of energy of activation (Table 6), the relative reactivity of these catalysts was in the order: Ru(III) + Os(VIII) > Os(VIII) > Ru(III). The reactivity was faster in the case of Os(VIII) compared with Ru(III). This may be attributed to the d-electronic configuration of the metal ions. Osmium having d^0 electronic configuration has greater catalytic efficiency to oxidize the substrate compared to the Ru(III), which is of d^5 electronic configuration. The reactivity in the case of

Ru(III) + Os(VIII) catalysis was maximum. This trend may be due to the formation of a transient complex between Ru(III) and Os(VIII). Ru(III) effectively forms a complex with Os(VIII) which increases the catalytic property of Os(VIII) and thereby the catalytic effect of the mixture of these is greater when compared with the individual catalysts and thus the positive synergistic catalytic activity of Ru(III) + Os(VIII) mixture. Therefore the observed reactivity in the present study follows the order: Ru(III) + Os(VIII) > Os(VIII) > Ru(III). Consequently, we can conclude that the synergistic effect of Ru(III) + Os(VIII) mixture is greater than the sum of their separate effects in the present study.

Conclusion

Ru(III), Os(VIII) and a mixture of both catalyzed synthesis of anthranilic acids from indoles has been performed efficiently using bromamine-B. The aforementioned metal ions were found to be efficient catalysts for the facile synthesis of anthranilic acids in aqueous alkaline acetonitrile medium. Ru(III) + Os(VIII) serves as an efficient catalyst when compared with the individual catalysts and shows positive synergetic catalytic activity. The stoichiometry and anthranilic acids products of Ru(III), Os(VIII) and Ru(III) + Os(VIII) catalysis were same but their observed kinetic patterns and mechanistic aspects were different, obeying the underlying rate law:

$$\text{rate} = k[\text{BAB}]_r[\text{Indole}][\text{Ru(III)}]^x[\text{OH}^-]^y$$

$$\text{rate} = k[\text{BAB}]_r[\text{Indole}][\text{Os(VIII)}][\text{OH}^-]^y$$

$$\text{rate} = k[\text{BAB}]_r[\text{Indole}]^0[\text{Ru(III)} + \text{Os(VIII)}][\text{OH}^-]^y$$

A comparison of the kinetics of the reactions catalyzed by these catalysts, under identical experimental conditions, was found to be in the order: Ru(III) + Os(VIII) > Os(VIII) > Ru(III). Further, under identical experimental conditions, the kinetics of all the three catalyzed reactions were compared with uncatalyzed reactions and were found to proceed 6- to 42-fold faster than the uncatalyzed reactions. Based on the observed experimental results, detailed mechanistic interpretations and the related kinetic modelings were worked out for each catalyst. It can be concluded that Ru(III), Os(VIII) and Ru(III) + Os(VIII) act as efficient catalysts in the synthesis of anthranilic acids from indoles brought about by BAB in alkaline medium and predominantly the synergism of Ru(III) + Os(VIII) mixed catalyst is prodigious.

Supporting information

Supporting information may be found in the online version of this article.

References

- [1] J. K. Joshi, V. R. Patel, K. Patel, *Indian J. Chem. Pharm. Sci.* **2007**, *69*, 697.

- [2] A. Thorarensen, J. Li, B. D. Wakefield, D. L. Romero, K. R. Marotti, M. T. Sweeney, G. E. Zurenko, *Bioorg. Med. Chem. Lett.* **2007**, *17*, 3113.
- [3] H. Jan, S. Jan, *Collect. Czech. Chem. Commun.* **1996**, *61*, 941.
- [4] J. F. M. Silva da, S. J. Garden, A. C. Pinto, *J. Braz. Chem. Soc.* **2001**, *12*, 273.
- [5] E. Chapman, H. Stephen, *J. Chem. Soc. Trans.* **1925**, 127, 1791.
- [6] M. M. Campbell, G. Johnson, *Chem. Rev.* **1978**, *78*, 65.
- [7] E. Kolvari, A. G. Choghmarani, P. Salehi, F. Shirini, M. A. Zolfigol, *J. Iran. Chem. Soc.* **2007**, *4*, 126.
- [8] R. V. Jagadeesh, Puttaswamy, *J. Phys. Org. Chem.* **2008**, *21*, 844.
- [9] Puttaswamy, R. V. Jagadeesh, *Appl. Catal. A: Gen.* **2005**, *292*, 259.
- [10] K. K. Baneji, B. Jayaram, D. S. Mahadevappa, *J. Sci. Ind. Res.* **1987**, *46*, 65.
- [11] D. H. Bremner, *Synth. Reag.* **1984**, *6*, 9.
- [12] K. S. Rangappa, M. P. Ragavendra, D. S. Mahadevappa, D. C. Gowda, *J. Org. Chem.* **1998**, *63*, 531.
- [13] F. A. Cotton, G. Wilkinson, P. L. Gaus, *Basic Inorganic Chemistry*, 3rd edn, John Wiley & Sons Inc.: New York, **1995**, p. 600.
- [14] W. P. Griffith, *The Chemistry of Rare Platinum Metals*, Interscience: New York, **1967**.
- [15] K. R. Bhat, K. Jyothi, B. T. Gowda, *Oxid. Commun.* **2002**, *25*, 117 and references therein.
- [16] M. S. Ahmed, D. S. Mahadevappa, *Talanta* **1980**, *27*, 669.
- [17] A. I. Vogel, *Textbook of Quantitative Inorganic Analysis*, 4th edn. ELBS-Longman: London, **1978**, p. 427.
- [18] J. C. Morris, J. R. Salazar, M. A. Winemann, *J. Am. Chem. Soc.* **1948**, *70*, 2036.
- [19] E. Bishop, V. J. Jennings, *Talanta* **1958**, *1*, 197.
- [20] B. G. Pryde, F. G. Soper, *J. Chem. Soc.* 1931, 1514; **1926**, 1582.
- [21] F. F. Hardy, J. P. Johnston, *J. Chem. Soc. Perkin Trans. 2* **1973**, 742.
- [22] D. S. Mahadevappa, K. S. Rangappa, N. M. M. Gowda, B. T. Gowda, *Int. J. Chem. Kinet.* **1982**, *14*, 1183.
- [23] S. Meenakshisundaram, R. M. Sockalingam, *J. Mol. Catal. A Chem.* **2000**, *160*, 269.
- [24] Puttaswamy, Nirmala Vaz, *Stud. Surf. Sci. Catal.* **2001**, *133*, 535.
- [25] Puttaswamy, R. V. Jagadeesh, *Cent. Eur. J. Chem.* **2005**, *3*, 482.
- [26] Puttaswamy, Nirmala Vaz, *Transition Met. Chem.* **2003**, *28*, 409.
- [27] M. H. Kondarasaiah, S. Ananda, Puttaswamy, N. M. M. Gowda, *Synth. React.* **2003**, *33*, 1145.
- [28] Puttaswamy, R. V. Jagadeesh, *Int. J. Chem. Kinet.* **2005**, *37*, 201.
- [29] Puttaswamy, R. V. Jagadeesh, *Ind. Eng. Chem. Res.* **2006**, *45*, 1563.
- [30] D. C. Bilehal, R. M. Kulkarni, S. T. Nandibewoor, *Can. J. Chem.* **2001**, *79*, 1926.
- [31] R. T. Mahesh, M. B. Bellaki, S. T. Nandibewoor, *Catal. Lett.* **2004**, *97*, 91.
- [32] C. P. Kathari, R. M. Mulla, S. T. Nandibewoor, *Oxid. Commun.* **2005**, *28*, 579.
- [33] S. A. Chimatdar, T. Basavaraj, S. T. Nandibewoor, *Inorg. React. Mechanism.* **2002**, *4*, 209.
- [34] M. Ardon, *J. Chem. Soc.* **1957**, 1811.
- [35] N. Kambo, S. K. Upadhyay, *Indian J. Chem.* **2004**, *43A*, 1210.
- [36] C. J. Collins, N. S. Bowman, *Isotope Effects in Chemical Reactions*, Van Nostrand Reinhold: New York, **1970**, p. 267.
- [37] K. B. Wiberg, *Chem. Rev.* **1955**, *55*, 713; b) K. B. Wiberg, *Physical Organic Chemistry*, Wiley: New York, **1964**.
- [38] E. A. Moelwyn-Hughes, *Kinetics of Reaction in Solutions*, Oxford University Press: London, **1947**, pp. 297.
- [39] A. M. Mackay, R. A. Mackay, *Introduction to Modern Inorganic Chemistry*, 4th edn, Prentice Hall: Englewood Cliffs, NJ, **1989**, p. 259.
- [40] R. D. Saverbrum, E. B. Sandell, *J. Am. Chem. Soc.* **1953**, *75*, 4170.
- [41] K. J. Laidler, *Chemical Kinetics*, 2nd edn. McGraw-Hill: New York, **1965**, pp. 219.