

## § 14. Formulation of Lagrangian Neoclassical Transport Theory

Satake, S. (Grad. Univ. Advanced Studies)  
Okamoto, M., Sugama, H.

Recently, neoclassical transport in the core region of tokamaks again attracts much attention. It is well-known that there appear non-standard guiding-center orbits near the magnetic axis called “potato” orbits[1]. Typical orbit width of potato particles is as large as  $(q^2\rho^2R_0)^{1/3}$ . The standard neoclassical transport theory[2,3] constructed in the small-orbit-width (SOW) approximation is not applicable to the near-axis region, and the orbital properties of potato particles need to be considered in analyzing transport in this region. In this purpose, we adopt a Lagrangian formulation[4] which is suitable for a collisionless (banana-regime) plasma with finite-width orbits. In this approach, transport phenomena are described by a reduced drift-kinetic equation in the space of three constants-of-motion (COM) along a collisionless particle orbit in a tokamak.

We apply for the first time Lagrangian formulation to the near-axis region[5] in which the finite-orbit-width (FOW) effect becomes really important. To utilize Lagrangian transport theory, we improve the treatment of like-particle collision term by adding a momentum-restoring term to a simple Lorentz operator. This transport theory reflects quantitatively the properties of all types of particles appearing near the magnetic axis.

The Lagrangian formulation is derived from the drift kinetic equation in an Eulerian representation in the  $(\mathbf{x}, \mathcal{E}, \mu)$  space,

$$\frac{\partial}{\partial t} f_a(\mathbf{x}, \mathcal{E}, \mu, t) + \dot{\mathbf{x}} \cdot \frac{\partial f_a}{\partial \mathbf{x}} = C_{ab}, \quad (1)$$

where  $\dot{\mathbf{x}} = d/dt$ ,  $\mathbf{x}$  is the guiding-center position and  $C_{ab}$  is a collision operator. The independent variables in eq. (1) are transformed into three constants of motion in the collisionless limit  $(z_1, z_2, z_3) = (\mathcal{E}, \mu, \langle\psi\rangle)$ , and the other three variables  $(\tilde{z}_4, \tilde{z}_5, \tilde{z}_6)$ . Here,  $\mathcal{E}$  is the energy of a particle,  $\mu$  is the magnetic momentum, and  $\langle\psi\rangle$  represents the averaged radial position of a particle orbit. The orbit average operator for any function  $a(\mathbf{z}, \tilde{\mathbf{z}})$  is defined as

$$\langle a \rangle \equiv \frac{1}{4\pi^2\tau_p} \oint \frac{d\theta}{\dot{\theta}} d\zeta d\phi a(\mathbf{z}, \tilde{\mathbf{z}}), \quad (2)$$

where  $\tau_p \equiv \oint d\theta/\dot{\theta}$  is the poloidal period of an particle orbit. By using the set of variables  $(\mathbf{z}, \tilde{\mathbf{z}})$ , eq. (1) is transformed into

$$\frac{\partial}{\partial t} f_a(\mathbf{z}, \theta, t) + \dot{\theta} \frac{\partial f_a}{\partial \theta} = C_{ab}, \quad (3)$$

where the property  $\partial/\partial\phi = \partial/\partial\zeta = 0$  is used.

In the collisionless regime, averaging both sides of eq. (3) yields the reduced drift-kinetic equation in the COM space,

$$\frac{\partial \bar{f}}{\partial t} = \frac{1}{J_c} \frac{\partial}{\partial \mathbf{z}} \cdot \left( J_c \left\langle \frac{\partial \mathbf{z}}{\partial \mathbf{v}} \cdot \Gamma(\bar{f}) \right\rangle \right) = \bar{C}, \quad (4)$$

where  $\bar{f} = f(\mathcal{E}, \mu, \langle\psi\rangle, t)$  and  $J_c$  is Jacobian.

Equation (4) is expanded in  $\delta_b = \Delta_b/L \ll 1$  and then solved order by order. Here,  $\Delta_b$  and  $L$  are typical potato width and typical gradient scale, respectively. By taking moments of  $O(\delta_b^2)$  part of eq. (4) with  $\mathcal{E}$  and  $\mu$ , one obtains the particle and heat flux equations,

$$\begin{aligned} \frac{\partial \bar{n}_i}{\partial t} + \frac{\lambda_n}{V'} \frac{\partial}{\partial \langle\psi\rangle} (V' \Gamma_i) &= 0, \quad (5) \\ \frac{\partial}{\partial t} \left( \frac{3}{2} \bar{n}_i T_i \right) + \frac{\lambda_q}{V'} \frac{\partial}{\partial \langle\psi\rangle} \left[ V' \left( q_i + \frac{3}{2} \Gamma_i T_i \right) \right] \\ &= -\lambda_q e_i \Gamma_i \frac{d\Phi}{d\langle\psi\rangle}, \quad (6) \end{aligned}$$

where

$$\begin{bmatrix} \Gamma_i \\ \frac{q_i}{T_i} \end{bmatrix} = \begin{bmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{bmatrix} \cdot \begin{bmatrix} \frac{d \ln \bar{n}_i}{d\langle\psi\rangle} + \frac{e_i}{T_i} \frac{d\Phi}{d\langle\psi\rangle} \\ \frac{d \ln T_i}{d\langle\psi\rangle} \end{bmatrix}. \quad (7)$$

These equations are in the comparable form to those of standard Eulerian representation. The difference arose by using  $\langle\psi\rangle$  instead of  $\psi$  can be seen to be factorized by  $\lambda_n$  and  $\lambda_q$ . These factors tend to be unity away from the axis and go infinity as  $\langle\psi\rangle \rightarrow 0$ . The transport coefficients  $A_{jk}$  have a convenient form to calculate numerically by means of Monte Carlo integration[5]. Fig. 1 is an example of calculated ion thermal conductivity  $\chi_i$ . Near the magnetic axis,  $\chi_i$  is found to decrease from the value predicted by the conventional theory in the SOW limit. The reductive tendency of  $\chi_i$  is common feature in recent experimental diagnostics and  $\delta f$  simulations. Our approach has first succeeded to explain this tendency by proper analytic model.




Fig. 1 :  $\chi_i$  in the near-axis region. Solid: numerical result, dashed: Conventional theory from [2].

- 1) Satake, S. *et al.*, **77**, 573 (2001).
- 2) Hinton, F. L. *et al.*, *Rev. Mod. Phys.* **48**, 239 (1976).
- 3) Hirshman, S. P. *et al.*, *Nuclear Fusion* **21**, 1079 (1981).
- 4) Bernstein, I. B. *et al.*, *Phys. Fluids* **26**, 1488 (1983).
- 5) Satake, S. *et al.*, *Phys. Plasmas* **9**, 3946 (2002).