

§35. Dependence of Local Transport on Beta and Configurations in $\gamma = 1.20$ Plasmas

Funaba, H., Watanabe, K.Y.,
LHD Experiment Group

High-beta plasmas of more than 5% of $\langle\beta\rangle$ were produced in a magnetic configuration with the optimal aspect ratio $A_p = 6.6$ ($\gamma = 1.20$) [1], while the usual A_p value is 5.8 ($\gamma = 1.254$) in the low-beta plasmas on LHD. From the results of the previous local transport analysis for the high-beta plasmas with $A_p = 5.8$, the local transport property, which is evaluated by referring to ISS04, is preserved or improved in the core region, while it is degraded in the peripheral region in the $A_p = 5.8$ configuration [2]. In this study, the local transport analysis for the high-beta plasmas with $A_p = 6.6$ are made and the results are compared with those of the $A_p = 5.8$ case. In order to evaluate the degradation in the local transport which is caused from reasons other than the change of the magnetic configuration, a new factor $g_{\text{ren}\chi}^{\text{int}}$ is introduced. This factor is derived by interpolating the renormalization factor for the local transport at each ρ . Since the effects of the change of the magnetic configurations on the local transport coefficients in the high beta region are represented by $g_{\text{ren}\chi}^{\text{int}}$, the effects of the beta increment or the increment in gradient beta are appear in $\chi^{\text{eff}}/(g_{\text{ren}\chi}^{\text{int}}\chi^{\text{ISS04}})$. Here, χ^{ISS04} is the model transport coefficient which has the same non-dimensional parameter dependence as ISS04.

Figures 1 and 2 show the beta dependences of the normalized local transport coefficient $\chi^{\text{eff}}/\chi^{\text{ISS04}}$ in the $A_p = 5.8$ and 6.6 configurations at (a) $\rho = 0.5$ and (b) $\rho = 0.9$, respectively. The outlines of the data region more than 1% of $\langle\beta\rangle$ in the configuration of $A_p = 5.8$ are shown by the dashed lines in Figs 1 (a) and (b). In Fig. 1 (b), as $\chi^{\text{eff}}/(g_{\text{ren}\chi}^{\text{int}}\chi^{\text{ISS04}})$ becomes large with the increment in $\langle\beta\rangle$, transport degradation due to effects other than the change of the magnetic configuration exists at $\rho = 0.9$. The results in the $A_p = 6.6$ configuration are shown in Fig. 2. The dashed lines in Figs 2 (a) and (b) are the same as those in Fig. 2 (a) and (b), respectively. Figure 2 (b) shows that the degradation of the local transport is larger than the effect of the change of the configuration in the $\langle\beta\rangle > 1\%$ region at $\rho = 0.9$. However, it is different from the case of $A_p = 5.8$ that the ratio $\chi^{\text{eff}}/(g_{\text{ren}\chi}^{\text{int}}\chi^{\text{ISS04}})$ seems to decrease in $\langle\beta\rangle > 2\%$. As shown in Fig. 2 (a), $\chi^{\text{eff}}/(g_{\text{ren}\chi}^{\text{int}}\chi^{\text{ISS04}})$ also seems to be improved in the $\langle\beta\rangle > 2\%$ region at $\rho = 0.5$ in $A_p = 6.6$.




Fig. 1. Beta dependence of the normalized local transport coefficient in the $A_p = 5.8$ configuration at (a) $\rho = 0.5$ and (b) $\rho = 0.9$.




Fig. 2. Beta dependence of the normalized local transport coefficient in the $A_p = 6.6$ configuration at (a) $\rho = 0.5$ and (b) $\rho = 0.9$.

[1] S.Sakakibara, *et al.*, Plasma Phys. Control. Fusion **50** (2008) 124014.

[2] H.Funaba, K.Y.Watanabe, *et al.*, Plasma Fusion Res. **3** (2008) 022.