



INTERNATIONAL ATOMIC ENERGY AGENCY

18th IAEA Fusion Energy Conference  
Sorrento, Italy, 4 to 10 October 2000

IAEA-CN-77/THP2/08

## NATIONAL INSTITUTE FOR FUSION SCIENCE

### Improved Stability due to Local Pressure Flattening in Stellarators

K. Ichiguchi, M. Wakatani, T. Unemura,  
T. Tatsuno and B.A. Carreras

(Received - Sep. 13, 2000 )

NIFS-649

Sep. 2000

This report was prepared as a preprint of work performed as a collaboration research of the National Institute for Fusion Science (NIFS) of Japan. This document is intended for information only and for future publication in a journal after some rearrangements of its contents.

Inquiries about copyright and reproduction should be addressed to the Research Information Center, National Institute for Fusion Science, Oroshi-cho, Toki-shi, Gifu-ken 509-02 Japan.

## RESEARCH REPORT NIFS Series

This is a preprint of a paper intended for presentation at a scientific meeting. Because of the provisional nature of its content and since changes of substance or detail may have to be made before publication, the preprint is made available on the understanding that it will not be cited in the literature or in any way be reproduced in its present form. The views expressed and the statements made remain the responsibility of the named author(s); the views do not necessarily reflect those of the government of the designating Member State(s) or of the designating organization(s). In particular, neither the IAEA nor any other organization or body sponsoring this meeting can be held responsible for any material reproduced in this preprint.

TOKI, JAPAN

# Improved Stability due to Local Pressure Flattening in Stellarators

K. Ichiguchi 1), M. Wakatani 2), T. Unemura 2), T. Tatsuno 3), B. A. Carreras 4)

1) National Institute for Fusion Science, Oroshi-cho 322-6, Toki, Japan 509-5292

2) Graduate School of Energy Science, Kyoto University, Gokasho, Uji, Japan 611-0011

3) Graduate School of Frontier Science, University of Tokyo, Tokyo, Japan 113-8656

4) Oak Ridge National Laboratory, Oak Ridge, Tennessee 37831, USA

e-mail: wakatani@energy.kyoto-u.ac.jp

**Abstract.** It is demonstrated that the stability of low  $n$  pressure gradient driven modes is improved by introducing local pressure flattening at low order rational surfaces in LHD (Large Helical Device) with the inward magnetic axis shift of 25cm, where  $n$  is a toroidal mode number.

## 1. Introduction

The largest stellarator/heliotron device called LHD (Large Helical Device) started physics experiments successfully [1]. The electron and ion temperatures,  $T_e \sim 3.8\text{keV}$  and  $T_i \sim 2.8\text{keV}$ , were obtained in the low density range,  $\bar{n}_e \sim 1.5 \times 10^{19}\text{m}^{-3}$  [2]. The obtained energy confinement time was about 50% better than the International Stellarator Scaling of energy confinement [3]. The maximum average beta value,  $\bar{\beta} \simeq 2.4\%$ , exceeded 2% already, which was the highest beta obtained in stellarator/heliotron devices [4]. Since  $\bar{\beta} \simeq 2.4\%$  is not limited by MHD instabilities and the target beta value of LHD is 5%, a higher power heating is expected in an optimized magnetic configuration.

In this paper we will discuss effects of local pressure profile flattening on interchange modes which may affect stability and confinement properties of LHD. For studying MHD stability in stellarator/heliotron devices, Mercier criterion [5] is valuable. For three-dimensional MHD equilibria under the assumption of existence of flux surfaces, Mercier criterion is usable for evaluating the beta limit [6]. Another important ingredient for three-dimensional MHD equilibrium and stability is the formation of magnetic islands [7]. This problem is related to the existence of three-dimensional nested flux surfaces [8]. The magnetic islands may be produced by resonant perturbed magnetic fields which are generated by internal resistive MHD instabilities or external error fields. If the magnetic islands appear at low order rational surfaces, it is expected that the pressure profile becomes flat in the island regions. It is shown that the MHD stability based on Mercier criterion changes significantly, although the pressure flattening is highly localized in the neighborhood of rational surface [9] [10] [11]. For this situation stability limit of low mode number interchange mode becomes important.

## 2. Reduced MHD Equations and a Model Pressure Profile with Locally Flat Regions at Rational Surfaces

For analyzing pressure driven instabilities in stellarator/heliotron devices, we use the ideal reduced MHD equations [6] [10] [12], which are written as

$$\frac{\partial \psi}{\partial t} = -(R/R_0)^2 \mathbf{B} \cdot \nabla u \quad (1)$$

$$\rho \frac{d}{dt} \nabla_{\perp}^2 u = -\mathbf{B} \cdot \nabla (\nabla_{\perp}^2 \psi) + R_0^2 \nabla \Omega \times \nabla P \cdot \nabla \zeta \quad (2)$$

$$\frac{dP}{dt} = 0 \quad (3)$$

where

$$\mathbf{B} \cdot \nabla = \frac{R_0 B_0}{R^2} \frac{\partial}{\partial \zeta} - \nabla \psi \times \nabla \zeta \cdot \nabla \quad (4)$$

$$\frac{d}{dt} = \frac{\partial}{\partial t} + (R/R_0)^2 \nabla u \times \nabla \zeta \cdot \nabla \quad (5)$$

$$\Omega = \frac{1}{2\pi} \int_0^{2\pi} d\zeta \left( \frac{R}{R_0} \right)^2 \left( 1 + \frac{|\mathbf{B} - \bar{\mathbf{B}}|^2}{B_0^2} \right). \quad (6)$$

Here  $\psi$ ,  $u$  and  $P$  denote the poloidal flux function, the stream function and the plasma pressure, respectively. An axisymmetric component of the magnetic field is given by  $\bar{\mathbf{B}}$  and  $R$  and  $\zeta$  denote the major radius and the toroidal angle, respectively. The magnetic axis is  $R = R_0$ , and the toroidal field at  $R = R_0$  is  $B_0$ . Since the free-boundary effect is not significant for the stability of currentless plasmas, a perfectly conducting wall is usually placed at the plasma boundary [10]. It is noted that the equilibrium state of  $\psi$  is consistent with the rotational transform profile due to stellarator fields.

For describing the locally flat pressure profile,

$$P(\rho) = C[P_0(\rho) + P_{ax}(\rho) + P_{res}(\rho) - A] \quad (7)$$

is assumed, where  $P_0(\rho)$  denotes a smooth and standard pressure profile,  $P_{ax}(\rho)$  corresponds to a flattening of pressure profile near the magnetic axis given by

$$P_{ax}(\rho) = [P_0(0) - P_0(\rho)] \exp \left[ -\frac{1}{2} \left( \frac{\rho}{w_a} \right)^4 \right], \quad (8)$$

and  $P_{res}(\rho)$  plays a role to flatten the pressure at rational surfaces

$$P_{res}(\rho) = \sum_m \{ [P_0(\rho_m) + P_{ax}(\rho_m)] - [P_0(\rho) + P_{ax}(\rho)] \} \\ \times \exp \left[ -\frac{1}{2} \left( \frac{\rho - \rho_m}{w_m} \right)^4 \right]. \quad (9)$$

Here  $\rho$  denotes a square root of normalized toroidal flux. In the expression of (7),  $A$  and  $C$  are numerical factors to fix pressures at both the magnetic axis and the plasma surface. In the expression of (8),  $w_a$  denotes a width of a region to make the pressure profile flat near the magnetic axis. Also in the expression of (9),  $\rho_m$  denotes a position of  $m$ -th rational surface and  $w_m$  denotes a width of a region to make the pressure profile flat at the rational surface  $\rho = \rho_m$ .

In order to calculate fixed boundary MHD equilibria for the LHD configuration with the pressure shown by the expression (7), VMEC code was applied. In the following calculations the LHD configuration with the inward magnetic axis shift of 25cm is picked up. Ideal MHD stability against pressure driven interchange modes was studied with the RESORM code [13], which solves linearized equations of eqs.(1) - (3) as an initial value problem.

### 3. Stabilization of Low- $n$ Interchange Modes with Flat Pressure Regions at Rational Surfaces in Toroidal Plasmas

For the pressure profile given by Eq.(7) in the LHD model configuration, global pressure driven

modes with  $n = 1, 2, 3$  are examined with the RESORM code [13], where  $n$  is a toroidal mode number.

(I)  $n = 1$  mode

The smooth pressure profile  $P_0(\rho)$  and the rotational transform profile  $\iota(\rho)$  are plotted in Fig.1. The rational surfaces for the  $n = 1$  mode,  $\iota = 1/1$ , and  $\iota = 1/2$  are shown with the dotted lines in Fig.1. Here the central beta value is assumed 2%. The unstable  $n = 1$  mode exists with the growth rate of  $4.846 \times 10^{-2}$ , which is destabilized at the  $\iota = 1/2$  surface. Here the growth rate is normalized by the poloidal Alfvén time. When the flat pressure region with the width  $w$  is increased at the  $\iota = 1/2$  surface, the growth rate decreases. The marginally stable pressure profile against the  $n = 1$  with  $w = 0.045$  is shown in Fig.2.

(II)  $n = 2$  mode

Here the same pressure and rotational transform profiles as shown in Fig.1 are used for the stability analysis of the  $n = 2$  mode. However, the relevant rational surfaces increases;  $\iota = 2/2$ ,  $\iota = 2/3$ ,  $\iota = 2/4$  and  $\iota = 2/5$ . The RESORM code shows that the  $n = 2$  mode is destabilized at the two rational surfaces,  $\iota = 2/4$  and  $\iota = 2/5$ . Here the growth rate is  $\gamma = 7.187 \times 10^{-2}$  at  $\beta(0) = 2\%$ . Thus it is required to introduce two locally flat pressure regions with different widths at  $\iota = 2/4$  and  $\iota = 2/5$  for stabilizing the  $n = 2$  mode. When  $w = 0.02$  at  $\iota = 2/4$  and  $w = 0.04$  at  $\iota = 2/5$ , the instability is suppressed completely and the obtained pressure profile is shown in Fig.3.

(III)  $n = 3$  mode

For the LHD configuration with the pressure and rotational transform profiles shown in Fig.1, there are six rational surfaces;  $\iota = 3/3, 3/4, 3/5, 3/6, 3/7, 3/8$ . For the case of Fig.1 the RESORM code gives the growth rate  $\gamma = 8.233 \times 10^{-2}$  at  $\beta(0) = 2\%$ , and the unstable mode is localized at the central region with a ballooning structure. These are typical characteristics of the toroidal non-resonant pressure-driven mode [14]. In order to suppress this non-resonant mode, the central pressure profile is flattened first with  $w_a = 0.6$  in Eq.(8). Then the growth rate decreases to  $\gamma = 6.975 \times 10^{-2}$ , and the unstable mode has a typical interchange mode structure destabilized at  $\iota = 3/4, 3/5, 3/6$  and  $3/7$ . For suppressing the pressure driven interchange mode with  $n = 3$  completely, flat pressure regions are generated at the four rational surfaces with  $w = 0.03$  at  $\iota = 3/7$ ,  $w = 0.03$  at  $\iota = 3/6$ ,  $w = 0.02$  at  $\iota = 3/5$  and  $w = 0.02$  at  $\iota = 3/4$ . The obtained pressure profile with  $\beta(0) = 2\%$  is shown in Fig.4. The average beta value is changed from  $\bar{\beta} = 0.632\%$  (see Fig.1) to  $\bar{\beta} = 1\%$  (see Fig.4).

It was demonstrated that the pressure driven modes with  $n = 1, n = 2$  and  $n = 3$  can be stabilized by generating the locally flat pressure regions at the relevant rational surfaces separately. Furthermore it is confirmed that the  $n = 1, 2, 3$  modes become stable simultaneously when the pressure profile is described with  $w_a = 0.6$  and locally flat regions with  $w = 0.025, 0.03, 0.065, 0.03, 0.025, 0.02$  at  $\iota = 0.4, 3/7, 0.5, 0.6, 2/3, 0.75$ , respectively.

#### 4. Concluding Remarks

It is expected that the locally flat pressure regions are produced by the non-linear evolution of resistive interchange modes which become unstable for beta values less than the Mercier limit. The other possibility to produce the locally flat pressure regions is external application of resonant helical magnetic fields. It has been demonstrated that the pressure driven instabilities with low toroidal mode numbers are stabilized by modifying the pressure profile to make the locally

flat pressure regions in the LHD model configuration. It is noted that large pressure gradients near the magnetic axis destabilize the non-resonant modes with medium toroidal mode numbers such as  $n = 3$  or 4. Therefore, in order to increase the ideal beta limit, broad pressure profiles with several locally flat pressure regions at dominant rational surfaces may be appropriate in LHD.

There are some indications that the experimental beta values exceed the Mercier limit, when smooth and monotonic pressure profiles are assumed in CHS [15] [16] and Heliotron E [17]. Formation of the above mentioned locally flattened pressure profiles at low order rational surfaces may explain the discrepancy. The important assumption is that the resistive interchange instabilities unstable in the magnetic hill region are responsible for generating such profiles. The other possibility to explain the discrepancy is that high- $n$  pressure driven modes do not play a role due to the finite Larmor radius stabilization [6]. It is noted that the LHD high beta plasma already obtained  $\bar{\beta} \simeq 2.4\%$ , which seems to exceed the Mercier limit. Future experiments for the LHD high beta plasmas will be expected for checking of our conjecture.

## References

- [1] MOTOJIMA, O., et al., Phys. Plasmas **6** (1999) 1843.
- [2] OHYABU, N., et al., Phys. Plasmas **7** (2000) 1802.
- [3] YAMADA, H., et al., Phys. Rev. Lett. **84** (2000) 1216.
- [4] SAKAKIBARA, S., et al., this conferece, paper EXP3/12
- [5] MERCIER, C., Nucl. Fusion **1** (1960) 47.
- [6] WAKATANI, M., *Stellarator and Heliotron Devices* (Oxford University Press, 1998)
- [7] HEGNA, C.C., and BHATTACHARJEE, A., Phys. Fluids **B 1** (1989) 392.
- [8] NAKAMURA, Y., et al., Phys. Plasmas **B 2** (1990) 2528.
- [9] TATSUNO, T., et al., Nucl. Fusion **39** (1999) 1391.
- [10] ICHIGUCHI, K., et al., J. Plasma Fusion Res. SERIES **2** (1999) 286.
- [11] CARRERAS, B.A., et al., Plasma Phys. Rep. **25** (1999) 958.
- [12] STRAUSS, H.R., Plasma Phys. **22** (1980) 2733.
- [13] ICHIGUCHI, K., et al., Nucl. Fusion **29** (1989) 2093.
- [14] CARRERAS, B.A., et al., Phys. Plasmas **5** (1998) 3700.
- [15] OKAMURA, S., et al., Nucl. Fusion **35** (1995)
- [16] OKAMURA, S., et al., Nucl. Fusion **39** (1999) 1337.
- [17] WAKATANI, M., et al., Proc. IAEA Conf. on Plasma Phys. and Controlled Nucl. Fusion (Washington, 1990) vol.2, p.567.
- [18] HIRSHMAN, S.P., et al., Comp. Phys. Comm. **43** (1986) 143.




FIG.1. An assumed pressure profile and a rotational transform profile in the LHD configuration with the inward magnetic axis shift of 25cm obtained by VMEC code[18]. Dotted lines show  $\iota = 1/1$  and  $\iota = 1/2$ , respectively. The central beta value is  $\beta(0) = 2\%$  and the average beta value is  $\bar{\beta} = 0.632\%$ . The radius  $\rho_p$  denotes a square root of normalized poloidal flux, which is also used in FIG.2 - FIG.4.




FIG.2. A marginally stable pressure profile against the  $n = 1$  pressure driven mode with the width of flat pressure region  $w = 0.045$ . The rotational transform profile obtained by VMEC code, and rational surfaces with  $\iota = 1/1$  and  $\iota = 1/2$  are also shown.




FIG.3. A marginally stable pressure profile against the  $n = 2$  pressure driven mode with the flat pressure regions given by  $w = 0.04$  at  $\iota = 2/5$  and  $w = 0.02$  at  $\iota = 2/4$ . The rotational transform profile obtained by VMEC code, and rational surfaces with  $\iota = 2/2, 2/3, 2/4, 2/5$  are shown.




FIG.4. A marginally stable pressure profile against the  $n = 3$  pressure driven mode with the flat pressure regions given by  $w = 0.03$  at  $\iota = 3/7, w = 0.03$  at  $\iota = 3/6, w = 0.02$  at  $\iota = 3/5$  and  $w = 0.02$  at  $\iota = 3/4$ . The rotational transform profile obtained by VMEC code, and rational surfaces with  $\iota = 3/3, 3/4, 3/5, 3/6, 3/7, 3/8$  are shown.

## Recent Issues of NIFS Series

- NIFS-587 T Nakamura and T Yabe,  
Cubic Interpolated Propagation Scheme for Solving the Hyper-Dimensional Vlasov-Poisson Equation in Phase Space Mar 1999
- NIFS-588 W.X. Wnag, N Nakajima, S Murakami and M. Okamoto,  
An Accurate  $\delta f$  Method for Neoclassical Transport Calculation, Mar 1999
- NIFS-589 K. Kishida, K. Araki, S. Kishiba and K Suzuki,  
Local or Nonlocal? Orthonormal Divergence-free Wavelet Analysis of Nonlinear Interactions in Turbulence; Mar 1999
- NIFS-590 K. Araki, K. Suzuki, K. Kishida and S. Kishiba,  
Multiresolution Approximation of the Vector Fields on  $T^3$ , Mar 1999
- NIFS-591 K Yamazaki, H. Yamada, K.Y. Watanabe, K. Nishimura, S. Yamaguchi, H. Nakanishi, A. Komori, H. Suzuki, T. Mito, H. Chikaraishi, K. Murai, O. Motojima and the LHD Group,  
Overview of the Large Helical Device (LHD) Control System and its First Operation, Apr 1999
- NIFS-592 T. Takahashi and Y Nakao,  
Thermonuclear Reactivity of D-T Fusion Plasma with Spin-Polarized Fuel; Apr 1999
- NIFS-593 H. Sugama,  
Damping of Toroidal Ion Temperature Gradient Modes, Apr 1999
- NIFS-594 Xiaodong Li,  
Analysis of Crowbar Action of High Voltage DC Power Supply in the LHD ICRF System, Apr 1999
- NIFS-595 K. Nishimura, R. Honuchi and T. Sato,  
Drift-kink Instability Induced by Beam Ions in Field-reversed Configurations; Apr. 1999
- NIFS-596 Y Suzuki, T-H. Watanabe, T. Sato and T. Hayashi,  
Three-dimensional Simulation Study of Compact Toroid Plasmoid Injection into Magnetized Plasmas; Apr. 1999
- NIFS-597 H. Sanuki, K. Itoh, M. Yokoyama, A. Fujisawa, K. Ida, S. Toda, S-I. Itoh, M. Yagi and A. Fukuyama,  
Possibility of Internal Transport Barrier Formation and Electric Field Bifurcation in LHD Plasma;  
May 1999
- NIFS-598 S. Nakazawa, N. Nakajima, M. Okamoto and N. Ohyaibu,  
One Dimensional Simulation on Stability of Detached Plasma in a Tokamak Divertor; June 1999
- NIFS-599 S. Murakami, N. Nakajima, M. Okamoto and J. Nhrenberg,  
Effect of Energetic Ion Loss on ICRF Heating Efficiency and Energy Confinement Time in Heliotrons;  
June 1999
- NIFS-600 R. Honuchi and T. Sato,  
Three-Dimensional Particle Simulation of Plasma Instabilities and Collisionless Reconnection in a Current Sheet,, June 1999
- NIFS-601 W. Wang, M. Okamoto, N. Nakajima and S. Murakami,  
Collisional Transport in a Plasma with Steep Gradients; June 1999
- NIFS-602 T. Mutoh, R. Kumazawa, T. Saki, K. Saito, F. Simpo, G. Nomura, T. Watari, X. Jikang, G. Cattanei, H. Okada, K. Ohkubo, M. Sato,

S. Kubo, T. Shimozuma, H. Idei, Y. Yoshimura, O. Kaneko, Y. Takeiri, M. Osakabe, Y. Oka, K. Tsumori, A. Komori, H. Yamada, K. Watanabe, S. Sakakibara, M. Shoji, R. Sakamoto, S. Inagaki, J. Miyazawa, S. Morita, K. Tanaka, B.J. Peterson, S. Murakami, T. Minami, S. Ohdachi, S. Kado, K. Narihara, H. Sasao, H. Suzuki, K. Kawahata, N. Ohyabu, Y. Nakamura, H. Funaba, S. Masuzaki, S. Muto, K. Sato, T. Monsaki, S. Sudo, Y. Nagayama, T. Watanabe, M. Sasao, K. Ida, N. Noda, K. Yamazaki, K. Akaishi, A. Sagara, K. Nishimura, T. Ozaki, K. Toi, O. Motojima, M. Fujiwara, A. Iyoshi and LHD Exp. Group 1 and 2,  
**First ICRF Heating Experiment in the Large Helical Device;** July 1999

- NIFS-603 P.C. de Vries, Y. Nagayama, K. Kawahata, S. Inagaki, H. Sasao and K. Nagasaki,  
**Polarization of Electron Cyclotron Emission Spectra in LHD;** July 1999
- NIFS-604 W. Wang, N. Nakajima, M. Okamoto and S. Murakami,  
 **$\delta f$  Simulation of Ion Neoclassical Transport;** July 1999
- NIFS-605 T. Hayashi, N. Mizuguchi, T. Sato and the Complexity Simulation Group,  
**Numerical Simulation of Internal Reconnection Event in Spherical Tokamak;** July 1999
- NIFS-606 M. Okamoto, N. Nakajima and W. Wang,  
**On the Two Weighting Scheme for  $\delta f$  Collisional Transport Simulation;** Aug. 1999
- NIFS-607 O. Motojima, A.A. Shishkin, S. Inagaki, K.Y. Watanabe,  
**Possible Control Scenario of Radial Electric Field by Loss-Cone-Particle Injection into Helical Device;** Aug. 1999
- NIFS-608 R. Tanaka, T. Nakamura and T. Yabe,  
**Constructing Exactly Conservative Scheme in Non-conservative Form;** Aug 1999
- NIFS-609 H. Sugama,  
**Gyrokinetic Field Theory;** Aug. 1999
- NIFS-610 M. Takechi, G. Matsunaga, S. Takagi, K. Ohkuni, K. Toi, M. Osakabe, M. Isobe, S. Okamura, K. Matsuoka, A. Fujisawa, H. Iguchi, S. Lee, T. Minami, K. Tanaka, Y. Yoshimura and CHS Group,  
**Core Localized Toroidal Alfvén Eigenmodes Destabilized By Energetic Ions in the CHS Heliotron/Torsatron;** Sep. 1999
- NIFS-611 K. Ichiguchi,  
**MHD Equilibrium and Stability in Heliotron Plasmas;** Sep. 1999
- NIFS-612 Y. Sato, M. Yokoyama, M. Wakatani and V. D. Puzovtsov,  
**Complete Suppression of Pfirsch-Schluter Current in a Toroidal  $l=3$  Stellarator;** Oct. 1999
- NIFS-613 S. Wang, H. Sanuki and H. Sugama,  
**Reduced Drift Kinetic Equation for Neoclassical Transport of Helical Plasmas in Ultra-low Collisionality Regime;** Oct. 1999
- NIFS-614 J. Miyazawa, H. Yamada, K. Yasui, S. Kato, N., Fukumoto, M. Nagata and T. Uyama,  
**Design of Spheromak Injector Using Conical Accelerator for Large Helical Device;** Nov. 1999
- NIFS-615 M. Uchida, A. Fukuyama, K. Itoh, S.-I. Itoh and M. Yagi,  
**Analysis of Current Diffusive Ballooning Mode in Tokamaks;** Dec. 1999
- NIFS-616 M. Tanaka, A. Yu. Grosberg and T. Tanaka,  
**Condensation and Swelling Behavior of Randomly Charged Multichain Polymers by Molecular Dynamics Simulations;** Dec. 1999
- NIFS-617 S. Goto and S. Kida,  
**Sparseness of Nonlinear Coupling;** Dec. 1999
- NIFS-618 M.M. Skoric, T. Sato, A. Maluckov and M.S. Jovanovic,  
**Complexity in Laser Plasma Instabilities** Dec. 1999



- NIFS-619 T.H. Watanabe, H Sugama and T Sato,  
**Non-dissipative Kinetic Simulation and Analytical Solution of Three-mode Equations of Ion Temperature Gradient Instability, Dec 1999**
- NIFS-620 Y. Oka, Y Takeiri, Yu I Belchenko, M Hamabe, O Kaneko, K Tsumori, M Osakabe, E Asano, T Kawamoto, R. Akiyama,  
**Optimization of Cs Deposition in the 1/3 Scale Hydrogen Negative Ion Source for LHD-NBI System ,Dec 1999**
- NIFS-621 Yu I Belchenko, Y Oka, O Kaneko, Y Takeiri, A Krivenko, M Osakabe, K Tsumori, E Asano, T Kawamoto, R Akiyama,  
**Recovery of Cesium in the Hydrogen Negative Ion Sources, Dec 1999**
- NIFS-622 Y Oka, O Kaneko, K Tsumori, Y Takeiri, M Osakabe, T Kawamoto, E Asano, and R Akiyama.  
**H- Ion Source Using a Localized Virtual Magnetic Filter in the Plasma Electrode: Type I LV Magnetic Filter: Dec 1999**
- NIFS-623 M Tanaka, S Kida, S Yanase and G Kawahara,  
**Zero-absolute-vorticity State in a Rotating Turbulent Shear Flow, Jan. 2000**
- NIFS-624 F Leuterer, S Kubo,  
**Electron Cyclotron Current Drive at  $\omega \approx \omega_c$  with X-mode Launched from the Low Field Side; Feb 2000**
- NIFS-625 K Nishimura,  
**Wakefield of a Charged Particulate Influenced by Emission Process of Secondary Electrons; Mar 2000**
- NIFS-626 K Itoh, M. Yagi, S.-I. Itoh, A Fukuyama,  
**On Turbulent Transport in Burning Plasmas; Mar. 2000**
- NIFS-627 K Itoh, S.-I. Itoh, L Giannone,  
**Modelling of Density Limit Phenomena in Toroidal Helical Plasmas, Mar. 2000**
- NIFS-628 K Akaishi, M Nakasuga and Y. Funato,  
**True and Measured Outgassing Rates of a Vacuum Chamber with a Reversibly Absorbed Phase; Mar 2000**
- NIFS-629 T Yamagishi,  
**Effect of Weak Dissipation on a Drift Orbit Mapping, Mar. 2000**
- NIFS-630 S. Toda, S.-I. Itoh, M. Yagi, A Fukuyama and K Itoh,  
**Spatial Structure of Compound Dither in L/H Transition; Mar. 2000**
- NIFS-631 N Ishihara and S. Kida,  
**Axial and Equatorial Magnetic Dipoles Generated in a Rotating Spherical Shell; Mar 2000**
- NIFS-632 T. Kuroda, H Sugama, R. Kanno and M Okamoto,  
**Ion Temperature Gradient Modes in Toroidal Helical Systems, Apr 2000**
- NIFS-633 V.D Pustovitov,  
**Magnetic Diagnostics: General Principles and the Problem of Reconstruction of Plasma Current and Pressure Profiles in Toroidal Systems, Apr 2000**
- NIFS-634 A.B. Mikhailovskii, S.V Kononov, V.D Pustovitov and V.S. Tsypin,  
**Mechanism of Viscosity Effect on Magnetic Island Rotation, Apr 2000**
- NIFS-635 H. Naitou, T Kuramoto, T Kobayashi, M. Yagi, S. Tokuda and T. Matsumoto,  
**Stabilization of Kinetic Internal Kink Mode by Ion Diamagnetic Effects; Apr. 2000**

- NIFS-636 A. Kageyama and S. Kida,  
**A Spectral Method in Spherical Coordinates with Coordinate Singularity at the Origin;**  
 Apr. 2000
- NIFS-637 R. Honuchi, W. Pei and T. Sato,  
**Collisionless Driven Reconnection in an Open System;** June 2000
- NIFS-638 K. Nagaoka, A. Okamoto, S. Yoshimura and M.Y. Tanaka,  
**Plasma Flow Measurement Using Directional Langmuir Probe under Weakly Ion-  
 Magnetized Conditions;** July 2000
- NIFS-639 Alexei Ivanov,  
**Scaling of the Distribution Function and the Critical Exponents near the Point of a  
 Marginal Stability under the Vlasov-Poisson Equations;** Aug. 2000
- NIFS-640 K. Ohi, H. Naitou, Y. Tauchi, O. Fukumasa,  
**Observation of the Limit Cycle in the Asymmetric Plasma Divided by the Magnetic  
 Filter;** Aug. 2000
- NIFS-641 H. Momota, G.H. Miley and J. Nadler,  
**Direct Energy Conversion for IEC Propulsions;** Aug. 2000
- NIFS-642 Y. Kondoh, T. Takahashi and H. Momota,  
**Revisit to the Helicity and the Generalized Self-organization Theory;** Sep 2000
- NIFS-643 H. Soltwisch, K. Tanaka,  
**Changes of the Electron Density Distribution during MHD Activity in CHS;** Sep. 2000
- NIFS-644 Fujisawa, A., Iguchi, H., Minami, T., Yoshimura, Y., Sanuki, H., Itoh, K., Isobe, M., Nishimura, S., Tanaka, K.,  
 Osakabe, M., Nomura, I., Ida, K., Okamura, S., Toi, K., Kado, S., Akiyama, R., Shimizu, A., Takahashi, C.,  
 Kojima, M., Matsuoka, K., Hamada, Y., Fujiwara, M.,  
**Observation of Bifurcation Property of Radial Electric Field Using a Heavy Ion Beam  
 Probe;** Sep. 2000  
 (IAEA-CN-77/EX6/6)
- NIFS-645 Todo, Y., Watanabe, T.-H., Park, H.-B., Sato, T.,  
**Fokker-Planck Simulation Study of Alfvén Eigenmode Burst** Sep. 2000  
 (IAEA-CN-77/TH6/2)
- NIFS-646 Muroga, T., Nagasaka, T.,  
**Development of Manufacturing Technology for High Purity Low Activation Vanadium  
 Alloys;** Sep 2000  
 (IAEA-CN-77/FTP1/09)
- NIFS-647 Itoh, K., Sanuki, H., Toda, S., Itoh, S.-I., Yagi, M., Fukuyama, A., Yokoyama, M.,  
**Theory of Dynamics in Long Pulse Helical Plasmas;** Sep. 2000  
 (IAEA-CN-77/THP1/19)
- NIFS-648 R. Sakamoto, H. Yamada, K. Tanaka, K. Narihara, S. Morita, S. Sakakibara, S. Masuzaki, L.R. Baylor, P.W. Fisher, S.K. Combs,  
 M.J. Gouge, S. Kato, A. Komori, O. Kaneko, N. Ashikawa, P. de Varies, M. Emoto, H. Funaba, M. Goto, K. Ida, H. Idei, K. Ikeda, S.  
 Inagaki, M. Isobe, S. Kado, K. Kawahata, K. Khlopenkov, S. Kubo, R. Kumazawa, T. Minami, J. Miyazawa, T. Morisaki, S.  
 Murakami, S. Muto, T. Mutoh, Y. Nagayama, Y. Nakamura, H. Nakanishi, K. Nishimura, N. Noda, T. Notake, T. Kobuchi, Y. Liang,  
 S. Ohdachi, N. Ohyabu, Y. Oka, M. Osakabe, T. Ozaki, R.O. Pavlichenko, B.J. Peterson, A. Sagara, K. Saito, H. Sasao, M. Sasao,  
 K. Sato, M. Sato, T. Seki, T. Shimozuma, M. Shoji, S. Sudo, H. Suzuki, M. Takechi, Y. Takein, N. Tamura, K. Toi, T. Tokuzawa, Y.  
 Toni, K. Tsumori, I. Yamada, S. Yamaguchi, S. Yamamoto, Y. Yoshimura, K.Y. Watanabe, T. Watari, and O. Motojima,  
**Impact of Pellet Injection on Extension of Operational Region in LHD;** Sep. 2000  
 (IAEA-CN-77/EXP4/18/)
- NIFS-649 K. Ichiguchi, M. Wakatani, T. Umemura, T. Tatsuno and B.A. Carreras  
**Improved Stability due to Local Pressure Flattening in Stellarators;** Sep. 2000  
 (IAEA-CN-77/THP2/08)