

Journal of the Arkansas Academy of Science

Volume 64

Article 16

2010

Seasonal Activity of the Ozark Highlands Leech, *Macrobdella diplotertia*, (Annelida: Hirudinea) in North-central Arkansas

M. Connior

South Arkansas Community College, mconnior@southark.edu

Stanley E. Trauth

Arkansas State University

Follow this and additional works at: <http://scholarworks.uark.edu/jaas>

 Part of the [Zoology Commons](#)

Recommended Citation

Connior, M. and Trauth, Stanley E. (2010) "Seasonal Activity of the Ozark Highlands Leech, *Macrobdella diplotertia*, (Annelida: Hirudinea) in North-central Arkansas," *Journal of the Arkansas Academy of Science*: Vol. 64 , Article 16.
Available at: <http://scholarworks.uark.edu/jaas/vol64/iss1/16>

This article is available for use under the Creative Commons license: Attribution-NoDerivatives 4.0 International (CC BY-ND 4.0). Users are able to read, download, copy, print, distribute, search, link to the full texts of these articles, or use them for any other lawful purpose, without asking prior permission from the publisher or the author.

This Article is brought to you for free and open access by ScholarWorks@UARK. It has been accepted for inclusion in Journal of the Arkansas Academy of Science by an authorized editor of ScholarWorks@UARK. For more information, please contact scholar@uark.edu.

Seasonal Activity of the Ozark Highlands Leech, *Macrobdella diplotertia*, (Annelida: Hirudinea) in North-central Arkansas

M.B. Connior¹ and S.E. Trauth²

¹Department of Biology, South Arkansas Community College, 300 S. West Avenue, El Dorado, AR 71730

²Department of Biological Sciences, Arkansas State University, PO Box 599, State University, AR 72467

¹Correspondence: mconnior@southark.edu

Abstract

The Ozark Highlands Leech, *Macrobdella diplotertia*, occurs intermittently throughout Arkansas, Kansas, and Missouri. Limited natural history of this species is known. Herein, we report a new county occurrence for this leech from a cattle pond in southern Marion County, Arkansas. We also report on the seasonal activity and novel hosts of this species. We surveyed the inhabited pond monthly to determine activity and collected natural history data on this species. This study indicated that this species appears to be most active in spring, summer, and early fall, but inactive during the winter. We also documented five new anuran hosts for this species. The Central Newt, *Notophthalmus viridescens*, was also present in this pond, which further supports a mimetic relationship previously proposed between these two animals.

Introduction

There are currently 77 described species of leeches occurring in North American north of Mexico (Klemm et al. 2009). Of these 77 species, 22 are known to occur in Arkansas (Moser et al. 2006). The distribution and natural histories of many of these species have yet to be determined completely (Klemm 1985). Herein, we report on the natural history of one of these species, the Ozark Highlands Leech (*Macrobdella diplotertia*). *Macrobdella* is a monophyletic genus with four species in North America (Phillips and Siddall 2005). Two species of *Macrobdella* occur in Arkansas, *M. diplotertia* in the north and *M. ditetra* in the south (Moser et al. 2006). *M. diplotertia* has been previously reported from two counties in Missouri (Meyer 1975, Trauth and Neal 2004), three counties in Kansas (Klemm et al. 1979) and two counties in Arkansas (Turbeville and Briggler 2003). No additional specimens, however, were discovered during a leech survey of northern Arkansas (Moser et al. 2006).

Macrobdella diplotertia is an omnivore known to be an amphibian egg predator and a sanguivore (Turbeville and Briggler 2003, Trauth and Neal 2004). It is also known to have a mimetic relationship with the Central Newt (McCallum et al. 2008). Interestingly, the leech *Placobdella picta* has been observed attached to the Ozark Highlands leech (Turbeville and Briggler 2003). Currently, no reports of these leeches feeding on adult amphibians have been published.

Methods

Leeches were observed and collected in a manmade cattle pond in southern Marion County, Arkansas (UTM 15N 0535878E, 3992944N). In order to obtain data on seasonal activity of the leeches, the pond was surveyed monthly during daylight hours for a year from June 2009 to May 2010. One of the authors (MBC) slowly walked around a section of the pond perimeter (~25 m) at a water depth of ~30 cm for 10 min each month and collected all observed leeches with a ~6 mm mesh dipnet. To supplement the survey, 5 single dipnet collections of the benthic substrate were collected every month and sifted through to collect any leeches in this material. We also opportunistically captured amphibians and examined them for leech parasitism. We also sampled the pond for the presence of Central Newts to provide further support of the proposed mimetic relationship between these two species (McCallum et al. 2008). All collected leeches were deposited in the Invertebrate Zoology collections of the National Museum of Natural History, Smithsonian Institution, Washington, DC, for further studies.

Results

Marion County (Arkansas) represents a new county occurrence for *M. diplotertia* (Fig. 1). We recorded these leeches to be most active in spring, summer, and early fall, but inactive during the winter




Figure 1: Distribution of the Ozark Highlands Leech (*Macrobdeella diploptertia*) in Kansas, Missouri, and Arkansas. Previous records (dots); new record (star).

(Fig. 2). During the warmer months, leeches were collected while freely swimming in the pond. The leeches were documented to feed on the eggs of gray treefrogs (*Hyla versicolor*) *in situ*. We also observed these leeches to feed from five novel adult anuran host species: Spring Peeper (*Pseudacris crucifer*); Gray Treefrog; Pickerel Frog (*Lithobates palustris*); Green Frog (*L. clamitans*); and American Bullfrog (*L. catesbeianus*). An adult leech was attached to the gular region of the Spring Peeper. An adult leech was attached adjacent to the orbit of the eye on the treefrog. A juvenile was attached to the inner thigh of the bullfrog. While in captivity directly after collection, we observed an adult leech attach to the dorsal cephalic region of a Spotted Salamander (*Ambystoma maculatum*) larva and feed there. Additionally, while in captivity we observed two different individuals attach to the web of the hind foot of a Pickerel frog and Green frog. We also observed on several occasions, we observed the leech *Placobdeella picta* attached to *Macrobdeella diploptertia*. The Central Newt was very common within the pond providing further evidence for potential interactions between these two species.

Discussion

The discovery of this population of *Macrobdeella diploptertia* in Marion County expands the distribution of this species eastward in Arkansas. Other populations may occur elsewhere within the Ozark Highlands region in the state. Amphibians that either live or reproduce within ponds that contain populations of these leeches may sustain substantial parasitism; however, the extent of this parasitism is currently

unknown. The observations of both amphibian adult and egg predation suggest that *M. diploptertia* affect the fitness of adult amphibians as well as survival of developing larvae.




Figure 2: Seasonal activity of the Ozark Highlands Leech (*Macrobdeella diploptertia*) in a manmade cattle pond in Marion County, Arkansas.

The genus *Macrobdeella* typically feed on amphibian adults and their larvae. *M. decora* has been reported feeding on brook trout, *Salvelinus fontinalis* (Rupp and Meyer 1954) and amphibian eggs, including *Ambystoma maculatum* (Cargo 1960), *Bufo terrestris* (Travis and Trexler 1986), *Lithobates catesbeianus* (Howard 1978), and *L. sylvatica* (Cory and Manion 1953). *M. ditetra* has been reported feeding on sirens, *Siren intermedia*, (Graham and Borda 2010) and southern leopard frogs, *L. sphenoccephala* (Beckerdite and Corkum 1973). An unusual case of hirudiniasis was reported with *M. ditetra* feeding on cattle, *Bos taurus* (Meyer 1959). Moore (1953) reported that *M. ditetra* readily consumed the eggs of *L. catesbeianus*. Because *M. sestertia* is quite rare, host preferences have not been determined, although Smith (1977) stated that beach patrons reported that these leeches commonly fed on humans. By combining our results with those of Trauth and Neal (2004) and Turbeville and Briggler (2003), *M. diploptertia* is now known to feed on adult

Seasonal Activity of the Ozark Highlands Leech, *Macrobdella diploptertia* in North-central Arkansas

amphibians, including Spring Peepers, Gray Treefrogs, Pickerel Frogs, Green Frogs, and American Bullfrogs, as well as the eggs of five amphibian species (*A. maculatum*, *H. versicolor*, *L. clamitans*, *L. sphenoccephala*, and *L. sylvatica*). Our results have augmented the known food sources for *M. diploptertia*, although further studies are required to assess the impact this species has on amphibian fitness and survival.

Acknowledgments

We wish to thank J. and R. Connior for access to the pond and I. Guenther for assistance with fieldwork. W. Moser provided comments on an earlier version of this manuscript as well as curatorial assistance.

Literature Cited

- Beckerdite FW** and **KC Corkum**. 1973. Observations on the life history of the leech *Macrobdella ditetra* (Hirudinea: Hirudinidae). Proceedings of the Louisiana Academy of Sciences 36:61-3.
- Cargo DG**. 1960. Predation of eggs of the spotted salamander *Ambystoma maculatum* by the leech *Macrobdella decora*. Chesapeake Science 1:119-20.
- Cory L** and **JL Manion**. 1953. Predation on eggs of the woodfrog, *Rana sylvatica*, by leeches. Copeia 1953:66.
- Graham SP** and **E Borda**. 2010. First report of leech parasitism in the amphibian family Sirenidae. Comparative Parasitology 77:105-7.
- Howard RD**. 1978. The influence of male-defended oviposition sites on early embryo mortality in Bullfrogs. Ecology 59:789-98.
- Klemm DJ**. 1985. Freshwater leeches (Annelida: Hirudinea). In: Klemm DJ, ed. A Guide to the Freshwater Annelida (Polychaeta, Naidid and Tubificid Oligochaeta, and Hirudinea) of North America. Dubuque: Kendall/Hunt Publishing Co. 198 p.
- Klemm DJ, DG Huggins** and **MJ Wetzel**. 1979. Kansas leeches (Annelida: Hirudinea) with notes on distribution and ecology. Technical Publications of the State Biological Survey of Kansas. 8:38-46.
- Klemm DJ, WE Moser** and **MJ Wetzel**. 2009. Classification and checklist of the leeches (Phylum Annelida: Class Clitellata: Subclass Hirudinida) occurring in North America north of Mexico. <<http://www.inhs.uiuc.edu/~mjwetzel/FWLeechesNA.html>>. 14 April 2009.
- McCallum ML, S Beharry** and **SE Trauth**. 2008. A complex mimetic relationship between the central newt and the Ozark Highlands leech. Southeastern Naturalist 7:173-9.
- Meyer MC**. 1959. Another unusual case of erratic hirudiniasis. Journal of Parasitology 45:39.
- Meyer MC**. 1975. A new leech, *Macrobdella diploptertia* sp. n. (Hirudinea: Hirudinidae), from Missouri. Proceedings of the Helminthological Society of Washington 42:82-5.
- Moore JP**. 1953. Three undescribed North American leeches (Hirudinea). Notulae Naturae of the Academy of Natural Sciences of Philadelphia 250:1-13.
- Moser WE, DJ Klemm, DJ Richardson, BA Wheeler, SE Trauth** and **BA Daniels**. 2006. Leeches (Annelida: Hirudinida) of northern Arkansas. Journal of the Arkansas Academy of Science 60:84-95.
- Phillips AJ** and **ME Siddall**. 2005. Phylogeny of the New World medicinal leech family Macrobdellidae (Oligochaeta: Hirudinida: Arynchobdellida). Zoological Scripta 34:559-64.
- Rupp RS** and **MC Meyer**. 1954. Mortality among brook trout, *Salvelinus fontinalis*, resulting from attacks of freshwater leeches. Copeia 1954:294-5.
- Smith DG**. 1977. The rediscovery of *Macrobdella sestertia* Whitman (Hirudinea: Hirudinidae). The Journal of Parasitology 63:759-60.
- Trauth SE** and **RG Neal**. 2004. Feeding response by the leech *Macrobdella diploptertia* (Annelida: Hirudinea) to Wood Frog and Spotted Salamander egg masses. Journal of the Arkansas Academy of Science 58:139-41.
- Travis J** and **JC Trexler**. 1986. Interactions among factors affecting growth, development and survival in experimental populations of *Bufo terrestris* (Anura:Bufonidae). Oecologia 69:110-6.
- Turbeville JM** and **JT Briggler**. 2003. The occurrence of *Macrobdella diploptertia* (Annelida: Hirudinea) in the Ozarks Highlands of Arkansas and preliminary observations on its feeding habits. Journal of Freshwater Ecology 18:155-9.