

Santa Clara University Scholar Commons

Biology

College of Arts & Sciences

10-1992

Two mechanisms for growth inhibition by elevated transport of sugar phosphates in *Escherichia coli*

Robert J. Kadner

G. Patrick Murphy

Craig Stephens

Santa Clara University, cstephens@scu.edu

Follow this and additional works at: <http://scholarcommons.scu.edu/bio>

 Part of the [Biology Commons](#)

Recommended Citation

Kadner R. J., Murphy G. P., Stephens C. M. (1992). Two mechanisms for growth inhibition by elevated transport of sugar phosphates in *Escherichia coli*. *J. Gen. Microbiol.* 138, 2007-2014 10.1099/00221287-138-10-2007

This Article is brought to you for free and open access by the College of Arts & Sciences at Scholar Commons. It has been accepted for inclusion in Biology by an authorized administrator of Scholar Commons. For more information, please contact rscroggin@scu.edu.

Two mechanisms for growth inhibition by elevated transport of sugar phosphates in *Escherichia coli*

ROBERT J. KADNER,* G. PATRICK MURPHY and CRAIG M. STEPHENS†

Department of Microbiology, School of Medicine, University of Virginia, Charlottesville, VA 22908, USA

(Received 24 February 1992; revised 24 June 1992; accepted 2 July 1992)

The *Escherichia coli* *uhpT* gene encodes an active transport system for sugar phosphates. When the *uhpT* gene was carried on a multicopy plasmid, amplified levels of transport activity occurred, and growth of these strains was inhibited upon the addition of various sugar phosphates. Two different mechanisms for this growth inhibition were distinguished. Exposure to glucose-6-phosphate, fructose-6-phosphate or mannose-6-phosphate, which enter directly into the glycolytic pathway, resulted in cessation of growth and substantial loss of viability. Cell killing was correlated with the production of the toxic metabolite, methylglyoxal. In contrast, addition of 2-deoxyglucose-6-phosphate, galactose-6-phosphate, glucosamine-6-phosphate or arabinose-5-phosphate, which do not directly enter the glycolytic pathway, resulted in growth inhibition without engendering methylglyoxal production or cell death. Inhibition of growth could result from excessive accumulation of organophosphates in the cell or depletion of inorganic phosphate pools as a result of the sugar-P/P_i exchange process catalysed by UhpT. The phosphate-dependent uptake of glycerol-3-phosphate by the GlpT antiporter was strongly inhibited under conditions of elevated sugar-phosphate transport. There are thus two separate toxic effects of elevated sugar-phosphate transport, one of which was lethal and related to increased flux through glycolysis. It is likely that the control of *uhpT* transcription by catabolite repression exists to limit the level of UhpT transport activity and thereby prevent the toxic events that result from elevated uptake of its substrates.

Introduction

The *uhpT* gene of *Escherichia coli* encodes an active transport system for sugar phosphates (Weston & Kadner, 1988). UhpT acts as an electroneutral phosphate antiporter to mediate the entry of a variety of organophosphates, including the phosphate esters of many sugars and sugar alcohols ranging in size from three to seven carbons, and glucosamine-6-phosphate (reviewed in Dietz, 1976; Maloney *et al.*, 1990). The driving force for active transport of sugar phosphates is the outward movement of P_i down the transmembrane P_i gradient, which is normally maintained by the proton-motive-force-dependent Pit transport system (Sonna *et al.*, 1988). Production of the Uhp transporter is induced specifically by extracellular glucose-6-phosphate (Glu6P) or 2-deoxyglucose-6-phosphate (2dGlu6P) (Dietz & Heppel, 1971) in a process mediated by the products of three regulatory genes, *uhpABC* (Weston & Kadner, 1987). UhpA, which bears sequence relatedness

to many transcriptional activators, including OmpR and NarL, appears to activate transcription of *uhpT*. Expression of UhpA from multicopy plasmids subverts the normal regulatory control by Glu6P and UhpBC function, and results in elevated and constitutive production of UhpT (Shattuck-Eidens & Kadner, 1983).

Expression of UhpT is subject to catabolite repression and is reduced 2- to 3-fold when cells are grown in the presence of both Glu6P and glucose. This decrease is reversed by addition of cyclic AMP, and it has been found that the *uhpT* promoter contains a typical binding site for the cyclic AMP receptor protein (CAP) (Merkel *et al.*, 1992). Deletion of the CAP-binding site in the *uhpT* promoter or of the *crp* gene resulted in an 8- to 10-fold decrease in *uhpT* transcription and loss of the response to glucose. The susceptibility of *uhpT* expression to catabolite repression was surprising. Glu6P itself is a potent catabolite repressor of unknown mechanism for numerous other systems (McGinnis & Paigen, 1969) and it seems unlikely that its utilization should be repressed in preference for some other carbon source. A likely reason for the catabolite repression of *uhpT* expression is to limit the amount of transport activity to

* Author for correspondence. Tel. (804) 924 2532; fax (804) 982 1071.

† Current address: Department of Developmental Biology, Stanford University School of Medicine, Stanford, CA 94305, USA.

prevent overloading the metabolic pathways with substrate. Ackerman *et al.* (1974) showed that the presence of Glu6P was lethal for wild-type *E. coli* when cyclic AMP was added to overcome catabolite repression. They showed that the toxicity was associated with elevated levels of methylglyoxal, as observed during deregulated metabolism of other carbon sources.

It was of interest to see whether similar toxicity could result from elevation of Uhp transport activity without interfering with catabolite repression. During studies of the genetic basis of *uhpT* regulation, we observed that strains carrying *uhpT*⁺ plasmids were sensitive to the presence of Glu6P in the medium. On solid media containing Glu6P, these strains grew poorly in the densely inoculated portion of streak plates but well as isolated colonies. Growth inhibition was even more dramatic in liquid media. This paper describes the deleterious effects of addition of sugar phosphates to strains overexpressing *uhpT*, and examines the genetic and environmental parameters influencing susceptibility to inhibition by sugar phosphates. Two distinctive effects of sugar phosphates are described: namely, lethal production of the toxic metabolite methylglyoxal due to overloading of the glycolytic pathway; and growth stasis elicited by sugar phosphates that are not directly metabolized through glycolysis. These results demonstrate the physiological importance of maintaining a limiting rate of transport during sugar phosphate catabolism and provide a clear rationale for the negative regulation of *uhpT* expression by catabolite repression.

Methods

Bacterial strains. The *Escherichia coli* K12 strains were derived from strain RK4353 [$\Delta(\argF-lac)U169\ araD139\ relA1\ rpsL150\ thi\ gyrA219\ non$]; strain RK5000 also carried the $\Delta(ilvB-uhpABCT)2056$ deletion (Kadner & Shattuck-Eidens, 1983). All host strains were *recA56*, unless otherwise specified. Plasmids were derived from pBR322. Plasmid pDSE1 carries a 19 kb insert from the Clarke-Carbon plasmid pLC17-47, which includes pColE1 and the entire *uhp* region; pDSE6 carries the 9.5 kb insert from plasmid pLC40-33 and encodes only *uhpC* and *uhpT* (Shattuck-Eidens & Kadner, 1983). Plasmid pRJK10 carries a 6.5 kb *HindIII-BamHI* fragment with a portion of the *ilvBN* operon and the entire *uhp* region (Weston & Kadner, 1988). Four derivatives of pRJK10 were made by insertion of a 1.5 kb kanamycin-resistance cassette into a restriction site in each of the four *uhp* genes.

Growth conditions. Cells were grown in MOPS medium (Neidhardt *et al.*, 1974) or medium A (Davis & Mingioli, 1950), with 0.5% glycerol as carbon source, thiamine (1 $\mu\text{g ml}^{-1}$), and required amino acids (100 $\mu\text{g ml}^{-1}$). The nominal P, contents of MOPS medium and A medium are 1.3 mM and 93.3 mM, respectively. When specified, casein hydrolysate (Difco) was added to 0.5%. The salts media of different pH contained the same total concentration of phosphate and other ingredients as in medium A, but were adjusted to the desired initial pH with KOH. Ampicillin was added to 25 $\mu\text{g ml}^{-1}$ to maintain plasmid presence. Cells were grown at 37 °C with rotary shaking. Cell growth was

measured as OD₄₂₀. Viable cell counts were measured by dilution and plating on LB agar plates.

Measurement of methylglyoxal production. Production of a toxic factor by cells was detected by suspending cells in water at the time of addition of 300 μM -Glu6P. After 60 min at 37 °C, the cells were removed by centrifugation and the medium was passed through a 0.22 μm filter and then concentrated by lyophilization. Serial dilutions of this material were pipetted onto a lawn of RK4353 on minimal agar plates, and zones of killing were observed after overnight incubation. Portions of the lethal extracts were incubated for 60 min with glutathione or glyoxylase I (Sigma) or both, and were then likewise pipetted onto RK4353 plates. The amount of S-lactyl-glutathione produced in the glyoxylase reaction was measured colorimetrically (Gawehn, 1983).

Methylglyoxal production and release from cells was measured by reaction of the growth medium with 2,4-dinitrophenylhydrazine (Cooper, 1975). Cells were incubated for various periods of time with sugar phosphates and the cells were removed by centrifugation. 2,4-Dinitrophenylhydrazine (665 μl of 0.25 mg ml⁻¹ in 2 M-HCl) was added to a 500 μl sample containing various volumes of the growth medium (2.5–500 μl). Following incubation for 15 min at 30 °C, 835 μl 10% (w/v) NaOH was added and *A*₅₅₀ determined. An absorbance of 16.4 corresponds to 1 μmol methylglyoxal (Cooper, 1975). The chromatographic mobility of the 2,4-dinitrophenylhydrazone formed was compared to that derived from commercial methylglyoxal by ascending chromatography on dimethylformamide-soaked Whatman #1 paper with a solvent system of n-butyl ether/ethanol/water (50:5:50).

Assay of transport activity. Transport of [¹⁴C]-labelled glycerol-3-phosphate or glucose-6-phosphate was measured as previously described (Shattuck-Eidens & Kadner, 1981). Uptake is expressed in units of nmol substrate accumulated per μl of cell water.

Results

Effect of glucose-6-phosphate on cell growth and viability

E. coli strains carrying plasmids pDSE1 or pRJK10, which contain the entire *uhpABCT* gene cluster with various extents of flanking sequence, exhibit high-level constitutive expression of *uhpT*. This amplification was shown by assays of Glu6P transport and of *uhpT*:*lacZ* reporter fusions *in vivo* (Weston & Kadner, 1988), and by determination of the level of Uhp transport activity following *in vitro* reconstitution into proteoliposomes (Sonna *et al.*, 1988). When 3.5 mM-Glu6P was added to cultures of these strains in minimal salts/glycerol medium, growth was strongly inhibited after a short lag and viability began to decline within 30 min (Fig. 1*a*). Cell killing proceeded with a half-time of about 12 min and led to a reduction in viable cell count by at least four orders of magnitude within 3 h. Growth of the same host strain bearing the vector plasmid was unaffected by addition of Glu6P. The same cessation of growth and loss of viability was seen upon addition of 1.75 mM-Glu6P (Fig. 1*b*), whereas 0.7 mM-Glu6P caused an inhibition of growth for 30–60 min followed by resumption at normal rate.




Fig. 1. Growth of *E. coli* strains carrying *uhp* plasmids and exposed to sugar phosphates. Culture OD_{420} was measured at the indicated times after addition of the specified compounds at zero time. Experiments (a–c) used medium A; (d–g) used MOPS medium with 1.3 mM- P_i , unless otherwise specified. (a) Effect of Glu6P on growth and viability. Strain 4353 carrying pDSE1; turbidity after \circ , no addition or \bullet , 3.5 mM-Glu6P addition; \blacksquare , colony-forming units after 3.5 mM-Glu6P addition. (b) Effect of different concentrations of Glu6P. Strain RK4353 carrying pDSE1: \circ , control; \bullet , Glu6P at 0.7 mM; \triangle , 1.75 mM; or \blacktriangledown , 3.5 mM. (c) Effect of Glu6P on an inducible strain. Strain RK4353 carrying pDSE6; symbols as (b). (d) Effect of different sugar phosphates. Strain RK5000 carrying pRJK10; \circ , control; \bullet , Glu6P or Gal6P; ∇ , Fru6P; \blacktriangledown , glucosamine-6P; \square , 2-deoxyGlu6P; \blacksquare , ribose-5P; all sugar phosphates at 2.5 mM. (e) Effect of Gal6P on strains carrying transposon insertions in the *uhp* plasmid. Strain RK5000 carrying pRJK10 with no addition (\circ) or with 2.5 mM-Gal6P (\bullet); or carrying a kanamycin resistance cassette in *uhpA* (∇), *uhpB* (\blacktriangledown), *uhpC* (\square) or *uhpT* (\blacksquare) and exposed to Gal6P. (f) Effect of P_i growth medium concentration on response to 2.5 mM-Gal6P. Strain RK5000 carrying pRJK10 in MOPS medium; \circ , control; plus Gal6P in P_i at \bullet , 0.26 mM; ∇ , 1.3 mM; \blacktriangledown , 6.5 mM; or \square , 13 mM. (g) Effect of growth medium on response to 2.5 mM-Gal6P. Strain RK5000 carrying pRJK10 in MOPS medium; \circ , control; \bullet , 2.5 mM-Glu6P in 1.3 mM- P_i ; ∇ , 2.5 mM-Gal6P in 1.3 mM- P_i ; \square , 2.5 mM-Gal6P in 2.6 mM- P_i ; \blacksquare , 2.5 mM-Gal6P in 7.8 mM- P_i ; \triangle , 2.5 mM-Gal6P in 1.3 mM- P_i , grown with 0.25% glucose; \blacktriangledown , 2.5 mM-Gal6P in 1.3 mM- P_i , grown with 0.25% galactose.

A different response to Glu6P was seen in cells carrying plasmid pDSE6, which contains *uhpT* but not *uhpA*. Expression of the plasmid-borne *uhpT* gene is dependent on the chromosomal copy of *uhpA* and is inducible by Glu6P (Shattuck-Eidens & Kadner, 1983). The addition of 1.75 or 3.5 mM-Glu6P resulted in complete inhibition of growth (Fig. 1c) and the same degree of cell killing as in RK4353(pDSE1) (data not shown). However, there was a lag of 60–80 min before the onset of killing and a longer lag before the cessation of cell growth. The additional lag presumably represents the time required for induction of transporter expression to the level at which inhibitory levels of Glu6P accumulated in the cell. This view was supported by the effect of the inhibition of protein synthesis by chloram-

phenicol (data not shown). Chloramphenicol (100 $\mu\text{g ml}^{-1}$) caused a rapid cessation of growth in strains carrying either pDSE1 or pDSE6, but no loss of viability for at least 4 h. Simultaneous addition of chloramphenicol and Glu6P (3.5 mM) to constitutively expressing strains carrying pRJK10 or pDSE1 resulted in cell killing which was identical in extent and kinetics to that seen upon addition of Glu6P alone. In contrast, the inducible strain carrying pDSE6 was completely protected from killing by Glu6P when formation of the transport system was prevented by chloramphenicol. These results indicated that the observed lethality occurs in response to amplified transport of Glu6P, rather than to the amplified production of the transport protein itself.

Table 1. *Effect of sugar phosphates on growth, viability, and methylglyoxal production*

Cells of strain RK5000(pRJK10) were grown in MOPS minimal medium with glycerol as carbon source and 1.32 mM-P_i. Sugars and sugar phosphates were added to 2.5 mM, and after 120 min, the relative change of culture turbidity or of colony-forming units was determined. Methylglyoxal was determined at this time as described in Methods.

Addition	Culture turbidity (relative change)	Viable counts (relative change)	Methylglyoxal (mM)
Control	2.36	2.3	0.06
Glucose-6P	1.21	0.06	3.45
Fructose-6P	1.33	0.06	3.31
Mannose-6P	1.89	0.27	0.67
Glucosamine-6P	1.08	1.97	0.13
2-Deoxyglucose-6P	< 1.05	1.65	0.06
Ribose-5P	3.18	2.71	0.09
Galactose-6P	< 1.05	< 1.15	0.07
Arabinose-5P	< 1.05	< 1.05	0.06
Glucose	3.78	ND	0.06

ND, not determined.

Response to other sugar phosphates

The effect of other sugar phosphates on growth and viability was examined (Fig. 1*d*; Table 1). In RK5000(pRJK10), 2.5 mM-fructose-6-phosphate (Fru6P) killed cells at the same rate and to the same extent as did Glu6P. Cells exposed to 2.5 mM-mannose-6-phosphate showed a similar increase in culture turbidity to the control culture, but there was a decrease in the number of colony-forming units that showed more daily variability and a lesser degree of killing than was seen with Glu6P or Fru6P. In contrast, 2-deoxyglucose-6-phosphate (2dGlu6P) and galactose-6-phosphate (Gal6P) at 2.5 mM inhibited growth but did not cause cell death. Addition of 2.5 mM-glucosamine-6-phosphate stopped growth for about 120 min but with no apparent cell killing, and ribose-5-phosphate stimulated growth. Arabinose-5-phosphate also inhibited growth without cell killing. If the cells had been previously grown in the presence of arabinose, the growth inhibition by arabinose-5-phosphate was reduced or eliminated (data not shown). These results suggest that the effect on cell viability of the elevated transport of a sugar phosphate depends on the metabolic fate of that compound, such that lethality occurs only if the sugar phosphate enters the glycolytic pathway directly.

In the inducible strain RK4353(pDSE6), 3.5 mM Fru6P, which is a poor inducer of *uhpT* expression, did not inhibit growth over the time period of this experiment (data not shown). 2-Deoxyglucose-6-phosphate is an effective inducer and caused a moderate growth inhibition but no killing. Arabinose-5-phosphate, which is not an inducer of the *uhp* system, had no effect on this strain.

To determine the genetic requirements for this response to sugar phosphates, the effect of 2.5 mM-Gal6P

on *uhpABCT* deletion strain RK5000 containing pRJK10 derivatives with mini-Tn10 insertion mutations in each of the *uhp* genes was examined. Gal6P caused growth inhibition in the *uhpB* and *uhpC* mutants, but not in strains with *uhpA* or *uhpT* insertions (Fig. 1*e*). Identical behaviour was seen in response to Glu6P, except that cell killing occurred along with growth inhibition (data not shown). The requirement for UhpA is consistent with previous observations that it is necessary for *uhpT* expression, as transport activity is virtually undetectable in the absence of *uhpA* even when *uhpT* is in multicopy (Weston & Kadner, 1988). When *uhpA* is present on a multicopy plasmid, *uhpT* expression in the absence of inducer is greatly increased and the *uhpB* and *uhpC* products are no longer necessary (Weston & Kadner, 1988), consistent with the observation that Glu6P was lethal to *uhpB* and *uhpC* insertion mutants. Amplified expression of UhpT transport activity is clearly necessary for the toxic effect of sugar phosphates.

Effect of medium conditions

The effect of the composition of the growth medium on the response to Glu6P was examined. Considerable protection was provided by 0.5% casein hydrolysate at low Glu6P levels (data not shown). In its presence, 0.7 mM-Glu6P did not inhibit growth to any obvious degree, and 1.75 mM-Glu6P elicited only a 90 min lag, in contrast to the extensive killing in the absence of casein hydrolysate. The addition of 3.5 mM-Glu6P resulted in a similar degree of killing irrespective of the presence of casein hydrolysate. The protective effect could result from a reaction of amino acids with a toxic agent generated upon metabolism of sugar phosphates.

A possible mechanism for the growth inhibition could be the acidification of the cytoplasm as a result of




Fig. 2. Level of methylglyoxal in culture medium of strains exposed to sugar phosphates. The cell-free culture medium was obtained at the indicated times after addition of sugar phosphates and methylglyoxal measured by the 2,4-dinitrophenylhydrazine assay ($1 \mu\text{mol} = 16.4 A_{550}$ units). Strain RK5000 carrying pRJK10 was grown in MOPS medium. (a) Glu6P was added at \circ , 0 mM; \bullet , 0.9 mM; ∇ , 1.8 mM; and \blacktriangledown , 3.5 mM. These concentrations of Glu6P resulted in killing after 120 min of 82%, 97.3%, and 97.5% of the cells. (b) Cells were exposed to sugar phosphates at 2.5 mM: \circ , control; \bullet , Glu6P; ∇ , Fru6P; \square , mannose-6P; or \blacktriangledown , any of glucosamine-6P, 2-deoxyglucose-6P, Gal6P, or ribose-5P.

elevated proton entry coupled directly or indirectly to uptake of sugar phosphates. However, the initial pH of the medium had only a slight effect of the susceptibility of strains to growth inhibition. The lag in growth that resulted after the addition of 0.7 mM-Glu6P was about 40 min longer at pH 5.5 than at pH 7.5, but growth was completely inhibited by 1.75 mM-Glu6P at both pHs (data not shown). The concentration of P_i in MOPS medium had a significant effect on susceptibility to sugar phosphates. Increasing $[P_i]_{\text{out}}$ from 1.3 mM to 6.8 mM reduced the inhibitory effect of 2.5 mM-Gal6P (Fig. 1f) and 2dGlu6P, but did not prevent the growth inhibition. Further increases to 13 mM or 65 mM- P_i did not provide any additional protection. This result could indicate that external P_i blocks entry of the growth-inhibitory sugar phosphates, or that higher external P_i levels could partially overcome the depletion of internal P_i pools by the action of the UhpT antiporter.

The presence of glucose in the medium blocked killing by Glu6P (data not shown) and reduced the inhibition caused by Gal6P (Fig. 1g), most likely as a result of catabolite repression of *uhpT* expression. Pregrowth of the cells with galactose did not relieve the inhibition caused by Gal6P.

Formation of methylglyoxal

Exposure of *uhp*-plasmid-bearing cells to Glu6P under conditions leading to lethality resulted in the production of a toxic factor which could be assayed by the killing of a lawn of sensitive cells. The toxic factor was inactivated by incubation with glyoxylase I plus glutathione but not with either alone. This result suggested that the toxic factor was a ketoaldehyde, most likely methylglyoxal (MG). In support of this proposal, cells exposed to Glu6P released material reactive with 2,4-dinitrophenylhydrazine (DNPH). The derivative formed with the growth medium of Glu6P-treated cells was identical in absorp-

tion spectrum and chromatographic mobility with the corresponding derivative formed with commercial MG (data not shown). MG has been shown to accumulate in cells in which the metabolic flux through the glycolytic pathway has been increased through some mechanism to subvert the usual regulation of metabolism (reviewed in Cooper, 1984).

To demonstrate the linkage between high-level sugar phosphate transport, MG production and cell death, MG levels in cultures were measured at various times after addition of Glu6P. MG levels rose rapidly after Glu6P addition, and the amount of MG produced was related to the concentration of Glu6P added (Fig. 2a). Even non-lethal concentrations of Glu6P engendered MG excretion, but the levels in the medium decreased to a basal level, at which time growth resumed. Incubation with 3.5 mM-Fru6P resulted in the production of MG at the same rate and extent as in cells incubated with Glu6P (Fig. 2b; Table 1). In contrast, there was no detectable synthesis of MG upon incubation with sugar phosphates (2dGlu6P, Gal6P, glucosamine-6-phosphate, or ribose-5-phosphate) that inhibited growth but did not cause killing (Table 1). Thus, cell killing following amplified uptake of some sugar phosphates was correlated with the formation of MG.

Function of *GlpT* under conditions of high-level *UhpT* activity

The basis for the growth inhibition resulting from excessive transport of the sugar phosphates that did not engender MG production is less obvious. It was possible that excessive transport of sugar phosphates resulted in depletion of the intracellular P_i pool to a growth-limiting level. To examine this possibility, the transport of $[^{14}\text{C}]$ glycerol-3-phosphate by *GlpT*, a phosphate antiporter related in sequence and mechanism to *UhpT* (Eiglmeier *et al.*, 1987), was examined with or without




Fig. 3. Effect of 2deoxyGlu6P on glycerol-3P uptake and of P_i on Glu6P uptake. Cells of strain RK5000 carrying pRJK10 were grown in MOPS/glycerol medium. (a) To one of duplicate cultures was added 2deoxyGlu6P to 2.5 mM for 88 min before assay of uptake of [^{14}C]glycerol-3P (200 μM ; 0.2 $\mu\text{Ci ml}^{-1}$); \circ , no addition; \bullet , 2dGlu6P. (b) Cells grown in MOPS/glycerol medium with P_i concentrations of \circ , 1.3 mM or \bullet , 26 mM. Uptake of [^{14}C]Glu6P (200 μM ; 0.2 $\mu\text{Ci ml}^{-1}$) was measured.

exposure of cells to Gal6P or 2dGlu6P. Transport of glycerol-3-phosphate was strongly inhibited by addition of 2dGlu6P (Fig. 3a); similar inhibition was seen upon addition of Gal6P (data not shown). Inhibition of transport occurred within 10 min after addition of the hexose phosphate and was relieved when cells resumed growth following prolonged incubation (data not shown).

The partial reduction in growth inhibition that results from increased P_i in the growth medium (Fig. 1f) was also consistent with the view that amplified sugar-phosphate transport might deplete intracellular P_i pools. However, it was possible that high external P_i might competitively inhibit sugar phosphate uptake and thereby relieve the growth inhibition by decreasing the accumulation of sugar phosphate in the cell. To investigate this explanation, the rate of uptake of [^{14}C]Glu6P was measured in media with 1.3 mM and 26 mM- P_i (Fig. 3b). High external P_i had only a slight effect if any on the rate of Glu6P uptake under these conditions. These results are consistent with the proposal that the nonlethal inhibitory effect of amplified sugar phosphate transport is dependent on the accumulation of the internal sugar phosphate, acting by depletion of the internal P_i pool or accumulation of osmotically active, charged metabolites.

Discussion

The results presented here show that elevated transport of sugar phosphates in *E. coli* strains overproducing the UhpT hexose-phosphate transport protein is deleterious to the cell by at least two mechanisms. Some sugar

phosphates caused extensive cell killing, whereas others resulted only in growth inhibition. Lethality was correlated with the ability of the transported sugar phosphate to enter the glycolytic pathway of metabolism and to engender MG synthesis. Glu6P and Fru6P are intermediates in glycolysis and rapidly caused a drastic drop in cell viability. In contrast, 2dGlu6P, which cannot be converted to Fru6P and is not metabolized further, and other sugar phosphates (galactose-, arabinose-, and ribose-phosphates), which are utilized by alternative catabolic pathways, did not lead to MG synthesis and were not lethal.

Previous studies have shown that elevated metabolic fluxes through the glycolytic pathway can engender cell killing by generating excessive levels of methylglyoxal (Freedberg *et al.*, 1971; Ackerman *et al.*, 1974). These abnormally high metabolic rates were achieved in two ways. Growth of wild-type cells in the presence of an inducing sugar, such as xylose or Glu6P, and cyclic AMP overcame catabolite repression and resulted in high level synthesis of the corresponding transport system (Ackerman *et al.*, 1974). High rates of glycerol metabolism required use of mutant strains which both expressed the glycerol-catabolic enzymes at a high, constitutive level, owing to loss of the GlpR repressor, and also possessed an altered glycerol kinase activity that was insensitive to feedback inhibition by fructose-1,6-bis-phosphate (Freedberg *et al.*, 1971). Amplified copy number of the *uhpT* gene provides another mechanism to increase glycolytic flux. MG production can explain some of the curious phenotypic properties of strains carrying Uhp plasmids. Growth on agar plates containing sugar phosphates as sole carbon source was healthy in regions of lower inoculum size but was sparse in the regions of high cell density, and growth in liquid medium in the presence of Glu6P was very poor. The growth of isolated colonies is probably explained by the diffusion of MG away from cells into the agar medium; at high cell density in liquid or solid medium, MG accumulates to toxic concentrations.

MG is formed from dihydroxyacetone phosphate by the enzyme methylglyoxal synthase and is converted to lactate by glyoxylase I in the presence of glutathione (reviewed in Cooper, 1984). A physiological role for MG synthesis has yet to be firmly established, however, MG toxicity in *E. coli* has been extensively documented. Toxicity appears to result primarily from inhibition of protein synthesis, though DNA synthesis may also be affected (Fraval & McBrien, 1980). It appears from this and previous work (Freedberg *et al.*, 1971; Ackerman *et al.*, 1974) that 'proper' regulation of metabolic flow through glycolysis (i.e. that which is sufficient to avoid overproduction of MG) is dependent on the rate of entry of substrates into the pathway being limited at the

transport step, and can be subverted by elevated transport activities. It should also be noted that the potentially low intracellular levels of P_i developed during Glu6P/Fru6P transport as a result of the phosphate antiport activity of UhpT would also favor MG production as MG synthase is inhibited by P_i and the alternate, normal route of glycolytic formation of glyceraldehyde-3-phosphate is P_i -dependent.

The observation that mannose-6-phosphate resulted in lower MG production and less cell death than seen with Glu6P and Fru6P, while glucosamine-6-phosphate resulted only in transient growth inhibition is interesting, considering that both sugars can be enzymically converted into Fru6P, and thus could potentially enter glycolysis. Both compounds also have biosynthetic roles, with mannose-6-phosphate being involved in capsular polysaccharide synthesis and glucosamine-6-phosphate in peptidoglycan and LPS synthesis. It is possible that glucosamine-6-phosphate is shunted preferentially into the synthetic pathway, which is essential for cell growth, rather than the dissimilatory pathway. Capsule synthesis, in contrast, is a dispensable function which may not be expressed under the growth conditions employed here. The delayed effect of mannose-6-phosphate may have been due to limiting levels of phosphomannoisomerase, the enzyme which converts mannose-6-phosphate into Fru6P.

The physiological significance of the regulation of *uhpT* expression by catabolite repression must be to reduce the flux of sugar phosphate transport to limiting levels. Rates of transport must be limited because the feedback controls of glycolytic activity are obviously not sufficient to avoid deleterious consequences of elevated substrate presentation. The detrimental effects resulting from excessive transport cannot be ascribed solely to toxic levels of MG arising from excessive production of dihydroxyacetone phosphate. Substrates that do not elicit MG production still caused very significant growth inhibition. This inhibition could be related to the very familiar growth inhibition caused by the accumulation of sugar phosphates in mutant strains that are blocked in their further metabolism. The basis for this effect is not clear, but might involve a salting-out effect owing to the sequestration of intracellular water by the high levels of organophosphates (> 10 mM).

The fact that increased transport of sugar phosphates strongly inhibited glycerol-3-phosphate transport is consistent with a depletion of the internal P_i pool by the antiport action of UhpT. This model is consistent with the protective effect of increased $[P_i]_{out}$ against the addition of Gal6P or 2dGlu6P. This protective effect of higher $[P_i]_{out}$ was not due simply to competition of P_i with Glu6P for entry on UhpT, since the directly measured rate of Glu6P uptake was not significantly inhibited at

this concentration of external P_i . Assuming that the K_m for P_i for GlpT is similar to that for UhpT (1.2 mM), it is likely that the level of internal P_i could be reduced to a level low enough to affect antiporter activity. Consistent with this view is the puzzling finding that the level of Glu6P uptake in *uhpT*-plasmid-bearing cells is only increased 2–4 fold over that of an induced haploid strain, despite the fact that reconstitution assays showed that the amount of active UhpT protein was increased more than 20-fold (Sonna *et al.*, 1988). This discrepancy may be explained by the suggestion that sugar phosphate transport in whole cells is not limited by the number of transporters, but by some physiological factor, potentially the capacity of the P_i pool. The inhibition of GlpT by UhpT substrates is consistent with this view. However, given the K_m values observed for UhpT, it is unlikely that the level of intracellular P_i could be reduced by the action of UhpT to below 5 μ M, at which growth inhibition occurs. Thus, the mechanism for inhibition of growth by accumulated sugar phosphates remains to be defined. These findings provide evidence for multiple mechanisms of growth inhibition by elevated rates of carbohydrate transport and reinforce the need for careful regulation of transport activity as a primary site of metabolic control.

This work was supported by research grant GM38681 from the National Institute of General Medical Sciences.

References

- ACKERMAN, R. S., COZZARELLI, N. R. & EPSTEIN, W. (1974). Accumulation of toxic concentrations of methylglyoxal by wild-type *Escherichia coli* K-12. *Journal of Bacteriology* **119**, 357–362.
- COOPER, R. A. (1975). Methylglyoxal synthase. *Methods in Enzymology* **41**, 502–508.
- COOPER, R. A. (1984). Metabolism of methylglyoxal in microorganisms. *Annual Review of Microbiology* **38**, 49–68.
- DAVIS, B. D. & MINGIOLI, E. S. (1950). Mutants of *Escherichia coli* requiring methionine or vitamin B₁₂. *Journal of Bacteriology* **60**, 17–28.
- DIETZ, G. W. (1976). The hexose phosphate transport system of *Escherichia coli*. *Advances in Enzymology* **44**, 237–259.
- DIETZ, G. W. & HEPPEL, L. A. (1971). Studies on the uptake of hexose phosphates. II. The induction of the glucose-6-phosphate transport system by exogenous but not by endogenously formed glucose-6-phosphate. *Journal of Biological Chemistry* **246**, 2885–2890.
- EIGLMEIER, K., BOOS, W. & COLE, S. T. (1987). Nucleotide sequence and transcriptional start point of the *glpT* gene of *Escherichia coli*: extensive sequence homology of the glycerol-3-phosphate transport protein with components of the hexose-6-phosphate transport system. *Molecular Microbiology* **1**, 251–258.
- FRAVAL, H. N. A. & MCBRIEN, D. C. H. (1980). The effect of methyl glyoxal on cell division and the synthesis of protein and DNA in synchronous and asynchronous cultures of *Escherichia coli* B/r. *Journal of General Microbiology* **117**, 127–134.
- FREEDBERG, W. B., KISTLER, W. S. & LIN, E. C. C. (1971). Lethal synthesis of methylglyoxal by *Escherichia coli* during unregulated glycerol metabolism. *Journal of Bacteriology* **108**, 137–144.
- GAWEHN, K. (1983). Methylglyoxal. In *Methods of Enzymatic Analysis*, vol. 6, pp. 593–596. Edited by H.-U. Bergmeyer. Weinheim: Verlag Chemie.

- KADNER, R. J. & SHATTUCK-EIDENS, D. M. (1983). Genetic control of the hexose phosphate transport system of *Escherichia coli*: mapping of deletion and insertion mutations in the *uhp* region. *Journal of Bacteriology* **155**, 1052–1061.
- MCGINNIS, J. F. & PAIGEN, K. (1969). Catabolite inhibition: a general phenomenon in the control of carbohydrate utilization. *Journal of Bacteriology* **100**, 902–913.
- MALONEY, P. C., AMBUDKAR, S. V., ANANTHARAM, V., SONNA, L. A. & VARADHACHARY, A. (1990). Anion-exchange mechanisms in bacteria. *Microbiological Reviews* **54**, 1–17.
- MERKEL, T. J., NELSON, D. M., BRAUER, C. L. & KADNER, R. J. (1992). Promoter elements required for positive control of transcription of the *Escherichia coli uhpT* gene. *Journal of Bacteriology* **174**, 2763–2770.
- NEIDHARDT, F. C., BLOCH, P. L. & SMITH, D. F. (1974). Culture medium for enterobacteria. *Journal of Bacteriology* **119**, 736–747.
- SHATTUCK-EIDENS, D. M. & KADNER, R. J. (1981). Exogenous induction of the *Escherichia coli* hexose phosphate transport system defined by *uhp-lac* operon fusions. *Journal of Bacteriology* **148**, 203–209.
- SHATTUCK-EIDENS, D. M. & KADNER, R. J. (1983). Molecular cloning of the *uhp* region and evidence for a positive activator for expression of the hexose phosphate transport system of *Escherichia coli*. *Journal of Bacteriology* **155**, 1060–1070.
- SONNA, L. A., AMBUDKAR, S. V. & MALONEY, P. C. (1988). The mechanism of glucose 6-phosphate transport by *Escherichia coli*. *Journal of Biological Chemistry* **263**, 6625–6630.
- WESTON, L. A. & KADNER, R. J. (1987). Identification of Uhp polypeptides and evidence for their role in exogenous induction of the sugar phosphate transport system of *Escherichia coli*. *Journal of Bacteriology* **169**, 3546–3555.
- WESTON, L. A. & KADNER, R. J. (1988). Role of *uhp* genes in expression of the *Escherichia coli* sugar-phosphate transport system. *Journal of Bacteriology* **170**, 3375–3383.