

CRYSTALLOGRAPHIC
COMMUNICATIONS

ISSN 2056-9890

Received 29 July 2016

Accepted 3 August 2016

Edited by H. Stoeckli-Evans, University of
Neuchâtel, Switzerland**Keywords:** crystal structure; C,N-disubstituted
acetamides; molecular structure; molecular
conformation; hydrogen bonding; C—
halogen $\cdots\pi$ bonding.**CCDC references:** 1497360; 1497359**Supporting information:** this article has
supporting information at journals.iucr.org/e

Crystal structures of two C,N-disubstituted acetamides: 2-(4-chlorophenyl)-N-(2-iodophenyl)acetamide and 2-(4-chlorophenyl)-N-(pyrazin-2-yl)acetamide

Badiadka Narayana,^a Hemmige S. Yathirajan,^{b*} Ravindranath Rathore^c and
Christopher Glidewell^d^aDepartment of Chemistry, Mangalore University, Mangalagangothri 574 199, India, ^bDepartment of Studies in
Chemistry, University of Mysore, Manasagangothri, Mysuru 570 006, India, ^cCentre for Biological Sciences
(Bioinformatics), Central University of South Bihar, BIT Campus, PO, B.V. College, Patna 800 014, Bihar State, India, and
^dSchool of Chemistry, University of St Andrews, Fife KY16 9ST, Scotland. *Correspondence e-mail:
yathirajan@hotmail.com


In the crystal of 2-(4-chlorophenyl)-N-(2-iodophenyl)acetamide, C₁₄H₁₁ClINO, molecules are linked by a combination of N—H \cdots O and C—H \cdots O hydrogen bonds to form a C(4)C(4)[R₂¹(7)] chain of rings and chains of this type are linked by a combination of C—Cl $\cdots\pi$ (arene) and C—I $\cdots\pi$ (arene) interactions to form deeply puckered twofold interwoven sheets. In the crystal of 2-(4-chlorophenyl)-N-(pyrazin-2-yl)acetamide, C₁₂H₁₀ClN₃O, molecules are linked into complex sheets by N—H \cdots N, C—H \cdots N and C—H \cdots O hydrogen bonds, and by C—H $\cdots\pi$ (arene) interactions.

1. Chemical context


Substituted acetamides of the type R¹CH₂CONHR², where R¹ and R² are aromatic or hetero-aromatic substituents, are of interest as they have some resemblance to benzyl penicillins (Pitt, 1952; Csöregy & Palm, 1977; Kojić-Prodić & Ružić-Toroš, 1978; Mijin & Marinković, 2006; Mijin *et al.*, 2008). Here we report on the molecular structures and supra-molecular assembly of two such amides, compounds (I) and (II). The compounds were prepared by the reaction between (4-chlorophenyl)acetic acid and either 2-iodoaniline for (I), or 2-aminopyrazine for (II), using 1-ethyl-3-(3-dimethylamino-propyl)-carbodiimide hydrochloride as the coupling agent.

2. Structural commentary

The molecular conformations of compounds (I) and (II), illustrated in Figs. 1 and 2, respectively, can be defined in terms of the torsional angles N1—C1—C2—C21, 141.8 (3) and 129.22 (18)° respectively, and by the dihedral angles between the central spacer unit, atoms N1,C1,O1,C2, and the two independent rings. The dihedral angles to the chlorinated ring (C21—C26) are 80.02 (11) and 61.74 (6)° in (I) and (II); those to the iodinated ring in (I) and the pyrazinyl ring in (II) are 67.48 (11) and 5.86 (11)°, respectively. This difference is probably associated with the participation in the inter-molecular hydrogen bond of both N atoms of the pyrazinyl ring in (II), as discussed below. The molecules of (I) and (II) do not therefore exhibit any internal symmetry, so that they are conformationally chiral: the centrosymmetric space groups

OPEN  ACCESS

confirm that each compound has crystallized as a conformational racemate.



In the pyrazine ring of compound (II) the four independent C–N distances span a range of only *ca* 0.01 Å, indicating that this ring is fully aromatic.

3. Supramolecular interactions

The hydrogen-bonded assembly in compound (I) is very simple: a combination of N–H···O and C–H···O hydrogen bonds (Table 1) links the molecules into a $C(4)C(4)[R_2^1(6)]$ chain of rings. This chain contains molecules which are related by a *c*-glide plane, producing a chain running parallel to the [001] direction (Fig. 3). There is also a C–H··· π (arene) contact in compound (I) (Table 1), lying within the [001] chain,




Figure 1
The molecular structure of compound (I), showing the atom-labelling scheme. Displacement ellipsoids are drawn at the 30% probability level.




Figure 2
The molecular structure of compound (II), showing the atom-labelling scheme. Displacement ellipsoids are drawn at the 30% probability level.

Table 1
Hydrogen-bond geometry (Å, °) for (I).

*Cg*2 is the centroid of the C21–C26 ring.

<i>D</i> –H··· <i>A</i>	<i>D</i> –H	H··· <i>A</i>	<i>D</i> ··· <i>A</i>	<i>D</i> –H··· <i>A</i>
N1–H1···O1 ⁱ	0.86	2.06	2.908 (3)	167
C2–H2A···O1 ⁱ	0.97	2.58	3.420 (4)	145
C2–H2B··· <i>Cg</i> 2 ⁱ	0.97	2.99	3.589 (3)	121

Symmetry code: (i) $x, -y + \frac{1}{2}, z + \frac{1}{2}$.

but the dimensions make it unlikely that this has any structural significance. Two chains of this type, which are related to one another by inversion, pass through each unit cell, and a combination of C–I··· π (arene) and C–Cl··· π (arene) interactions links the chains into a sheet in the form of a (4,4) net lying parallel to (100) (Fig. 4). The dimensions of these interactions are: for C12–I12···*Cg*1ⁱ [symmetry code: (i) $x, \frac{3}{2} - y, \frac{1}{2} + z$, where *Cg*1 represents the centroid of the C11–C16] ring, I···*Cg* 3.7977 (14), C···*Cg* 5.082 (3) Å and C–I···*Cg* 116.34 (8)°; for C24–Cl24···*Cg*2ⁱⁱ [symmetry code: (ii) $x, -\frac{1}{2} - y, -\frac{1}{2} + z$, where *Cg*2 represents the centroid of the C21–C26 ring], Cl···*Cg* 3.4557 (8), C···*Cg* 4.504 (3) Å and C–Cl···*Cg* 116.19 (11)°. The metrics of the C–Cl···*Cg* interaction are well within the normal range, as deduced using database analysis (Imai *et al.*, 2008).

Because the repeat unit of this sheet in the [010] direction spans two unit cells, there are in fact two such sheets present, related to one another by a unit translation along [010]: the deep puckering of the sheets (Fig. 5) means that the two independent sheets are interwoven. The structure of (I) also contains a short I···O contact with dimension I12···O1ⁱ




Figure 3
Part of the crystal structure of compound (I) showing the formation of a hydrogen-bonded chain of rings running parallel to the [001] direction. Hydrogen bonds are shown as dashed lines and, for the sake of clarity, the H atoms bonded to the C atoms which are not involved in the motif shown have been omitted. The atoms marked with an asterisk (*) or a hash (#) are at the symmetry positions $(x, \frac{1}{2} - y, \frac{1}{2} + z)$ and $(x, \frac{1}{2} - y, -\frac{1}{2} + z)$, respectively.

3.058 (2) Å and C12—I12···O1ⁱ 170.88 (8)° [symmetry code: (i) $x, \frac{3}{2} - y, \frac{1}{2} + z$] which complements the C—Cl···C_g contact. The I···O distance here is significantly shorter than the sum of the van der Waals radii, 3.56 Å (Rowland & Taylor, 1996), or 3.30 Å if account is taken of the polar flattening model (Nyburg & Faerman, 1985).

The hydrogen-bonded supramolecular assembly in compound (II) is more complex than that in compound (I): molecules of (II) are linked into complex sheets by a combination of N—H···N, C—H···N and C—H···O hydrogen bonds, weakly augmented by two C—H···π(arene) hydrogen bonds (Table 2): hydrogen bonds of N—H···O type, often observed in the structures of amides, are absent, however. The formation of this structure can readily be analysed in terms of two simple sub-structures in one- and two-dimensions (Ferguson *et al.*, 1998*a,b*; Gregson *et al.*, 2000). In the simpler of the sub-structures, a combination of N—H···N and C—H···N hydrogen bonds links molecules which are related by the 2_1 screw axis along $(x, \frac{3}{4}, \frac{1}{2})$ into a $C(4)C(5)[R_2^2(7)]$ chain of rings running parallel to the [100] direction (Fig. 6). A more complex one-dimensional sub-




Figure 4

A projection down [100] of part of the crystal structure of compound (I) showing the formation of a sheet built from C—Cl···π(arene) and C—I···π(arene) interactions, shown as thin tapered lines. For the sake of clarity, the H atoms have all been omitted.




Figure 5

A projection down [001] of one of the (100) sheets in the crystal structure of compound (I) showing the deep puckering of the sheet enabling interweaving. The C—X···π(arene) interactions (X = Cl or I) are shown as thin tapered lines, and for the sake of clarity, the H atoms have all been omitted.




Figure 6

Part of the crystal structure of compound (II) showing the formation of a hydrogen-bonded chain of rings running parallel to the [010] direction and built from N—H···N and C—H···N hydrogen bonds, shown as dashed lines. For the sake of clarity, the C-bound H atoms which are not involved in the motifs shown have been omitted. The atoms marked with an asterisk (*) or a hash (#) are at the symmetry positions $(\frac{1}{2} + x, \frac{3}{2} - y, 1 - z)$ and $(-\frac{1}{2} + x, \frac{3}{2} - y, 1 - z)$, respectively.

Table 2

Hydrogen-bond geometry (Å, °) for (II).

Cg2 is the centroid of the C21–C26 ring.

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
N1–H1 \cdots N14 ⁱ	0.85 (2)	2.23 (2)	3.077 (2)	175 (2)
C2–H2A \cdots O1 ⁱⁱ	0.97	2.57	3.461 (3)	153
C13–H13 \cdots N11 ⁱⁱⁱ	0.93	2.50	3.277 (2)	142
C22–H22 \cdots Cg2 ^{iv}	0.93	2.99	3.6416 (17)	129
C25–H25 \cdots Cg2 ^{iv}	0.93	2.89	3.743 (2)	154

Symmetry codes: (i) $x + \frac{1}{2}, -y + \frac{3}{2}, -z + 1$; (ii) $-x + \frac{1}{2}, y - \frac{1}{2}, z$; (iii) $x - \frac{1}{2}, -y + \frac{3}{2}, -z + 1$; (iv) $-x + 1, y + \frac{1}{2}, -z + \frac{3}{2}$.

structure results from the combination of the N–H \cdots N, C–H \cdots N and C–H \cdots O hydrogen bonds, in the form of a ribbon containing alternating $R_2^2(7)$ and $R_4^4(22)$ rings (Fig. 7). The combination of these two chains along [100] and [010] generates a sheet lying parallel to (001) in the domain $\frac{1}{4} < z < \frac{3}{4}$, and a second such sheet, related to the first by inversion, lies in the domain $\frac{3}{4} < z < \frac{5}{4}$. The C–H \cdots π (arene) interactions both lie within the sheet.

4. Database survey

The structures of a number of 2-aryl-*N*-aryl acetamides related to compounds (I) and (II) have been reported recently. We note in particular the structure of 2-(4-chlorophenyl)-*N*-(2,6-dimethylphenyl)acetamide (III) (Narayana *et al.*, 2016), where the molecules are linked by a combination of N–H \cdots O and C–H \cdots O hydrogen bonds to form a $C(4)C(4)[R_2^2(7)]$ chain of rings very much like that in compound (I), except that the molecules comprising the chain in (III) are related by translation along [100], whereas those in (I) are related by a *c*-glide plane. Other recently reported structures include those of *N*-(4-bromophenyl)-2-(4-chlorophenyl)acetamide (IV) (Fun, Shahani *et al.*, 2012), 2-(4-bromophenyl)-*N*-(pyrazin-2-yl)acetamide (V) (Nayak *et al.*, 2013) and 2-(4-chlorophenyl)-*N*-(2,6-dimethylphenyl)acetamide (VI) (Fun, Quah *et al.*, 2012), which are related to compounds (I)–(III), respectively. In addition, the structures of some compounds related to (I), but carrying more than one substituent in the *N*-aryl ring have been reported (Praveen *et al.*, 2013*a,b*; Nayak *et al.*, 2014).

5. Synthesis and crystallization

For the synthesis of compounds (I) and (II), equimolar quantities (1.0 mmol of each component) of (4-chlorophenyl)acetic acid and either 2-iodoaniline for (I), or 2-aminopyrazine for (II), were dissolved in dichloromethane (20 ml) in the presence of 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride (0.01 mol) and triethylamine (0.02 mol) at 273 K. The mixtures were stirred at 273 K for 3 h, and then poured with stirring into an excess of aqueous hydrochloric acid (4 mol dm^{−3}). The aqueous mixtures were exhaustively extracted with dichloromethane and in each case, the combined organic extracts were washed first with saturated aqueous sodium hydrogencarbonate solution and then with




Figure 7

Part of the crystal structure of compound (II) showing the formation of a hydrogen-bonded ribbon of $R_2^2(7)$ and $R_4^4(22)$ rings running parallel to the [100] direction and built from N–H \cdots N, C–H \cdots N and C–H \cdots O hydrogen bonds, shown as dashed lines. For the sake of clarity, the C-bound H atoms which are not involved in the motifs shown have been omitted.

brine. The solutions were dried with anhydrous sodium sulfate and then the solvent was removed under reduced pressure, to give the products. Compound (I): yield 78%, m. p. 441–443 K; analysis found C 45.4, H 2.9, N 3.9%, $C_{14}H_{11}ClINO$ requires C 45.2, H 3.0, N 3.8%. Compound (II): yield 85%, m. p. 421–423 K; analysis found C 58.3, H 4.2, N 16.9%, $C_{12}H_{10}ClN_3O$ requires C 58.2, H 4.1, N 17.0%. Crystals suitable for single-crystal X-ray diffraction analysis were grown by slow evaporation, at ambient temperature and in the presence of air, of solutions in dichloromethane.

6. Refinement

Crystal data, data collection and structure refinement details are summarized in Table 3. All H atoms were located in

Table 3
Experimental details.

	(I)	(II)
Crystal data		
Chemical formula	C ₁₄ H ₁₁ ClINO	C ₁₂ H ₁₀ ClN ₃ O
<i>M_r</i>	371.59	247.68
Crystal system, space group	Monoclinic, <i>P</i> 2 ₁ / <i>c</i>	Orthorhombic, <i>Pbca</i>
Temperature (K)	295	295
<i>a</i> , <i>b</i> , <i>c</i> (Å)	24.001 (1), 6.2369 (3), 9.3266 (4)	10.7041 (4), 7.5724 (3), 28.6619 (11)
α , β , γ (°)	90, 99.621 (2), 90	90, 90, 90
<i>V</i> (Å ³)	1376.48 (11)	2323.21 (15)
<i>Z</i>	4	8
Radiation type	Mo <i>K</i> α	Mo <i>K</i> α
μ (mm ⁻¹)	2.51	0.32
Crystal size (mm)	0.30 × 0.18 × 0.12	0.40 × 0.30 × 0.20
Data collection		
Diffractometer	Bruker APEXII area detector	Bruker APEXII area detector
Absorption correction	Multi-scan (<i>SADABS</i> ; Sheldrick, 2003)	Multi-scan (<i>SADABS</i> ; Sheldrick, 2003)
<i>T</i> _{min} – <i>T</i> _{max}	0.528, 0.740	0.739, 0.939
No. of measured, independent and observed [<i>I</i> > 2 σ (<i>I</i>)] reflections	15082, 3960, 3183	24592, 3380, 2287
<i>R</i> _{int}	0.026	0.029
(<i>sin</i> θ / λ) _{max} (Å ⁻¹)	0.703	0.703
Refinement		
<i>R</i> [<i>F</i> ² > 2 σ (<i>F</i> ²)], <i>wR</i> [<i>F</i> ²], <i>S</i>	0.037, 0.074, 1.07	0.047, 0.136, 1.02
No. of reflections	3960	3380
No. of parameters	163	157
H-atom treatment	H-atom parameters constrained	H-atom parameters constrained
$\Delta\rho_{\text{max}}$, $\Delta\rho_{\text{min}}$ (e Å ⁻³)	1.49, −0.60	0.47, −0.51

Computer programs: *APEX2* and *SAINT-Plus* (Bruker, 2012), *SHELXS97* (Sheldrick, 2008), *SHELXL2014* (Sheldrick, 2015) and *PLATON* (Spek, 2009).

difference Fourier maps. The C-bound H atoms were then treated as riding atoms in geometrically idealized positions with C–H distances 0.93 Å (aromatic and hetero-aromatic) or 0.97 Å (CH₂) and with *U*_{iso}(H) = 1.2*U*_{eq}(C). For the H atoms bonded to N atoms in compound (II), the atomic coordinates were refined with *U*_{iso}(H) = 1.2*U*_{eq}(N) giving the N–H distance shown in Table 2; an attempt to refine similarly the corresponding H-atom coordinates in compound (I) led to an unsatisfactorily low value, 0.74 (3) Å for the N–H distance, possibly associated with the presence of the strongly scattering iodene atom: accordingly this distance was thereafter fixed at 0.86 Å. A small number of low-angle reflections, which had been attenuated by the beam stop [(100) and (200) for (I); (002) for (II)] were omitted from the final cycles of refinement. In the final analysis of variance for compound (I), there was a large value, 4.245, of $K = [\text{mean}(F_o^2)/\text{mean}(F_c^2)]$ for the group of 428 very weak reflections having *F_o*/*F_c*(max) in the range 0.000 < *F_o*/*F_c*(max) < 0.008.

Acknowledgements

BN thanks the UGC (India) for financial assistance through a BSR one-time grant for the purchase of chemicals. The authors thank P. S. Nayak of Mangalore University for the synthesis of the compounds. RSR thanks the Head of the Sophisticated Analytical Instrument Facility (SAIF), IIT, Chennai, for X-ray data collection.

References

- Bruker (2012). *APEX2* and *SAINT-Plus*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Csöregi, I. & Palm, T.-B. (1977). *Acta Cryst.* **B33**, 2169–2175.
- Ferguson, G., Glidewell, C., Gregson, R. M. & Meehan, P. R. (1998*a*). *Acta Cryst.* **B54**, 129–138.
- Ferguson, G., Glidewell, C., Gregson, R. M. & Meehan, P. R. (1998*b*). *Acta Cryst.* **B54**, 139–150.
- Fun, H.-K., Quah, C. K., Nayak, P. S., Narayana, B. & Sarojini, B. K. (2012). *Acta Cryst.* **E68**, o2678.
- Fun, H.-K., Shahani, T., Nayak, P. S., Narayana, B. & Sarojini, B. K. (2012). *Acta Cryst.* **E68**, o519.
- Gregson, R. M., Glidewell, C., Ferguson, G. & Lough, A. J. (2000). *Acta Cryst.* **B56**, 39–57.
- Imai, Y. N., Inoue, Y., Nakanishi, I. & Kitaura, K. (2008). *Protein Sci.* **17**, 1129–1137.
- Kojić-Prodić, B. & Ružić-Toroš, Ž. (1978). *Acta Cryst.* **B34**, 1271–1275.
- Mijin, D. & Marinković, A. (2006). *Synth. Commun.* **36**, 193–198.
- Mijin, D. Z., Prascovic, M. & Petrovic, S. D. (2008). *J. Serb. Chem. Soc.* **73**, 945–950.
- Narayana, B., Yathirajan, H. S., Rathore, R. S. & Glidewell, C. (2016). Private communication (CCDC 1491671). CCDC, Cambridge, England.
- Nayak, P. S., Jasinski, J. P., Golen, J. A., Narayana, B., Kaur, M., Yathirajan, H. S. & Glidewell, C. (2014). *Acta Cryst.* **C70**, 889–894.
- Nayak, P. S., Narayana, B., Jasinski, J. P., Yathirajan, H. S. & Sarojini, B. K. (2013). *Acta Cryst.* **E69**, o891.
- Nyburg, S. C. & Faerman, C. H. (1985). *Acta Cryst.* **B41**, 274–279.
- Pitt, G. J. (1952). *Acta Cryst.* **5**, 770–775.

- Praveen, A. S., Yathirajan, H. S., Jasinski, J. P., Keeley, A. C., Narayana, B. & Sarojini, B. K. (2013*a*). *Acta Cryst.* **E69**, o900–o901.
- Praveen, A. S., Yathirajan, H. S., Jasinski, J. P., Keeley, A. C., Narayana, B. & Sarojini, B. K. (2013*b*). *Acta Cryst.* **E69**, o996–o997.
- Rowland, R. S. & Taylor, R. (1996). *J. Phys. Chem.* **100**, 7384–7391.
- Sheldrick, G. M. (2003). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Sheldrick, G. M. (2015). *Acta Cryst.* **C71**, 3–8.
- Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.

supporting information

Acta Cryst. (2016). E72, 1270-1275 [doi:10.1107/S2056989016012512]

Crystal structures of two C,N-disubstituted acetamides: 2-(4-chlorophenyl)-N-(2-iodophenyl)acetamide and 2-(4-chlorophenyl)-N-(pyrazin-2-yl)acetamide

Badiadka Narayana, Hemmige S. Yathirajan, Ravindranath Rathore and Christopher Glidewell

Computing details

For both compounds, data collection: *APEX2* (Bruker, 2012); cell refinement: *APEX2* (Bruker, 2012); data reduction: *SAINT-Plus* (Bruker, 2012); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL2014* (Sheldrick, 2015); molecular graphics: *PLATON* (Spek, 2009); software used to prepare material for publication: *SHELXL2014* (Sheldrick, 2015) and *PLATON* (Spek, 2009).

(I) 2-(4-Chlorophenyl)-N-(2-iodophenyl)acetamide

Crystal data

C₁₄H₁₁ClINO

M_r = 371.59

Monoclinic, *P2₁/c*

a = 24.001 (1) Å

b = 6.2369 (3) Å

c = 9.3266 (4) Å

β = 99.621 (2)°

V = 1376.48 (11) Å³

Z = 4

F(000) = 720

D_x = 1.793 Mg m⁻³

Mo *Kα* radiation, λ = 0.71073 Å

Cell parameters from 5030 reflections

θ = 0.9–33.5°

μ = 2.51 mm⁻¹

T = 295 K

Block, colourless

0.30 × 0.18 × 0.12 mm

Data collection

Bruker APEXII area detector
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

φ and ω scans

Absorption correction: multi-scan
(SADABS; Sheldrick, 2003)

T_{min} = 0.528, *T_{max}* = 0.740

15082 measured reflections

3960 independent reflections

3183 reflections with *I* > 2σ(*I*)

R_{int} = 0.026

θ_{max} = 30.0°, θ_{min} = 2.6°

h = -33→33

k = -8→8

l = -13→9

Refinement

Refinement on *F*²

Least-squares matrix: full

R[*F*² > 2σ(*F*²)] = 0.037

wR(*F*²) = 0.074

S = 1.07

3960 reflections

163 parameters

0 restraints

Hydrogen site location: inferred from
neighbouring sites

H-atom parameters constrained

w = 1/[σ²(*F_o*²) + (0.0189*P*)² + 1.9949*P*]

where *P* = (*F_o*² + 2*F_c*²)/3

(Δ/σ)_{max} = 0.001

Δρ_{max} = 1.49 e Å⁻³

Δρ_{min} = -0.60 e Å⁻³

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.25595 (11)	0.3149 (5)	0.2858 (3)	0.0344 (6)
O1	0.24811 (9)	0.3574 (4)	0.1558 (2)	0.0496 (6)
N1	0.21438 (10)	0.3174 (4)	0.3652 (2)	0.0340 (5)
H1	0.2223	0.2839	0.4557	0.041*
C2	0.31392 (12)	0.2634 (6)	0.3678 (3)	0.0445 (7)
H2A	0.3101	0.1900	0.4574	0.053*
H2B	0.3342	0.3962	0.3935	0.053*
C11	0.15786 (11)	0.3726 (4)	0.3066 (3)	0.0325 (6)
C12	0.13408 (11)	0.5596 (4)	0.3479 (3)	0.0350 (6)
I12	0.18141 (2)	0.77640 (3)	0.48960 (2)	0.04554 (8)
C13	0.07856 (13)	0.6103 (6)	0.2902 (4)	0.0506 (8)
H13	0.0623	0.7349	0.3192	0.061*
C14	0.04773 (14)	0.4773 (7)	0.1907 (4)	0.0610 (10)
H14	0.0107	0.5131	0.1511	0.073*
C15	0.07091 (15)	0.2923 (6)	0.1492 (5)	0.0626 (10)
H15	0.0497	0.2025	0.0815	0.075*
C16	0.12596 (14)	0.2378 (5)	0.2076 (4)	0.0456 (7)
H16	0.1415	0.1106	0.1802	0.055*
C21	0.34782 (11)	0.1252 (5)	0.2816 (3)	0.0363 (6)
C22	0.39910 (13)	0.1937 (5)	0.2504 (4)	0.0441 (7)
H22	0.4129	0.3279	0.2821	0.053*
C23	0.43017 (13)	0.0644 (6)	0.1724 (4)	0.0472 (8)
H23	0.4646	0.1119	0.1512	0.057*
C24	0.41012 (12)	-0.1323 (6)	0.1268 (3)	0.0440 (7)
Cl24	0.44931 (4)	-0.2968 (2)	0.03026 (13)	0.0754 (3)
C25	0.35920 (13)	-0.2053 (5)	0.1558 (4)	0.0460 (7)
H25	0.3457	-0.3397	0.1237	0.055*
C26	0.32848 (12)	-0.0756 (6)	0.2333 (3)	0.0440 (7)
H26	0.2940	-0.1241	0.2536	0.053*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0379 (14)	0.0370 (15)	0.0273 (14)	0.0082 (11)	0.0026 (11)	-0.0040 (11)
O1	0.0484 (12)	0.0757 (16)	0.0244 (11)	0.0167 (11)	0.0056 (9)	0.0077 (11)
N1	0.0421 (12)	0.0387 (13)	0.0214 (11)	0.0094 (10)	0.0057 (9)	0.0008 (9)
C2	0.0408 (15)	0.065 (2)	0.0257 (14)	0.0123 (14)	-0.0011 (11)	-0.0074 (14)
C11	0.0379 (14)	0.0331 (14)	0.0268 (13)	0.0007 (11)	0.0059 (10)	0.0026 (11)
C12	0.0385 (14)	0.0322 (14)	0.0349 (15)	-0.0005 (11)	0.0082 (11)	-0.0002 (11)

I12	0.05868 (14)	0.03484 (11)	0.04157 (12)	-0.00133 (9)	0.00388 (9)	-0.00538 (9)
C13	0.0365 (16)	0.0478 (19)	0.067 (2)	0.0083 (13)	0.0081 (15)	-0.0018 (17)
C14	0.0366 (17)	0.068 (2)	0.075 (3)	-0.0006 (16)	-0.0011 (16)	0.002 (2)
C15	0.0490 (19)	0.064 (2)	0.068 (3)	-0.0141 (17)	-0.0083 (17)	-0.011 (2)
C16	0.0515 (18)	0.0405 (17)	0.0431 (18)	0.0003 (13)	0.0031 (14)	-0.0052 (14)
C21	0.0332 (13)	0.0469 (17)	0.0271 (14)	0.0082 (12)	0.0003 (10)	0.0029 (12)
C22	0.0397 (15)	0.0464 (18)	0.0441 (18)	-0.0023 (13)	0.0013 (13)	-0.0022 (14)
C23	0.0350 (15)	0.062 (2)	0.0460 (19)	-0.0005 (14)	0.0100 (13)	0.0001 (16)
C24	0.0380 (15)	0.060 (2)	0.0328 (16)	0.0157 (14)	0.0030 (12)	-0.0055 (14)
Cl24	0.0581 (5)	0.0985 (8)	0.0703 (7)	0.0263 (5)	0.0122 (5)	-0.0270 (6)
C25	0.0452 (17)	0.0417 (17)	0.0487 (19)	0.0025 (13)	0.0008 (14)	-0.0069 (14)
C26	0.0304 (14)	0.0551 (19)	0.0466 (18)	-0.0005 (13)	0.0063 (12)	0.0012 (15)

Geometric parameters (Å, °)

C1—O1	1.225 (3)	C14—H14	0.9300
C1—N1	1.338 (4)	C15—C16	1.385 (5)
C1—C2	1.506 (4)	C15—H15	0.9300
N1—C11	1.418 (3)	C16—H16	0.9300
N1—H1	0.8600	C21—C22	1.379 (4)
C2—C21	1.507 (4)	C21—C26	1.385 (4)
C2—H2A	0.9700	C22—C23	1.385 (5)
C2—H2B	0.9700	C22—H22	0.9300
C11—C12	1.381 (4)	C23—C24	1.360 (5)
C11—C16	1.381 (4)	C23—H23	0.9300
C12—C13	1.388 (4)	C24—C25	1.373 (5)
C12—I12	2.089 (3)	C24—Cl24	1.741 (3)
C13—C14	1.366 (5)	C25—C26	1.378 (4)
C13—H13	0.9300	C25—H25	0.9300
C14—C15	1.365 (5)	C26—H26	0.9300
O1—C1—N1	122.7 (3)	C14—C15—C16	120.2 (3)
O1—C1—C2	121.7 (3)	C14—C15—H15	119.9
N1—C1—C2	115.6 (2)	C16—C15—H15	119.9
C1—N1—C11	122.9 (2)	C11—C16—C15	120.0 (3)
C1—N1—H1	118.5	C11—C16—H16	120.0
C11—N1—H1	118.5	C15—C16—H16	120.0
C1—C2—C21	112.7 (2)	C22—C21—C26	118.3 (3)
C1—C2—H2A	109.0	C22—C21—C2	121.0 (3)
C21—C2—H2A	109.0	C26—C21—C2	120.6 (3)
C1—C2—H2B	109.0	C21—C22—C23	120.6 (3)
C21—C2—H2B	109.0	C21—C22—H22	119.7
H2A—C2—H2B	107.8	C23—C22—H22	119.7
C12—C11—C16	119.4 (3)	C24—C23—C22	119.8 (3)
C12—C11—N1	120.7 (2)	C24—C23—H23	120.1
C16—C11—N1	119.8 (3)	C22—C23—H23	120.1
C11—C12—C13	119.9 (3)	C23—C24—C25	121.2 (3)
C11—C12—I12	121.1 (2)	C23—C24—Cl24	120.0 (3)

C13—C12—I12	118.9 (2)	C25—C24—C124	118.8 (3)
C14—C13—C12	120.1 (3)	C24—C25—C26	118.7 (3)
C14—C13—H13	120.0	C24—C25—H25	120.6
C12—C13—H13	120.0	C26—C25—H25	120.6
C15—C14—C13	120.4 (3)	C25—C26—C21	121.5 (3)
C15—C14—H14	119.8	C25—C26—H26	119.3
C13—C14—H14	119.8	C21—C26—H26	119.3
O1—C1—N1—C11	-1.1 (5)	N1—C11—C16—C15	-179.4 (3)
C2—C1—N1—C11	177.0 (3)	C14—C15—C16—C11	-1.0 (6)
O1—C1—C2—C21	-40.1 (4)	C1—C2—C21—C22	121.2 (3)
N1—C1—C2—C21	141.8 (3)	C1—C2—C21—C26	-59.7 (4)
C1—N1—C11—C12	-111.7 (3)	C26—C21—C22—C23	0.2 (5)
C1—N1—C11—C16	68.6 (4)	C2—C21—C22—C23	179.3 (3)
C16—C11—C12—C13	0.1 (4)	C21—C22—C23—C24	-0.4 (5)
N1—C11—C12—C13	-179.6 (3)	C22—C23—C24—C25	0.4 (5)
C16—C11—C12—I12	-177.1 (2)	C22—C23—C24—C124	-179.3 (3)
N1—C11—C12—I12	3.2 (4)	C23—C24—C25—C26	-0.3 (5)
C11—C12—C13—C14	-1.1 (5)	C124—C24—C25—C26	179.4 (3)
I12—C12—C13—C14	176.2 (3)	C24—C25—C26—C21	0.2 (5)
C12—C13—C14—C15	1.1 (6)	C22—C21—C26—C25	-0.1 (5)
C13—C14—C15—C16	0.0 (6)	C2—C21—C26—C25	-179.2 (3)
C12—C11—C16—C15	0.9 (5)		

Hydrogen-bond geometry (Å, °)

Cg2 is the centroid of the C21—C26 ring.

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
N1—H1...O1 ⁱ	0.86	2.06	2.908 (3)	167
C2—H2 <i>A</i> ...O1 ⁱ	0.97	2.58	3.420 (4)	145
C2—H2 <i>B</i> ...Cg2 ⁱ	0.97	2.99	3.589 (3)	121

Symmetry code: (i) *x*, -*y*+1/2, *z*+1/2.**(II) 2-(4-Chlorophenyl)-*N*-(pyrazin-2-yl)acetamide***Crystal data*C₁₂H₁₀ClN₃O*M_r* = 247.68Orthorhombic, *Pbca**a* = 10.7041 (4) Å*b* = 7.5724 (3) Å*c* = 28.6619 (11) Å*V* = 2323.21 (15) Å³*Z* = 8*F*(000) = 1024*D_x* = 1.416 Mg m⁻³Mo *Kα* radiation, λ = 0.71073 Å

Cell parameters from 3760 reflections

θ = 1.4–32.3°

μ = 0.32 mm⁻¹*T* = 295 K

Block, colourless

0.40 × 0.30 × 0.20 mm

Data collection

Bruker APEXII area detector diffractometer	24592 measured reflections 3380 independent reflections
Radiation source: fine-focus sealed tube	2287 reflections with $I > 2\sigma(I)$
Graphite monochromator	$R_{\text{int}} = 0.029$
φ and ω scans	$\theta_{\text{max}} = 30.0^\circ$, $\theta_{\text{min}} = 2.4^\circ$
Absorption correction: multi-scan (SADABS; Sheldrick, 2003)	$h = -15 \rightarrow 14$
$T_{\text{min}} = 0.739$, $T_{\text{max}} = 0.939$	$k = -9 \rightarrow 10$
	$l = -40 \rightarrow 40$

Refinement

Refinement on F^2	Hydrogen site location: mixed
Least-squares matrix: full	H-atom parameters constrained
$R[F^2 > 2\sigma(F^2)] = 0.047$	$w = 1/[\sigma^2(F_o^2) + (0.0547P)^2 + 0.9776P]$
$wR(F^2) = 0.136$	where $P = (F_o^2 + 2F_c^2)/3$
$S = 1.02$	$(\Delta/\sigma)_{\text{max}} < 0.001$
3380 reflections	$\Delta\rho_{\text{max}} = 0.47 \text{ e } \text{\AA}^{-3}$
157 parameters	$\Delta\rho_{\text{min}} = -0.51 \text{ e } \text{\AA}^{-3}$
0 restraints	

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.34192 (16)	0.6027 (3)	0.57916 (6)	0.0514 (4)
O1	0.23005 (12)	0.6165 (3)	0.58396 (5)	0.0760 (5)
N1	0.40586 (13)	0.6699 (2)	0.54222 (5)	0.0488 (4)
H1	0.484 (2)	0.654 (3)	0.5421 (7)	0.059*
C2	0.42190 (17)	0.5037 (3)	0.61387 (7)	0.0561 (5)
H2A	0.4031	0.3787	0.6118	0.067*
H2B	0.5091	0.5195	0.6057	0.067*
N11	0.44426 (13)	0.8104 (2)	0.47344 (6)	0.0522 (4)
C12	0.35844 (15)	0.7566 (2)	0.50359 (6)	0.0430 (4)
C13	0.23119 (15)	0.7832 (3)	0.49531 (6)	0.0472 (4)
H13	0.1730	0.7436	0.5170	0.057*
N14	0.19234 (13)	0.8640 (2)	0.45707 (5)	0.0507 (4)
C15	0.27929 (17)	0.9194 (3)	0.42717 (7)	0.0531 (4)
H15	0.2550	0.9773	0.4000	0.064*
C16	0.40366 (18)	0.8925 (3)	0.43564 (7)	0.0562 (5)
H16	0.4617	0.9335	0.4141	0.067*
C21	0.40218 (15)	0.5639 (2)	0.66329 (6)	0.0440 (4)
C22	0.29842 (16)	0.5106 (2)	0.68824 (6)	0.0470 (4)
H22	0.2383	0.4408	0.6738	0.056*
C23	0.28265 (16)	0.5594 (2)	0.73413 (6)	0.0492 (4)
H23	0.2124	0.5228	0.7506	0.059*

C24	0.37099 (18)	0.6618 (2)	0.75533 (6)	0.0486 (4)
Cl24	0.35395 (7)	0.71602 (9)	0.81390 (2)	0.0808 (2)
C25	0.47361 (18)	0.7207 (3)	0.73145 (8)	0.0575 (5)
H25	0.5325	0.7925	0.7460	0.069*
C26	0.48787 (16)	0.6715 (3)	0.68544 (7)	0.0555 (5)
H26	0.5570	0.7118	0.6689	0.067*

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0365 (8)	0.0669 (12)	0.0507 (9)	0.0063 (8)	0.0034 (7)	-0.0025 (8)
O1	0.0359 (7)	0.1235 (14)	0.0685 (9)	0.0117 (8)	0.0105 (6)	0.0210 (9)
N1	0.0278 (6)	0.0693 (10)	0.0492 (8)	0.0028 (7)	0.0001 (6)	-0.0025 (7)
C2	0.0434 (9)	0.0685 (13)	0.0563 (10)	0.0119 (9)	0.0050 (8)	0.0023 (9)
N11	0.0331 (7)	0.0684 (11)	0.0549 (8)	0.0003 (7)	0.0030 (6)	-0.0011 (8)
C12	0.0315 (7)	0.0526 (9)	0.0449 (8)	-0.0001 (7)	-0.0004 (6)	-0.0100 (7)
C13	0.0303 (7)	0.0622 (11)	0.0489 (9)	0.0015 (7)	-0.0005 (6)	-0.0074 (8)
N14	0.0373 (7)	0.0616 (10)	0.0532 (8)	0.0057 (7)	-0.0032 (6)	-0.0080 (7)
C15	0.0474 (10)	0.0615 (12)	0.0503 (9)	0.0049 (9)	-0.0019 (7)	-0.0021 (9)
C16	0.0437 (9)	0.0686 (13)	0.0563 (10)	0.0005 (9)	0.0070 (8)	0.0014 (9)
C21	0.0337 (7)	0.0449 (9)	0.0534 (9)	0.0055 (7)	0.0011 (7)	0.0054 (7)
C22	0.0401 (8)	0.0454 (10)	0.0556 (10)	-0.0074 (7)	-0.0003 (7)	0.0003 (8)
C23	0.0453 (9)	0.0463 (10)	0.0560 (9)	-0.0047 (8)	0.0067 (7)	0.0046 (8)
C24	0.0536 (10)	0.0421 (9)	0.0499 (9)	0.0035 (8)	-0.0046 (8)	0.0009 (7)
Cl24	0.1048 (5)	0.0848 (5)	0.0529 (3)	-0.0046 (4)	-0.0051 (3)	-0.0082 (3)
C25	0.0456 (10)	0.0533 (11)	0.0738 (12)	-0.0094 (8)	-0.0114 (9)	-0.0054 (9)
C26	0.0341 (8)	0.0615 (12)	0.0709 (12)	-0.0066 (8)	0.0044 (8)	0.0059 (10)

Geometric parameters (Å, °)

C1—O1	1.210 (2)	C15—C16	1.368 (3)
C1—N1	1.360 (2)	C15—H15	0.9300
C1—C2	1.511 (3)	C16—H16	0.9300
N1—C12	1.384 (2)	C21—C22	1.381 (2)
N1—H1	0.85 (2)	C21—C26	1.382 (3)
C2—C21	1.503 (3)	C22—C23	1.377 (3)
C2—H2A	0.9700	C22—H22	0.9300
C2—H2B	0.9700	C23—C24	1.366 (3)
N11—C16	1.323 (3)	C23—H23	0.9300
N11—C12	1.325 (2)	C24—C25	1.369 (3)
C12—C13	1.397 (2)	C24—Cl24	1.7378 (19)
C13—N14	1.322 (2)	C25—C26	1.379 (3)
C13—H13	0.9300	C25—H25	0.9300
N14—C15	1.333 (2)	C26—H26	0.9300
O1—C1—N1	123.67 (18)	C16—C15—H15	119.4
O1—C1—C2	121.92 (18)	N11—C16—C15	122.33 (18)
N1—C1—C2	114.39 (15)	N11—C16—H16	118.8

C1—N1—C12	128.03 (14)	C15—C16—H16	118.8
C1—N1—H1	116.6 (14)	C22—C21—C26	117.92 (17)
C12—N1—H1	115.3 (14)	C22—C21—C2	120.83 (17)
C21—C2—C1	112.98 (15)	C26—C21—C2	121.24 (16)
C21—C2—H2A	109.0	C23—C22—C21	121.00 (17)
C1—C2—H2A	109.0	C23—C22—H22	119.5
C21—C2—H2B	109.0	C21—C22—H22	119.5
C1—C2—H2B	109.0	C24—C23—C22	119.51 (17)
H2A—C2—H2B	107.8	C24—C23—H23	120.2
C16—N11—C12	116.79 (15)	C22—C23—H23	120.2
N11—C12—N1	114.40 (14)	C23—C24—C25	121.21 (18)
N11—C12—C13	121.38 (16)	C23—C24—Cl24	119.43 (15)
N1—C12—C13	124.19 (16)	C25—C24—Cl24	119.35 (15)
N14—C13—C12	120.94 (16)	C24—C25—C26	118.61 (18)
N14—C13—H13	119.5	C24—C25—H25	120.7
C12—C13—H13	119.5	C26—C25—H25	120.7
C13—N14—C15	117.31 (15)	C25—C26—C21	121.70 (17)
N14—C15—C16	121.23 (18)	C25—C26—H26	119.2
N14—C15—H15	119.4	C21—C26—H26	119.2
O1—C1—N1—C12	-2.8 (3)	N14—C15—C16—N11	0.3 (3)
C2—C1—N1—C12	175.70 (18)	C1—C2—C21—C22	77.6 (2)
O1—C1—C2—C21	-52.2 (3)	C1—C2—C21—C26	-103.5 (2)
N1—C1—C2—C21	129.22 (18)	C26—C21—C22—C23	-1.7 (3)
C16—N11—C12—N1	179.33 (17)	C2—C21—C22—C23	177.25 (17)
C16—N11—C12—C13	1.0 (3)	C21—C22—C23—C24	-0.1 (3)
C1—N1—C12—N11	178.81 (18)	C22—C23—C24—C25	1.7 (3)
C1—N1—C12—C13	-2.9 (3)	C22—C23—C24—Cl24	-177.50 (14)
N11—C12—C13—N14	-0.3 (3)	C23—C24—C25—C26	-1.4 (3)
N1—C12—C13—N14	-178.47 (17)	Cl24—C24—C25—C26	177.80 (16)
C12—C13—N14—C15	-0.4 (3)	C24—C25—C26—C21	-0.5 (3)
C13—N14—C15—C16	0.4 (3)	C22—C21—C26—C25	2.0 (3)
C12—N11—C16—C15	-1.0 (3)	C2—C21—C26—C25	-176.93 (19)

Hydrogen-bond geometry (\AA , $^\circ$)

Cg2 is the centroid of the C21—C26 ring.

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
N1—H1 \cdots N14 ⁱ	0.85 (2)	2.23 (2)	3.077 (2)	175 (2)
C2—H2A \cdots O1 ⁱⁱ	0.97	2.57	3.461 (3)	153
C13—H13 \cdots N11 ⁱⁱⁱ	0.93	2.50	3.277 (2)	142
C22—H22 \cdots Cg2 ⁱⁱ	0.93	2.99	3.6416 (17)	129
C25—H25 \cdots Cg2 ^{iv}	0.93	2.89	3.743 (2)	154

Symmetry codes: (i) $x+1/2, -y+3/2, -z+1$; (ii) $-x+1/2, y-1/2, z$; (iii) $x-1/2, -y+3/2, -z+1$; (iv) $-x+1, y+1/2, -z+3/2$.