

# PCCP

Accepted Manuscript



This article can be cited before page numbers have been issued, to do this please use: R. A. Cormanich, N. S. Keddie, R. Rittner, D. O'Hagan and M. Buhl, *Phys. Chem. Chem. Phys.*, 2015, DOI: 10.1039/C5CP04537A.



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

*Accepted Manuscripts* are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.



Journal Name

COMMUNICATION

## Particularly strong C-H $\cdots\pi$ interactions between benzene and all-*cis* 1,2,3,4,5,6-hexafluorocyclohexane

Received 00th January 20xx,  
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

Rodrigo A. Cormanich,<sup>a,b</sup> Neil Keddie,<sup>a</sup> Roberto Rittner,<sup>b</sup> David O'Hagan\*<sup>a</sup> and Michael Bühl<sup>a†</sup>

www.rsc.org/


We present the first high-level *ab initio* benchmark study of the interaction energy between fluorocyclohexanes and benzene. These compounds form CH $\cdots\pi$  interactions with aromatic solvents which causes notable shielding of the axial cyclohexane protons. For the recently synthesised all-*cis* 1,2,3,4,5,6-hexafluorocyclohexane the interaction energy with benzene amounts to -7.9 kcal/mol and -6.4 kcal/mol at the MP2 and SCS-MP2 levels, respectively (extrapolated to the complete basis set limit), which according to dispersion-corrected density functional calculations, is largely due to dispersion.

When fluorine atoms are added to organic compounds, interesting physicochemical properties may arise, which cannot be obtained using other elements from the periodic table.<sup>[1]</sup> Indeed, it was recently shown both experimentally<sup>[2]</sup> by <sup>1</sup>H NMR and theoretically<sup>[3]</sup> that the all-*cis* 1,2,4,5- (**1**) and the 1,2,3,4-tetrafluorocyclohexanes (**2**, Figures 1a and 1b) form CH $\cdots\pi$  interactions with molecules of aromatic solvents. These interactions arise from the high polarity of these all-*cis* tetrafluoro species, which have a "negative face" on the side of the axial fluorine atoms and "a positive" face on the hydrogen side (Figure 1c). Such CH $\cdots\pi$  interactions lead to a close contact between the *axial* hydrogens and the arene  $\pi$  electrons. Thus, the axial hydrogen atoms show unusually large upfield shifts caused by anisotropic diamagnetism originating from the arene ring current (Figure 1d).


Recently, the synthesis of all-*cis* 1,2,3,4,5,6-hexafluorocyclohexane (**3**) was reported.<sup>[4]</sup> This compound (Figure 2a) was found to bear a very high dipole moment for a non-ionic organic compound, calculated to be 6.2 D at the M11/6-311G(2d,p) level. This hexafluorocyclohexane will reasonably have an even higher interaction energy with benzene compared to the tetrafluorocyclohexanes **1** and **2**.

Our previous DFT analysis of those CH $\cdots\pi$  interactions for the

complex formed between **1** or **2** and benzene indicated a binding energy of ca. -1.5 kcal mol<sup>-1</sup> at the B3LYP/def2-TZVP level and a CH $\cdots\pi$  distance of ca. 3.10-3.25 Å.<sup>[3]</sup> When one includes Grimme's DFT-D or DFT-D3 dispersion corrections<sup>[5]</sup> the binding energies for those complexes rises to ca. -6 kcal mol<sup>-1</sup> and much closer C-H $\cdots\pi$  contacts of ca. 2.6-2.7 Å result. However, the NMR chemical shifts computed for the B3LYP-optimised complexes appeared to reproduce the upfield shifts observed in aromatic solvents better than those using the tighter geometries obtained with dispersion corrections.



**Figure 1:** Schematic representations of **a)** all-*cis* 1,2,3,4-tetrafluorocyclohexane (**1**), **b)** all-*cis* 1,2,3,4-tetrafluorocyclohexane (**2**); **c)** the "negative" and "positive" faces for compound **1**; and **d)** the induced ring-current effect on axial <sup>1</sup>H atoms of **1** interacting with a benzene molecule.



**Figure 2:** **a)** Structural representation of all-*cis* 1,2,3,4,5,6-hexafluorocyclohexane (**3**). **b)** Electrostatic potential (ESP) of **3** [B3LYP-D3/def2-TZVP] color-coded on a scale from -0.03 au (red) to +0.03 au (blue) and mapped onto an isodensity surface  $\rho = 0.001$  au.

<sup>a</sup>EastChem School of Chemistry, University of St Andrews, North Haugh, St Andrews, Fife, KY16 9ST, UK Fax: 01334 463800; Tel: 01334 467171;


<sup>b</sup>Chemistry Institute, University of Campinas, Campinas, SP, 13083-970, Brazil, Tel (19) 3521 3150; Fax. (19) 3521 3023

E-mail: mb105@st-andrews.ac.uk

† Electronic Supplementary Information (ESI) available: Computational details, NMR spectra and experimental details, Extrapolation details and data, calculated isotropic shielding data.

In order to validate these DFT results and to arrive at a more confident value for the interaction energy between these polar fluorocyclohexanes and aromatics, high-level *ab initio* benchmarks are necessary. The newly synthesised hexafluorocyclohexane **3** is an ideal target for this purpose, because its complex with the simplest aromatic, benzene, can have high symmetry ( $C_{3v}$ , Figure 3), all  $CH\cdots\pi$  contacts being equivalent. We now present such a high-level *ab initio* benchmark for the  $CH\cdots\pi$  binding energies between compound **3** and benzene at MP2 and SCS-MP2 levels, extrapolated to the complete basis set (CBS). In addition, the  $^1H$  chemical shift values for **3** were recorded in  $CD_2Cl_2$  and benzene- $d^6$ , and the resulting upfield shifts were compared with those computed on going from free **3** to the complex with benzene.

In order to more fully evaluate the effect of fluorine atoms on the binding energy of **3** with benzene, two additional model systems were calculated, namely all-*cis* 1,3,5-trifluorocyclohexane (**4**) and cyclohexane (**5**) (Figure 3).



**Figure 3:** Calculated geometries of benzene with: **a**) all-*cis* 1,2,3,4,5,6-hexafluorocyclohexane (**3**), **b**) all-*cis* 1,3,5-trifluorocyclohexane (**4**) and **c**) cyclohexane (**5**). Geometries shown have  $C_{3v}$  symmetry.

Initial calculations were performed at B3LYP, B3LYP-D3 and MP2 levels in conjunction with the def2-TZVP basis set, similar to the levels used in our previous studies of compounds **1** and **2.3**. With tight optimisation criteria and an "ultrafine" integration grid (see Computational Details in the ESI), the complexes of **3** and **4** with benzene are true minima at the B3LYP level in  $C_{3v}$  symmetry. At B3LYP-D3 a very small imaginary frequency appears, which describes rotation of the two rings relative to each other about the  $C_3$  axis. This rotation is indicated to lower the energy by just fractions of a kcal/mol, implying essentially free rotation of the two parallel rings. Thus, we kept  $C_{3v}$  symmetry imposed throughout.

Consistent with our previous work on **1** and **2**,<sup>[3]</sup> the B3LYP functional showed much longer contacts between the cyclohexanes and benzene, as well as smaller energy values than B3LYP-D3 and also MP2 (Table 1). Indeed, as expected, B3LYP fails to find an interaction between the parent cyclohexane (**5**) and benzene. On the other hand, B3LYP-D3 and MP2 find strong  $CH\cdots\pi$  binding energies with short distances for all compounds **3-5**. When converted into enthalpies and Gibbs free energies using standard thermodynamic corrections from the frequency calculations from each level, the binding energy becomes weaker for enthalpies and even endergonic for Gibbs free energies (Table S1 in the ESI). Still, for the complexes between benzene and **3** or **4**, B3LYP-D3 and MP2 indicate binding energies approaching and even exceeding strengths of typical hydrogen bonds (e.g. ca. 5 kcal/mol for the water dimer<sup>6</sup>). This binding energy increases steadily with the number of fluorines, from ca. -3 kcal/mol for  $n = 0$  via -5 kcal/mol and -6 kcal/mol for  $n = 3$  and 4, respectively, to -7 kcal/mol for  $n = 6$  (B3LYP-D3/def2-TZVP level, Table 1). From these data, it is evident that it is not only the bond dipoles from the axial fluorine atoms in  $3\cdot C_6H_6$  that are responsible for the strong binding, but that both equatorial and axial fluorine atoms are important. Comparison of

B3LYP and B3LYP-D3 results (Table 1) indicates that the largest fraction of this interaction stems from dispersion rather than from electrostatic (e.g. dipole-quadrupole) interactions.

**Table 1:** Calculated distances and binding energies obtained at B3LYP/def2-TZVP, B3LYP-D3/def2-TZVP and MP2/aug-cc-pVDZ levels for complexes of **1-5** and benzene. Complexes for compounds **3-5** have  $C_{3v}$  symmetry. Optimised C-H $\cdots\pi$  distance in angstroms were obtained with basis set superposition error (BSSE) corrections included through the counterpoise method.

		C-H $\cdots\pi$ distance	Binding energy (kcal mol <sup>-1</sup> )
B3LYP	<b>1</b>	3.27 Å <sup>[a]</sup>	-1.26
	<b>2</b>	3.35 Å <sup>[a]</sup>	-1.28
	<b>3</b>	3.11 Å	-2.12
	<b>4</b>	3.44 Å	-0.75
	<b>5</b>	n.a. <sup>[b]</sup>	n.a. <sup>[b]</sup>
B3LYP-D3	<b>1</b>	2.77 Å	-5.76
	<b>2</b>	2.90 Å <sup>[a]</sup>	-6.05
	<b>3</b>	2.69 Å	-7.06
	<b>4</b>	2.79 Å	-4.84
	<b>5</b>	2.83 Å	-3.40
MP2	<b>1</b>	2.78	-5.71
	<b>2</b>	2.90 Å <sup>[a]</sup>	-6.34
	<b>3</b>	2.71 Å	-6.95
	<b>4</b>	2.81 Å	-4.88
	<b>5</b>	2.87 Å	-3.17

<sup>[a]</sup> Average of three C-H $\cdots\pi$  distances.

<sup>[b]</sup> Unbound (no minimum found)

The close correspondence between B3LYP-D3 and MP2 data in Table 1 is noteworthy. Because MP2 results tend to be much more basis-set dependent than DFT, we decided to perform extrapolations to the CBS limit for compounds **3** and **4** following a protocol by Helgaker *et al.*<sup>[7]</sup> This protocol involves single-point calculations with correlation-consistent basis sets (up to aug-cc-pVQZ; for details see ESI, Tables S2-S5 and Figure S1). It has recently been reported that while standard MP2 may overestimate weak intermolecular interactions relative to CCSD(T) benchmarks (e.g. for the benzene dimer),<sup>[8]</sup> the parametrised spin-component-scaled (SCS) variant<sup>[9]</sup> performs much better. We therefore performed both MP2/CBS and SCS-MP2/CBS extrapolations (Table 2).

**Table 2:** Binding energies (kcal mol<sup>-1</sup>) obtained at the HF, MP2 and SCS-MP2 *ab initio* methods with the complete basis set (CBS) for compounds **3** and **4**.

	<b>3</b>	<b>4</b>
HF/CBS	-0.13	+1.41
MP2/CBS <sup>[a]</sup>	-7.93	-5.75
SCS-MP2/CBS <sup>[a]</sup>	-6.39	-4.33

<sup>[a]</sup> Estimated uncertainty  $\pm 0.4$  kcal mol<sup>-1</sup> (see ESI for details).

At the MP2-optimised distance, the HF method shows essentially vanishing or even repulsive interactions for both

compounds **3** and **4** with benzene.<sup>[10]</sup> Predicted binding energies for **3** at MP2/CBS and SCS-MP2/CBS levels are ca. -8 kcal/mol and -6 kcal/mol, respectively, (ca. -6 kcal/mol and -4 kcal/mol, respectively, for **4**), bracketing the B3LYP-D3 values in Table 1, thus reinforcing the reliability of that level. The SCS-MP2 predicted binding energy of -6.4 kcal/mol for **3**·C<sub>6</sub>H<sub>6</sub> is, to our knowledge, the largest C-H... $\pi$  interaction energy between benzene and an aliphatic hydrocarbon, larger than that between benzene and chloroform.<sup>[11]</sup> In view of the dominance of dispersion discussed above one may argue that the overall interaction energy is not arising from three local C-H... $\pi$  interactions,<sup>[12]</sup> but should rather be attributed to the large molecular size of **3**. Irrespective of the individual contributions to the total interaction energy, however, topological analysis indicates the presence of three distinct bond paths between the axial H atoms of **3** and C atoms of benzene, and a weakly attractive noncovalent interaction between them (MP2/aug-cc-pVDZ density, see Figure S2 in the ESI).

<sup>1</sup>H NMR chemical shift ( $\delta$ ) values for compound **3** have been obtained theoretically at the BHandH/6-311+G(2d,p) level<sup>[13]</sup> and are compared to experimentally obtained data in Table 3 (More details in Tables S6-S8 in the ESI). Observed solvent shifts  $\Delta\delta$  on going from dichloromethane to benzene are modelled as the difference between pristine **3** and its complex with benzene. Irrespective of the source geometry, optimised at either B3LYP, B3LYP-D3 or MP2 levels, the computed trends are in qualitative agreement with experimental values (Table 3) accounting for roughly half of the observed upfield shift. In order to simulate the entire shielding effect exerted by the arene solvent, more solvent molecules would have to be included in a dynamic description. Calculations for a single benzene molecule placing "ghost atoms" at the positions of axial and equatorial H atoms in the complex with **3** indicate that the observed shifts are largely (but not exclusively) due to the anisotropy (ring current) effect in the aromatic solvent (Table S9 in the ESI).


**Table 3:** Theoretical chemical shift ( $\delta$ ) values obtained from BHandH/6-311+G(2d,p) calculations on B3LYP, B3LYP-D3 and MP2 optimised geometries with the def2-TZVP basis set for pristine compound **3** and its complex with benzene, as well as the experimental values in dichloromethane and benzene.

	$\delta(\text{Hax})$		$\delta(\text{Heq})$		$\Delta\delta(\text{Hax})$	$\Delta\delta(\text{Heq})$
	gas /CH <sub>2</sub> Cl <sub>2</sub>	C <sub>6</sub> H <sub>6</sub>	gas /CH <sub>2</sub> Cl <sub>2</sub>	C <sub>6</sub> H <sub>6</sub>		
<b>B3LYP</b>	4.90 <sup>[a]</sup>	3.79 <sup>[b]</sup>	6.10 <sup>[a]</sup>	5.45 <sup>[b]</sup>	-1.11	-0.65
<b>B3LYP-D3</b>	3.74 <sup>[a]</sup>	2.86 <sup>[b]</sup>	4.92 <sup>[a]</sup>	4.39 <sup>[b]</sup>	-0.88	-0.53
<b>MP2</b>	4.01 <sup>[a]</sup>	3.17 <sup>[b]</sup>	5.17 <sup>[a]</sup>	4.67 <sup>[b]</sup>	-0.84	-0.50
<b>Exp.</b>	4.53	2.88	5.32	4.41	-1.65	-0.91

[a] Pristine **3**. [b] **3**·C<sub>6</sub>H<sub>6</sub> complex

As observed previously for the benzene complexes of **1** and **2**, the results for the B3LYP geometry for **3** seem to fit better to the observed solvent shifts than those using the more optimal B3LYP-D3 structure. Apparently, the shorter CH... $\pi$  separations in the B3LYP-D3 minimum (Table 1) bring the axial H atoms out of the shielding cone. From the anharmonicity of the **3**·C<sub>6</sub>H<sub>6</sub> stretching potential (Figure 4a), thermal averaging might be expected to increase the intermolecular distance somewhat (approaching the B3LYP value), but from the small computed variation of the chemical shifts in the distance range of interest, 2.6 Å - 3.1 Å, little effect on the  $\Delta\delta$  values is expected upon thermal averaging over this single coordinate. Again, full dynamics and more solvent molecules would have to be included for quantitative modelling. In

any event, the upfield shifts of the axial H atoms for compound **3** in aromatic solvents are of similar magnitude (ca. -1.7 ppm) to those observed in the all-*cis* tetrafluoro derivatives **1** and **2**.<sup>[3a]</sup>



**Figure 4:** a) Binding energies of the **3**·C<sub>6</sub>H<sub>6</sub> complex vs C-H... $\pi$  distances. b) Dependence of the chemical shifts in the **3**·C<sub>6</sub>H<sub>6</sub> complexation vs the C-H... $\pi$  distance. Energies calculated at B3LYP-D3/def2-TZVP level and shielding tensors on BHandH/6-311+G(2d,p) levels. BSSE corrections included.

## Conclusions

In summary, we have presented the first high-level *ab initio* benchmark study (MP2/CBS and SCS-MP2/CBS) for the CH... $\pi$  interaction energies between fluorocyclohexanes and benzene. The interaction energies proved to be strong (ca. 6-8 kcalmol<sup>-1</sup> in total for **3**), and are well described by dispersion-corrected DFT functionals. The affinity toward aromatic solvents is reflected in notable changes in <sup>1</sup>H chemical shifts that are rationalised by way of ring current effects. This affinity could clearly be exploited in crystal engineering or liquid crystal design.

## Theoretical and Experimental Procedures

Geometries were fully optimised in C<sub>3v</sub> symmetry at the B3LYP/def2-TZVP, B3LYP-D3/def2-TZVP and MP2/aug-cc-pVDZ levels including BSSE corrections; single point energies for the MP2 geometries refined at MP2 and SCS-MP2 levels using auc-cc-pVxZ basis sets (X = D, T, Q) and extrapolated to the CBS limit.<sup>[7]</sup> <sup>1</sup>H NMR measurements were carried out on a Bruker Avance III 500 spectrometer, operating at 500 MHz, using the deuterated solvent as the reference for internal deuterium lock. See ESI for further details and references.

## Acknowledgements

We thank EaStCHEM and the University of St Andrews for access to a computing facility managed by Dr H. Fruchtl, CNPq and FAPESP for the studentship for R.A.C. (#2011/01170-1, FAPESP) and for financial support (#2014/25903-6), as well as CNPq for the fellowship for R.R. DO'H thanks the Royal Society for a Wolfson Research Merit Award.

## Notes and references

## COMMUNICATION

Journal Name

- 1 (a) D. O'Hagan, *Chem. Soc. Rev.*, 2008, **37**, 308; (b) P. A. Champagne, J. Desroches, J.-D. Hamel, M. Vandamme, J.-F. Paquin, *Chem. Rev.*, 2015 (In press; DOI: 10.1021/cr500706a).
- 2 (a) A. J. Durie, A. M. Z. Alexandra, T. Lebl, P. Kirsch, D. O'Hagan, *Chem. Commun.*, 2011, **47**, 8265; (b) A. J. Durie, A. M. Z. Slwein, T. Lebl, P. Kirsch, D. O'Hagan, *Chem. Commun.*, 2012, **48**, 9643; (c) A. J. Durie, T. Fujiwara, R. Cormanich, M. Bühl, A. M. Z. Slawin, D. O'Hagan, *Chem. – Eur. J.*, 2014, **20**, 6259.
- 3 (a) R. A. Cormanich, A. J. Durie, R. Bjornsson, R. Rittner, O'Hagan, D.; Bühl, M., *Helv. Chim. Acta.*, 2014, **97**, 797. (b) R. A. Cormanich, R. Rittner, D. O'Hagan, M. Bühl, *J. Phys. Chem. A*, 2014, **118**, 7901.
- 4 N. S. Keddie, A. M. Z. Slawin, T. Lebl, D. Philp, D. O'Hagan, *Nature Chem.*, 2015, **7**, 483.
- 5 (a) S. Grimme, S. Ehrlich, L. Goerigk *J. Comput. Chem.*, 2011, **32**, 1456. (b) S. Grimme, J. Antony, S. Ehrlich, H. Krieg *J. Chem. Phys.*, 2010, **132**, 154104.
- 6 M. W. Feyereisen, D. Feller, D. A. Dixon *J. Phys. Chem.*, 1996, **100**, 2993.
- 7 T. Helgaker, W. Klopper, H. Koch, J. Noga, *J. Chem. Phys.*, 1997, **106**, 9639.
- 8 E. Miliordos, E. Apra, S. S. Xantheas, *J. Phys. Chem. A*, 2014, **118**, 7658.
- 9 S. J. Grimme *Chem. Phys.*, 2003, **118**, 9095.
- 10 A small net binding should prevail at the HF level upon full optimisation, given the large dipole moments of the fluorocyclohexanes (calculated dipole moments for **3** and **4** are 5.15 D and 5.94 D, respectively, at B3LYP-D3/def2-TZVP).
- 11 -5.2 kcal/mol (expt.: [a] A. Fujii, K. Shibasaki, T. Kazama, R. Itaya, N. Mikami, S. Tsuzuki, *Phys. Chem. Chem. Phys.*, 2008, **10**, 2836); stronger individual C-H... $\pi$  interactions exceeding 6 kcal/mol are known, but these involve larger aromatic systems (Review: [b] M. Nishio, *Phys. Chem. Chem. Phys.*, 2011, **13**, 13873).
- 12 In line with observations that number of C–H groups of aliphatic hydrocarbons in contact with a benzene ring does not correlate with the magnitude of the interaction energy: A. Fujii, H. Hayashi, J. W. Park, T. Kazama, N. Mikamia, S. Tsuzuki, *Phys. Chem. Chem. Phys.*, 2011, **13**, 14131.
- 13 A. D. Becke, *J. Chem. Phys.*, 1993, **98**, 5648.