



The androgen receptor gene CAG repeat in relation to 4-year changes in androgen-sensitive endpoints in community-dwelling older European men

DOI:
[10.1530/EJE-16-0447](https://doi.org/10.1530/EJE-16-0447)

Document Version
Accepted author manuscript

[Link to publication record in Manchester Research Explorer](#)

Citation for published version (APA):

Eendebak, R. J. A. H., Huhtaniemi, I., Pye, S., Ahern, T., O'Neill, T. W., Bartfai, G., Casanueva, F., Maggi, M., Forti, G., Alston, R., Giwercman, A., Han, T. S., Kula, K., Lean, M. E. J., Punab, M., Pendleton, N., Keevil, B., Vanderschueren, D., Rutter, M., ... Wu, F. (2016). The androgen receptor gene CAG repeat in relation to 4-year changes in androgen-sensitive endpoints in community-dwelling older European men. *European Journal of Endocrinology*, 175, 583-593 . <https://doi.org/10.1530/EJE-16-0447>

Published in:
European Journal of Endocrinology

Citing this paper

Please note that where the full-text provided on Manchester Research Explorer is the Author Accepted Manuscript or Proof version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version.

General rights

Copyright and moral rights for the publications made accessible in the Research Explorer are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Takedown policy

If you believe that this document breaches copyright please refer to the University of Manchester's Takedown Procedures [<http://man.ac.uk/04Y6Bo>] or contact uml.scholarlycommunications@manchester.ac.uk providing relevant details, so we can investigate your claim.



1 **The androgen receptor gene CAG repeat in relation to 4-year changes in androgen-**
 2 **sensitive endpoints in community-dwelling older European men**

3 Robert J.A.H. Eendebak M.Sc. ^{‡1}, Ilpo T. Huhtaniemi M.D. Ph.D.², Stephen R. Pye Ph.D.³, Tomas Ahern MB
 4 ChB¹, Terence W. O'Neill M.D.³, György Bartfai M.D. Ph.D. D.S.⁴, Felipe F. Casanueva M.D. Ph.D.⁵, Mario Maggi
 5 M.D.⁶, Gianni Forti M.D.⁶, Robert D. Alston Ph.D.¹, Aleksander Giwercman M.D. Ph.D.⁷, Thang S. Han Ph.D.⁸,
 6 Krzysztof Kula M.D. Ph.D.⁹, Michael E. J. Lean M.D.¹⁰, Margus Punab M.D. Ph.D.¹¹, Neil Pendleton M.D.¹², Brian
 7 G. Keevil M.Sc.¹³, Dirk Vanderschueren M.D. Ph.D.¹⁴, Martin K. Rutter M.D.^{15,16}, Gindo Tampubolon Ph.D.¹⁷,
 8 Royston Goodacre Ph.D.¹⁸ and Frederick C.W. Wu M.D.¹, for the EMAS Group

9
 10 [‡]Correspondence and reprint requests can be directed to: Robert J.A.H. Eendebak M.Sc. Andrology Research
 11 Unit, Centre for Endocrinology and Diabetes, Institute of Human Development, Faculty of Medical and Human
 12 Sciences, University of Manchester (United Kingdom), Old St. Mary's building, Hathersage Road, M13 9WL, or to
 13 roberteendebak@gmail.com

14
 15 ¹University of Manchester, Manchester Academic Health Sciences Centre, Faculty of Medical and Human Sciences, Institute of
 16 Human Development, Centre for Endocrinology and Diabetes, Andrology Research Unit

17 ²Institute of Reproductive and Developmental Biology, Department of Surgery and Cancer, Imperial College London

18 ³Arthritis Research UK Centre for Epidemiology, Centre for Musculoskeletal Health, The University of Manchester, ,
 19 Manchester Academic Health Sciences Centre, Manchester, United Kingdom

20 ⁴Albert Szent György Medical University, Department of Obstetrics and Gynaecology & Andrology

21 ⁵University Santiago de Compostela, Department of Medicine

22 ⁶University of Florence, Department of Clinical Physiopathology, Andrology Unit

23 ⁷Lund University, Malmö University Hospital, Scania Andrology Centre, Department of Urology

24 ⁸University College London, Department of Endocrinology

25 ⁹Medical University Lodz, Department of Andrology and Reproductive Endocrinology

26 ¹⁰University of Glasgow, Department of Human Nutrition

27 ¹¹Tartu University Clinic, United Labs, Andrology Unit

28 ¹²University of Manchester, School of Community Based Medicine, Salford Royal NHS Trust

29 ¹³University South Manchester Hospital, Department of Clinical Biochemistry

30 ¹⁴Catholic University Leuven, Department of Andrology and Endocrinology

31 ¹⁵University of Manchester, Manchester Academic Health Sciences Centre, Central Manchester University Hospitals NHS
 32 Foundation Trust, Manchester Diabetes Centre

33 ¹⁶University of Manchester, Faculty of Medical and Human Sciences, Institute of Human Development, Endocrinology and
 34 Diabetes Research Group

35 ¹⁷University of Manchester, Cathie Marsh Institute for Social Research, Faculty of Humanities

36 ¹⁸University of Manchester, School of Chemistry, Manchester Institute for Biotechnology

37
 38 **European Journal of Endocrinology (EJE) manuscript reference number: EJE-16-0447**

39 **Title:** 19 words (maximum: 20 words)

40 **Short title:** AR CAG repeat and longitudinal changes in ASEs (46 characters including spaces;
 41 maximum number allowed)

42 **Total word count article (excluding abstract):** 4770 words (maximum approximately 4500 words
 43 including references)

44 **Key words:** androgen receptor, androgen action, gonadal axis, ageing, genetics

45
 46
 47
 48

49 **Abstract total: 243 words (maximum number of words: 250)**
50

51 **Context:** The *Androgen Receptor (AR)* gene exon 1 CAG repeat length has been proposed to be a
52 determinant of between-individual variations in androgen action in target tissues, which might regulate
53 phenotypic differences of human ageing. However, findings on its phenotypic effects are inconclusive.

54 **Objective:** To assess whether the *AR* CAG repeat length is associated with longitudinal changes in
55 endpoints which are influenced by testosterone (T) levels in middle-aged and elderly European men.

56 **Design:** Multinational European observational prospective cohort study

57 **Participants:** 1887 men (mean±sd age: 63±11 years; median follow-up: 4.3 years) from centres of 8
58 European countries comprised the analysis sample after exclusion of those with diagnosed diseases
59 of the hypothalamic-pituitary-testicular (HPT) axis.

60 **Main outcome measures:** Longitudinal associations between the *AR* CAG repeat and changes in
61 androgen-sensitive endpoints (ASEs) and medical conditions were assessed using regression
62 analysis adjusting for age and centre. The *AR* CAG repeat length was treated both as a continuous
63 and categorical (6-20; 21-23; 24-39 repeats) predictor. Additional analysis investigated whether results
64 were independent of baseline T or oestradiol (E2) levels.

65 **Results:** The *AR* CAG repeat, when used as a continuous or categorical predictor, was not
66 associated with longitudinal changes in ASEs or medical conditions after adjustments. These results
67 were independent of T and E2 levels.

68 **Conclusion:** Within a 4-year timeframe, variations in the *AR* CAG repeat do not contribute to the rate
69 of phenotypic ageing, over and above, that, which might be associated with the age-related decline in
70 T levels.

71

72

73

74 **Introduction**

75 The length of the *Androgen Receptor (AR)* tri-nucleotide CAG repeat in exon 1, encoding a poly-
76 glutamine tract, has been proposed to regulate androgen action in target tissue. An inverse
77 association between the *AR* CAG repeat length and androgen action may exist. The *AR* CAG repeat
78 might regulate androgen action in response to testosterone (T) and dihydrotestosterone (1-4) in target
79 tissues, and affect androgen-sensitive endpoints (ASEs), such as body composition and metabolic
80 parameters (leptin and insulin levels) (5), cardiovascular risk factors (HDL cholesterol and arterial
81 vasoreactivity) (6), bone density (7), and treatment response to T supplementation (8).

82 Previously, results from the Massachusetts Male Aging Study (MMAS) have indicated that shorter *AR*
83 CAG repeats are associated with a greater decline in T levels over time (9). Others have indicated that
84 the presence of either extreme short or long *AR* CAG repeat (<9 or ≥38) length is associated with
85 increased risk for prostate cancer and Kennedy's disease, respectively (4, 10, 11-14). In addition,
86 Nenonen and colleagues (15) reported a non-linear association between the *AR* CAG repeat length
87 and fertility status across 33 studies, whereby men with either <22 or >23 CAG repeats were at
88 increased risk of reduced fertility.

89 In contrast, Van Pottelbergh and co-workers did not observe an association between the *AR* CAG
90 repeat and androgen levels, androgen insensitivity index (LHxTT product) or bone markers within a
91 cross-sectional cohort study consisting of ambulatory elderly men (16). In addition, Bentmar-
92 Holgersson and co-workers (17) did not observe an association between the *AR* CAG repeat and PSA
93 levels or prostate cancer risk within cross-sectional data from the European Male Ageing Study
94 (EMAS). However, additional cross-sectional results from EMAS have led Huhtaniemi and colleagues
95 (18) to propose that the potential downstream consequences of longer *AR* CAG repeat length and the
96 concomitant decreased androgen action may be modified by compensatory increased oestradiol (E2)
97 levels. However, most previous studies have been cross-sectional in design or were performed within
98 single centres and hence do not allow for assessment of longitudinal changes and may have limited
99 external validity. The potential importance of androgen action in ageing men remains unclear. Clinical
100 features developing with ageing may at least in part be a consequence of the age-related decline in T
101 levels modified by variations in tissue response to androgens. Longitudinal cohort studies may provide
102 the opportunity to discern how genetic markers, such as the *AR* CAG repeat, are related to changes in
103 features of ageing, which are believed to be regulated by androgen action.

104 The aim of the current study was to assess whether the *AR CAG* repeat length was associated with
105 changes in ASEs, independent of circulating T or E2 levels, in community-dwelling middle-aged and
106 elderly European men. In addition, longitudinal associations between the *AR CAG* repeat length and
107 the development of medical conditions, common in the elderly, were assessed in a similar manner.

108 We hypothesized that the *AR CAG* repeat length is associated with longitudinal changes in some
109 ASEs that may contribute to the phenotype of ageing men.

110

111 **Methods**

112 **Participants and study design**

113 The European Male Ageing Study (EMAS), as described elsewhere (19-21), is a multi-centre,
114 prospective, population-based cohort study of the endocrine and metabolic determinants of male
115 ageing. The eight participating centres are: Florence (Italy), Leuven (Belgium), Lodz (Poland), Malmö
116 (Sweden), Manchester (United Kingdom), Santiago de Compostela (Spain), Szeged (Hungary), and
117 Tartu (Estonia). Ethics approval for the study was obtained in each centre according to local
118 requirements. The number of men recruited ranged from 396 to 451 per centre (total n=3369). DNA
119 extraction and *AR CAG* repeat analysis were carried out on 267 to 368 samples per centre (total
120 n=2659). The protocols used for blood processing and sampling, DNA extraction and determination of
121 *AR CAG* repeat length within EMAS have been described, previously (18). The protocols used for
122 assessment of body composition (lean and fat mass), ultrasound of the heel, blood pressure,
123 hematological, biochemical, lipid and carbohydrate metabolism, sexual, physical, psychological and
124 prostate function, vitality and cognitive function in EMAS have been described, previously (18) (19,
125 20). Follow-up assessment was performed a median of 4.3 years (95% CI: 4.23 – 4.36 years) after the
126 baseline assessment using the same protocols as the baseline assessment.

127

128 **Exclusion criteria**

129 As depicted in the flow chart (figure 1), participants (of the total n=3369) were excluded if they
130 reported treatment for pituitary, testicular or adrenal disorders and/or use of medication affecting
131 hypothalamic-pituitary-testicular (HPT) axis function at baseline (n=179) or follow-up (n=132).
132 Participants were also excluded if they died (n=168), were lost to follow-up (n=407), missing total T

133 data (n=78), if their genotyping failed quality control standards (n=177) or if missing *AR* CAG repeat
134 data (n=341) was recorded. This lead to an analytical sample size of 1887 men.

135

136 **Hormone assays**

137 T was measured by liquid chromatography–tandem mass spectrometry, with paired baseline and
138 follow-up samples analysed simultaneously (22). LH, FSH, and SHBG were measured by the E170
139 platform electrochemiluminescence immunoassay (Roche Diagnostics) (23). E2 was measured by
140 both radio-immunoassay (at both phases) and by mass spectrometry (at baseline). Free (F) T was
141 calculated using the Vermeulen formula (24). Intra- and interassay coefficients of variation (CVs) were:
142 T, 4.0% and 5.6%; SHBG, 1.7% and 3.2%; LH, 1.9% and 3.0%; FSH, 1.8% and 5.3%, and (radio-
143 immunoassay) E2 5.2% and 9.1% and (GC-MS) E2 3.5% and 3.7%, respectively. The detection limit
144 for the reproductive hormones were: total T (TT) [0.55 nmol/L or 0.16 µg/L], SHBG [8.80 nmol/L or
145 10.00 µg/L], LH [0.10 U/L], FSH [0.61 U/L], radio-immunoassay E2 [18.14 pmol/L or 4.94 ng/L] and
146 GC-MS E2 [9.91 pmol/L or 2.70 ng/L].

147

148 **Other measures**

149 Participants provided information on their self-rated general health (SF36 questionnaire) and were
150 asked if they were currently being treated for the following medical conditions: heart problems, stroke,
151 hypertension, diabetes, bronchitis, cancer, kidney or liver disease. The presence of heart problems,
152 stroke or hypertension was indicative of cardiovascular disease. The responses from the participants
153 were further classified as either 'none' or 'one or more' or 'two or more' reported comorbidities from the
154 eight chronic conditions. Self-reported poor health status was assessed using responses from
155 participants on the SF36 questionnaire concerning how the participants rated their overall general
156 health. Self-reported poor health status was considered if responses included 'fair' or 'poor'.

157

158 **Statistical analyses**

159 The relationship between the *AR* CAG repeat and outcomes (ASEs) was assessed using the *AR* CAG
160 repeat both as a continuous predictor, as well as a tertiled categorical [tertile1: 6-20 (n=581), tertile2:
161 21-23 (n=667) and tertile3: 24-39 (n=639) CAG repeats] predictor.

162 Outcomes such as changes in blood pressure, body composition, heel ultrasound, physical activity,
163 carbohydrate and lipid metabolism, cognitive processing speed (as measured via the DSST) and
164 biochemical parameters, as well as the international prostate symptom score (IPSS), prostate specific
165 antigen (PSA) and reproductive hormone levels were treated as continuous outcomes. In addition, in
166 order to assess the relationship between the *AR* CAG repeat and changes in sexual, physical and
167 psychological function, individual scores on the EMAS sexual function questionnaire, as well as the
168 SF36 and BDI, were used as continuous outcomes. Changes in ASEs, were defined by the absolute
169 change of an ASE (i.e. follow-up ASE value - baseline ASE value) adjusted for the baseline ASE
170 value. The relationship between the *AR* CAG repeat (predictor) and the development of medical
171 conditions or self-reported poor health status (outcome variables) was also assessed. The
172 development of a medical condition was defined as subjects who reported being treated for a medical
173 condition at follow-up who did not report having the condition at baseline. The development of self-
174 reported poor health status was defined in a similar manner.

175 Linear regression was used to determine the longitudinal associations between the *AR* CAG repeat
176 and each of the ASEs with results expressed as absolute differences (β -coefficients) and 95%
177 confidence intervals (CI). Logistic regression analysis was used to assess the relationships between
178 the *AR* CAG repeat and the development of medical conditions or self-reported poor health status
179 (binary outcomes) with results expressed as odds ratios and 95% CI. For both linear and logistic
180 regression analyses, adjustments were made for age, centre and baseline TT and (GC-MS) E2 levels.
181 The cut off value for statistical significance was set to $p < 0.01$, when using the *AR* CAG repeat as a
182 continuous linear predictor, in order to account for potential false-positive results, as used in our
183 baseline cross-sectional analysis (18). In order to account for the multiple comparisons performed
184 when using the *AR* CAG repeat as a tertiled predictor, a Bonferroni correction was applied, which
185 lowered the threshold for statistical significance to $p < 0.003$. Statistical thresholds of $p < 0.05$, $p < 0.01$
186 and $P < 0.003$ are included in each of the tables, but only statistical thresholds of $p < 0.01$ (tables 2a-2e)
187 and $p < 0.003$ (S2a-S2f) are deemed significant. All statistical analyses were performed using STATA
188 13 SE (<http://www.stata.com>).

189
190
191

192 **Results**193 **Characteristics of the study subjects at baseline and follow-up (including the distribution of the**
194 **AR CAG repeat length)**

195 Men with complete AR CAG repeat data were middle-aged, often overweight, with a relatively low
196 prevalence of comorbidity burden, and with reproductive hormone levels within the eugonadal range.
197 Most clinical endpoints changed over time in men with complete AR CAG repeat data except for
198 glucose, triglycerides, hemoglobin, heel bone mineral density (US-BMD), androgen insensitivity index
199 (LHxTT product), E2 levels, aromatase activity (E2:TT ratio), mental function (SF36 mental function),
200 inability to bend, sadness, and prevalence of prostate disease in unadjusted analysis (Table 1). The
201 distribution of the AR CAG repeat length approximated a normal distribution (data not shown) with
202 mean±SD = 22±3 CAG repeats and a range of 6 - 39 CAG repeats. The distribution of the AR CAG
203 repeat was similar in men with complete AR CAG repeat data, when compared to men who were
204 excluded (S1) indicating a low risk from selection bias.

205

206 **Changes in body composition, heel ultrasound, physical, prostate and cognitive function**

207 The AR CAG repeat was not associated with changes in body composition parameters, such as BMI,
208 waist circumference and mid-upper arm circumference (MUAC), and heel ultrasound parameters (US-
209 BMD, US-BUA, US-SOS) (Table 1a). The AR CAG repeat was not associated with changes in
210 physical activity (PASE) or physical performance (50 feet walk test and PPT-rating) scores. The AR
211 CAG repeat was not associated with changes in indices of prostate function, such as PSA levels or
212 IPSS scores. In addition, the AR CAG repeat was not associated with changes in cognitive processing
213 speed, as assessed via DSST scores (Table 2a). Adjustment for baseline TT or E2 levels did not
214 change the results obtained. Results were similar when using the AR CAG repeat as a tertiled
215 categorical predictor (S2a). However, when using a less stringent p-value threshold, the AR CAG
216 repeat was associated with changes in 50ft walking distance, limited walking, decreased vigorous
217 activity and IPSS-scores.

218

219

220

221

222 Changes in carbohydrate and lipid metabolism, blood pressure and hematological parameters

223 The *AR* CAG repeat was not associated with changes in fasting plasma glucose levels or a measure
224 of insulin resistance (HOMA-IR). In addition, the *AR* CAG repeat was not associated with changes in
225 total cholesterol, HDL-cholesterol, LDL-cholesterol or triglyceride levels.

226 The *AR* CAG repeat was not associated with changes in blood pressure or hemoglobin levels (Table
227 2a). Adjustment for baseline TT or E2 levels did not change the results obtained. Results were similar
228 when using the *AR* CAG repeat as a tertiled categorical predictor (S2a). However, when using a less
229 stringent p-value threshold, the *AR* CAG repeat was associated with changes in fasting glucose, HDL-
230 cholesterol and triglyceride levels.

231

232 Changes in reproductive hormone levels and phase 2 reproductive hormone levels

233 The *AR* CAG repeat was not associated with changes in either TT or FT levels (Table 2b). The *AR*
234 CAG repeat was not associated with changes in LH, FSH, TT:LH ratio or the LHxTT product.
235 However, the *AR* CAG repeat was positively associated with TT, FT and E2 levels, but not the E2:TT
236 ratio, at follow-up, in a cross-sectional manner (Table 2c). After adjustment for baseline E2 levels the
237 cross-sectional relationship between the *AR* CAG and TT levels became non-significant. Results were
238 similar when using the *AR* CAG repeat as a tertiled categorical predictor (S2b). However, when using
239 a less stringent p-value threshold, the *AR* CAG repeat was associated with changes in TT, FT, E2, LH
240 levels and the LHxTT product.

241

242 Changes in sexual, physical, psychological, mental and quality of life questionnaire scores

243 The *AR* CAG repeat was not associated with changes in sexual, physical or psychological function
244 questionnaire scores. In addition, the *AR* CAG repeat was not associated with changes in overall
245 sexual function (SFQ-OSF), overall physical function (SF36 physical function), psychological (BDI-
246 total) and mental function (SF36 mental function), and quality of life (SF36 vitality) scores (Table 2c).
247 Adjustment for baseline TT or E2 levels did not change the results obtained. Results were similar
248 when using the *AR* CAG repeat as a tertiled categorical predictor (S2c, S2d and S2e). However, when
249 using a less stringent p-value threshold, the *AR* CAG repeat was associated with changes in overall
250 sexual function, fatigue, mental function, and vitality scores.

251

252 **Changes in medical conditions**

253 The *AR* CAG repeat was not associated with the development of self-reported poor health status,
254 comorbidity or multi-morbidity burden, or any other medical conditions (Table 2e). Adjustment for
255 baseline TT or E2 levels did not change the results obtained. Results were similar when using the *AR*
256 CAG repeat as a tertiled categorical predictor (S2f). However, when using a less stringent p-value
257 threshold, the *AR* CAG repeat was associated with the development of poor health.

258

259 **Discussion**

260 The main finding from this longitudinal study was the lack of association between the *AR* CAG repeat
261 and changes in a wide variety of putative ASEs and medical conditions potentially important in the
262 phenotype of ageing in men. *AR* CAG repeat lengths, treated as a continuous variable (Tables 2a-2e)
263 or separated into tertiles (S2a-2f), showed similar results.

264

265 Our findings differed from those presented by Krithivas and colleagues (9). They reported that the *AR*
266 CAG repeat was associated with the magnitude of the longitudinal decline in T levels within the
267 MMAS, which we did not observe in our study. However, their study had a longer follow-up period
268 (approximately 8 vs. 4 years) than EMAS, but contained a smaller sample size than EMAS (n=1709
269 men vs. n=3369 men). Their study used the radio-immunoassay to measure T, which is known to have
270 a sub-optimal performance at low levels (11, 25). In their study, the relationship between reproductive
271 hormone levels and the *AR* CAG repeat was investigated per 3 *AR* CAG repeats, which may not
272 represent a clinically meaningful increase. Finally, in their study, quantification of the decline in T
273 levels over time in relation to the *AR* CAG repeat length was performed on pairing of just 4 individuals
274 based on identical baseline TT, age and waist to hip ratio.

275 Our findings agree with those from Trivison and colleagues (26), whom indicate that the *AR* CAG
276 repeat is not associated with changes in reproductive hormone levels within the MMAS. The study by
277 Trivison et al. (26) used similar methodology as Krithivas et al (9) and may suffer from similar
278 limitations. However, Trivison and colleagues studied the change in reproductive hormone levels
279 using the *AR* CAG repeat length as a continuous measure. Our study investigated the change in a
280 large number of endpoints in relation to the *AR* CAG repeat, assessed as either a continuous or
281 tertiled predictor, and might be more similar to the study by Trivison and colleagues.

282 Zitzmann and colleagues have proposed that the *AR* CAG repeat might be a putative biomarker for
283 'androgenicity' (1, 2, 10). Our results in men from the general population do not support this concept.
284 Our cohort consisted of community-dwelling middle-aged and elderly European men from the general
285 population, and the few subjects with diagnosed pituitary, testicular or adrenal disease were excluded
286 from the analysis. Men in the present study presumably have an intact HPT-axis, and thus the
287 potential consequences of any variations in the *AR* CAG repeat length are likely to be minimized or
288 rendered clinically insignificant by compensatory regulatory feedback changes involving gonadotropins
289 and E2, although no relationship between the *AR* CAG repeat length and longitudinal changes in
290 either could be observed. Our findings did not exclude the possibility that in men who have either
291 pituitary or testicular deficits, in whom the feedback regulation has been disrupted, the *AR* CAG repeat
292 may impact on the severity of symptoms associated with androgen deficiency or the response to T
293 replacement therapy.

294

295 The present cross-sectional results at follow-up confirmed our earlier finding at baseline (18) that
296 longer *AR* CAG repeat length was, associated with higher E2 levels. However, we did not observe that
297 the *AR* CAG repeat length was associated with longitudinal changes in E2 levels. Our longitudinal
298 results indicate that variations in *AR* CAG repeat length may not contribute to the phenotype of
299 ageing, over and above, that, which could be associated with the age-related decline in T levels. Our
300 findings have to be interpreted with caution, since a large number of endpoints were assessed in
301 relation to a single genetic marker. The relationship between the *AR* CAG repeat and changes in
302 ASEs was unclear prior to this study. Although we have reported all results, we are cautious in
303 interpreting associations, which are above our p-value thresholds ($p > 0.01$ and $p > 0.003$). We have
304 chosen a more stringent p-value threshold in line with recommendations proposed to account for
305 multiplicity (27). However, our findings do suggest that the effect of the *AR* CAG repeat on changes in
306 phenotypic endpoints in ageing men is small.

307

308 **Strengths and limitations**

309 EMAS is a multi-centre European longitudinal cohort study, which investigates the endocrine and
310 metabolic determinants of male ageing, such as alterations in androgenic and anabolic hormone
311 levels. The present analysis examined the temporal associations of within-subject differences in ASEs

312 and medical conditions in relation to variations in a genetic marker of androgen action, which should
313 not be influenced by reverse causality, since the *AR* CAG repeat length is fixed throughout life.
314 Both baseline and follow-up T levels from EMAS men were measured via liquid chromatography-
315 tandem mass spectrometry, which minimized any potential method-related variation (25). The EMAS
316 questionnaires related to sexual, physical and psychological function were carefully standardized and
317 translated into local languages in 8 centres (21, 28-32). Finally, men with missing *AR* CAG repeat data
318 showed minor differences in baseline FT and follow-up FT, follow-up insulin resistance and follow-up
319 vigorous activity scores, as compared to the analytical sample after age and centre adjustment (S2g).
320 Thus, potential bias due to missing *AR* CAG repeat data is likely to be minimal.
321 However, measurement of E2 levels by the radio-immunoassay should be considered a limitation, due
322 to its suboptimal performance at low hormone concentrations (33). Another limitation of the current
323 study is that it contains only middle-aged and elderly men of European origin. Thus, the findings might
324 not extend to younger men or individuals of a non-European background, since the *AR* CAG repeat
325 length is known to differ across ethnic groups (34, 35). The median duration to follow-up was 4.3
326 years, which may be too short to discern slower longitudinal changes associated with variation in the
327 *AR* CAG repeat length.

328

329 **Conclusions**

330 We demonstrate in community-dwelling middle-aged and elderly men of European origin that
331 variations in *AR* CAG repeat length are not associated longitudinally with short-term changes in ASEs
332 or the development of medical conditions. The *AR* CAG repeat as a genetic marker of androgen action
333 is unlikely to contribute to major changes in the phenotype of ageing men.

334

335 **Declaration of interest**

336 The authors declare no conflicts of interest.

337

338 **Funding**

339 The European Male Ageing Study (EMAS) is funded by the Commission of the European
340 Communities Fifth Framework Program "Quality of Life and Management of Living Resources" Grant
341 QLK6-CT-2001-00258 and facilitated by the Manchester Biomedical Research Centre and the NIHR

342 Greater Manchester: Clinical Research Network. Additional support was also provided by Arthritis
 343 Research UK Centre for Epidemiology and the National Institute for Health Research and the
 344 Manchester Biomedical Research Centre. The principal investigator of EMAS is Professor Frederick
 345 Wu, M.D. Andrology Research Unit, University of Manchester, Manchester, U.K.

346

347 **Author contributions**

348 RJAHE and FCWW wrote the analysis plan for the study. BGK provided lab measurements required to
 349 perform the analysis. RJAHE performed the analysis. ITH, SRP, TA, RDA, AG, NP, MKR, GT, RG and
 350 FCWW supervised the analysis. RJAHE wrote the paper. ITH, SRP, TA, TWO, GB, FFC, MM, RDA,
 351 AG, TSH, KK, MEJ, MP, NP, BGK, DV, MKR, GT, RG and FCWW supervised the writing process.

352

353 **Acknowledgements**

354 R.J.A.H.E. is grateful for support received from the Biotechnology and Biological Sciences Research
 355 Council Doctoral Training Partnership (BBSRC-DTP), as well as the Fundatie van de Vrijvrouwe van
 356 Renswoude and Scholten-Cordes scholarship foundations. The authors wish to thank the men who
 357 participated in the eight countries and the research/nursing staff in the eight centres: C Pott
 358 (Manchester), E Wouters (Leuven), M Nilsson (Malmö), M del Mar Fernandez (Santiago de
 359 Compostela), M Jedrzejowska (Łódź), H-M Tabo (Tartu), A Heredi (Szeged) for their data collection,
 360 and C Moseley (Manchester) for data entry and project co-ordination.

361

362 **References**


- 363 1. Zitzmann M. Mechanisms of disease: pharmacogenetics of testosterone therapy in
 364 hypogonadal men. *Nature Clinical Practice Urology* 2007 **4** 161-166.
- 365 2. Zitzmann M. The Role of the CAG Repeat Androgen Receptor Polymorphism in Andrology.
 366 *Advances in the Management of Testosterone Deficiency* 2009 **37** 52-61.
- 367 3. Gottlieb B, Trifiro M, Lombroso R & Pinsky L. The androgen receptor gene mutations
 368 database. *Nucleic Acids Research* 1997 **25** 158-162.
- 369 4. La Spada AR, Wilson EM, Lubahn DB, Harding AE & Fischbeck KH. Androgen receptor gene
 370 mutations in X-linked spinal and bulbar muscular atrophy. *Nature* 1991 **352** 77-79.
- 371 5. Zitzmann M, Gromoll J, von Eckardstein A & Nieschlag E. The CAG repeat polymorphism in
 372 the androgen receptor gene modulates body fat mass and serum concentrations of leptin and
 373 insulin in men. *Diabetologia* 2003 **46** 31-39.
- 374 6. Zitzmann M, Brune M, Kornmann B, Gromoll J, von Eckardstein S, von Eckardstein A &
 375 Nieschlag E. The CAG repeat polymorphism in the AR gene affects high density lipoprotein
 376 cholesterol and arterial vasoreactivity. *Journal of Clinical Endocrinology & Metabolism* 2001 **86**
 377 4867-4873.
- 378 7. Zitzmann M, Brune M, Kornmann B, Gromoll J, Junker R & Nieschlag E. The CAG repeat
 379 polymorphism in the androgen receptor gene affects bone density and bone metabolism in
 380 healthy males. *Clinical Endocrinology (Oxf)* 2001 **55** 649-657.

- 381 8. Zitzmann M & Nieschlag E. Androgen receptor gene CAG repeat length and body mass index
382 modulate the safety of long-term intramuscular testosterone undecanoate therapy in
383 hypogonadal men. *Journal of Clinical Endocrinology & Metabolism* 2007 **92** 3844-3853.
- 384 9. Krithivas K, Yurgalevitch SM, Mohr BA, Wilcox CJ, Batter SJ, Brown M, Longcope C,
385 McKinlay JB & Kantoff PW. Evidence that the CAG repeat in the androgen receptor gene is
386 associated with the age-related decline in serum androgen levels in men. *Journal of*
387 *Endocrinology* 1999 **162** 137-142.
- 388 10. Zitzmann M & Nieschlag E. The CAG repeat polymorphism within the androgen receptor gene
389 and maleness. *International Journal of Andrology* 2003 **26** 76-83.
- 390 11. Hsing AW, Stanczyk FZ, Belanger A, Schroeder P, Chang L, Falk RT & Fears TR.
391 Reproducibility of serum sex steroid assays in men by RIA and mass spectrometry. *Cancer*
392 *Epidemiology Biomarkers & Prevention* 2007 **16** 1004-1008.
- 393 12. Hakimi JM, Schoenberg MP, Rondinelli RH, Piantadosi S & Barrack ER. Androgen receptor
394 variants with short glutamine or glycine repeats may identify unique subpopulations of men
395 with prostate cancer. *Clinical Cancer Research* 1997 **3** 1599-1608.
- 396 13. Ingles SA, Ross RK, Yu MC, Irvine RA, LaPera G, Haile RW & Coetzee GA. Association of
397 prostate cancer risk with genetic polymorphisms in vitamin D receptor and androgen receptor.
398 *Journal of the National Cancer Institute* 1997 **89** 166-170.
- 399 14. Heinlein CA & Chang C. Androgen receptor in prostate cancer. *Endocrine Reviews* 2004 **25**
400 276-308.
- 401 15. Nenonen HA, Giwercman A, Hallengren E & Giwercman YL. Non-linear association between
402 androgen receptor CAG repeat length and risk of male subfertility - a meta-analysis.
403 *International Journal of Andrology* 2011 **34** 327-332.
- 404 16. Van Pottelbergh I, Lumbroso S, Goemaere S, Sultan C & Kaufman JM. Lack of influence of
405 the androgen receptor gene CAG-repeat polymorphism on sex steroid status and bone
406 metabolism in elderly men. *Clinical Endocrinology (Oxf)* 2001 **55** 659-666.
- 407 17. Holgersson MB, Giwercman A, Bjartell A, Wu FCW, Huhtaniemi IT, O'Neill TW, Pendleton N,
408 Vanderschueren D, Lean MEJ, Han TS, et al. Androgen Receptor Polymorphism-Dependent
409 Variation in Prostate-Specific Antigen Concentrations of European Men. *Cancer Epidemiology*
410 *Biomarkers & Prevention* 2014 **23** 2048-2056.
- 411 18. Huhtaniemi IT, Pye SR, Limer KL, Thomson W, O'Neill TW, Platt H, Payne D, John SL, Jiang
412 M, Boonen S, et al. Increased Estrogen Rather Than Decreased Androgen Action Is
413 Associated with Longer Androgen Receptor CAG Repeats. *Journal of Clinical Endocrinology &*
414 *Metabolism* 2009 **94** 277-284.
- 415 19. Lee DM, O'Neill TW, Pye SR, Silman AJ, Finn JD, Pendleton N, Tajar A, Bartfai G,
416 Casanueva F, Forti G, et al. The European Male Ageing Study (EMAS): design, methods and
417 recruitment. *International Journal of Andrology* 2009 **32** 11-24.
- 418 20. Lee DM, Pye SR, Tajar A, O'Neill TW, Finn JD, Boonen S, Bartfai G, Casanueva FF, Forti G,
419 Giwercman A, et al. Cohort profile: the European Male Ageing Study. *International Journal of*
420 *Epidemiology* 2013 **42** 391-401.
- 421 21. O'Connor DB, Corona G, Forti G, Tajar A, Lee DM, Finn JD, Bartfai G, Boonen S, Casanueva
422 FF, Giwercman A, et al. Assessment of sexual health in aging men in Europe: Development
423 and validation of the European Male Ageing Study sexual function questionnaire. *Journal of*
424 *Sexual Medicine* 2008 **5** 1374-1385.
- 425 22. Gallagher LM, Owen LJ & Keevil BG. Simultaneous determination of androstenedione and
426 testosterone in human serum by liquid chromatography-tandem mass spectrometry. *Annals of*
427 *Clinical Biochemistry* 2007 **44** 48-56.
- 428 23. Wu FC, Tajar A, Pye SR, Silman AJ, Finn JD, O'Neill TW, Bartfai G, Casanueva F, Forti G,
429 Giwercman A, et al. Hypothalamic-pituitary-testicular axis disruptions in older men are
430 differentially linked to age and modifiable risk factors: the European Male Aging Study. *Journal*
431 *of Clinical Endocrinology & Metabolism* 2008 **93** 2737-2745.
- 432 24. Vermeulen A, Verdonck L & Kaufman JM. A critical evaluation of simple methods for the
433 estimation of free testosterone in serum. *Journal of Clinical Endocrinology & Metabolism* 1999
434 **84** 3666-3672.
- 435 25. Huhtaniemi IT, Tajar A, Lee DM, O'Neill TW, Finn JD, Bartfai G, Boonen S, Casanueva FF,
436 Giwercman A, Han TS, et al. Comparison of serum testosterone and estradiol measurements
437 in 3174 European men using platform immunoassay and mass spectrometry; relevance for
438 the diagnostics in aging men. *European Journal of Endocrinology* 2012 **166** 983-991.
- 439 26. Travison TG, Shackelton R, Araujo AB, Morley JE, Williams RE, Clark RV, McKinlay JB.
440 Frailty, Serum Androgens, and the CAG repeat Polymorphism: Results from the

- 441 Massachusetts Male Aging Study. *Journal of Clinical Endocrinology & Metabolism* 2010 **95**
 442 2746-2754.
- 443 27. Streiner DL & Norman GR. Correction for Multiple Testing Is There a Resolution? *Chest* 2011
 444 **140** 16-18.
- 445 28. Corona G, Lee DM, Forti G, O'Connor DB, Maggi M, O'Neill TW, Pendleton N, Bartfai G,
 446 Boonen S, Casanueva FF, et al. Age-Related Changes in General and Sexual Health in
 447 Middle-Aged and Older Men: Results from the European Male Ageing Study (EMAS). *Journal*
 448 *of Sexual Medicine* 2010 **7** 1362-1380.
- 449 29. Tajar A, Forti G, O'Neill TW, Lee DM, Silman AJ, Finn JD, Bartfai G, Boonen S, Casanueva
 450 FF, Giwercman A, et al. Characteristics of Secondary, Primary, and Compensated
 451 Hypogonadism in Aging Men: Evidence from the European Male Ageing Study. *Journal of*
 452 *Clinical Endocrinology & Metabolism* 2010 **95** 1810-1818.
- 453 30. Tajar A, Huhtaniemi IT, O'Neill TW, Finn JD, Pye SR, Lee DM, Bartfai G, Boonen S,
 454 Casanueva FF, Forti G, et al. Characteristics of Androgen Deficiency in Late-Onset
 455 Hypogonadism: Results from the European Male Aging Study (EMAS). *Journal of Clinical*
 456 *Endocrinology & Metabolism* 2012 **97** 1508-1516.
- 457 31. Wu FC, Tajar A, Beynon JM, Pye SR, Silman AJ, Finn JD, O'Neill TW, Bartfai G, Casanueva
 458 FF, Forti G, et al. Identification of late-onset hypogonadism in middle-aged and elderly men.
 459 *New England Journal of Medicine* 2010 **363** 123-135.
- 460 32. Lee DM, Tajar A, O'Neill TW, O'Connor DB, Bartfai G, Boonen S, Bouillon R, Casanueva FF,
 461 Finn JD, Forti G, et al. Lower vitamin D levels are associated with depression among
 462 community-dwelling European men. *Journal of Psychopharmacology* 2011 **25** 1320-1328.
- 463 33. Taylor AE, Keevil B, Huhtaniemi IT. Mass spectrometry and immunoassay: how to measure
 464 steroid hormones today and tomorrow. *European Journal of Endocrinology* 2015 **173** D1-D12
- 465 34. Ackerman CM, Lowe LP, Lee H, Hayes MG, Dyer AR, Metzger BE, Lowe WL, Urbanek M.
 466 Ethnic Variation in Allele Distribution of the Androgen Receptor (AR) (CAG)(n) Repeat.
 467 *Journal of Andrology* 2012 **33** 210-215.
- 468 35. Nelson KA & Witte JS. Androgen receptor CAG repeats and prostate cancer. *American*
 469 *Journal of Epidemiology* 2002 **155** 883-890.
- 470

471

1 **Figure 1, flow chart**



2
3
4

Table 1. Baseline and follow-up candidate ageing-related parameters of men with complete AR CAG repeat data

Parameter	Baseline	Follow-up
	Men with AR CAG repeat data present (n=1887)	Men with AR CAG repeat data present (n=1887)
Study age, (years)	58.3 ±10.5	62.7 ±10.5***
Systolic blood pressure, (mmHg)	144.4 ±19.8	146.2 ±19.7***
Diastolic blood pressure, (mmHg)	87.0 ±11.8	84.6 ±11.6***
BMI, (kg/m ²)	27.6 ±3.9	27.8 ±4.2***
Waist circumference, (cm)	98.0 ±10.6	99.5 ±11.3***
MUAC (cm)	27.8 ±2.6	27.2 ±2.7***
PASE	205.0 ±89.1	181.4 ±96.0***
50ft walk, (sec)	13.1 ±2.5	14.0 ±3.7***
PPT-rating	24.3 ±2.4	23.7 ±2.5***
Fasting plasma glucose, (mmol/L)	5.6 ±1.2	5.5 ±1.3
HOMA-IR	3.1 ±4.1	3.0 ±2.9**
Total cholesterol, (mmol/L)	5.6 ±1.0	5.2 ±1.1***
HDL-cholesterol, (mmol/L)	1.4 ±0.4	1.4 ±0.4***
LDL-cholesterol, (mmol/L)	3.5 ±0.9	3.2 ±1.0***
Triglycerides, (mmol/L)	1.6 ±1.2	1.5 ±2.0
PSA, (ng/mL)	1.6 ±2.6	2.1 ±6.5**
IPSS scores	5.2 ±5.7	6.3 ±6.2***
Hb, (g/L)	150.3 ±10.4	149.8 ±11.6
DSST	28.9 ±8.4	27.8 ±9.0***
US BMD, (g/cm ²)	0.6 ±0.1	0.9 ±14.9
US BUA, (dBIMHz/cm)	81.0 ±19.0	83.1 ±18.2***
US SOS, (kHz)	1552.6 ±34.2	1550.8 ±32.9***
Testosterone, (nmol/L)	17.0 ±5.9	16.6 ±6.0***
Free testosterone, (pmol/L)	305.1 ±85.7	289.6 ±86.7***
SHBG, (nmol/L)	41.6 ±18.4	44.1 ±19.7***
LH, (U/L)	5.8 ±3.8	6.2 ±4.5***
FSH, (U/L)	7.8 ±7.5	8.2 ±8.6***
TT:LH ratio	3.6 ±1.9	3.5 ±2.0***
LHxTT product	101.2 ±76.5	103.3 ±76.9
Oestradiol (pmol/L; radio-immunoassay)	91.3 ±27.9	90.6 ±35.2
Oestradiol (pmol/L; GC-MS)	73.6 ±24.9	---
E2:TT ratio	6.0 ±4.1	6.1 ±3.9
Overall sexual function	21.0 ±6.5	21.1 ±6.9***
SF36 physical function	51.1 ±7.5	50.5 ±8.1**
BDI total	6.5 ±6.1	6.2 ±6.4*
SF36 mental function	52.2 ±8.7	52.1 ±9.0
SF36 vitality	15.2 ±2.8	15.0 ±2.9*
Morning erection frequency score	3.5 ±1.9	3.4 ±1.9*
Frequency of sexual thoughts score	5.2 ±2.0	4.8 ±2.1***
Erectile function score	1.9 ±1.0	2.1 ±1.0***
Vigorous activity score	2.2 ±0.7	2.1 ±0.8**
Limited walking score	2.9 ±0.4	2.8 ±0.5***
Unable to bend score	2.7 ±0.6	2.6 ±0.6
Sadness score	4.2 ±0.9	4.3 ±0.9
Loss of energy score	0.6 ±0.6	0.6 ±0.6*
Fatigue score	0.5 ±0.6	0.5 ±0.6*
Poor health, n(%)	366 (19.6)	429 (23.7)***
≥1 illnesses, n(%)	763 (40.4)	1062 (56.3)***
≥2 illnesses, n(%)	290 (20.5)	586 (41.5)***
Diabetes, n(%)	115 (6.2)	154 (8.4)***
CVD, n(%)	596 (32.0)	781 (44.2)***
Cancer, n(%)	85 (4.5)	148 (8.3)***
Prostate disease, n(%)	27 (8.0)	170 (9.3)

Data are expressed as unadjusted mean ±SD for continuous variables and number (percentages) for categorical variables.

Abbreviations: CardioVascular Disease; BDI, Beck Depression Inventory score; SF-36, medical outcome study short form 36 item questionnaire; BMI, Body Mass Index; WC, waist circumference; MUAC, Mid Upper Arm Circumference; PASE, Physical Activity Scale for the Elderly; PPT, Physical Performance Test; DSST, Digital Symbol Substitution Test; FPG, Fasting Plasma Glucose concentration; HOMA-IR, HOmeostatic Model of Insulin Resistance; PSA, serum Prostate Specific Antigen concentration; IPSS, international prostate symptom score; Hb, plasma HaemogloBin concentration; SOS, Speed of Sound; BUA, Broadband Ultrasound Attenuation. BMD, Bone Mineral Density.

Longitudinal (unadjusted) within-group differences:

* = p < 0.05 as assessed by paired T-test for continuous variables or McNemar test for binary variables

** = p < 0.01 as assessed by paired T-test for continuous variables or McNemar test for binary variables

*** = p < 0.001 as assessed by paired T-test for continuous variables or McNemar test for binary variables

Table 2a. Longitudinal changes in candidate androgen-sensitive parameters associated with the number of CAG repeats in the *Androgen Receptor* (linear regression)

Parameter (difference)	N	Model 1 Unadjusted	Model 2 Adjusted for age and centre	Model 3 Adjusted for age, centre and baseline total testosterone	Model 4 Adjusted for age, centre and baseline oestradiol
Systolic blood pressure, (mmHg)	1836	-0.02 (-0.06, 0.02)	-0.02 (-0.06, 0.02)	-0.02 (-0.06, 0.02)	-0.02 (-0.06, 0.02)
Diastolic blood pressure, (mmHg)	1835	-0.03 (-0.07, 0.01)	-0.03 (-0.07, 0.01)	-0.03 (-0.07, 0.01)	-0.03 (-0.07, 0.01)
BMI, (kg/m ²)	1812	0.00 (-0.04, 0.05)	0.00 (-0.04, 0.05)	0.00 (-0.04, 0.05)	0.01 (-0.04, 0.05)
Waist circumference, (cm)	1830	-0.03 (-0.08, 0.02)	-0.03 (-0.08, 0.01)	-0.03 (-0.08, 0.01)	-0.03 (-0.07, 0.02)
MUAC, (cm)	1827	-0.01 (-0.06, 0.03)	-0.01 (-0.06, 0.03)	-0.01 (-0.06, 0.03)	-0.02 (-0.06, 0.02)
PASE	1584	-0.00 (-0.05, 0.04)	-0.00 (-0.05, 0.04)	-0.00 (-0.05, 0.04)	-0.00 (-0.05, 0.04)
50ft walk, (sec)	1807	0.04 (-0.01, 0.08)	0.04 (-0.01, 0.08)	0.04 (0.00, 0.09)^a	0.04 (-0.00, 0.08)
PPT-rating	1735	-0.01 (-0.06, 0.03)	-0.02 (-0.06, 0.02)	-0.02 (-0.06, 0.02)	-0.02 (-0.06, 0.02)
Fasting plasma glucose, (mmol/L)	1772	0.02 (-0.02, 0.06)	0.02 (-0.02, 0.06)	0.02 (-0.02, 0.07)	0.02 (-0.02, 0.06)
HOMA-IR	1557	0.01 (-0.03, 0.04)	0.01 (-0.03, 0.04)	0.01 (-0.02, 0.05)	0.00 (-0.03, 0.04)
Total cholesterol, (mmol/L)	1771	0.00 (-0.04, 0.04)	0.00 (-0.04, 0.04)	-0.00 (-0.04, 0.04)	0.01 (-0.04, 0.05)
HDL-cholesterol, (mmol/L)	1763	-0.01 (-0.06, 0.03)	-0.02 (-0.06, 0.03)	-0.02 (-0.06, 0.02)	-0.02 (-0.06, 0.03)
LDL-cholesterol, (mmol/L)	1696	0.02 (-0.02, 0.06)	0.02 (-0.02, 0.06)	0.02 (-0.02, 0.06)	0.02 (-0.02, 0.06)
Triglycerides, (mmol/L)	1773	-0.03 (-0.08, 0.01)	-0.03 (-0.08, 0.01)	-0.03 (-0.07, 0.01)	-0.03 (-0.07, 0.02)
PSA, (ng/mL)	1475	0.01 (-0.04, 0.06)	0.01 (-0.04, 0.06)	0.01 (-0.04, 0.06)	0.01 (-0.04, 0.06)
IPSS scores	1719	-0.04 (-0.09, 0.00)	-0.04 (-0.09, 0.00)	-0.04 (-0.09, 0.00)	-0.05 (-0.09, -0.00)^a
Hb, (g/L)	1539	0.00 (-0.04, 0.05)	0.00 (-0.04, 0.05)	-0.00 (-0.05, 0.05)	-0.01 (-0.05, 0.04)
DSST	1796	-0.02 (-0.07, 0.02)	-0.03 (-0.07, 0.02)	-0.03 (-0.07, 0.02)	-0.02 (-0.07, 0.02)
US BMD, (g/cm ²)	1773	-0.02 (-0.07, 0.02)	-0.02 (-0.07, 0.02)	-0.02 (-0.07, 0.02)	-0.02 (-0.07, 0.02)
US BUA, (dBIMHz/cm)	1742	0.02 (-0.02, 0.07)	0.02 (-0.02, 0.07)	0.03 (-0.02, 0.07)	0.03 (-0.02, 0.07)
US SOS, (kHz)	1739	0.02 (-0.03, 0.06)	0.02 (-0.03, 0.06)	0.02 (-0.02, 0.07)	0.02 (-0.02, 0.07)

Data are expressed as standardized beta regression coefficients (95% confidence intervals).

Abbreviations: BDI, Beck Depression Inventory score; SF-36, medical outcome study short form 36 item questionnaire; BMI, Body Mass Index; WC, waist circumference; MUAC, Mid Upper Arm Circumference; PASE, Physical Activity Scale for the Elderly; PPT, Physical Performance Test; DSST, Digital Symbol Substitution Test; FPG, Fasting Plasma Glucose concentration; HOMA-IR, HOmeostatic Model of Insulin Resistance; HDL-cholesterol, High Density Lipid cholesterol; LDL-cholesterol, Low Density Lipid cholesterol; PSA, serum Prostate Specific Antigen concentration; IPSS, international prostate symptom score; Hb, plasma HaemogloBin concentration; SOS, Speed of Sound; BUA, Broadband Ultrasound Attenuation. BMD, Bone Mineral Density.

a = p<0.05, b = p<0.01 (all p>0.01)

Table 2b. Longitudinal changes in reproductive hormone levels associated with the number of CAG repeats in the *Androgen Receptor* (linear regression)

Parameter (difference)	N	Model 1 Unadjusted	Model 2 Adjusted for age and centre	Model 3 Adjusted for age, centre and baseline total testosterone	Model 4 Adjusted for age, centre and baseline oestradiol
Total testosterone, (nmol/L)	1887	0.02 (-0.02, 0.07)	0.02 (-0.02, 0.07)	---	0.02 (-0.02, 0.07)
Free testosterone, (pmol/L)	1866	0.03 (-0.01, 0.07)	0.03 (-0.01, 0.07)	---	0.03 (-0.01, 0.07)
SHBG, (nmol/L)	1866	0.01 (-0.03, 0.06)	0.01 (-0.03, 0.06)	0.01 (-0.04, 0.05)	0.01 (-0.03, 0.06)
LH, (U/L)	1864	-0.02 (-0.07, 0.02)	-0.02 (-0.06, 0.03)	-0.02 (-0.06, 0.03)	-0.02 (-0.07, 0.02)
FSH, (U/L)	1865	-0.02 (-0.06, 0.03)	-0.02 (-0.06, 0.03)	-0.02 (-0.06, 0.03)	-0.02 (-0.07, 0.02)
E2, (pmol/L; radio-immunoassay)	1859	0.02 (-0.02, 0.06)	0.02 (-0.02, 0.07)	0.02 (-0.03, 0.06)	---
E2:TT ratio	1859	0.02 (-0.02, 0.06)	0.02 (-0.02, 0.06)	0.02 (-0.02, 0.06)	0.02 (-0.03, 0.06)
TT:LH ratio	1864	0.00 (-0.04, 0.05)	0.01 (-0.04, 0.05)	0.00 (-0.04, 0.04)	0.01 (-0.03, 0.05)
LHxTT product	1864	0.01 (-0.03, 0.05)	0.01 (-0.03, 0.05)	0.01 (-0.03, 0.05)	0.01 (-0.04, 0.05)

Data are expressed as standardized beta regression coefficients (95% confidence intervals).

a = p<0.05, b = p<0.01 (all p>0.01)

Table 2c. Reproductive hormone levels at follow-up associated with the number of CAG repeats in the *Androgen Receptor* (linear regression)

Parameter (Phase 2)	N	Model 1 Unadjusted	Model 2 Adjusted for age and centre	Model 3 Adjusted for age, centre and baseline total testosterone	Model 4 Adjusted for age, centre and baseline oestradiol
Total testosterone, (nmol/L)	1887	0.07 (0.02, 0.11)^b	0.07 (0.02, 0.11)^b	---	0.04 (-0.00, 0.08)
Free testosterone, (pmol/L)	1871	0.08 (0.04, 0.13)^c	0.08 (-0.04, 0.12)^c	---	0.06 (0.02, 0.10)^b
SHBG, (nmol/L)	1871	0.00 (-0.04, 0.05)	0.01 (-0.03, 0.05)	-0.03 (-0.06, 0.00)	-0.01 (-0.05, 0.03)
LH, (U/L)	1871	-0.01 (-0.06, 0.03)	-0.01 (-0.05, 0.03)	-0.01 (-0.06, 0.03)	-0.02 (-0.06, 0.03)
FSH, (U/L)	1871	-0.04 (-0.08, 0.01)	-0.03 (-0.08, 0.01)	-0.03 (-0.07, 0.01)	-0.03 (-0.08, 0.01)
E2, (pmol/L; radio-immunoassay)	1865	0.07 (0.03, 0.12)^b	0.07 (0.03, 0.12)^b	0.06 (0.01, 0.10)^a	---
E2:TT ratio	1865	0.02 (-0.03, 0.06)	0.02 (-0.03, 0.06)	0.04 (-0.00, 0.08)	0.01 (-0.03, 0.06)
TT:LH ratio	1871	0.04 (-0.00, 0.09)	0.04 (-0.00, 0.08)	0.02 (-0.02, 0.06)	0.03 (-0.01, 0.08)
LHxTT product	1871	0.03 (-0.02, 0.07)	0.03 (-0.01, 0.07)	0.00 (-0.04, 0.04)	0.01 (-0.03, 0.05)

Data are expressed as standardized beta regression coefficients (95% confidence intervals)

a = p<0.05, b = p<0.01

Table 2d. Longitudinal changes in sexual, physical, psychological, mental and quality of life questionnaire scores associated with the number of CAG repeats in the *Androgen Receptor (AR)* (linear regression)

Parameter (difference)	N	Model 1 Unadjusted	Model 2 Adjusted for age and centre	Model 3 Adjusted for age, centre and baseline total testosterone	Model 4 Adjusted for age, centre and baseline oestradiol
Morning erection scores	1704	-0.02 (-0.07, 0.02)	-0.02 (-0.06, 0.02)	-0.02 (-0.06, 0.02)	-0.02 (-0.06, 0.02)
Sexual thoughts scores	1715	-0.02 (-0.06, 0.02)	-0.02 (-0.06, 0.02)	-0.02 (-0.06, 0.02)	-0.02 (-0.06, 0.02)
Erectile function scores	1652	-0.00 (-0.05, 0.04)	-0.00 (-0.05, 0.04)	-0.00 (-0.05, 0.04)	-0.01 (-0.05, 0.04)
SFQ-OSF scores	1167	-0.02 (-0.08, 0.03)	-0.01 (-0.07, 0.04)	-0.02 (-0.07, 0.03)	-0.01 (-0.07, 0.04)
Vigorous activity score	1797	-0.00 (-0.04, 0.04)	-0.00 (-0.04, 0.04)	-0.00 (-0.04, 0.04)	-0.00 (-0.04, 0.04)
Limited walking score	1781	-0.02 (-0.07, 0.02)	-0.02 (-0.06, 0.02)	-0.02 (-0.07, 0.02)	-0.02 (-0.06, 0.02)
Unable to bend	1794	-0.02 (-0.06, 0.02)	-0.01 (-0.05, 0.02)	-0.01 (-0.05, 0.03)	-0.01 (-0.05, 0.03)
SF36 physical function	1721	-0.01 (-0.05, 0.03)	-0.01 (-0.05, 0.03)	-0.01 (-0.05, 0.03)	-0.01 (-0.05, 0.03)
Sadness score	1773	0.02 (-0.02, 0.06)	0.02 (-0.02, 0.06)	0.02 (-0.02, 0.06)	0.02 (-0.02, 0.06)
Loss of energy score	1814	-0.01 (-0.05, 0.03)	-0.01 (-0.05, 0.03)	-0.01 (-0.05, 0.03)	-0.02 (-0.06, 0.02)
Fatigue score	1816	-0.01 (-0.05, 0.03)	-0.01 (-0.05, 0.03)	-0.01 (-0.05, 0.03)	-0.01 (-0.05, 0.03)
BDI-total	1793	-0.03 (-0.07, 0.01)	-0.03 (-0.07, 0.01)	-0.03 (-0.07, 0.01)	-0.03 (-0.07, 0.01)
SF36 mental function	1720	0.03 (-0.01, 0.08)	0.04 (-0.01, 0.08)	0.04 (-0.01, 0.08)	0.04 (-0.01, 0.08)
SF36 vitality	1792	0.02 (-0.03, 0.06)	0.02 (-0.03, 0.06)	0.01 (-0.03, 0.06)	0.02 (-0.02, 0.06)

Data are expressed as standardized beta regression coefficients (95% confidence interval).

a = p<0.05, b = p<0.01 (all p>0.01)

Table 2e. Development of medical conditions associated with the number of CAG repeats in the *Androgen Receptor* (logistic regression)

Parameter (development)	N	Model 1 Unadjusted	Model 2 Adjusted for age and centre	Model 3 Adjusted for age, centre and baseline total testosterone	Model 4 Adjusted for age, centre and baseline oestradiol
Poor Health	1445	1.05 (1.00, 1.11)^a	1.05 (1.00, 1.11)^a	1.06 (1.00, 1.11)^a	1.05 (1.00, 1.11)^a
≥1 illness	1124	0.99 (0.95, 1.03)	0.99 (0.95, 1.04)	1.00 (0.95, 1.04)	1.00 (0.96, 1.04)
≥2 illness	939	1.00 (0.95, 1.06)	1.00 (0.95, 1.06)	1.01 (0.95, 1.06)	1.01 (0.96, 1.07)
Diabetes	1687	0.97 (0.89, 1.06)	0.97 (0.89, 1.06)	0.99 (0.90, 1.08)	0.98 (0.89, 1.07)
Cardiovascular disease	1172	1.01 (0.96, 1.06)	1.01 (0.97, 1.06)	1.02 (0.97, 1.07)	1.01 (0.97, 1.06)
Cancer	1703	1.04 (0.97, 1.13)	1.05 (0.97, 1.13)	1.04 (0.97, 1.12)	1.05 (0.97, 1.13)
Prostate disease	1646	1.01 (0.95, 1.08)	1.01 (0.95, 1.08)	1.02 (0.95, 1.09)	1.02 (0.95, 1.09)

Data are expressed as odds ratios (95% confidence intervals).

a = p<0.05, b = p<0.01 (all p>0.01)