



Tummeltshammer, K. and Amso, D. and French, R. and Kirkham, Natasha (2016) Across space and time: infants learn from backward and forward visual statistics. *Developmental Science* , ISSN 1363-755x. (In Press)

Downloaded from: <http://eprints.bbk.ac.uk/15047/>

Usage Guidelines

Please refer to usage guidelines at <http://eprints.bbk.ac.uk/policies.html> or alternatively contact lib-eprints@bbk.ac.uk.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Running Head: *Infant visual statistical learning*

Title:

Across space and time:

Infants learn from backward and forward visual statistics

Word Count: 3,995 [revised 4,482]

Review Copy Only

Research Highlights

- Eight-month-old infants demonstrate sensitivity to both backward and forward conditional probabilities, looking longer at unrelated part-pairs than at statistically related base pairs taken from a familiarized visual stream.
- Infants demonstrate this sensitivity whether familiarized with the conditional probabilities in a temporal sequence of looming shapes or in a spatial array of streaming shapes.
- Infants are sensitive to the direction of the conditional relationship between paired items (i.e., whether one item should be preceded or followed by another) when presented in a temporal sequence, but not when presented in a spatial array.

Abstract

This study investigates whether infants are sensitive to backward and forward transitional probabilities within temporal and spatial visual streams. Two groups of 8-month-old infants were familiarized with an artificial grammar of shapes, comprised of backward and forward base pairs (i.e., two shapes linked by strong backward or forward transitional probability) and part-pairs (i.e., two shapes with weak transitional probabilities in both directions). One group viewed the continuous visual stream as a temporal sequence, while the other group viewed the same stream as a spatial array. Following familiarization, infants looked longer at test trials containing part-pairs than base pairs, though they had appeared with equal frequency during familiarization. This pattern of looking time was evident for both forward and backward pairs, in both the temporal and spatial conditions. Further, differences in looking time to part-pairs that were consistent or inconsistent with the predictive direction of the base pairs (forward or backward) indicated that infants were indeed sensitive to direction when presented with temporal sequences, but not when presented with spatial arrays. These results suggest that visual statistical learning is flexible in infancy and depends on the nature of visual input.

Keywords: Statistical learning, visual learning, infant cognitive development

1
2
3 To accurately represent the structure of their visual environment, infants must detect and
4 extract meaningful signals from the massive amounts of noisy input encountered in the first few
5 years of life. A growing number of studies suggest that infants are sensitive to statistical
6 regularities in the visual domain, which enables them to extract patterns for further processing
7 (Fiser & Aslin, 2002; Kirkham, Slemmer & Johnson, 2002; Kirkham, Slemmer, Richardson, &
8 Johnson, 2007; Tummeltshammer & Kirkham, 2013; Wu, Gopnik, Richardson, & Kirkham,
9 2011). This visual statistical sensitivity is thought to facilitate many perceptual and cognitive
10 processes such as action prediction, spatial cueing, causation, object recognition, and category
11 learning (Baldwin, Andersson, Saffran, & Meyer, 2008; Sobel & Kirkham, 2006;
12 Tummeltshammer & Kirkham, 2013; Wu, et al., 2011; Younger & Cohen, 1986).
13
14
15
16
17
18
19
20
21
22
23
24
25
26

27 This research begs the question of whether infants' early emerging statistical sensitivities
28 may be input general or input specific. There has been limited inquiry into the bounds of visual
29 statistical learning in infancy and the extent to which the mechanism may be dependent on
30 stimulus characteristics, context, or manner of presentation. The current study investigates the
31 flexibility of infants' sensitivity to visual statistical regularities by manipulating two critical
32 features of the input stream: visual context (i.e., temporal sequence or spatial array) and
33 statistical properties (i.e., backward or forward transitional probability).
34
35
36
37
38
39
40
41
42

43 ***1. Temporal and spatial statistics***

44
45
46 Much of the research on infant statistical learning has been undertaken in relation to early
47 language comprehension, considering which statistical signals are utilized by infants to discover
48 word boundaries, phonetic contrasts, and grammar structure in a speech stream (Gomez &
49 Gerken, 1999; Maye, Werker, & Gerken, 2002; Pelucchi, Hay, & Saffran, 2009b; Saffran, Aslin,
50 & Newport, 1996). Paradigms using artificial language sequences have been adapted to pure
51
52
53
54
55
56
57
58
59
60

1
2
3 auditory, visual, and motor domains to investigate the possibility of similar processes or even a
4 domain-general statistical learning mechanism (Baldwin et al., 2008; Kirkham et al., 2002;
5 Saffran, et al., 1999; Thiessen, 2011). For example, Kirkham and colleagues (2002) familiarized
6 infants with a temporal stream of six colored shapes that loomed onscreen. The continuous
7 sequence could be segmented into three pairs using transitional probability: The first shape in a
8 pair always predicted the second (within-pair TP=1.0) while the second shape did not predict the
9 next pair (between-pair TP=0.33). After familiarization, infants as young as 2 months looked
10 longer at sequences that violated this statistical pattern, providing the first evidence that infants
11 make use of transitional probability to parse a visual sequence. Fiser and Aslin (2002) conducted
12 a visual statistical learning investigation using spatial arrays, in which they familiarized infants
13 with two items that reliably co-occurred beside a third item that occurred randomly. They found
14 that infants looked longer at arrays containing co-occurring shapes than at arrays made up of
15 items that did not reliably co-occur, indicating that conditional statistics may also play a role in
16 infants' parsing of the spatial world.
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35

36 An accepted interpretation of these findings is that infants isolate visual regularities by
37 comparing statistical distributions, such as item frequency and conditional probability, via a
38 'statistical learning' mechanism (e.g., Aslin & Newport, 2012). This view suggests that infants
39 perform statistical computations over a set of input, making rapid and automatic predictions
40 about individual items within the set. For example, when watching others' actions, they may
41 compute the transitional probability of subsequent movements, in order to anticipate which
42 movement is likely to occur next and identify the action's endpoints (e.g., Baldwin et al., 2008).
43
44
45
46
47
48
49
50
51
52
53 Indeed infants' statistical learning capabilities have been demonstrated experimentally with a
54
55
56
57
58
59
60

1
2
3 multiple statistics across a variety of inputs (e.g., Marcovitch & Lewkowicz, 2009, Thiessen &
4
5 Erickson, 2013).
6

7
8 Another possible interpretation is that infants' sensitivity to statistical distributions may be
9
10 the by-product of an 'associative learning' strategy (or 'chunking', e.g., Miller, 1956). According
11
12 to this view, individual elements that frequently co-occur are extracted and stored as a structured
13
14 chunk (Perruchet & Peereman, 2004). Infants' responses, then, reflect a memory-based
15
16 recognition of regularities encountered in the environment without ascribing sophisticated
17
18 computational abilities to infants (see models, *PARSER* [Perruchet & Vintner, 1998] and *TRACX*
19
20 [French, Addyman, & Mareschal, 2011]). As noted by Perruchet and colleagues, statistical
21
22 learning and chunking explanations converge at many points, as both are considered automatic,
23
24 incidental, pre-attentive, domain-general processes that largely lead to the same behavioral
25
26 responses (Perruchet & Peereman, 2004; Perruchet and Pacton, 2006). Further, these processes
27
28 may not be mutually exclusive (e.g., chunk-based representations may arise from a sensitivity to
29
30 statistical regularities) and may even be selectively recruited when relevant.
31
32
33
34
35

36 Although the Kirkham, et al. (2002) and Fiser & Aslin (2002) studies provide some
37
38 converging evidence for extending infants' use of conditional probabilities to the visual domain,
39
40 it is unclear whether the same learning mechanism can be implicated across all types of input.
41
42 For example, while auditory input is temporal in nature and relies on temporal cues to highlight
43
44 meaningful features (e.g., pitch, prosody, semantics), visual input is spatial in nature and
45
46 emphasizes spatial relations at many levels of processing (e.g., retinotopic mapping, object
47
48 tracking, navigation)¹. With different features and neural architectures implicated in their
49
50
51
52

53
54
55 ¹ Although it is worth pointing out that many visual events can be represented both temporally and spatially, such
56
57 as action sequences and chains of causally related events. Similarly, spatial relations do play a role in the auditory
58
59 domain, such as in localizing and orienting to sounds.
60

1
2
3 processing, it is not obvious that auditory and visual information would elicit the same
4
5 sensitivities from infant learners. Even within a single domain, it remains inconclusive whether
6
7 the same mechanism can be implicated for learning both temporal and spatial inputs.
8
9

10 ***II. Backward and forward probabilities***

11
12 Visual streams containing both backward and forward conditional probabilities provide an
13
14 interesting opportunity to evaluate the input-specificity of infant statistical learning. While some
15
16 statistics, such as pair frequency, do not contain any information about order or direction,
17
18 conditional probabilities can differ when computed with respect to the forward direction (i.e., X
19
20 followed by Y) or backward direction (i.e., Y preceded by X). Consider the following example,
21
22 illustrated by Perruchet and Desauty (2008): In English the letter q is always followed by u , but
23
24 u can be preceded by many letters other than q . Therefore, the forward probability $q \rightarrow u$ is 1.0,
25
26 while the backward probability $q \leftarrow u$ is actually much lower. Recent research in the domain of
27
28 language comprehension has demonstrated that infants and adults are sensitive to statistical
29
30 regularities defined in the backward as well as the forward direction (Jones & Pashler, 2007;
31
32 Pelucchi, Hay, & Saffran, 2009a; Perruchet & Desauty, 2008). However, language is inherently
33
34 a temporal input, which suggests a need to be receptive to temporal order. Thus sensitivity to
35
36 backward and forward statistics could be input-specific, not domain-general.
37
38
39
40
41
42
43

44 It is currently unknown whether infants are able to detect backward probabilities in a
45
46 continuous visual stream, and further, it is unclear whether infants can learn from both backward
47
48 and forward statistics within the same set of input. Is a visual event represented with respect to
49
50 what came before it, as much as it may strongly predict the next event? Since backward and
51
52 forward probabilities within a single pair of events can differ, sensitivity to this difference would
53
54 lend support to the idea that infants encode events with respect to their conditional statistical
55
56
57
58
59
60

1
2
3 relationships. However, failing to discriminate between backward and forward probabilities
4
5 would rather suggest that infants bind co-occurring events into associative chunks independent of
6
7 their predictive direction.
8
9

10 The goals of the present study are: 1) to determine whether infants are sensitive to both
11
12 backward and forward transitional probabilities in a visual stream, and 2) to test whether infants
13
14 extract the same statistical regularities from temporal and spatial presentations. Two groups of 8-
15
16 month-old infants were familiarized to an artificial grammar of shapes, comprised of backward
17
18 and forward base pairs (i.e., two shapes linked by strong backward or forward transitional
19
20 probability) and part-pairs (i.e., two shapes with weak transitional probabilities in both
21
22 directions). One group viewed a temporal sequence, with shapes looming one at a time in the
23
24 center of the screen, while the other group viewed a spatial array, with shapes moving across the
25
26 screen in horizontal rows. Following familiarization, all infants saw test trials of base pairs and
27
28 part-pairs, which had appeared with equal frequency in the familiarized pattern. Consistent with
29
30 previous findings using natural speech, we predicted that infants would discriminate both
31
32 backward and forward base pairs from part-pairs, demonstrated by differences in looking time
33
34 (Pelucchi, Hay, & Saffran, 2009a). Further, we hypothesized that infants would show evidence
35
36 of encoding the backward or forward direction of the base pairs, looking longer at part-pairs that
37
38 were inconsistent with the predictive direction of the base pair (e.g., longer looking to XA than to
39
40 DZ if $A \rightarrow D$ were a forward pair, since A should predict D). While differences in looking times in
41
42 either direction would indicate that infants indeed discriminated between base pairs and part-
43
44 pairs on the basis of their transitional probability, we expected a novelty preference (i.e., longer
45
46 looking at the part-pairs) due to previous findings using similar visual stimuli with
47
48
49
50
51
52
53
54
55
56
57
58
59
60

familiarization/test procedures in our laboratory (e.g., Kirkham, et al., 2002; Kirkham, et al., 2007; Wu & Kirkham, 2011)².

Method

Participants

Forty-four full-term 8-month-old infants participated in this between-subjects experiment: 22 infants in the *Temporal* condition (10 females, $M=7$ months 29.8 days, Range = 7m 7d – 9m 1d), with three additional infants excluded due to fussiness, and 22 infants in the *Spatial* condition (11 females, $M=7$ months 28.5 days, Range=7m 9d – 8m 28d), with three additional infants excluded due to fussiness. Infants were recruited on a voluntary basis via local advertisements. Informed consent was received from all caregivers, and all babies received a small gift.

Apparatus & Stimuli

Visual stimuli were displayed on a 23” screen and sounds were played through stereo external speakers on either side of the display. Infants and their eye movements were monitored using a Tobii TX300 eye-tracker with a video camera built-in below the display. Infants viewed streams of 11 unique colorful shapes (see a sample in Figure 1). The shapes were animated to loom in the center or to stream across the screen using Adobe Director software. The video clips were shown via Tobii’s Studio presentation software.

[Insert Figure 1 approximately here]

Design & Procedure

Infants were tested individually in a quiet room, seated on their caregiver’s lap approximately 60 cm away from the screen. All infants viewed a continuous stream of colorful

² However, it is not uncommon for infants to exhibit preferences for related items, as familiarity and novelty preferences may relate to task complexity (Fiser & Aslin, 2002; Roder, Bushnell, & Sasseville, 2000).

1
2
3 shapes, presented as either a temporal sequence (*Temporal* condition) or a spatial array (*Spatial*
4 condition). A familiarization/test procedure was used: Infants saw the probabilistic pattern
5 repeated several times, and their looking time was subsequently measured as they viewed
6 specific test items taken from the familiarized pattern.
7
8
9
10
11

12 The continuous stream of shapes was comprised of three forward base pairs (*AD*, *BD*, and
13 *CD*), three backward base pairs (*EF*, *EG*, and *EH*) and three single shapes (*X*, *Y*, and *Z*). The
14 forward base pairs were linked by strong forward probabilities, as the first shape always
15 predicted the second shape (e.g., $C \rightarrow D = 1.0$). The backward base pairs were linked by strong
16 backward probabilities, as the first shape always preceded the second shape (e.g., $E \leftarrow F = 1.0$).
17 Forward base pairs had weak backward probabilities, as three different shapes could precede the
18 second shape (i.e., $A, B, C \leftarrow D$). Backward base pairs had weak forward probabilities, as three
19 different shapes could follow the first shape (i.e., $E \rightarrow F, G, H$). Finally, the single shapes (*X*, *Y*,
20 and *Z*) had weak transitional probabilities in both directions and could precede or follow any
21 base pairs (e.g., $Z \rightarrow A, B, C, E$ and $D, F, G, H \leftarrow Z$). These unpaired shapes formed weak “part-
22 pairs” with the shapes adjacent to them. Following familiarization, looking times to base pairs
23 and part-pairs were compared to determine whether infants were sensitive to the transitional
24 probabilities in the stream. Shapes were randomly assigned to the pairings for each infant, so that
25 looking time measures would not be biased by any arbitrary shape or shape pair preferences.
26
27 Two pairs (*AD* and *EH*) and one single shape (*X*) appeared twice as often as the others, in order
28 to balance the frequencies of base pairs and part-pairs compared at test. If infants parsed the
29 stream using pair-frequency, then base pairs would be indistinguishable from part-pairs. Thus
30 any differences in looking time to base pairs and part-pairs at test could be taken as evidence of
31 infants’ sensitivity to the transitional probabilities.
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

[Insert Table 1 approximately here]

Temporal condition. Infants in the *Temporal* condition viewed this stream as a temporal sequence, with forward and backward transitional probabilities indicating which shapes appeared before and after a given shape in *time*. The shapes loomed one at a time in the center of the screen for one second, resulting in a 48-second block of 48 shapes (see Figure 1, as well as Supporting Information, clip 1). Infants viewed three blocks of temporal sequences, as determined by an independent pilot test using infant-controlled habituation. This length of familiarization was sufficient for infants to become familiar with the shape transitions, but not to become fussy or inattentive. Thus across the entire *Temporal* familiarization, infants viewed the more frequent base and part-pairs 24 times each and the less frequent base and part-pairs 12 times each. A short attention-grabbing video clip (Elmo from Sesame Street) was inserted between each familiarization block and abstract sounds (e.g. *bring, ding-dong, honk*) were inserted randomly to attract less attentive infants back to the screen, but offer no additional cues for segmenting the stream.

Spatial condition. Infants in the *Spatial* condition viewed this stream as a spatial array that moved across the screen from left-to-right or right-to-left (between subjects) for 48 seconds (see Figure 1, as well as Supporting Information, clip 2). The same base pairs and part-pairs were contained in the stream (see Table 1); but in this case, the forward and backward transitional probabilities indicated which shapes appeared on the left or right of a given shape in *space*. Infants viewed four blocks of the spatial arrays, as an independent pilot test using infant-controlled habituation determined that infants took longer to habituate to the stream when presented as a spatial array than when presented as a temporal sequence. Thus across the entire spatial familiarization, infants viewed the more frequent base and part-pairs 32 times each and

1
2
3 the less frequent base and part-pairs 16 times each. The same attention-grabbing video clip and
4
5 abstract sounds were used to maintain infants' attention to the screen.
6
7

8 Following *Temporal* or *Spatial* familiarization, all infants viewed test trials of base pairs
9
10 and part-pairs taken from the familiarized pattern. Each test pair was repeated six times in a
11
12 single trial that lasted 12 seconds. There were 8 test trials, the order of which was counter-
13
14 balanced across participants: 4 base pairs, 2 forward (i.e., *BD* and *CD*) and 2 backward (i.e., *EF*
15
16 and *EG*); and 4 part-pairs, 2 forward (i.e., *DZ* and *XA*) and 2 backward (*YE* and *HX*). Since base
17
18 pairs and part-pairs presented at test had appeared with equal frequency during familiarization
19
20 (e.g., *CD* appeared as often as *DZ*), any differences in infants' mean looking times could be
21
22 attributed to their sensitivity to the transitional probabilities in the stream. Importantly, two part-
23
24 pairs were inconsistent with the predictive direction of a forward or backward base pair (i.e., *XA*
25
26 was inconsistent with forward base pair $A \rightarrow D$ as *A* should always predict *D*; and *HX* was
27
28 inconsistent with backward base pair $E \leftarrow H$ as *H* should always be preceded by *E*). The other two
29
30 part-pairs were consistent with the base pair statistics (i.e., *DZ* and *YE*, since *D* and *E* could be
31
32 preceded or followed by multiple shapes). Thus, if infants were sensitive to whether base pairs
33
34 were forward or backward, then they should look longer at the inconsistent part-pairs than the
35
36 consistent part-pairs.
37
38
39
40
41
42

43 **Data Analysis.** Infants' looking times to base pairs and part-pairs at test were compared
44
45 between the *Temporal* and *Spatial* conditions. Following from previous visual statistical learning
46
47 studies, we hypothesized that infants would demonstrate longer mean looking times for part-
48
49 pairs than base pairs (e.g., Kirkham, et al., 2002; Kirkham, et al., 2007; Marcovitch &
50
51 Lewkowicz, 2009; Wu, et al., 2011). Further, we hypothesized that if they were sensitive to both
52
53 forward and backward transitional probabilities, then infants would look longer at part-pairs than
54
55
56
57
58
59
60

1
2
3 both backward and forward base pairs (i.e., no interaction with direction). Finally, we
4
5 hypothesized that infants would look longer at part-pairs that were inconsistent, rather than
6
7 consistent, with the predictive direction of the forward and backward base pair probabilities,
8
9 indicating that they had encoded the base pairs' directions (e.g., if $A \rightarrow D$ is encoded as a forward
10
11 pair, then XA should be more surprising than DZ , since A should predict D but D must not be
12
13 preceded by A).
14
15
16

17 Results

18
19 During test trials, infants looked longer at part-pairs than base pairs. This was apparent for
20
21 infants in the *Temporal* condition (part-pairs $M=7.90$ s, $SD=1.99$ s; base pairs $M=6.64$ s,
22
23 $SD=2.60$ s) as well as for infants in the *Spatial* condition (part-pairs $M=7.44$ s, $SD=1.95$ s; base
24
25 pairs $M=6.83$ s, $SD=2.19$ s). Eighteen out of 22 infants showed a preference for part-pairs in the
26
27 *Temporal* condition (Wilcoxin Signed Ranks test, $z=3.49$, $p<0.001$), and 16 out of 22 infants in
28
29 the *Spatial* condition ($z=2.18$, $p=0.03$). As looking times were not normally distributed,
30
31 parametric tests using log-transformed looking time data were conducted to confirm this result:
32
33 *Temporal* $t(21)=3.98$, $p=0.001$; *Spatial* $t(21)=2.41$, $p=0.02$.
34
35
36
37
38

39 Log-transformed looking times were further examined in a 2(Condition: *Temporal*,
40
41 *Spatial*) x 2(Direction: backward, forward) x 2(Type: base pairs, part-pairs) mixed ANOVA,
42
43 which showed a main effect of Type, $F(1,42)=17.27$, $p<0.001$, $\eta_p^2=0.29$, confirming that infants
44
45 indeed looked longer at part-pairs than base pairs. A marginal effect of Direction was also
46
47 apparent, $F(1,42)=3.80$, $p=0.06$, $\eta_p^2=0.08$, as infants looked slightly longer at forward direction
48
49 test trials compared to backward direction test trials. There were no other effects or interactions,
50
51 all $p>0.11$, indicating that longer looking at part-pairs than base pairs was evident for both
52
53 backward and forward probabilities and persisted in both the *Spatial* and *Temporal* conditions.
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Planned separate post-hoc comparisons confirmed that the preference for part-pairs was significant in both the forward direction, $t(43)=3.67, p=0.001$, and the backward direction, $t(43)=2.82, p=0.007$. Raw looking times for these variables are shown in Figure 2.

[Insert Figure 2 approximately here]

Finally, the four part-pair test trials were divided according to whether the part-pairs were consistent or inconsistent with the predictive direction of the base pair from which the part-pair item was taken (i.e., whether the part-pair item should have predicted another item with a backward or forward probability of 1.0). Mean looking times were compared in separate 2 (Direction: backward, forward) x 3 (Type: base pairs, consistent part-pairs, inconsistent part-pairs) repeated measures ANOVAs. In the *Temporal* condition, there was a main effect of Type, $F(2,42)=9.12, p<0.001$, and no effect of Direction or interaction ($p>0.490$). A post-hoc Friedman Ranks test showed that infants looked longest at inconsistent part-pairs, less at consistent part-pairs, and least at base pairs, $\chi^2=12.09, p=0.002$. However, in the *Spatial* condition, no main effect of Type emerged, $F(2,42)=0.144, p=0.866$, nor any effect of Direction or interaction, indicating that infants did not discriminate between the consistent and inconsistent part-pairs. This result suggests that infants who saw the *temporal* presentation were sensitive to the predictive directions of the base pairs, while infants who saw the *spatial* presentation were not. Raw looking times for these comparisons are shown in Figure 3.

[Insert Figure 3 approximately here]

Discussion

The results of the present study provide the first evidence that infants are sensitive to both backward and forward transitional probabilities in a continuous stream of visual input. Following familiarization with sequences containing both backward and forward transitional probabilities,

1
2
3 infants looked longer at part-pairs than base pairs, though base and part-pairs were equally
4
5 familiar and frequent. This result is consistent with previous studies of visual statistical learning,
6
7 in which infants also showed preferences for unrelated items (Marcovitch & Lewkowicz, 2009;
8
9 Wu, et al., 2011). Infants' looking preferences were apparent for both forward and backward
10
11 directions, indicating that they were sensitive to both forward and backward transitional
12
13 probabilities when parsing the visual stream. This result suggests that both directions are of
14
15 critical importance to the infant visual learner. Indeed, forward and backward statistical relations
16
17 may serve different, but equally important, functions: Forward relations facilitate prediction and
18
19 planning, while backward relations support explanation and causal attribution. Future studies
20
21 may investigate whether infants use both forward and backward statistics to modify their visual
22
23 behavior. A few studies have shown that infants make use of forward statistical regularities in
24
25 planning saccades, showing faster latencies to statistically predictable locations (Amso &
26
27 Davidow, 2012; Kirkham, et al., 2007; Tummeltshammer & Kirkham, 2013).

28
29 Further, differences in sensitivity to violations of the forward or backward transitional
30
31 probabilities indicate that infants may not encode temporal and spatial visual regularities in the
32
33 same way. When presented with temporal sequences, infants looked longer at inconsistent,
34
35 compared to consistent, part-pairs, indicating that they had encoded the predictive direction of
36
37 the base pairs (e.g., represented $A \rightarrow D$ as a forward pair and looked longer when A appeared
38
39 without D). This result was apparent for both backward and forward pairs (i.e., no Type x
40
41 Direction interactions), such that infants showed no advantage in sensitivity to one direction of
42
43 transitional probability over the other. This finding is consistent with a study by Turk-Browne
44
45 and Scholl (2009), who showed that adults encode visual triplets with their temporal order,
46
47 though they can also recognize the triplets independent of order. However, when presented with
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 spatial arrays, infants showed no differences in looking time to consistent and inconsistent part-
4
5
6 pairs, suggesting that they had not encoded the base pairs as forward or backward (e.g., looked
7
8 equally long when *A* appeared without *D* or *D* appeared without *A*). Rather, infants who saw the
9
10 spatial presentations may have bound the paired shapes into associative chunks without
11
12 preserving information about their direction (e.g., representing $A \rightarrow D$ as single unit *AD*).
13
14

15 It has been suggested that infants' responses on statistical learning tasks may emerge from
16
17 an 'associative learning' or 'chunking' strategy, reflecting memory-based recognition of
18
19 regularities in the environment without demanding statistical computations (French, Addyman,
20
21 & Mareschal, 2011; Perruchet & Peereman, 2004; Perruchet & Pacton, 2006). Though many
22
23 statistical learning tasks can indeed be accomplished using associative learning, sensitivity to
24
25 more subtle statistical features (such as whether pairs are linked by backward or forward
26
27 conditional probabilities) supports the engagement of a more computational strategy. Further
28
29 work is needed to disentangle the roles of memory-based chunking and statistical learning
30
31 accounts in infants' deconstruction and representation of their visual experience. The present
32
33 results indicate that infants' sensitivity to visual regularities depends on the nature of the input
34
35 stream, suggesting that statistical learning mechanisms may be selectively recruited when
36
37 relevant.
38
39
40
41
42

43 There are a number of fundamental distinctions between temporal and spatial presentations
44
45 that may account for the observed differences in infants' sensitivities across the two conditions.
46
47 First, temporal and spatial presentations place different demands on the cognitive resources
48
49 necessary to accomplish statistical learning, namely attention and working memory. In temporal
50
51 sequences, items are presented one-by-one such that the order of information is fixed and the
52
53 targets of attention are pre-selected for the learner. Spatial arrangements, on the other hand,
54
55
56
57
58
59
60

1
2
3 present multiple relations at the same time and attention can be freely allocated, such that items
4
5 may be fixated in many different orders. Equivalent learning, then, would depend on infants'
6
7 ability to distribute attention evenly across all items in the spatial display. Studies with adults
8
9 have suggested that success in statistical learning may indeed depend on the efficient deployment
10
11 of attention (Baker, Olson, & Behrmann, 2004; Fiser & Aslin, 2001; Turk-Browne, Junge, &
12
13 Scholl, 2005). Further, temporal information is transient and can only be reviewed in memory,
14
15 whereas spatial information is readily available for visual inspection. This difference in
16
17 perceptual access may affect the depth and detail of encoding, with temporal relations encoded
18
19 more robustly than spatial relations. Pilot work using infant-controlled habituation indicated that
20
21 infants required more trials to habituate to the spatially presented stream than the temporally
22
23 presented stream, justifying the difference in length of familiarization used in the current
24
25 experiment. This pilot result is consistent with the idea that spatial and temporal relations may
26
27 have different processing and encoding demands, as the speed of habituation is thought to reflect
28
29 information processing time (e.g., Dember & Earl, 1957; Fantz, 1964; Hunter & Ames, 1988;
30
31 Sokolov, 1963). The difference in familiarization length is unlikely to have produced the present
32
33 results, as one would expect longer familiarization in the spatial condition to yield greater
34
35 sensitivity, which is the opposite of what we observed.
36
37
38
39
40
41
42

43
44 Second, statistical direction may not be equally salient or relevant in temporal and spatial
45
46 relations, as these dimensions emphasize different features of a visual event. Fiser and Aslin
47
48 (2001) suggest that higher-order statistical information may only be represented when it is
49
50 necessary to solve a particular task; otherwise visual feature processing would be
51
52 computationally intractable. Accordingly, infants may only show sensitivity to the direction of
53
54 statistical relations when it signals a meaningful distinction. While one could argue that
55
56
57
58
59
60

1
2
3 directionality is inherent to temporal positions (i.e., before and after) and critically informative to
4
5 temporally defined events (e.g., identifying cause and effect), it is unclear whether the same is
6
7 true of spatial positions (i.e., above, below, beside). It may be informative to represent two items
8
9 as adjacent in space, but their absolute position is likely to be flexible or to depend on vantage
10
11 point. For the developing child, it may be particularly adaptive to employ flexible, context-
12
13 relevant learning mechanisms that are sensitive to the nature of visual input, as these
14
15 mechanisms will operate more efficiently over the young learner's rapidly changing
16
17 representations. For example, the physical layout of a room, inherently spatial in nature, may
18
19 take on a highly relevant temporal quality when the infant begins to self-locomote and needs to
20
21 plan her path and navigate obstacles. Conversely, language, which is primarily temporal when
22
23 spoken to the infant, may become spatially represented once the child learns to read and becomes
24
25 highly familiar with the left-to-right (or some other coherent) structure of written words³. In this
26
27 way, such developmental milestones can profoundly alter the child's environmental input, with a
28
29 likely consequence of tuning their sensitivities to the consistent and relevant features of that
30
31 input.
32
33
34
35
36
37

38
39 In conclusion, the present study has demonstrated that infants are sensitive to both forward
40
41 and backward transitional probabilities within continuous visual streams. Importantly, infants'
42
43 sensitivity to conditional statistics depends on the nature of visual input. That is, infants may
44
45 encode and represent statistical information differently across domains, depending on its salience
46
47 or relevancy, providing evidence for a flexible statistical learning strategy with a variety of
48
49 applications in infant cognitive development.
50
51

52
53
54
55
56
57 ³ We thank an anonymous reviewer for this suggestion.
58
59
60

References

- 1
2
3
4
5
6 Amso, D., & Davidow, J. (2012). The development of implicit learning from infancy to
7
8 adulthood: Item relations, salience, and cognitive flexibility. *Developmental*
9
10 *Psychobiology*, *54*, 664-673. doi: 10.1002/dev.20587
11
12
13 Aslin, R.N., & Newport, E.L. (2012). Statistical learning: From acquiring specific items to
14
15 forming general rules. *Current Directions in Psychological Science*, *21*, 170-176. doi:
16
17 10.1177/0963721412436806
18
19
20 Baker, C.I., Olson, C.R., & Behrmann, M. (2004). Role of attention and perceptual grouping in
21
22 visual statistical learning. *Psychological Science*, *15*, 460-466. doi: 10.1111/j.0956-
23
24 7976.2004.00702.x
25
26
27 Baldwin, D., Andersson, A., Saffran, J., & Meyer, M. (2008). Segmenting dynamic human
28
29 action via statistical structure. *Cognition*, *106*, 1382-1407. doi:
30
31 10.1016/j.cognition.2007.07.005
32
33
34 Dember, W.N., & Earl, R.W. (1957). Analysis of exploratory, manipulatory, and curiosity
35
36 behaviors. *Psychological Review*, *64*, 91-96.
37
38
39 Fantz, R.L. (1964). Visual experience in infants: Decreased attention to familiar patterns relative
40
41 to novel ones. *Science*, *146*, 668.
42
43
44 Fiser, J., & Aslin, R.N. (2001). Unsupervised statistical learning of higher-order spatial
45
46 structures from visual scenes. *Psychological Science*, *12*, 499-504. doi: 10.1111/1467-
47
48 9280.00392
49
50
51 Fiser, J., & Aslin, R. N. (2002). Statistical learning of new visual feature combinations by
52
53 infants. *Proceedings of the National Academy of Sciences, USA*, *99*, 15822-15826. doi:
54
55 10.1073/pnas.232472899
56
57
58
59
60

- 1
2
3 French, R.M., Addyman, C., & Mareschal, D. (2011). TRACX: a recognition-based
4
5 connectionist framework for sequence segmentation and chunk extraction. *Psychological*
6
7 *Review*, 118, 614-636. doi: 10.1037/a0025255
8
9
- 10 Gomez, R.L., & Gerken, L. (1999). Artificial grammar learning by 1-year-olds leads to specific
11
12 and abstract knowledge. *Cognition*, 70, 109-135. doi: 10.1016/S0010-0277(99)00003-7
13
14
- 15 Hunter, M.A., & Ames, E.W. (1988). A multifactor model of infant preferences for novel and
16
17 familiar stimuli. *Advances in Infancy Research*, 5, 69-95.
18
19
- 20 Jones, J., & Pashler, H. (2007). Is the mind inherently forward looking? Comparing prediction
21
22 and retrodiction. *Psychonomic Bulletin & Review*, 14, 295-300. doi: 10.3758/BF03194067
23
24
- 25 Kirkham, N. Z., Richardson, D. C., Slemmer, J. A., & Johnson, S. P. (2007). Location, location,
26
27 location: Development of spatiotemporal sequence learning in infancy. *Child Development*,
28
29 78, 1559-1571. doi: 10.1111/j.1467-8624.2007.01083.x
30
31
- 32 Kirkham, N.Z., Slemmer, J. A., & Johnson, S. P. (2002). Visual statistical learning in infancy.
33
34 *Cognition*, 83, B35-B42. doi: 10.1016/S0010-0277(02)00004-5
35
36
- 37 Marcovitch, S., & Lewkowicz, D.L. (2009). Sequence learning in infancy: the independent
38
39 contributions of conditional probability and pair frequency information. *Developmental*
40
41 *Science*, 12, 1020-1025. doi: 10.1111/j.1467-7687.2009.00838.x
42
43
- 44 Maye, J., Werker, J.F., & Gerken, L. (2002). Infant sensitivity to distributional information can
45
46 affect phonetic discrimination. *Cognition*, 82, B101-B111. doi: 10.1016/S0010-
47
48 0277(01)00157-3
49
- 50 Miller, G.A. (1956). The magical number seven, plus or minus two: Some limits on our capacity
51
52 for processing information. *Psychological Review*, 63, 81-97.
53
54
55
56
57
58
59
60

- 1
2
3 Pelucchi, B., Hay, J.F., & Saffran, J.R. (2009a). Learning in reverse: Eight-month-old infants
4 track backward transitional probabilities. *Cognition*, *113*, 244-247. doi:
5
6 10.1016/j.cognition.2009.07.011
7
8
9
10 Pelucchi, B., Hay, J.F., & Saffran, J.R. (2009b). Statistical learning in a natural language by 8-
11 month-old infants. *Child Development*, *80*, 674-685. doi: 10.1111/j.1467-
12 8624.2009.01290.x
13
14
15
16
17 Perruchet, P., & Desauty, S. (2008). A role for backward transitional probabilities in word
18 segmentation? *Memory & Cognition*, *36*, 1299-1305. doi: 10.3758/MC.36.7.1299
19
20
21
22 Perruchet, P. & Pacton, S. (2006). Implicit learning and statistical learning: one phenomenon,
23 two approaches. *TRENDS in Cognitive Sciences*, *10*, 233-238. doi:
24
25 10.1016/j.tics.2006.03.006
26
27
28
29 Perruchet, P., & Peereman, R. (2004). The exploitation of distributional information in syllable
30 processing. *Journal of Neurolinguistics*, *17*, 97-119. doi: 10.1016/S0911-6044(03)00059-9
31
32
33
34 Perruchet, P., & Vintner, A. (1998). PARSER: A model for word segmentation. *Journal of*
35 *Memory and Language*, *39*, 246-263. doi: 10.1006/jmla.1998.2576
36
37
38
39 Roder, B.J., Bushnell, E.W., & Sassville, A.M. (2000). Infants' preferences for familiarity and
40 novelty during the course of visual processing. *Infancy*, *1*, 491-507.
41
42
43
44 Saffran, J.R., Aslin, R.N., & Newport, E.L. (1996). Statistical learning by 8-month-old infants.
45 *Science*, *274*, 1926-1928. doi: 10.1126/science.274.5294.1926
46
47
48
49 Saffran, J.R., Johnson, E.K., Aslin, R.N., & Newport, E.L. (1999). Statistical learning of tone
50 sequences by human infants and adults. *Cognition*, *70*, 27-52. doi: 10.1016/S0010-
51 0277(98)00075-4
52
53
54
55
56
57
58
59
60

- 1
2
3 Sobel, D.M., & Kirkham, N.Z. (2006). Blickets and babies: the development of causal reasoning
4
5 in toddlers and infants. *Developmental Psychology*, *42*, 1103-1115. doi: 10.1037/0012-
6
7 1649.42.6.1103
8
9
- 10 Sokolov, E. (1963). *Perception and the Conditioned Reflex*. Oxford, England: Pergamon.
11
12
- 13 Thiessen, E.D. (2011). Domain general constraints on statistical learning. *Child Development*,
14
15 *82*, 462-470. doi: 10.1111/j.1467-8624.2010.01522.x
16
17
- 18 Thiessen, E.D., & Erickson, L.C. (2013). Beyond word segmentation: A two-process account of
19
20 statistical learning. *Current Directions in Psychological Science*, *22*, 239-243. doi:
21
22 10.1177/0963721413476035
23
24
- 25 Tummeltshammer, K.S., & Kirkham, N.Z. (2013). Learning to look: probabilistic variation and
26
27 noise guide infants' eye movements. *Developmental Science*, *16*, 760-771. doi:
28
29 10.1111/desc.12064
30
31
- 32 Turk-Browne, N.B., Jungé, J., & Scholl, B.J. (2005). The automaticity of visual statistical
33
34 learning. *Journal of Experimental Psychology: General*, *134*, 552-564. doi: 10.1037/0096-
35
36 3445.134.4.552
37
38
- 39 Turk-Browne, N.B., & Scholl, B.J. (2009). Flexible visual statistical learning: Transfer across
40
41 space and time. *Journal of Experimental Psychology: Human Perception and Performance*,
42
43 *35*, 195-202. doi: 10.1037/0096-1523.35.1.195
44
45
- 46 Wu, R., Gopnik, A., Richardson, D.C., & Kirkham, N.Z. (2011). Infants learn about objects from
47
48 statistics and people. *Developmental Psychology*, *47*, 1220-1229. doi: 10.1037/a0024023
49
50
- 51 Younger, B.A., & Cohen, L.B. (1986). Developmental change in infants' perception of
52
53 correlations among attributes. *Child Development*, *57*, 803-815. doi: 10.2307/1130356
54
55
56
57
58
59
60

Figure Captions

Figure 1. Sample visual streams presented to infants in the Temporal (top) and Spatial (bottom) conditions. During familiarization, the stream comprised of base pairs and part-pairs was presented continuously (left), and during test trials, a single base pair or part-pair was presented multiple times (right). It is worth noting that although shapes were displayed at the same rates, in comparison to the temporal condition, the spatial condition presented more shapes onscreen at the same time and each shape was visible for a longer duration.

Figure 2. Mean looking times (in seconds) at test for base pairs and part-pairs; separated by *Temporal* and *Spatial* conditions.


Figure 3. Mean looking times (in seconds) at test for base pairs, consistent part-pairs (i.e., *DZ* and *YE*), and inconsistent part-pairs (i.e., *XA* and *HX*); separated by *Temporal* and *Spatial* conditions.

Table Legends


Table 1. Elements comprising the probabilistic visual stream with their transitional probabilities and pair frequencies. In order to balance the pair frequencies of the base pairs and part-pairs presented at test, some shapes differed in their individual frequencies (i.e., one forward base pair, one backward base pair, and one single shape appeared twice as often as the others).

Review Copy Only



Familiarization




Test Trials




Spatial





1
2
3
4
5
6
7
8
9
10
11
12
13
14
15



developmentalscience@wiley.com

	Base Pairs	Single Shapes	Transitional Probability		Test Pair Frequency	
			Pairs	Part-Pairs	Pairs	Part-Pairs
Forward	A D A D B D C D	X X Y Z	$TP_{for} = 1.0$ $C \rightarrow D$ $TP_{back} = 0.25$ $C \leftarrow D$	$TP_{for} = 0.25$ $D \rightarrow Z$ $TP_{back} = 0.5$ $D \leftarrow Z$	BD = 1/24 CD = 1/24	DZ = 1/24 XA = 1/24
Backward	E F E G E H E H	X X Y Z	$TP_{for} = 0.25$ $E \rightarrow F$ $TP_{back} = 1.0$ $E \leftarrow F$	$TP_{for} = 0.5$ $Y \rightarrow E$ $TP_{back} = 0.25$ $Y \leftarrow E$	EF = 1/24 EG = 1/24	YE = 1/24 HX = 1/24
<p>Sample sequence of 48 shapes: ADXEGYADZEHXEFZCDXADYEFXBDZADXEHYBDXEHXCDYEHZEGX</p> <p>Parsed using within-pair probabilities: AD-X-EG-Y-AD-Z-EH-X-EF-Z-CD-X-AD-Y-EF-X-BD-Z-AD-X-EH-Y-BD-X-EH-X-CD-Y-EH-Z-EG-X</p>						