



University of HUDDERSFIELD

University of Huddersfield Repository

Rahman, Rubina, Mirza, Naznin and Stephenson, John

Risk factors for elevated intraocular pressure on first day postoperative review following pars plana vitrectomy

Original Citation

Rahman, Rubina, Mirza, Naznin and Stephenson, John (2016) Risk factors for elevated intraocular pressure on first day postoperative review following pars plana vitrectomy. In: British and Eire VR (BEAVRS) 2016 meeting, Thursday 10th & Friday 11th of November 2016, Grand Harbour Hotel, Southampton. (Unpublished)

This version is available at <http://eprints.hud.ac.uk/30683/>

The University Repository is a digital collection of the research output of the University, available on Open Access. Copyright and Moral Rights for the items on this site are retained by the individual author and/or other copyright owners. Users may access full items free of charge; copies of full text items generally can be reproduced, displayed or performed and given to third parties in any format or medium for personal research or study, educational or not-for-profit purposes without prior permission or charge, provided:

- The authors, title and full bibliographic details is credited in any copy;
- A hyperlink and/or URL is included for the original metadata page; and
- The content is not changed in any way.

For more information, including our policy and submission procedure, please contact the Repository Team at: E.mailbox@hud.ac.uk.

<http://eprints.hud.ac.uk/>

Risk factors for elevated intraocular pressure on first day postoperative review following pars plana vitrectomy

Rubina Rahman MBBS, FRCS, FRCOphth

Naznin Mirza MBChB

John Stephenson PHD

Financial Disclosure: The authors have no financial or propriety interest in any material or method mentioned.

Objectives

- To assess the relationship between day 1 post-operative intra-ocular pressure and patient demographic information, lens status, tamponade medium (air, C2F6 and SF6) and laser treatment.


Methods

- Consecutive case study at Calderdale and Huddersfield NHS Foundation Trust.
- All patients undergoing pars plana vitrectomy were pseudophakic or underwent combined surgery.
- Indications for surgery: MH, ERM or RRD.
- None of the patients received prophylactic anti-glaucoma medication.



Results

- 164 patients were evaluated on day 1 post surgery.
- 10 patients had IOP >30mmHg, incidence 6%
- Range 30 – 39mmHg
- Mean 32.5mmHg
- Incidence of hypotony (< 8mmHg) : 5 (2.7%).



Results: Vitrectomy gauge & lens status


Indication	23g	25g	27g	Phakic	Pseudo
ERM	22	14	25	52	9
MH	20	13	14	42	5
RRD	33	3	20	48	8

- No association with postoperative intraocular pressure.



Results:

Tamponade	Number of patients	
Air	37	22%
C2F6(20%)	74	45%
SF6 (20%)	25	15%
SF6 (25%)	28	17%



- 67 patients had retinopexy during vitrectomy.
- 50 patients received laser treatment, 88% in RRD group, 17 had cryotherapy.
- Number of laser burns ranged from 11-1489 (average 712)
- **IOP elevated by 0.3mmHg for every additional 100 burns (p= 0.028)**

Conclusion:

- Tamponade with C2F6 or SF6 & increasing number of laser burns predisposes to increased IOP.
- Incidence of IOP > 30 mmhg is low (6%) in small gauge PPV and phacovitrectomy TSV without prophylactic antiglaucoma medications.
- Incidence of hypotony on day 1 is LOW (2.7%) in tamponade filled eyes.
- Prophylactic anti-glaucoma medication is only justified in high risk patients.