



Muhammad, N; Mpyet, C; Adamu, MD; William, A; Umar, MM; Goyol, M; Muazu, H; Onyebuchi, U; Isiyaku, S; Flueckiger, RM; Chu, BK; Willis, R; Pavluck, AL; Alhassan, A; Olobio, N; Gordon, BA; Solomon, AW (2016) Mapping Trachoma in Kaduna State, Nigeria: Results of 23 Local Government Area-Level, Population-Based Prevalence Surveys. *Ophthalmic epidemiology*, 23 (sup1). pp. 46-54. ISSN 0928-6586 DOI: <https://doi.org/10.1080/09286586.2016.1250918>


Downloaded from: <http://researchonline.lshtm.ac.uk/3197202/>

DOI: [10.1080/09286586.2016.1250918](https://doi.org/10.1080/09286586.2016.1250918)

Usage Guidelines

Please refer to usage guidelines at <http://researchonline.lshtm.ac.uk/policies.html> or alternatively contact researchonline@lshtm.ac.uk.

Available under license: <http://creativecommons.org/licenses/by-nc-nd/2.5/>



Mapping Trachoma in Kaduna State, Nigeria: Results of 23 Local Government Area-Level, Population-Based Prevalence Surveys

Nasiru Muhammad, Caleb Mpyet, Mohammed Dantani Adamu, Adamani William, Murtala Muhammad Umar, Musa Goyol, Habila Muazu, Uwaezuoke Onyebuchi, Sunday Isiyaku, Rebecca M. Flueckiger, Brian K. Chu, Rebecca Willis, Alexandre L. Pavluck, Abdullahi Alhassan, Nicholas Olobio, Bruce A. Gordon & Anthony W. Solomon

To cite this article: Nasiru Muhammad, Caleb Mpyet, Mohammed Dantani Adamu, Adamani William, Murtala Muhammad Umar, Musa Goyol, Habila Muazu, Uwaezuoke Onyebuchi, Sunday Isiyaku, Rebecca M. Flueckiger, Brian K. Chu, Rebecca Willis, Alexandre L. Pavluck, Abdullahi Alhassan, Nicholas Olobio, Bruce A. Gordon & Anthony W. Solomon (2016) Mapping Trachoma in Kaduna State, Nigeria: Results of 23 Local Government Area-Level, Population-Based Prevalence Surveys, *Ophthalmic Epidemiology*, 23:sup1, 46-54, DOI: [10.1080/09286586.2016.1250918](https://doi.org/10.1080/09286586.2016.1250918)

To link to this article: <https://doi.org/10.1080/09286586.2016.1250918>



Published with license by Taylor & Francis©
2016 Nasiru Muhammad, Caleb Mpyet, Mohammed Dantani Adamu, Adamani William, Murtala Muhammad Umar, Musa Goyol, Habila Muazu, Uwaezuoke Onyebuchi, Sunday Isiyaku, Rebecca M. Flueckiger, Brian K. Chu, Rebecca Willis, Alexandre L. Pavluck, Abdullahi Alhassan, Nicholas Olobio, Bruce A. Gordon, and Anthony W. Solomon, for the Global Trachoma Mapping Project.



Published online: 05 Dec 2016.



Submit your article to this journal [↗](#)



Article views: 549



View related articles [↗](#)



View Crossmark data [↗](#)



Citing articles: 2 View citing articles [↗](#)

Mapping Trachoma in Kaduna State, Nigeria: Results of 23 Local Government Area-Level, Population-Based Prevalence Surveys

Nasiru Muhammad^a, Caleb Mpyet^{b,c}, Mohammed Dantani Adamu^a, Adamani William^b, Murtala Muhammad Umar^d, Musa Goyol^e, Habila Muazu^f, Uwaezuoke Onyebuchi^g, Sunday Isiyaku^b, Rebecca M. Flueckiger^h, Brian K. Chu^h, Rebecca Willis^h, Alexandre L. Pavluck^h, Abdullahi Alhassaniⁱ, Nicholas Olobio^g, Bruce A. Gordon^j, and Anthony W. Solomon^{k,l}, for the Global Trachoma Mapping Project*

^aOphthalmology Unit, Surgery Department, Usmanu Danfodiyo University, Sokoto, Nigeria; ^bSightsavers, Kaduna, Nigeria; ^cOphthalmology Department, University of Jos, Jos, Nigeria; ^dNational Eye Centre, Kaduna, Nigeria; ^eNetherlands Leprosy Relief, Nigeria; ^fGambo Sawaba Hospital, Zaria, Nigeria; ^gFederal Ministry of Health, Abuja, Nigeria; ^hTask Force for Global Health, Decatur, GA, USA; ⁱKaduna State Ministry of Health, Kaduna, Nigeria; ^jWater Sanitation and Hygiene, Department of Public Health, Social and Environmental Determinants of Health, World Health Organization, Geneva, Switzerland; ^kClinical Research Department, London School of Hygiene & Tropical Medicine, London, UK; ^lLondon Centre for Neglected Tropical Disease Research, London, UK

ABSTRACT

Introduction: To prepare for global elimination of trachoma by 2020, the World Health Organization (WHO) recommends mapping of trachoma at district-level to enable planning of elimination activities in affected populations. The aim of our study was to provide data on trachoma for each local government area (LGA) of Kaduna State, Nigeria, as such data were previously unavailable.

Method: As part of the Global Trachoma Mapping Project (GTMP), a population-based cross-sectional trachoma survey was conducted in each of the 23 LGAs of Kaduna State, between May and June 2013. The protocols of the GTMP were used.

Results: The prevalence of trachomatous inflammation – follicular (TF) in children aged 1–9 years was between 0.03% and 8% across the LGAs, with only one LGA (Igabi) having a TF prevalence $\geq 5\%$. The LGA-level prevalences of trichiasis in persons aged 15 years and older were between 0.00% and 0.78%. Eleven LGAs had trichiasis prevalences of 0.2% and over in adults; a threshold equivalent to 1 case per 1000 total population. The LGA-level proportion of households with access to improved water sources ranged from 9% to 96%, while household access to latrines ranged from 5% to 99%.

Conclusion: Kaduna State has generally hypoendemic trachoma, but a few trichiasis surgeries are still required to attain the WHO elimination targets. Better access to improved water and sanitation is needed.

ARTICLE HISTORY

Received 31 March 2016
Revised 3 September 2016
Accepted 14 October 2016

KEYWORDS

Global Trachoma Mapping Project; Kaduna; neglected tropical diseases; Nigeria; trachoma; trichiasis; water and sanitation

Introduction

Trachoma, a neglected tropical disease, is caused by *Chlamydia trachomatis*, and is the leading infectious cause of blindness, accounting for 1.4% of the global burden of blindness.¹ It is characterized by chronic conjunctivitis that results in scarring of the conjunctivae, and, in some people, trichiasis (in-turned eyelashes) with consequent corneal scarring. The World Health Organization (WHO) recommends mapping of trachoma at district level² in areas where disease is suspected, for planning based on disease burden and/or documentation of elimination. Mapping data should help to focus efforts to achieve global elimination of trachoma as a public health problem by the year 2020.³

The Global Trachoma Mapping Project (GTMP) commenced in 2012.⁴ The GTMP's goal was to complete the global baseline trachoma map, in order to plan public health interventions and to mobilize resources for trachoma control.⁵

Kaduna State, located in north-western Nigeria, has a population of 7,474,369 (2013 projection)⁶ residing in an estimated 1,115,974 regular households located in 23 local government areas (LGAs).^{7,8} Regular households consist of a group of persons living under the same roof with a recognized head, sharing a common catering arrangement and functioning as a social unit.⁸ A previous population-based survey reported trachomatous corneal opacity as the 7th most common cause of low

CONTACT Nasiru Muhammad ✉ nasiru69@yahoo.com 📧 Ophthalmology Unit, Surgery Department, Usmanu Danfodiyo University, Sokoto, Nigeria.

*See Appendix

Color versions of one or more of the figures in the article can be found online at www.tandfonline.com/oi/pe.

Published with license by Taylor & Francis

© 2016 Nasiru Muhammad, Caleb Mpyet, Mohammed Dantani Adamu, Adamani William, Murtala Muhammad Umar, Musa Goyol, Habila Muazu, Uwaezuoke Onyebuchi, Sunday Isiyaku, Rebecca M. Flueckiger, Brian K. Chu, Rebecca Willis, Alexandre L. Pavluck, Abdullahi Alhassani, Nicholas Olobio, Bruce A. Gordon, and Anthony W. Solomon, for the Global Trachoma Mapping Project.

This is an Open Access article distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives License (<http://creativecommons.org/licenses/by-nc-nd/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited, and is not altered, transformed, or built upon in any way.

vision in the state (0.3% of low vision).⁹ Kaduna has an eye care program that has provided eye care services, including lid surgery for trichiasis, over the last 20 years, largely in health facilities or during cataract outreach campaigns.

The aim of this study was to provide prevalence data on trachoma in all 23 LGAs of Kaduna State, as no previous trachoma-specific surveys had been conducted here. This should provide data to assist in decision-making with respect to local trachoma elimination. The objectives were to estimate the prevalence of trachomatous inflammation – follicular (TF)¹⁰ in children aged 1–9 years in each Kaduna LGA, to estimate the prevalence of trichiasis in persons aged 15 years and older in each Kaduna LGA, and to assess the water and sanitation coverage in each Kaduna LGA.

Materials and methods

Between May and June 2013, in each LGA, we conducted a population-based cross-sectional prevalence survey using GTMP protocols.¹¹

Sample size

Each LGA was considered as a separate evaluation unit. LGA populations ranged from 127,581 to 500,464, with an estimated 50% of the population aged 15 years and older.⁷ As described elsewhere, for each LGA, the sample size for estimating TF prevalence required framing of sufficient households to reach an expected 1222 children aged 1–9 years.¹¹ To estimate trichiasis prevalence in persons 15 years and older, all eligible persons living in selected households were examined.

Ethical considerations

The Ethics Committee of the London School of Hygiene & Tropical Medicine (reference 6319) and the National Health Research Ethics Committee of Nigeria (reference NHREC/01/01/2007) each granted approval for the study. The Kaduna State Ministry of Health granted administrative permission for the work to go ahead. Consent of participants was obtained verbally, with that for minors granted by parents or guardians; consent was documented in the LINKS Android smartphone application (app).¹¹ Participants found to have trichiasis during the surveys were appropriately referred, while those with active trachoma (TF and/or trachomatous inflammation – intense, TI, in either eye) were offered two tubes of 1% tetracycline eye ointment each.

Survey teams

The survey teams comprised graders and recorders who were trained and certified according to GTMP protocols (version 1).¹¹ Training was conducted over 4 days, with the first 2 days focused on training and selecting graders. Graders who achieved a minimum kappa rating for TF of 0.7 against a GTMP-certified grader trainer were selected for the survey. Ophthalmologists provided supervision to the teams, giving feedback and crosschecking in-service diagnostic accuracy.

Sampling design

Study subjects were selected using a 3-stage sampling approach. In the first stage, we systematically selected 25 clusters, using probability proportional to size sampling, from the list of towns and villages in each LGA. Each cluster was then sub-divided into the lowest administrative unit (ward) and one unit was selected using simple random sampling. In each unit, we selected 25 households. Despite its epidemiological drawbacks, the random walk method was used to select households within wards, as the security situation was tenuous, and we felt that locally familiar field methods would place survey teams at less risk of generating unmerited suspicion than ones that would be novel to included communities.^{12–14} All residents of selected households who were aged 1 year and older and had lived in the area for at least the previous 6 months were enumerated, and invited to participate.

Data collection

Consent to proceed was obtained by the survey team from the head of household or his/her representative, who also provided information on water and sanitation access for the household. Global positioning system (GPS) coordinates for the household's location were then recorded. Trachoma grading was undertaken according to the WHO simplified grading scheme,¹⁰ using $\times 2.5$ magnifying loupes under sunlight illumination. Hands were cleaned with an alcohol-based hand gel between the examination of successive subjects.

Data management

All collected data were directly entered into the LINKS app on Android smartphones and uploaded to a cloud-based server. The GTMP Data Manager cleaned and analyzed the data, as described elsewhere.¹¹ The proportion of 1–9-year-old children seen to have TF in each cluster was adjusted for age in 1-year age bands, and the

proportion of ≥ 15 -year-olds seen to have trichiasis in each cluster was adjusted for age and sex in 5-year age bands, with the Kaduna population pyramid from the 2006 census used as the reference. The adjusted prevalences of TF, TI and trichiasis for each LGA were the arithmetic means of the adjusted cluster level proportions of each sign.

Results

Study sample

A total of 33,884 children aged 1–9 years were enumerated, of whom 32,882 (97.0%) were examined, and 43,127 adults aged 15+ years were enumerated, of whom 38,824 (90.0%) were examined, in the 23 LGAs combined. The number of children examined per LGA ranged from 1044 to 1991, while the number of adults examined ranged from 1076 to 2524 per LGA. Sabon Gari LGA had the highest absentee rate (19.8%) among children, while the refusal rate among children was highest in Kaduna North (6.4%; Table 1).

Prevalence of trachoma

The prevalence of TF in children aged 1–9 years ranged from 0.0% (95% confidence interval, CI, 0.0–0.1%) in Kaduna South LGA to 8.1% (95% CI 5.7–10.9%) in Igabi LGA; in this age group, the prevalence of TI

ranged from 0.0% in nine LGAs to 1.1% (95% CI 0.4–1.9%) in Makarfi LGA. The prevalence of trichiasis in those aged 15+ years ranged from 0.0% in Jema'a LGA to 0.8% (95% CI 0.4–1.2%) in Lere LGA (Table 1). Figure 1 shows the LGAs surveyed, and Figure 2 shows TF prevalences and Figure 3 shows trichiasis prevalences for each LGA.

Water and sanitation coverage

The proportion of households with access to an improved water source ranged from 9% in Kudan to 96% in Jema'a LGA (Table 2). Access to water inside the compound or within 1 km of the house was lowest in Zangon Kataf (53%) and universal in Giwa (100%). Household-level latrine access ranged from 5% in Giwa to 100% in Kudan and Soba (Table 2).

Trachoma elimination targets

All LGAs except Igabi recorded TF prevalences $< 5\%$. The majority of LGAs had prevalences of trichiasis below the elimination threshold, with ten LGAs needing to carry out more trichiasis surgeries to reach the elimination target of < 1 case of unmanaged trichiasis per 1000 total population (0.1%), equivalent to a prevalence of 0.2% in adults aged 15+ years.¹⁵

Table 1. Local government area-level prevalences of trachomatous inflammation – follicular (TF) and trichiasis in Kaduna State, Nigeria, Global Trachoma Mapping Project, 2013.

Local Government Area	Sample examined n		Refused n		Absent n		Trachoma prevalence % (95% CI)	
	Children (1-9 yrs)	Adults (15+ yrs)	Children (1-9 yrs)	Adults (15+ yrs)	Children (1-9 yrs)	Adults (15+ yrs)	TF (1-9 yrs)	Trichiasis (15+ yrs)
Birnin Gwari	1222	1225	0	0	1	2	0.1 (0.0-0.2)	0.40 (0.14-0.65)
Chikun	840	1007	4	19	1	321	2.7 (1.0-4.8)	0.03 (0.0-0.10)
Giwa	1500	1308	1	19	52	213	1.7 (1.0-2.6)	0.35 (0.15-0.61)
Igabi	1444	1379	4	23	21	76	8.1 (5.7-10.9)	0.34 (0.16-0.59)
Ikara	1169	803	7	92	3	286	1.7 (1.1-2.3)	0.20 (0.04-0.43)
Jaba	982	1302	1	3	4	39	0.5 (0.3-0.9)	0.08 (0.00-0.23)
Jema'a	1219	1882	11	8	10	16	1.0 (0.5-1.8)	0.00
Kachia	1059	1166	14	1	22	30	1.4 (0.9-1.9)	0.16 (0.04-0.33)
Kaduna North	827	1619	60	87	17	117	0.6 (0.2-1.1)	0.07 (0.02-0.14)
Kaduna South	1126	918	2	0	36	37	0.0 (0.0-0.1)	0.08 (0.0-0.18)
Kagarko	1353	1148	1	5	2	8	2.0 (1.2-2.9)	0.38 (0.11-0.75)
Kajuru	916	1184	9	12	5	120	2.1 (0.9-3.8)	0.11 (0.02-0.26)
Kaura	1108	1460	4	2	6	140	2.0 (1.2-3.1)	0.09 (0.00-0.20)
Kauru	870	1378	37	31	60	334	4.9 (3.1-7.3)	0.50 (0.23-0.79)
Kubau	1014	1168	5	32	52	143	1.0 (0.3-2.0)	0.41 (0.22-0.60)
Kudan	1531	1282	21	49	50	93	2.4 (1.7-3.2)	0.32 (0.06-0.56)
Lere	799	1118	1	0	4	28	1.4 (0.5-2.1)	0.78 (0.40-1.24)
Makarfi	1070	1362	2	32	0	104	1.1 (0.3-2.3)	0.33 (0.14-0.55)
Sabon Gari	794	1177	2	36	203	196	0.5 (0.0-1.3)	0.09 (0.03-0.17)
Sanga	978	1279	4	4	5	68	2.2 (1.3-3.1)	0.09 (0.01-0.20)
Soba	1219	1062	3	84	5	97	0.2 (0.0-0.4)	0.23 (0.02-0.41)
Zangon Kataf	1070	1280	3	0	7	132	2.9 (2.0-3.9)	0.03 (0.00-0.08)
Zaria	898	1940	4	4	6	126	0.0 (0.0-0.4)	0.08 (0.01-0.14)

Prevalences of trichiasis are displayed to two decimal places in order to provide clarity on whether or not the best estimate of prevalence was above or below the elimination threshold of 0.2% in adults ≥ 15 years. CI, confidence interval.




Figure 1. Local government areas surveyed in Kaduna State, Nigeria, Global Trachoma Mapping Project, 2013.

Related targets

Our water, sanitation and hygiene (WASH) data show that United Nations sustainable development goal 6 (“Ensure access to water and sanitation for all”)¹⁶ will require considerable work in most LGAs of Kaduna (Table 3). Providing universal access to water and sanitation will help to prevent future recrudescence of trachoma in this population and contribute to the control and elimination of other neglected tropical diseases, as noted in a recent WHO Global Strategy document.¹⁷

Discussion

The results of this study suggests that active trachoma is not of public health significance in most LGAs of Kaduna State, except Igabi, which could benefit from implementation of the SAFE strategy (surgery, antibiotics, facial cleanliness and environmental improvement), including a single round of mass azithromycin treatment, and implementation of the F and E components before re-survey.¹⁸ Alternatively, given Igabi’s relatively large population, consideration could be given to undertaking further surveys at a more granular level, to determine whether there is really a need for interventions throughout the entire LGA. This LGA has a densely populated urban slum that may explain the higher TF prevalence than seen in other LGAs of Kaduna State.

The prevalence of TI was low in all LGAs. The generally low level of trachoma endemicity in Kaduna State may be partly due to ongoing provision of a comprehensive eye care program for more than two decades and the social development efforts that have happened over the same period. Our findings contrast with reports from neighboring Katsina,^{12,19} Kano,¹³ Kebbi and Sokoto²⁰ States, and also from the Republic of Niger further to the north,²¹ where many surveyed evaluation units have recorded TF prevalences higher than 10% and trichomatous trichiasis prevalences higher than 1%.

Trichiasis, the blinding stage of trachoma, was below the elimination threshold (0.2% prevalence in adults) in most LGAs. However, ten of the 23 LGAs may need to undertake more trichiasis surgeries to reach the elimination threshold. The low precision of the trichiasis estimates generated needs to be taken into account when interpreting these data, as previously acknowledged.¹¹ Stakeholders should prioritize these ten LGAs by engaging already-available human resources to offer active trichiasis case finding and community-based surgeries. Eyelid surgery services will need to be continuously available even after elimination end points have been reached, as new cases are likely to continue to develop.

The United Nations Millennium Development Goal target 7C was to halve, by 2015, the proportion of people without sustainable access to safe drinking water and basic sanitation.²² The results of this study




Figure 2. Prevalence of trachomatous inflammation – follicular (TF) in 1–9-year-old children by local government area in Kaduna State, Nigeria, Global Trachoma Mapping Project, 2013.

provide a snapshot of the 2013 situation in Kaduna. A total of 12 of the 23 LGAs in Kaduna State had <60% of households with access to improved water sources and 15 of 23 LGAs lacked 60% household-level access to improved sanitation. This situation calls for all stakeholders to redouble current efforts at improving access to water and sanitation, if the targets enshrined in the Sustainable Development Goals are to be achieved by 2030.²³ We note that in this set of surveys, Igabi, which had the highest TF prevalence, had the second-lowest household-level access to improved sanitation. Even if not necessarily causal, both active trachoma and lack of access to sanitation are markers of deprivation, and prioritizing WASH sector actions to populations with higher trachoma prevalences would go some way towards putting the most vulnerable people first.¹⁷

Our data have some limitations. First, a number of LGAs (Table 2) had total populations that exceeded the 250,000 recommended as the upper limit for “districts” in trachoma surveys; it is possible that small foci of disease were missed by chance. Second, the response rate in some LGAs such as Ikara, where only 803 of 1181 resident adults aged 15+ years (68%) were examined, may have introduced bias in the prevalence estimates. In general, however, the response was excellent; examination rates for resident 1–9-year-olds, for example, ranged from 79–100%, with the majority exceeding 90% (Table 1). Third, the random-walk method has been justifiably criticized for potentially permitting biased household selection,^{24–26} unfortunately, we felt that security considerations here outweighed epidemiological ones. Fourth, because trichiasis is a rare condition,




Figure 3. Prevalence of trichiasis in adults aged 15+ years by local government area in Kaduna State, Nigeria, Global Trachoma Mapping Project, 2013.

the CIs for estimates of its prevalence are relatively wide.¹¹ Finally, we did not record the presence or absence of trachomatous conjunctival scarring in eyes that had trichiasis; the recognition that this might be critical for distinguishing trachomatous from non-trachomatous trichiasis did not emerge until 2015,²⁷ after these surveys were completed.

These surveys show that Kaduna State has relatively hypoendemic trachoma. To reach the trachoma elimination threshold (trichiasis prevalences <math><0.2\%</math> in adults aged 15+ years and TF prevalences <math><5\%</math> in 1–9-year-olds), provision of community-based trichiasis surgery for 11 LGAs, and implementation of the A, F and E components of the SAFE strategy in one LGA will be needed. Our data also demonstrate a particular need for attention towards improving access to appropriate water and sanitation facilities for the local population.

Declaration of interest

The authors report no conflicts of interest. The authors alone are responsible for the content and writing of the paper.

Funding

This study was principally funded by the Global Trachoma Mapping Project (GTMP) grant from the United Kingdom's Department for International Development (ARIES: 203145) to Sightsavers, which led a consortium of non-governmental organizations and academic institutions to support ministries of health to complete baseline trachoma mapping worldwide. The GTMP was also funded by the United States Agency for International Development (USAID) through the ENVISION project implemented by RTI International under cooperative agreement number AID-OAA-A-11-00048, and the END in Asia project implemented by FHI360 under cooperative agreement number OAA-A-10-00051. A committee established in

Table 2. Household access to water and sanitation facilities in Kaduna State, Nigeria, Global Trachoma Mapping Project, 2013.

Local Government Area	Households enumerated, <i>n</i>	Proportion of households, %		
		With access to improved water source	With water source within yard/1 km	With access to improved sanitation facilities
Birnin Gwari	612	45	78	96
Chikun	632	61	95	43
Giwa	614	50	100	5
Igabi	624	26	99	8
Ikara	613	79	96	19
Jaba	623	67	87	44
Jema'a	623	96	79	83
Kachia	616	56	86	24
Kaduna North	615	84	86	99
Kaduna South	613	85	94	95
Kagarko	622	41	93	20
Kajuru	607	64	89	41
Kaura	613	80	82	82
Kauru	625	27	92	26
Kubau	623	28	99	54
Kudan	616	9	96	99
Lere	600	66	98	47
Makarfi	624	59	95	31
Sabon Gari	626	64	99	49
Sanga	619	68	80	16
Soba	624	17	97	99
Zangon Kataf	613	48	53	8
Zaria	873	56	99	90

Table 3. Activities needed for trachoma elimination as a public health problem and achievement of United Nations sustainable development goal 6 in Kaduna State, Nigeria, as of 2013.

Local Government Area	2006 population, <i>n</i>	Households, <i>n</i> (2006 census) ⁸	Households in need of improved water sources, <i>n</i>	Households in need of latrines, <i>n</i>	Trichiasis surgeries to achieve elimination target (<0.2%), <i>n</i>	Need for mass antibiotic treatment to clear ocular <i>Chlamydia trachomatis</i> infection
Birnin Gwari	258,581	44,643	24,554	1786	290	Individualized
Chikun	372,272	76,289	29,936	43,698	0	Individualized
Giwa	292,384	49,552	24,533	46,891	246	Individualized
Igabi	430,753	77,243	56,944	70,809	338	Mass treatment (single round before re-survey), or re-survey at finer level
Ikara	194,723	34,017	7269	27,414	0	Individualized
Jaba	155,973	29,472	9602	16,557	0	Individualized
Jema'a	278,202	51,839	1913	8569	0	Individualized
Kachia	252,568	47,116	20,500	35,719	0	Individualized
Kaduna North	364,575	71,283	11,591	813	0	Individualized
Kaduna South	402,731	81,693	12,524	3995	0	Individualized
Kagarko	239,058	43,306	25,620	34,671	241	Individualized
Kajuru	109,810	19,183	6985	11,345	0	Individualized
Kaura	174,626	33,008	6730	6083	0	Individualized
Kauru	221,276	38,382	27,942	28,372	372	Individualized
Kubau	280,704	50,606	36,472	23,233	330	Individualized
Kudan	138,956	24,406	22,107	39	93	Individualized
Lere	339,740	57,962	19,898	30,529	1103	Individualized
Makarfi	146,574	26,785	10,859	18,543	107	Individualized
Sabon Gari	291,358	52,977	18,955	27,251	0	Individualized
Sanga	151,485	27,302	8734	22,846	0	Individualized
Soba	291,173	50,483	41,664	81	49	Individualized
Zangon Kataf	318,991	59,646	30,843	54,880	0	Individualized
Zaria	406,990	68,781	30,016	6775	0	Individualized

March 2012 to examine issues surrounding completion of global trachoma mapping was initially funded by a grant from Pfizer to the International Trachoma Initiative. BAG is an employee of the World Health Organization (WHO); the views expressed in this article are the views of the authors alone and do not necessarily reflect the views or policy of

WHO. AWS was a Wellcome Trust Intermediate Clinical Fellow (098521) at the London School of Hygiene & Tropical Medicine. None of the funders had any role in project design, in project implementation or analysis or interpretation of data, in the decisions on where, how or when to publish in the peer-reviewed press, or in the preparation of the manuscript.

References

- Bourne RRA, Stevens GA, White RA, et al. Causes of vision loss worldwide, 1990–2010: a systematic analysis. *Lancet Global Health* 2013;1:e339–349.
- Solomon AW, Zondervan M, Kuper H, et al. *Trachoma control: a guide for program managers*. Geneva: World Health Organization, 2006.
- Smith JL, Haddad D, Polack S, et al. Mapping the global distribution of trachoma: why an updated atlas is needed. *PLoS Negl Trop Dis* 2011;5:e973.
- Solomon AW, Kurylo E. The Global Trachoma Mapping Project. *Comm Eye Health* 2014; 27(85):18.
- Emerson P. The SAFE strategy for trachoma control: poised for rapid scale-up. *Comm Eye Health* 2014;27 (86):38.
- Kaduna State Government. *About Kaduna state*. Available at: www.kadunastate.gov.ng Accessed 22 December 2013.
- National Population Commission. *2006 National and state population and housing tables – Priority tables I*. Abuja, Nigeria: Author, 2010.
- National Population Commission. *2006 Housing characteristics and amenities tables – Priority tables (LGA) volume II*. Abuja, Nigeria: Author, 2010.
- Rabiu MM. Prevalence of blindness and low vision in north central, Nigeria. *West Afr J Med* 2008;27:238–244.
- Thylefors B, Dawson CR, Jones BR, et al. A simple system for the assessment of trachoma and its complications. *Bull World Health Organ* 1987;65: 477–483.
- Solomon AW, Pavluck AL, Courtright P, et al. The Global Trachoma Mapping Project: methodology of a 34-country population-based study. *Ophthalmic Epidemiol* 2015;22:214–225.
- Mpyet C, Muhammad N, Adamu MD, et al. Prevalence of trachoma in Katsina state, Nigeria: results of 34 district-level surveys. *Ophthalmic Epidemiol* 2016;23 (S1):55–62.
- Mpyet C, Muhammad N, Adamu MD, et al. Prevalence of trachoma in Kano state, Nigeria: results of 44 local government area-level surveys. *Ophthalmic Epidemiol*. doi:10.1080/09286586.2016.1265657
- Mpyet C, Muhammad N, Adamu MD, et al. Prevalence of trachoma in Bauchi state, Nigeria: results of 20 local government area-level surveys. *Ophthalmic Epidemiol* 2016;23(S1):39–45.
- World Health Organization. Report of the 2nd global scientific meeting on trachoma, Geneva, 25–27 August, 2003. Geneva: World Health Organization, 2003.
- United Nations General Assembly. Resolution adopted by the General Assembly on 25 September 2015 (A/70/L.1). Transforming our world: the 2030 agenda for sustainable development. New York: United Nations, 2015.
- Boisson S, Engels D, Gordon BA, et al. Water, sanitation and hygiene for accelerating and sustaining progress on neglected tropical diseases: a new Global Strategy 2015–20. *Intl Health* 2016;8(Suppl. 1):i19–i21.
- World Health Organization. *Technical Consultation on Trachoma Surveillance Meeting Report*. Decatur, GA, USA, 2014.
- Jip NF, King JD, Diallo MO, et al. Blinding trachoma in Katsina State, Nigeria: population-based prevalence survey in ten local government areas. *Ophthalmic Epidemiol* 2008;15:294–302.
- Muhammad N, Mohammed A, Isiyaku S, et al. Mapping trachoma in 25 local government areas of Sokoto and Kebbi states, northwestern Nigeria. *Br J Ophthalmol* 2014;98:432–437.
- Cromwell EA, Amza A, Kadri B, et al. Trachoma prevalence in Niger: results of 31 district-level surveys. *Trans R Soc Trop Med Hyg* 2014;108:42–48.
- International Agency for the Prevention of Blindness. Blindness, poverty and development: the impact of VISION 2020 on the U.N. Millennium Development Goals. 2006. 12 pp. Available from: <http://www.v2020.org> Accessed 26 April 2014.
- World Health Organization. Health in 2015 from MDGs to SDGs; 2015:216 pp.
- Turner AG, Magnani RJ, Shuaib M. A not quite as quick but much cleaner alternative to the Expanded Programme on Immunization (EPI) cluster survey design. *Int J Epidemiol* 1996, 25:198–203.
- Brogan D, Flagg EW, Deming M, et al. Increasing the accuracy of the Expanded Programme on Immunization's cluster survey design. *Ann Epidemiol* 1994;4:302–311.
- Grais RF, Rose AMC, Guthmann JP. Don't spin the pen: two alternative methods for second stage sampling in cluster surveys in urban zones. *Emerg Themes Epidemiol* 2007;4:8.
- World Health Organization Alliance for the Global Elimination of Trachoma by 2020. Second Global Scientific Meeting on Trachomatous Trichiasis, November 4–6, 2015, Cape Town, South Africa. Geneva: World Health Organization, 2016.

Appendix

The Global Trachoma Mapping Project Investigators are: Agatha Aboe (1,11), Liknaw Adamu (4), Wondu Alemayehu (4,5), Menbere Alemu (4), Neal D. E. Alexander (9), Berhanu Bero (4), Simon J. Brooker (1,6), Simon Bush (7,8), Brian K. Chu (2,9), Paul Courtright (1,3,4,7,11), Michael Dejene (3), Paul M. Emerson (1,6,7), Rebecca M. Flueckiger (2), Allen Foster (1,7), Solomon Gadisa (4), Katherine Gass (6,9), Teshome Gebre (4), Zelalem Habtamu (4), Danny Haddad (1,6,7,8), Erik Harvey (1,6,10), Dominic Haslam (8), Khumbo Kalua (5), Amir B. Kello (4,5), Jonathan D. King (6,10,11), Richard Le Mesurier (4,7), Susan Lewallen (4,11), Thomas M. Lietman (10), Chad MacArthur (6,11), Colin Macleod (3,9), Silvio P. Mariotti (7,11), Anna Massey (8), Els Mathieu (6,11), Siobhain McCullagh (8), Addis Mekasha (4), Tom Millar (4,8), Caleb Mpyet (3,5), Beatriz Muñoz (6,9), Jeremiah Ngondi (1,3,6,11), Stephanie Ogden (6), Alex Pavluck (2,4,10), Joseph Pearce (10), Serge Resnikoff (1), Virginia Sarah (4), Boubacar Sarr (5), Alemayehu Sisay (4), Jennifer L. Smith (11), Anthony W. Solomon (1,2,3,4,5,6,7,8,9,10,11), Jo Thomson (4); Sheila K. West (1,10,11), Rebecca Willis (2,9).

Key: (1) Advisory Committee, (2) Information Technology, Geographical Information Systems, and Data Processing, (3) Epidemiological Support, (4) Ethiopia Pilot Team, (5) Master Grader Trainers, (6) Methodologies Working Group, (7) Prioritisation Working Group, (8) Proposal Development, Finances and Logistics, (9) Statistics and Data Analysis, (10) Tools Working Group, (11) Training Working Group.