

Black, N. (2016) [Accepted Manuscript] Higher Mortality in Weekend Admissions to the Hospital: True, False, or Uncertain? JAMA. ISSN 0098-7484 DOI: https://doi.org/10.1001/jama.2016.16410

Downloaded from: http://researchonline.lshtm.ac.uk/3331450/

DOI: 10.1001/jama.2016.16410

Usage Guidelines

 $Please\ refer\ to\ usage\ guidelines\ at\ http://researchonline.lshtm.ac.uk/policies.html\ or\ alternatively\ contact\ researchonline@lshtm.ac.uk.$

Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners

Higher mortality in weekend admissions to hospital: true, false or uncertain?

Concern about patients having a a-higher risk of dying if admitted to an acute hospital on Saturday or Sunday, the so-called 'weekend effect', has taken on great political importance in England over the past year. From being a long-standing topic of interest only to health services researchers and policy-makers, this issueit assumed high media coverage when the Secretary of State for Health, Jeremy Hunt, employed it as the prime reason for increasing service provision at weekends and, in turn, why the junior doctors in the NHS should accept new terms and conditions of service. In support of his claims, the Department of Health for England assembled an overview of the topic and concluded that there was 'significant evidence demonstrating this 'weekend effect'.' (1) Attributing the 'weekend effect' to the junior doctors' working practices was felt by them to be unjustified and only served to redouble their opposition to the new contract they were being offered. This resulted in a series of strikes including, for the first time in England, physicians doctors withdrawing emergency care. ([2) - perhaps add ref, Goddard in JAMA, Aug 22, 2016] In light of these events While the dispute has rumbled on, researchers have reexamined the evidence of higher mortality for weekend admissions.

The first reports that hospital mortality was higher during at-weekends appeared in the 1970s. Studies in the USA, the UK and Australia found that deaths involving-in-neonates wereas higher among those born at the weekend. Such a 'weekend effect' was confirmed by Bell and Redelmeier (32) in a study of examining adult admissions to Canadian hospitals in the 1990s who reported that although. However, while they found-mortality was higher for a quarter of the 100 conditions that were the most frequent causes of death, for the majority of conditions, no such effect was apparent. This mixed picture was confirmed by subsequent studies of specific conditions.

The situation changed around 2010 when some ambitious attempts were made in England to compare the mortality for the entire all-patient populations admitted at weekends to the mortality for all that for weekday admissions, rather than focusing on specific diseases as previous studies had. (Nick — not sure I understand the difference — please clarify). (1) These studies reported mortality as being 10-15% higher among for weekend admissions. and formed the basis of politicians'

Commented [NB1]: Hope that's clearer.

subsequent claims. However, these analyses were based on administrative databases that which have little information abouton the clinical state of patients at the time of en admission. These were the same data, subjected to the same analytical techniques, that had been used over the preceding decade to derive hospital-wide standardized mortality ratios. Just as those metrics and comparisons have neveret been adequately validated (43), comparisons of mortality by day of admission havesuffer from the same methodological limitations. (43) The authors were inevitably restricted in their ability to considertake into account differences in case-mix between weekend and weekday admissions as the databases include incomplete information about patients'en their comorbid conditions and no indication aboutef the severity or acuity of the patient's primary and secondary diagnoses [AU:correct?] conditions. In addition, inconsistency in diagnostic coding between weekends and weekdays could not be ruled out. Thus, higher mortality observed during at-weekends could have been due to sicker patients with more severe illness or greater co-morbidities being admitted on those days. What was needed were more rigorous analyses using databases with more clinical information that which had been collected in a more consistent way (such as using agreed diagnostic definitions).

Recent studies using high quality clinical databases have indeed justified these concerns about the existence of a 'weekend effect' on mortality. Each of the three potential limitations of administrative databases have been shown to have contributed to potentially misleading findings. First, the problem of inconsistent coding was demonstrated in a study of admissions for stroke. (54) The authors found that non-emergency admissions after a previous stroke were more likely to be coded as a new stroke on weekdays than at weekends (41% v 27%). As 30 day mortality for admissions for non-emergency strokeies (4%) was much lower than admissions for new strokes (22%), the survival of weekday admissions appeared to be better than at weekend admissions. When this coding error was corrected, there was no evidence of a 'weekend effect'. The authors concluded that "similar biases are likely to occur in studies of the 'weekend effect' in acute admissions for other conditions for which administrative diagnostic coding is prone to inaccuracy." (5)

The second concern, insufficient consideration of comorbidity, was addressed in another study of stroke admissions. (65) The authors used a national clinical

Commented [NB2]: Fine

Commented [NB3]: I'd prefer to state the findings were misleading rather than just 'potentially'!

database of stroke care in England and Wales in which clinicians collect and validate detailed data <u>abouten</u> the clinical characteristics of new acute cases. <u>AfterHaving</u> adjust<u>mented</u> for case-mix differences, there was no evidence of a weekend effect on 30 day mortality.

Failure to considertake the severity or acuity of patients into account, the third concern, was addressed indirectly in a study that compared the two routes of emergency admissions: direct referrals (from primary care physicians) and via the emergency department room (EDR). (76) Whereas the daily number of admissions via the EDR duringat weekends was similar to that on weekdays, the number of direct admissions was 61% lower. Compared to mortality for admissions during weekdays, mWhile mortality for admissions via the EDR was only 5% higher at weekends, whereas for direct admissions mortality it—was 21% higher. Given that, apart from initial treatment in the EDR, both sets of patients received the same inpatient care, this finding provides suggestive circumstantial evidence that the mortality difference between weekend and weekday is more likely to be attributable to how sick-patients' severity of illness and co-morbid conditions at the time of are on admission, rather than the quality of hospital care.

Since then stronger evidence has emerged. Researchers have overcome the lack of severity measures in administrative data by using a recently introduced standard measure of risk of dying based on seven physiological variables, known as. Using the National Early Warning Score (NEWS). In a, they conducted a study in three general hospitals, theat authors and found that patients admitted onadmissions at weekends had greater severity of illness were sicker than than those admitted on weekdays and had a 10% higher risk of dying. (7) although when When th NEWS [AU:correct] is was considered in the analysis [AU:correct2] taken into account, the difference in mortality between weekend and weekday admissions disappeared.

<u>TAll-these</u> recent studies suggest that patients admitted <u>during at-weekends</u> are at no greater risk of dying than those admitted during the week. <u>Although the majority of the recent studies were conducted in England, research focused on this question in any country would have to deal with these same major limitations of using <u>administrative data noted above</u>, particularly case-mix adjustment and illness acuity of the patients at the time of admission.</u>

Commented [NB4]: Correct!

The results of the various recent studies is doe not sn't-mean there is no 'weekend effect', only that if care provided at weekends is poorer than weekdays it does not sn't result in patients' deaths. This should come as no surprise given that death is not a particularly sensitive measure of the overall quality of a-hospital carse because only about 4% of hospital deaths are thought to be avoidable. (98) Instead, to determine find out if the widely held concerns about a 'weekend effect' are justified, investigations need to focus on other dimensions of quality such as health outcomes (morbidity, health-related quality of life), safety (falls, hospital-acquired infections), patients' experience (delays in diagnosis, not receiving sufficient information) and educational quality (training of junior doctors at weekends). In addition, concerns about poorer operational efficiency (extended lengths of stay, delayed discharges) during weekends also need to be addressed. Teortunately, the development of various databases and surveys in these areas means investigation of these factors can be conducted readily and relatively inexpensively. cheaply conducted.

In the meantime, policy-makers need to avoid unjustified claims about an increased heightened-risk of death among patients ying if admitted duringat weekends as doing sothis only serves to undermine the public's confidence and trust in their hospitals and to damagenegatively affect to damage the morale of staff. Neither of these are in the interests of patients or the public.

Commented [NB5]: Personal view but not keen on 'negatively affect'. Prefer plain English of 'damage'!

References

- 1. Department of Health. Research into the 'weekend effect' on patient outcomes and mortality. 30 October 2015. https://www.gov.uk/government/publications/research-into-the-weekend-effect-on-hospital-mortality/research-into-the-weekend-effect-on-patient-outcomes-and-mortality
- 2. Goddard AF. Lessons to Be Learned From the UK Junior Doctors' Strike. JAMA August 22, 2016 doi:10.1001/jama.2016.12029
- 32. Bell CM, Redelmeier DA. Mortality among patients admitted to hospitals on weekends as compared with weekdays. N Engl J Med 2001; 345: 663–68.
- 43. Black N. Assessing the quality of hospitals. Hospital standardised mortality ratios should be abandoned. BMJ 2010;340:c2066

Formatted: Heading 1, Right: 0.53 cm, Font Alignment: Baseline, Pattern: Clear (White)

Formatted: Font: (Default) Arial, 12 pt, Not Bold, Font color: Auto

Formatted: Font: (Default) Arial, 12 pt

- $\underline{5}4$. Li L, Rothwell P on behalf of the Oxford Vascular Study. Biases in detection of apparent "weekend effect" on outcome with administrative coding data: population based study of stroke. BMJ 2016;353:i2648
- <u>6</u>5. Bray BD, Cloud GC, James MA, Hemingway H, Paley L, Stewart K, Tyrrell PJ, Wolfe CD, Rudd AG, SSNAP collaboration. Weekly variation in health-care quality by day and time of admission: a nationwide, registry-based, prospective cohort study of acute stroke care. Lancet 2016:388:170-7.
- <u>76</u>. Meacock R, Anselmi L, Kristensen SR, Doran T, Sutton M. Higher mortality rates amongst emergency patients admitted to hospital at weekends reflect a lower probability of admission. J Health Serv Res Policy 2016; DOI: 10.1177/1355819616649630.
- 8₹. Mohammed M, Faisal M, Richardson D, Howes R, Beaston K, Speed K, Wright J. Adjusting for illness severity shows there is no difference in patient mortality at weekends or weekdays for emergency medical admissions. Quarterly J Medicine 2016; http://dx.doi.org/10.1093/qjmed/hcw104
- 98. Hogan H, Zipfel R, Neuburger J, Hutchings A, Darzi A, Black N. Avoidability of hospital deaths and association with hospital-wide mortality ratios: retrospective case record review and regression analysis. BMJ 2015; 351: h3239.