


OPEN ACCESS



All articles published in the Journal of Threatened Taxa are registered under Creative Commons Attribution 4.0 International License unless otherwise mentioned. JoTT allows unrestricted use of articles in any medium, reproduction and distribution by providing adequate credit to the authors and the source of publication.



Journal of Threatened Taxa

The international journal of conservation and taxonomy

www.threatenedtaxa.org

ISSN 0974-7907 (Online) | ISSN 0974-7893 (Print)

NOTE

RANGE EXTENSION OF THE ENDANGERED SALIM ALI'S FRUIT BAT *LATIDENS SALIMALII* (CHIROPTERA: PTEROPODIDAE) IN THE ANAMALAI HILLS, TAMIL NADU, INDIA

Claire F.R. Wordley, Eleni K. Foui, Divya Mudappa, Mahesh Sankaran & John D. Altringham

26 October 2016 | Vol. 8 | No. 12 | Pp. 9486–9490

[10.11609/jott.2796.8.12.9486-9490](https://doi.org/10.11609/jott.2796.8.12.9486-9490)



For Focus, Scope, Aims, Policies and Guidelines visit http://threatenedtaxa.org/About_JoTT.asp

For Article Submission Guidelines visit http://threatenedtaxa.org/Submission_Guidelines.asp

For Policies against Scientific Misconduct visit http://threatenedtaxa.org/JoTT_Policy_against_Scientific_Misconduct.asp

For reprints contact info@threatenedtaxa.org

Partner



Publisher/Host





ISSN 0974-7907 (Online)
ISSN 0974-7893 (Print)

OPEN ACCESS



Latidens salimalii is the only species in the genus *Latidens*, and is endemic to India (Bates & Harrison 1997). It is the only fruit bat to be protected under Schedule I of the Indian Wildlife (Protection) Act, amended in 2006, and is listed as Endangered by the IUCN (Molur & Vanitharani 2008). It was first collected in the High Wavy Mountains in Tamil Nadu in 1948, and misclassified as the Greater Short-Nosed Fruit Bat *Cynopterus sphinx*, but was later recognised as a new genus and species by Kitti Thonglongya and named after Indian ornithologist Salim Ali in 1972 (Thonglongya 1972). The High Wavy Mountains were the only known location of this species until 1999, when its presence was recorded in the Kalakkad-Mundanthurai Tiger Reserve, Tamil Nadu (Ghosh et al. 1999).

Since then, *Latidens salimalii* has been reported from a few more localities in the southern Western Ghats of India; the Kalakkad-Mundanthurai Tiger Reserve, the Kardana Coffee Estate, Meghamalai, and the High Wavy Mountains in Tamil Nadu (Molur et al. 2002; Vanitharani et al. 2004, 2005) (Fig. 1). There is a report from Uppinangadi in Karnataka, which was described as unverified by Molur & Vanitharani (2008), and another unconfirmed report from the Periyar Tiger Reserve in Kerala (Molur et al. 2002).

RANGE EXTENSION OF THE ENDANGERED SALIM ALI'S FRUIT BAT *LATIDENS SALIMALII* (CHIROPTERA: PTEROPODIDAE) IN THE ANAMALAI HILLS, TAMIL NADU, INDIA

Claire F.R. Wordley¹, Eleni K. Foui², Divya Mudappa³, Mahesh Sankaran⁴ & John D. Altringham⁵

^{1,2,4,5} School of Biology, University of Leeds, Leeds LS2 9JT, UK

^{2,4} National Centre for Biological Sciences, Tata Institute of Fundamental Research, GKVK, Bellary Road, Bengaluru, Karnataka 560065, India

^{2,3} Nature Conservation Foundation, 3076/5, 4th Cross, Gokulam Park, Mysuru, Karnataka 570002, India

¹ c.wordley@live.com (corresponding author), ² e_foui@yahoo.com,

³ divya@ncf-india.org, ⁴ mahesh@ncbs.res.in,

⁵ j.d.altringham@leeds.ac.uk

This cave roosting species has been found at altitudes of 800–1,100 m, and was thought to occupy an area of about 1,100km², based on the location of known roosts (Vanitharani et al. 2004; Molur & Vanitharani 2008). It has been found in montane tropical evergreen forest and coffee and cardamom plantations (Molur et al. 2002; Vanitharani et al. 2004). Based on remains in their day-roosts, *L. salimalii* is thought to feed on *Prunus ceylanicus*, *Ficus glomerata*, *F. macrocarpa*, *F. beddomei*, *Elaeocarpus serratus*, *E. tuberculatus*, *E. oblongus*, *Diospyros ovaliflora* and *Dichapetalum gelonioides* (Singaravelan & Marimuthu 2003; Vanitharani et al.

DOI: <http://dx.doi.org/10.11609/jott.2796.8.12.9486-9490> | ZooBank: urn:lsid:zoobank.org:pub:A540CEF7-D751-42A7-8245-59A5AD0DBD7C

Editor: P.O. Nameer, Kerala Agricultural University, Thrissur, India.

Date of publication: 26 October 2016 (online & print)

Manuscript details: Ms # 2796 | Received 17 May 2016 | Final received 18 June 2016 | Finally accepted 01 September 2016

Citation: Wordley, C.F.R., E.K. Foui, D. Mudappa, M. Sankaran & J.D. Altringham (2016). Range extension of the endangered Salim Ali's Fruit Bat *Latidens salimalii* (Chiroptera: Pteropodidae) in the Anamalai Hills, Tamil Nadu, India. *Journal of Threatened Taxa* 8(12): 9486-9490; <http://dx.doi.org/10.11609/jott.2796.8.12.9486-9490>

Copyright: © Wordley et al. 2016. Creative Commons Attribution 4.0 International License. JoTT allows unrestricted use of this article in any medium, reproduction and distribution by providing adequate credit to the authors and the source of publication.

Funding: Funding was provided by CEPF (Critical Ecosystems Partnership Fund), IUCN-NL Ecosystems Grants Programme, NERC (Natural Environment Research Council) (NE/I528393/1), the British Ecological Society (4863 / 5903), the University of Leeds, and UKIERI (UK India Education and Research Initiative).

Conflict of Interest: The authors declare no competing interests.

Acknowledgements: We thank the Tamil Nadu Forest Department, the Tamil Nadu Electricity Board and managers of Peria Karamalai Tea Company, Bombay Burmah Trading Corporation, Tata Coffee Ltd. and Parry Agro Industries Ltd., for permissions and local support. We would like to thank the field assistants Satish Kumar A., Dinesh T., Pandi, Sunderaj, Karthik, Kannan C., Anand Raj and Anand Kumar. We are also grateful to Emma Rigby, Sarah Proctor, Aurelie Laurent, Ruth Angell, Kate Parker, Swati Sidhu, Dina Nisthar and Aditya Malgaonkar for their help in the field. Many thanks to R. Raghunath for mapping the locations where *L. salimalii* has been found. Finally, we thank T.R. Shankar Raman, M. Ananda Kumar, and Ganesh Raghunathan for logistical support and advice throughout the project.

CRITICAL ECOSYSTEM
PARTNERSHIP FUND



BRITISH
ECOLOGICAL
SOCIETY

NERC
SCIENCE OF THE
ENVIRONMENT

UKIERI
UK-India Education
and Research Initiative




Figure 1. Location of the Anamalai Tiger Reserve, new locations for *Latidens salimalii* and previously reported locations. Map by R. Raghunath.

2004; Agoramoorthy & Hsu 2005).

The study was conducted in the Valparai plateau in the state of Tamil Nadu in the southern Western Ghats, an area of approximately 220km² (10.2–10.4°N & 76.8–77.0°E), and in the adjacent Anamalai Tiger Reserve (958km², 10.12–11.07°N & 76.0–77.56°E) (Fig. 1). The vegetation is classified as mid-elevation tropical wet evergreen forest of the *Cullenia exarillata*—*Mesua fërrea*—*Palaquium ellipticum* type (Pascal 1988; Raman

et al. 2009). The Valparai plateau is an agricultural landscape dominated by tea plantations interspersed with shade grown coffee plantations, eucalyptus plantations, tropical rainforest fragments, streams, and riverine vegetation (Mudappa & Raman 2007). The site is adjacent to the Anamalai Tiger Reserve in Tamil Nadu, Eravikulam National Park, Vazhachal Reserve Forest and Parambikulam Tiger Reserve in Kerala. Elevation ranges from approximately 800–1600 m, and the

Table 1. Locations of *Latidens salimalii* found in the Anamalai range.

Site ID	Date of capture	Latitude	Longitude	Sex	Age	Weight (g)	Forearm (mm)	Reproductive status	Notes
1	17.v.2009	76.837109	10.305527	M	J	68.5	68.3	NSA	Lots of parasites
2	12.ii.2012	76.897300	10.268458	M	A	52	69.8	NSA	Lots of parasites
3	11.xii.2014	76.91542	10.37059	F	A	60	68.8	PL	

Site 1 - Candura; Site 2 - Sangli Road; Site 3 - Anali; NSA - Bat does not appear to be currently sexually active, i.e. testes are not swollen; PL - Bat has previously lactated.

average annual rainfall is 3500mm, of which about 70% falls during the southwest monsoon (June–September) (Raman et al. 2009).

EKF caught bats across the Valparai plateau using ground level mist nets from 2008 until 2010. From 2011–2014, CFRW caught bats in mist nets and harp traps in forest fragments, coffee, tea and cardamom plantations, along rivers, and at roosts in tunnels and caves. CFRW caught from February to May in 2011–2013, and from November to December 2014 inside the Anamalai Tiger Reserve with appropriate permits. All bats caught were identified using the most up to date guides and then released on site within 15 min of capture (Bates & Harrison 1997; Srinivasulu et al. 2010). Bats were caught in accordance with Natural England protocol, and their welfare was of the highest priority at all times (http://www.naturalengland.org.uk/Images/wmlg39_tcm6-35872.pdf). Forearm measurements were taken to the nearest 0.1mm using dial calipers. Bats were weighed in cotton bags using spring balances, with weights to the nearest 0.5g.

One individual of *L. salimalii* was caught by EKF in 2009 and two by CFRW, in 2012 and 2014 (Table 1). All three sites were riparian locations bordered on both sides by forest. Site 1, Candura, is an abandoned vanilla plantation separated by a river and a reservoir from the Vazhachal Reserve Forest, now a naturally regenerating site with some restoration efforts. Site 2, Sangli Road forest fragment, is a large (102.8 ha) forest fragment less than 1km from the Vazhachal Reserved Forest, with a small stream running through it. The bat was caught in a mist net set over the stream. Site 3, Anali, is located inside the Anamalai Tiger Reserve, within 1km of the border with a tea plantation. The bat was caught in a mist net set over a small river.

All bats caught fitted the recorded characteristics of *L. salimalii*; most notably all showed the single pair of upper incisors that distinguishes *Latidens* from all other species of fruit bat known from the Indian subcontinent (Bates & Harrison 1997; Srinivasulu et al. 2010). The forearm measurements (68.3–69.8 mm) were similar to those recorded for this species in the High Wavy Mountains (66.0–69.0 mm) (Bates & Harrison 1997;

Srinivasulu et al. 2010). Unlike the other fruit bats in the area (*Cynopterus brachyotis* and *Rousettus leschenaultii*), *L. salimalii* has no external tail (Image 1) (Bates & Harrison 1997; Srinivasulu et al. 2010). It is also smaller than *R. leschenaultii* (forearm 75.0–86.0 mm), but larger than *C. brachyotis* (forearm 57.3–63.3 mm) (Srinivasulu et al. 2010). The underside of the wings and interfemoral membranes had some fur in all individuals.

Finding this species in the Anamalai Tiger Reserve and Valparai plateau is not unexpected, given the relative proximity of this location to the Meghamalai Wildlife Sanctuary and the similarity in habitat type and altitude to the other locations where *L. salimalii* has been found. However, it is important to note the presence of an endemic, Endangered, Schedule I species in a new locality. Two of the locations are on private lands, where the forest fragments are not legally protected, although imminent destruction of these forests is unlikely. Despite being reported as roosting in colonies of hundreds (Vanitharani et al. 2004; Molur & Vanitharani 2008), we caught only three individuals during seven years of seasonal mist-netting and harp trapping in the area, so it is likely that it is not locally common. That we mostly caught it over water is likely to be due to it flying lower when coming to drink, and thus being easier to catch than when foraging.

L. salimalii is described as being threatened by hunting for local consumption in traditional medicine, human disturbance of roosting sites and tree cutting in coffee estates (Molur et al. 2002; Singaravelan & Marimuthu 2003; Vanitharani et al. 2004). As all other fruit bats in India are listed as vermin, it is likely that few people are aware of the protected status (Schedule I) of *Latidens salimalii*. Local education on the status of *L. salimalii* where it is found may be beneficial; but a deep change in the way that bats are viewed overall in India may be needed to protect these rare endemics.

References

- Agoramoorthy, G. & M. Hsu (2005). Population size, feeding, forearm length and body weight of a less known India fruit bat, *Latidens salimalii*. *Current Science* 88(3): 354–356.
- Bates, P.J.J. & D.L. Harrison (1997). *Bats of the Indian Subcontinent*.



Image 1. Photographs of all three bats caught. Note the lack of external tail. Photo credit: Bat one by EKF, bat two by CFRW, bat three by A. Satish Kumar.

Harrison Zoological Museum Publications, 258pp.

Ghosh, M.K., T.P Bhattacharyya & S.S. Saha (1999). Occurrence of Salim Ali's Fruit Bat (*Latidens salimalii* Thonglongya, 1972) in the Kalakkad-Mundanthurai Tiger Reserve, Tamil Nadu. *Tigerpaper* 26(2): 32.

Molur, S., G. Marimuthu, C. Srinivasulu, S. Mistry, A.M. Hutson, P.J.J. Bates, S. Walker, K.P. Priya & A.R.B. Priya (2002). Status of South Asian Chiroptera. Conservation Action Management Plan (CAMP) Workshop Report, Zoo Outreach Organisation, 320pp.

Molur, S. & J. Vanitharani (2008). *Latidens salimalii*. The IUCN Red List of Threatened Species 2008: e.T11374A3274238. Downloaded on 08 September 2016; <http://dx.doi.org/10.2305/IUCN.UK.2008>.

[RLTS.T11374A3274238.en](https://doi.org/10.1111/j.1526-RLTS.T11374A3274238.en)

Mudappa, D. & T.R.S. Raman (2007). Rainforest restoration and wildlife conservation on private lands in the Western Ghats, pp. 210–240. In: Shahabuddin, G. & M. Rangarajan (eds.). *Making Conservation Work*. Permanent Black, Ranikhet, India, 312pp.

Pascal, J.P. (1988). *Wet Evergreen Forests of The Western Ghats of India: Ecology, Structure, Floristic Composition and Succession*. Pondicherry Institute, Français Pondichéry, 345pp.

Raman, T.R.S., D. Mudappa & V. Kapoor (2009). Restoring rainforest fragments: survival of mixed-native species seedlings under contrasting site conditions in the Western Ghats, India. *Restoration Ecology* 17(1): 137–147; <http://dx.doi.org/10.1111/j.1526->

100X.2008.00367.x

- Singaravelan, N. & G. Marimuthu (2003).** Mist net captures of the rarest fruit bat *Latidens salimalii*. *Current Science* 84(1): 101–103.
- Srinivasulu, C., P.A. Racey & S. Mistry (2010).** A key to the bats (Mammalia: Chiroptera) of South Asia. *Journal of Threatened Taxa* 2(7): 1001–1076; <http://dx.doi.org/10.11609/JoTT.o2352.1001-76>
- Thonglongya, K. (1972).** A new genus and species of fruit bat from South India (Chiroptera: Pteropodidae). *Journal of the Bombay Natural History Society* 69(1): 151–158.
- Vanitharani, J., U.S.U. Malathi & K.A. Sundari (2005).** New records of bats from Kalakad Mundanthurai Tiger Reserve, India. *BatNet CCINSA Newsletter* 6(1): 13–14.
- Vanitharani, J., M. Pearch, L.J. Praba & R. Annamalai (2004).** A review of the distribution and status of *Latidens salimalii* (Chiroptera: Pteropodidae) with new records from the Western Ghats, India. *Lutra* 47(1): 21–32.





OPEN ACCESS



All articles published in the Journal of Threatened Taxa are registered under Creative Commons Attribution 4.0 International License unless otherwise mentioned. JoTT allows unrestricted use of articles in any medium, reproduction and distribution by providing adequate credit to the authors and the source of publication.

ISSN 0974-7907 (Online); ISSN 0974-7893 (Print)

October 2016 | Vol. 8 | No. 12 | Pages: 9397–9504
Date of Publication: 26 October 2016 (Online & Print)

DOI: 10.11609/jott.2016.8.12.9397-9504

www.threatenedtaxa.org

Article

Identifying orchid hotspots for biodiversity conservation in Laos: the limestone karst vegetation of Vang Vieng District, Vientiane Province

-- Pankaj Kumar, Stephan W. Gale, André Schuiteman, Somsanith Bouamanivong & Gunter A. Fischer, Pp. 9397–9417

Communications

On the occurrence of Common Baron (Lepidoptera: Nymphalidae: Limenitidinae: *Euthalia aconthea* Cramer, 1777) in the Delhi area and analysis of abiotic factors affecting its distribution in India

-- Rajiv K. Singh Bais, Pp. 9418–9433

Diversity and seasonality of polypore fungi in the moist deciduous forests of Peechi-Vazhani Wildlife Sanctuary, Kerala, India

-- A. Muhammed Iqbal, Kattany Vidyasagaran & P. Narayan Ganesh, Pp. 9434–9442

Short Communications

Camera trapping the Palawan Pangolin *Manis culionensis* (Mammalia: Pholidota: Manidae) in the wild

-- Paris N. Marler, Pp. 9443–9448

Migratory Pallas's Gull *Larus ichthyaetus* (Pallas, 1773): a new record from Sikkim, the eastern Himalaya, India

-- Santosh Sharma & Dinesh Bhatt, Pp. 9449–9453

An inventory of herpetofauna from Wadi Sayq, Dhofar, Oman

-- Lawrence Derek Ball & James Stefan Borrell, Pp. 9454–9460

Species diversity and spatial distribution of snakes in Jigme Dorji National Park and adjoining areas, western Bhutan

-- Bal Krisna Koirala, Dhan Bdr Gurung, Phurba Lhendup & Sonam Phuntsho, Pp. 9461–9466

New records of petiolate potter wasps (Hymenoptera: Vespidae: Eumeninae) from Bhutan

-- Tshering Nidup, Thinley Gyeltshen, P. Girish Kumar, Wim Klein & Phurpa Dorji, Pp. 9467–9472

Recent records of the Pale Jezebel *Delias sanaca sanaca* (Moore, 1857) (Lepidoptera: Pieridae) from Mussoorie hills, western Himalaya, India

-- Arun P. Singh, Pp. 9473–9478

An observation on the fruit feeding behavior of butterflies in some areas of Bangladesh

-- Tahsinur Rahman Shihan, Pp. 9479–9485

Notes

Range extension of the endangered Salim Ali's Fruit Bat *Latidens salimalii* (Chiroptera: Pteropodidae) in the Anamalai Hills, Tamil Nadu, India

-- Claire F.R. Wordley, Eleni K. Foui, Divya Mudappa, Mahesh Sankaran & John D. Altringham, Pp. 9486–9490

A checklist of butterflies of Dakshina Kannada District, Karnataka, India

-- Deepak Naik & Mohammed S. Mustak, Pp. 9491–9504