



UNIVERSITY OF LEEDS

This is a repository copy of *Evaluating the impact of trigeminal neuralgia*.

White Rose Research Online URL for this paper:

<http://eprints.whiterose.ac.uk/110714/>

Version: Accepted Version

Article:

Zakrzewska, JM, Wu, J orcid.org/0000-0001-6093-599X, Mon Williams, M et al. (2 more authors) (2017) Evaluating the impact of trigeminal neuralgia. *Pain*, 158 (6). pp. 1166-1174. ISSN 0304-3959

<https://doi.org/10.1097/j.pain.0000000000000853>

© 2017, International Association for the Study of Pain. This is an author produced version of a paper published in *Pain*. Uploaded in accordance with the publisher's self-archiving policy.

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>

PAIN

Evaluating the impact of trigeminal neuralgia.

--Manuscript Draft--

Manuscript Number:	PAIN-D-14-15450R2
Article Type:	Clinical Note
Section/Category:	Clinical
Keywords:	trigeminal neuralgia , disability, quality of life, sociodemographic
Corresponding Author:	Joanna Maria Zakrzewska, MD, FDSRCS, FFDRCSI, FFPMRCA,FHEA Eastman Dental Hospital London, UNITED KINGDOM
First Author:	Joanna Maria Zakrzewska, MD, FDSRCS, FFDRCSI, FFPMRCA,FHEA
Order of Authors:	Joanna Maria Zakrzewska, MD, FDSRCS, FFDRCSI, FFPMRCA,FHEA Jianhua Wu, BSc, PhD Mark Mon Williams, PhD Nick Phillips, BSc, PhD, FRCS, FRCS (SN) Sue Pavitt, BSc,PhD
Abstract:	<p>There is a lack of prospective systematic studies on the clinical characteristics of pain in trigeminal neuralgia (TN) as well as its 'psychosocial burden'.</p> <p>Patients with idiopathic TN were categorised into three sub-types (n = 225). Group 1 (n= 155, 68.9%) had TN without concomitant pain, Group 2 (n=32, 14.2%) had TN with intermittent concomitant pain and Group 3 (n=39, 16.9%) had TN with autonomic symptoms. We tested two hypotheses: (i) that different pain profiles would be associated with the different groups; (2) that the severe pain associated with TN would impact negatively on activities of daily living and thereby result in disability as defined by the World Health Organisation. A different pain profile was found across the groups. We obtained unequivocal evidence that TN causes disability with up to 45% of patients being absent from usual daily activities 15 days or more in the past 6 months. On the Hospital Anxiety and Depression Scale, 35.7% patients had mild to severe depression and over 50% were anxious. The Pain Catastrophizing Scale showed that 78% of patients had considerable negative thoughts with scores > 20 and a mean score of 36.4. Prior to referral only 54% had been prescribed carbamazepine whilst opioids had been prescribed in 14.6% of the patients. Prior to referral over 80% had already been to one specialist centre which had not provided appropriate management. Patients with TN report varied characteristics but all result in some degree of psychosocial disability especially before adequate therapy is attained .</p>

We have deleted the word significant from the last sentence in the abstract and we have added this phrase **in patients with trigeminal neuralgia and its variants** to tables 5 and 6.

Impact of TN

Abstract

There is a lack of prospective systematic studies on the clinical characteristics of pain in trigeminal neuralgia (TN) as well as its 'psychosocial burden'.

Patients with idiopathic TN were categorised into three sub-types (n = 225). Group 1 (n= 155, 68.9%) had TN without concomitant pain, Group 2 (n=32, 14.2%) had TN with intermittent concomitant pain and Group 3 (n=39, 16.9%) had TN with autonomic symptoms. We tested two hypotheses: (i) that different pain profiles would be associated with the different groups; (2) that the severe pain associated with TN would impact negatively on activities of daily living and thereby result in disability as defined by the World Health Organisation. A different pain profile was found across the groups. We obtained unequivocal evidence that TN causes disability with up to 45% of patients being absent from usual daily activities 15 days or more in the past 6 months. On the Hospital Anxiety and Depression Scale, 35.7% patients had mild to severe depression and over 50% were anxious. The Pain Catastrophizing Scale showed that 78% of patients had considerable negative thoughts with scores > 20 and a mean score of 36.4. Prior to referral only 54% had been prescribed carbamazepine whilst opioids had been prescribed in 14.6% of the patients. Prior to referral over 80% had already been to one specialist centre which had not provided appropriate management. Patients with TN report varied characteristics but all result in some degree of psychosocial disability especially before adequate therapy is attained.

Evaluating the impact of trigeminal neuralgia.

Joanna M. Zakrzewska*1; Jianhua Wu 2; Mark Mon Williams3;

Nick Phillips 4; Sue H. Pavitt2

*1 Corresponding author

Facial Pain Unit Eastman Dental Hospital,

UCLH NHS Foundation Trust

256 Gray's Inn Road

London WC1X 8LD

UK

tel: +44 (0) 20 345 61195

fax ; + 44 (0) 20 345 61105

email j.zakrzewska@ucl.ac.uk

UCL IRIS URL <https://iris.ucl.ac.uk/iris/browse/profile?upi=JZAKR77>

2 School of Dentistry, University of Leeds, LS2 9LU, UK

3 School of Psychology, University of Leeds, LS2 9JT, UK

4 Department of Neurosurgery, Leeds General Infirmary, Great George Street, LS1 3EX,UK

Key words: trigeminal neuralgia, quality of life, disability, sociodemographic

Running title: The impact of trigeminal neuralgia

Number of pages: 19

Number of tables: 6

Number of figures and legends: 3

Evaluating the impact of trigeminal neuralgia.

Joanna M. Zakrzewska*¹; Jianhua Wu ²; Mark Mon Williams³;

Nick Phillips ⁴; Sue H. Pavitt²

*¹ Corresponding author

Facial Pain Unit Eastman Dental Hospital,

UCLH NHS Foundation Trust

256 Gray's Inn Road

London WC1X 8LD

UK

tel: +44 (0) 20 345 61195

fax ; + 44 (0) 20 345 61105

email j.zakrzewska@ucl.ac.uk

UCL IRIS URL <https://iris.ucl.ac.uk/iris/browse/profile?upi=JZAKR77>

2 School of Dentistry, University of Leeds, LS2 9LU, UK

3 School of Psychology, University of Leeds, LS2 9JT, UK

4 Department of Neurosurgery, Leeds General Infirmary, Great George Street, LS1 3EX,UK

Key words: trigeminal neuralgia, quality of life, disability, sociodemographic

Running title: The impact of trigeminal neuralgia

Number of pages: 19

Number of tables: 6

Number of figures and legends: 3

Abstract

1 There is a lack of prospective systematic studies on the clinical characteristics of pain in
2 trigeminal neuralgia (TN) as well as its 'psychosocial burden'.
3

4
5 Patients with idiopathic TN were categorised into three sub-types (n = 225). Group 1 (n=
6 155, 68.9%) had TN without concomitant pain, Group 2 (n=32, 14.2%) had TN with
7 intermittent concomitant pain and Group 3 (n=39, 16.9%) had TN with autonomic
8 symptoms. We tested two hypotheses: (i) that different pain profiles would be associated
9 with the different groups; (2) that the severe pain associated with TN would impact
10 negatively on activities of daily living and thereby result in disability as defined by the
11 World Health Organisation. A different pain profile was found across the groups. We
12 obtained unequivocal evidence that TN causes disability with up to 45% of patients being
13 absent from usual daily activities 15 days or more in the past 6 months. On the Hospital
14 Anxiety and Depression Scale, 35.7% patients had mild to severe depression and over
15 50% were anxious. The Pain Catastrophizing Scale showed that 78% of patients had
16 considerable negative thoughts with scores > 20 and a mean score of 36.4. Prior to
17 referral only 54% had been prescribed carbamazepine whilst opioids had been prescribed
18 in 14.6% of the patients. Prior to referral over 80% had already been to one specialist
19 centre which had not provided appropriate management. Patients with TN report varied
20 characteristics but all result in some degree of psychosocial disability especially before
21 adequate therapy is attained.
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1. Introduction

Trigeminal neuralgia (TN) is defined by the International Association for the Study of Pain (IASP) as “a sudden, unilateral, severe brief stabbing recurrent pain in the distribution of one or more branches of the fifth cranial nerve”¹⁹. The disorder is often misdiagnosed, being commonly confused with toothache or temporomandibular disorders, with reports of patients undergoing unnecessary, aggressive and irreversible dental treatments before obtaining a correct diagnosis^{6,29;30}. The estimated misdiagnosis by general practitioners could be as high as 48% which led to a ‘neurologist validated’ diagnosis incidence rate of 12.6 per 100,000 person years in the Netherlands¹¹.

Although a relatively rare condition, the impact on the lives of patients is profound^{1, 31}. Allsop et al¹ provided a qualitative study suggesting TN places a large burden on patients and TN is known to be associated with severe pain¹⁶. Nevertheless, there are few studies that have determined the impact of the pain on the lives of the affected individuals. This is highly problematic as it prevents proper evaluation of the health economics associated with existing treatment for these patients. In turn, this prevents evaluation of alternative treatment regimens. We hypothesised that the pain associated with TN would cause a high degree of disability (as defined by the World Health Organisation, WHO) within the population. The framework described in the WHO’s international classification of functioning, disability and health (ICF) outlines how impairment (e.g. pain) relates to disability. The ICF identifies three levels: the body function and structure level; the activity level and the participation level (WHO 2001). The ICF framework suggests that a functional deficit does not automatically result in an activity limitation or participation restriction. Rather, these different levels are proposed to interact with each other in a complex manner (mediated by personal and family factors). There are three studies that support the general idea that the pain associated with TN will lead to disability. Tolle et al²⁴ found a significant impact of TN on activities of daily living in their study of 82 patients, mean pain interference of 3.6 ± 2.4 . Wu et al²⁷ reported that TN was associated with increased depression 2.2%, anxiety 1.8% and sleep disturbance 1.2% in a retrospective study of 3273 TN patients compared to controls 13,092 based on health insurance claims databases. Mačianskytė et al¹⁷ identified disability, anxiety and depression in 30 to 47% of 30 TN patients.

1 The aim of the current study was to collect detailed clinical characteristics from patients
2 attending a TN clinic. We wished to determine the burden of TN at the point of referral to the
3 specialist centre. Our goal was explore differences between three groups of patients. Group 1
4 - Type 1 or ‘classical’ without concomitant persistent pain, irrespective of neurovascular
5 compression findings; Group 2 - Type 2 with intermittent concomitant pain often associated
6 with ‘after-pain’ following an attack ²¹ ; Group 3 - TN with autonomic symptoms which are
7 either present consistently or intermittently. We wanted to explore whether these different
8 classifications produced different pain profiles (thereby supporting the clinical utility of this
9 categorisation).
10
11
12
13
14
15
16

17 **2. Material and Methods**

21 2.1 Study subjects

22 We enrolled 237 consecutively consenting patients with TN (and its variants - short-lasting
23 unilateral neuralgiform headache attacks with autonomic system symptoms (SUNA) or short-
24 lasting unilateral neuralgiform headache attacks with conjunctival injection and tearing
25 (SUNCT)) into a cohort study between April 2007 and December 2015. The patients were all
26 attending a national Facial Pain Unit within a London teaching hospital. Patients with
27 multiple sclerosis or tumour related TN (n = 12) were excluded but those who had undergone
28 previous surgery for TN were included if their pain was the same as pre-operatively. The
29 resulting cohort (n = 225) undertook a baseline assessment that comprised both physician
30 measures and patient-completed questionnaires.
31
32
33
34
35
36
37
38
39
40

41 2.2 Ethics approval

42 The project had ethical approval and all patients were provided with an information sheet and
43 gave written consent according to the Declaration of Helsinki. The study was approved by
44 South East Research Ethics Committee REC Reference Number 07/MRE01/38.
45
46
47
48
49
50

51 2.3 Patient history of TN

52 Basic demographic data including ethnicity and social status were recorded. A note was made
53 regarding the specialists seen prior to attendance. The medical history was ascertained from
54 the primary care physician (GP) and particular note was made of conditions associated with
55 TN and the presence of headaches, migraines and other chronic pain. All current and past
56 treatments for TN were recorded including maximum doses and, where possible, efficacy and
57
58
59
60
61
62
63
64
65

The impact of Trigeminal Neuralgia

1 tolerability. Patients were asked about any history of bruxism (teeth grinding), jaw clenching
2 habits and jaw clicking. Examination included a gross cranial nerve examination and sensory
3 testing with light touch, cotton wool and pin-prick. The muscles of mastication were
4 examined for tender spots to determine if a musculoskeletal temporomandibular disorder was
5 present and an oral examination was done.
6
7
8
9

10 A history of the patient's experience was collected: duration; onset, including whether this
11 was acute and memorable (i.e. could they remember the circumstances of the first attack);
12 length of attacks; whether these were single stabs, a series of stabs in quick succession or
13 more continuous 'saw tooth' ⁵; whether any 'after pain' remained and its duration and
14 characteristics; remission periods and their length; provoking factors (including whether
15 attacks were only evoked or could be spontaneous). The pain location for the third division
16 was noted and whether this remained only in the lower part of the face ie. Pre-auricular area
17 down to the mental area or extended to the temporal region.
18
19
20
21
22
23
24
25
26

27 2.4 Patient questionnaires

28 Pain ascertainment

29 Patients completed the following questionnaires at their first visit: The Brief Pain
30 Inventory ^{4,13} which uses a Lickert (1-10) scale to determine pain intensity and quality
31 of life (an extended one was introduced later); The Graded Chronic Pain Scale (GGPS)
32 ²⁶ which uses a Lickert (1-4) scale to determine pain over last six months; The McGill
33 pain questionnaire, where users select words that best describe their pain ¹⁸; The Pain
34 Catastrophizing Scale (PCS) ²³ to assess catastrophizing (scores > 20 deemed
35 'significant').
36
37
38
39
40
41
42
43
44

45 Depression and anxiety ascertainment

46 The Hospital Anxiety and Depression scale (HAD) ³² scores > 8 significant replaced the
47 DAPOS to assess depression which was used for the first year only; plus two NICE
48 questions to screen for depression ²².
49
50
51
52

53 Data collection

54 Each patient was assigned a unique study ID number. A standardised data extraction form
55 and simple UCLH/ University of Leeds TN patient data base was developed to facilitate the
56 data collection. The patient data were transferred from the paper case report form into the TN
57
58
59
60
61
62
63
64
65

1 database (quality of inputting audited). Patients were followed up via routine clinical
2 consultations and self-completed questionnaires. All patients were seen by one clinician (JZ),
3 with 58 (26%) also seen by a headache neurologist to independently validate the diagnosis.
4 All patients had an MRI to exclude a symptomatic cause and identify potential neurovascular
5 compression, some of which had been done prior to referral. Patients were managed
6 according to the Facial Pain Unit protocol, similar to the recently published Danish protocol
7
8
9
10
11⁹. Questionnaires were collected at each routine visit.
12
13

14 2.5 Statistical Analysis

15 A case series analysis was performed. Summary statistics were used to describe the sample:
16 means and standard deviations were provided for continuous variables and frequency
17 distributions for categorical variables. One-way ANOVAs were used for continuous
18 outcomes and chi-squared tests were used for categorical outcomes (significance level set at
19 5%). Pain severity was categorized as mild, moderate, or severe.
20
21
22
23
24
25
26

27 3. Results

28
29
30 A total of 237 patients were referred to the clinic between 2007 and 2015. Twelve of these
31 patients (5.1%) had TN secondary to other causes and were excluded from further analysis.
32
33 Nine of these 12 patients (6 women and 3 men) had multiple sclerosis before the onset of
34 their TN, three had tumours.. The remainder (n = 225) were divided into three groups: Group
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000

Table 1 provides details of type of surgery they had and its effect.

[TABLE 1 AROUND HERE]

3.1 Baseline characteristics

The baseline demographic and clinical characteristics of the 225 patients are presented in Table 2. The mean (SD) age was 60.9 (12.5) years, and the median age [IQR] at first attack was 57.0 [46.0, 65.0] years. The median duration of TN was 4.0 [2.0, 7.0] years. There were

1 no statistically significant differences in symptom duration or age between the three groups.
2 There was a predominance of females (63.6%) within the population. There was a bias
3 towards a higher socioeconomic position within the population, with over 70% of the patients
4 having an index of multiple deprivation of 3 or less (1 = least deprived and 5 = most
5 deprived). Two thirds of the patients (63.7%) had managerial or professional occupations
6 according to the standard occupational classification (Office for National Statistics 2010).
7
8
9

10
11
12 [TABLE 2 AROUND HERE]
13
14
15

16 A family history of TN was reported by 13 (5.8%) of the patients. Nearly half of the patients
17 (45.7%) were referred to the clinic by a primary care medical practitioner (GP), a fifth were
18 referred by a dentist with the rest referred by specialists. There was not a statistically higher
19 incidence of lower facial pain in those referred by a dentist. Prior to referral, the majority of
20 the patients had already consulted a GP (80.0%) or dentist (71.6%). Many had used
21 secondary dental (42.2%) or medical (58.7%) services and 32.5% had seen two or more
22 dental or medical specialists. One hundred and fifteen patients (87.1%) had seen a
23 neurosurgeon or neurologist or both.
24
25
26
27
28
29
30

31 32 33 3.2 Medical history and oral health

34 A quarter of patients had headaches and 44 had migraines or migraines with tension type
35 headache (10 of these were in Group 3). Patients in all groups had a similar incidence of
36 hypertension (37.3%) and other cardiovascular diseases (12.4%) and those proportions were
37 lower than the population prevalence (hypertension 52.0% and cardiovascular disease 22%)
38 of similar age (table 3). Fifty four patients (24.0%) reported bruxism but the frequency was
39 not different between groups. The quality of oral hygiene was evenly distributed among poor
40 (31.5%), moderate (28.9%) and good (31.5%) quality for 108 patients who were examined.
41 Two hundred and eleven patients had partial/full dentition and only 24 (14.3%) patients had
42 little conservation but all had evidence of some dental treatment. Oral health was not
43 associated either with the duration of TN or pain severity.
44
45
46
47
48
49
50
51
52
53

54 [TABLE 3 AROUND HERE]
55
56
57

58 3.3 Pain characteristics 59 60 61 62 63 64 65

The impact of Trigeminal Neuralgia

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

The pain locations for Group 1 (n = 155) and Group 2 (n = 32) were similar in all trigeminal divisions. In contrast, those patients in Group 3 (n = 38) had a significantly lower proportion of pain in the first division of the trigeminal nerve and third division (0% and 10.3%) but a higher proportion of pain in the second division (Table 4). Right-sided TN (65.6%) was twice as prevalent as left-sided TN (32.5%), while 1.9% of the patients had bilateral TN. All patients had extra-oral pain but 79.9% also reported intra-oral pain (whereas 68.6% reported only extra-oral pain) across all type of TN.

[TABLE 4 AROUND HERE]

Patients reported a variety of different types of attacks. The predominant type of attack was a single stab for all groups but a series of stabs was also frequently reported in Group 1 (36.4%) and Group 2 (31.3%) and patients would also report having a combination of these types. Pain severity was similar across the three types of TN. Less than 5% of patients in Group 1 and 3 could not recall the circumstances of the first attack while a fifth of Group 2 (79.3%) could not remember the circumstances (p-value = 0.001).

The pain frequency in TN is high with over 90% reporting daily pain attacks though only a small proportion reported attacks on an hourly basis. Very few patients reported attacks lasting more than a few minutes but 62% of Group 2 reported a prolonged after-pain.

Remission periods were reported consistently across the groups but were least likely to be in Group 3. Overall, the remission duration decreased with time. Nearly all patients had spontaneous pain but all had pain provoked by light touch on the face or intra-orally. A large proportion of patients across all TN types (30-60%; p-value = 0.05) could be provoked by cold wind or bodily movement. Four patients had attacks that could be provoked by noise or light and one patient had attacks that could be provoked by alcohol. Group 3 reported the most ipsilateral autonomic features (p-value = <0.001) and the most common ones were tearing, nasal stuffiness, redness of the cheek or eye, and these were all statistically different from the other groups (p-value <0.001) (see table 3). These were not observed but reported several times at follow ups.

The McGill pain questionnaire was fully completed by 193 of the 225 patients (85.8%). The words were analysed by sensory type (1-10), affective type (11-15), evaluative type (16) and

The impact of Trigeminal Neuralgia

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
miscellaneous type (17-20). The sensory words most commonly used by patients across all Groups were ‘shooting’ (84.5%), followed by ‘sharp’ (72.5%) and ‘stabbing’ (55.4%). More than 50% of patients choose a word from ‘fearful’, ‘frightful’ or ‘terrifying’. The evaluative and miscellaneous words most commonly chosen in all TN categories were ‘unbearable’ (45.1%) and ‘piercing’ (45.1%).

3.4 Treatment

Pharmacological management of TN was the most commonly reported treatment modality for TN. All patients received at least one medication for their facial pain before referral. A quarter of the patients (27.6%) had two medications while nearly half (46.7%) had used three or more medications. Prior to referral, the most frequently prescribed drug was carbamazepine (including retard type) 122/225 (54.2%) which also caused significant side effects. Opioids had been prescribed in 14.6% of the patients. Over 75% were put on anticonvulsants after the first consultation as shown in figure 1.

[FIGURE 1 AROUND HERE]

3.5 Impact of pain

Pain had a significant impact on health status, including functioning and well-being across all TN groups as shown in Table 4. On the NICE questions, 106/216 (49.1%) felt depressed and 101/216 (47.6%) had little pleasure in life. This correlated with the HAD results where 75/210 (35.7%) patients had mild to severe depression. More than 50% were anxious as indexed by the HAD. The Pain Catastrophizing Scale (PCS) showed that a significant proportion of patients with TN had considerable negative thoughts about their pain: 146/188 (77.7%) patients had a PCS score of 20 and over with a mean score of 36.4 (95% CI: 34.9-37.9).

Completion of the CGPS questionnaire (grading pain in the past 6 months) achieved 175/225 (77.8%) respondents. Ninety five of these patients (54.3%) had high and moderately or severely limiting disability (Grade III and IV). 89/198 (44.9%) patients were absent from usual daily activities (work, school or housework) 15 days or more in the past 6 months because of the facial pain. The CGPS was significantly associated with HAD -Anxiety ($p < 0.001$) and HAD -Depression ($p < 0.001$) using Pearson’s Chi-square test, table 5.

1
2 [TABLE 5 AROUND HERE]
3
4

5 On the Brief Pain Inventory (BPI)–Facial, reporting pain intensity and impact in the week
6 prior to assessment, two thirds (66.3%) of patients reported moderate (score of 4-7) or severe
7 (score of 7-10) overall pain within the prior 24 hours and the mean overall Pain Severity
8 Index scores for 194 patients was 3.9 (95% CI: 3.6-4.3), indicating a mild to moderate level
9 of pain. The pain severity was significantly associated with HAD-Depression ($p = 0.022$) but
10 not HAD -Anxiety ($p = 0.163$) using Pearson’s Chi-square test table 6.
11
12
13
14
15
16

17
18 [TABLE 6 AROUND HERE]
19
20

21 The mean pain interference on the BPI was higher (mean = 4.9) when the facial domains
22 were assessed. The pain interference was more salient for 110 patients who also reported pain
23 interference on seven face related daily activities (mean 4.9, 95% CI: 4.3-5.5). Pain severity
24 significantly affected general activity, mood and enjoyment of life (Figure 2). Sleep was
25 affected for those with severe pain, with thirty five patients reporting that pain affected their
26 sleep. Pain severely interfered with all daily activities, especially activities involving the face
27 (such as eating a meal, brushing or flossing teeth and eating hard food like apples) as seen in
28 Figure 3.
29
30
31
32
33
34
35

36
37
38 [FIGURE 2 AND 3 AROUND HERE]
39
40

41 42 **4. Discussion** 43 44

45 The reported research represents the largest and most comprehensive biopsychosocial
46 prospective study on TN that has been conducted to date.
47
48

49 50 51 4.1. Population characteristics 52

53 The study involved a population of patients with TN who attended a specialist clinic. In this
54 regard, the population may not be representative of the general TN population (as less severe
55 cases may not end up at such a specialist clinic). Indeed, the average socioeconomic position
56 of the population was skewed away from the greatest levels of deprivation. It seems likely
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

that TN (like most other diseases including multiple sclerosis) would be over represented in lower socioeconomic positions. Thus, the population reported within this paper are probably overrepresented with those who could utilise social capital to attend the clinic. Patients access many health care professionals prior to referral to a specialist clinic. These clinics can provide appropriate treatment but over 37% of those visiting a dentist get dental treatment which is often irreversible (such as root canal treatments and extractions) which contribute to the burden ¹⁶. Garvan and Siegfried ⁶ reported that in their series of 140 patients 67 patients had 680 teeth extracted. Patients with only intraoral pain may obtain continuing care from dental practitioners and not be referred whereas patients in Group 3 may be more complex and less responsive to carbamazepine and so referred in earlier. Repeated consultations and inappropriate use of medications (despite numerous guidelines) add to the economic costs to the providers and the patients.

4.2 TN variants

The participants were considered as falling within one of three groups. Group 1 had TN without concomitant pain, Group 2 had TN with concomitant pain and Group 3 had TN with autonomic symptoms. We hypothesised that the different groups would show disparate pain profiles. There were clear differences between the groups. Group 3 were more likely to have a lower proportion of pain in the lower part of the face and had prominent autonomic symptoms. Groups 1 and 2 were more likely to experience a series of stabbing pains, with Group 2 more likely to experience a prolonged after-pain and were less likely to recall the circumstances of their first attack. We observed in our cohort that 68.4% had TN without concomitant pain and 14.2% had TN with at least some concomitant pain and 30% had attacks lasting minutes rather than seconds – this was a distinctive difference to the Maarbjerger et al ¹⁵ cohort who had 49% with concomitant pain. This might be explained by differing clinical referral processes to a specialist headache neurology clinic compared to ours , a specialist facial pain clinic within a dental school¹⁵. Haviv et al ⁸ hypothesised that length of individual pain attacks correlates to presence of background pain - a similar finding to this cohort. They reported that 87% of those with attack duration of over 2 minutes (n= 20) had background pain whereas this was 30% in those with shorter attacks. Bowsher ³ reported that 50% of 50 patients who had prior surgery reported longer attack duration but, in the present study, the 10.2% who had undergone surgery were not significantly different from the rest of the population (except for having consulted more specialists).

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

In agreement with other studies, our work shows that autonomic features are often present and this is reflected in emerging data showing that SUNCT, SUNA and TN may be variants of the same disorder^{12,25}. Patients did report some altered sensation and this was highest in Group 2. This may be of importance as a recent study has suggested that classical TN (Group 1) may have sub-clinical hypoesthesia whereas those with concomitant pain (Group 2) were more likely to have more clinically detectable sensory changes but all had sensory changes not just on the side of pain but also on the opposite side and in other parts of the body suggesting central sensitisation²⁸. What still needs to be established is whether patients move between the different groups during the course of disorder.

4.3 Clinical characteristics

This study has provided useful information on the clinical characteristics of TN patients. Our results concur with a large UK population study that showed greater prevalence in women and a higher occurrence of right-sided symptoms and pain in second and third divisions that is consistent with the somatology relationship of sensory fibres in the trigeminal nerve Hall et al⁷. The study has highlighted that pain in the third division may only involve the lower branches and not the temporal branches does not always extend to the entire third division and is often present only in the lower division as proposed by Henderson¹⁰ who suggests a mouth –ear zone and nose-orbit zone. Other features, including the McGill pain descriptors, are similar to other studies reporting on phenotypic features^{3,8,16} and highlights the frequent use of words like “fearful” and “terrifying”. Fear could be a driver for the spontaneous pain rather than evoked pain as shown in a small fMRI study done with patients with classical TN, where patients were told that their pain would be triggered during the examination²⁰.

4.4 Management

Pharmacological management is the most commonly used treatment modality for patients with TN. The anticonvulsant class (particularly carbamazepine) is considered the gold standard treatment³⁰. It is clear though that a wide range of drugs are being provided to TN patients and over 75% have used more than one drug prior to referral with 21% having used four or more. The prescribed drugs include opioids despite the lack of any evidence that these drugs are effective in TN. Despite all the guideline recommendations only 54% of the patients were or had used carbamazepine which agrees with Hall et al’s⁷ survey of primary care management of TN (where the figure was 58%). An additional 37 patients were prescribed this drug after their attendance at the clinic and other non-effective drugs were

1 stopped. This suggests that large numbers of TN patients outside specialist clinics are not
2 being provided with the optimal drug regime which adds to the burden of their disease. The
3 effectiveness of carbamazepine/retard was witnessed by the fact that 91.8% of the patients
4 taking this drug reported that it was partially or completely effective. Unfortunately, 42.8% of
5 these patients reported significant side effects which is to be expected as all these drugs result
6 in side effects - especially cognitive side effects ².
7
8
9
10

11 12 4.5 Burden of disease

13 The results of the present study provide further empirical evidence that patients with TN
14 suffer considerable pain and disability – even when prescribed the optimal drug regimen. We
15 hypothesised that the high levels of pain would produce disability as defined by the WHO’s
16 International Classification of Disability. The data showed unambiguously that patients with
17 TN experience considerable activity limitation and the limitation is particularly pronounced
18 with activities that involve the face (e.g. eating a hard piece of food, such as an apple).
19 Moreover, there was a relationship between pain and the ability of the patients to participate.
20 We found evidence that over 50% of the patients had to take significant time off work which
21 has a significant economic impact as the median age for the start of the TN is 57 and over
22 50% are still in employment. This has never previously been reported. The disability
23 experienced by patients with TN is consistent with the high levels of anxiety and depression
24 recorded within this population. Fear, unpredictability of the pain attacks and lack of
25 confidence in dealing with flare ups results in high catastrophizing scores.
26
27
28
29
30
31
32
33
34
35
36
37
38
39

40 The quantitative data in the present study are consistent with the qualitative data reported in
41 Allsop et al ¹. The overall picture suggests a population who have a high chance of
42 experiencing excruciating pain, leading to activity limitation and participation restriction.
43 These findings suggest that patients with TN should be offered psychological support –this is
44 not routinely provided at present. A multidisciplinary approach for pain management
45 (including, for example, Cognitive Behavioural Therapy) can be effective for pain control,
46 developing coping skills and restoration of functional status. It is important that patients with
47 TN are provided with an individualised pain management programme that is not limited to
48 pharmacological or surgical interventions alone.
49
50
51
52
53
54
55
56
57

58 **5. Conclusions**

1 We conclude that patients with TN are experiencing a poor quality of life even when being
2 treated with the optimal drug regimen. This suggests an urgent need to evaluate alternative
3 treatment pathways (such as newer medications, psychological support, earlier surgical
4 interventions and access to newer surgical innovations such stereotactic radiosurgery,
5 neuromodulation). For example, Lee et al ¹⁴ have suggested that surgical management can
6 significantly improve quality of life in patients with TN when assessed using the Brief Pain
7 Inventory –Facial. The data presented in the current manuscript provides a starting point for
8 evaluations of the true health burden and cost of TN.
9
10
11
12
13
14

15 **Acknowledgments**

16 We thank the patients with TN and Teresa Belai who inputted the data. For input on
17 selection of questionnaires we are grateful to:, Dr Liam Hill , Dr Matthew Allsop, Prof Jenny
18 Hewison, Ms Carolyn Czoski Murray & Dr Maureen Twiddy. We thank Dr Paul Baxter and
19 Thomas Fleming for constructing the patient database and providing advice on analysis.
20
21
22
23
24
25
26

27 **Funding**

28 JZ undertook the work at UCL/UCLHT who received a proportion of funding from the
29 Department of Health's NIHR Biomedical Research Centre funding scheme. SP, NP and
30 MMW received funding from the University of Leeds Biomedical Health Research Centre,
31 Problem Solving funding scheme to support a collaboration with JZ and the development of
32 the UCHL/UoL prospective TN patient cohort database and some data entry.
33
34
35
36
37
38
39

40 **Conflict of interest:** The authors have no conflict of interest to declare
41
42
43
44

45 **Reference List**

- 46 [1] Allsop MJ, Twiddy M, Grant H, Czoski-Murray C, Mon-Williams M, Mushtaq F,
47 Phillips N, Zakrzewska JM, Pavitt S. Diagnosis, medication, and surgical
48 management for patients with trigeminal neuralgia: a qualitative study. Acta
49 Neurochir (Wien) 2015;1925-1933.
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
- [2] Besi E, Boniface DR, Cregg R, Zakrzewska JM. Comparison of tolerability and adverse symptoms in oxcarbazepine and carbamazepine in the treatment of trigeminal neuralgia and neuralgiform headaches using the Liverpool Adverse Events Profile (AEP). *J Headache Pain* 2015;16:563.
- [3] Bowsher D. Trigeminal neuralgia : a symptomatic study on 126 successive patients with and without previous intervention. *Pain Clinic* 2000;12:93-101.
- [4] Cleeland CS, Ryan KM. Pain assessment: global use of the Brief Pain Inventory. *Ann Acad Med Singapore* 1994;23:129-138.
- [5] Cohen AS, Matharu MS, Goadsby PJ. Short-lasting unilateral neuralgiform headache attacks with conjunctival injection and tearing (SUNCT) or cranial autonomic features (SUNA)--a prospective clinical study of SUNCT and SUNA. *Brain* 2006;129:2746-2760.
- [6] Garvan NJ, Siegfried J. Trigeminal neuralgia--earlier referral for surgery. *Postgraduate Medical Journal* 1983;59:435-437.
- [7] Hall GC, Carroll D, Parry D, McQuay HJ. Epidemiology and treatment of neuropathic pain: The UK primary care perspective. *Pain* 2006;122:156-162.
- [8] Haviv Y, Khan J, Zini A, Almoznino G, Sharav Y, Benoliel R. Trigeminal neuralgia (part I): Revisiting the clinical phenotype. *Cephalalgia* 2015;36:730-746.
- [9] Heinskou T, Maarbjerg S, RoCHAT P, Wolfram F, Jensen RH, Bendtsen L. Trigeminal neuralgia--a coherent cross-specialty management program. *J Headache Pain* 2015;16:66.

The impact of Trigeminal Neuralgia

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
- [10] Henderson WR. Trigeminal neuralgia: the pain and its treatment. *Br Med J* 1967;1:7-15.
- [11] Koopman JS, Dieleman JP, Huygen FJ, de MM, Martin CG, Sturkenboom MC. Incidence of facial pain in the general population. *Pain* 2009;147:122-127.
- [12] Lambro G, Matharu MS. SUNCT, SUNA and trigeminal neuralgia: different disorders or variants of the same disorder? *Curr Opin Neurol* 2014;27:325-331.
- [13] Lee JY, Chen HI, Urban C, Hojat A, Church E, Xie SX, Farrar JT. Development of and psychometric testing for the Brief Pain Inventory-Facial in patients with facial pain syndromes. *J Neurosurg* 2010;113:516-523.
- [14] Lee JY, Sandhu S, Miller D, Solberg T, Dorsey JF, Alonso-Basanta M. Higher dose rate Gamma Knife radiosurgery may provide earlier and longer-lasting pain relief for patients with trigeminal neuralgia. *J Neurosurg* 2015;1-8.
- [15] Maarbjerg S, Gozalov A, Olesen J, Bendtsen L. Concomitant persistent pain in classical trigeminal neuralgia--evidence for different subtypes. *Headache: The Journal of Head and Face Pain* 2014;54:1173-1183.
- [16] Maarbjerg S, Gozalov A, Olesen J, Bendtsen L. Trigeminal neuralgia--a prospective systematic study of clinical characteristics in 158 patients. *Headache: J Head Face Pain* 2014;54:1574-1582.
- [17] Maėianskyte D, Januėis G, Kubilius R, Adomaitiene V, Sciupokas A. Associations Between Chronic Pain and Depressive Symptoms in Patients With Trigeminal Neuralgia. *Medicina Kaunas* 2011;47:386-392.

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
- [18] Melzack R. The McGill Pain Questionnaire: major properties and scoring methods. *Pain* 1975;1:277-299.
- [19] Merskey H, Bogduk N. Classification of chronic pain. Descriptors of chronic pain syndromes and definitions of pain terms. Seattle: IASP Press, 1994.
- [20] Moisset X, Villain N, Ducreux D, Serrie A, Cunin G, Valade D, Calvino B, Bouhassira D. Functional brain imaging of trigeminal neuralgia. *Eur J Pain* 2011;15:124-131.
- [21] Olesen J. The International Classification of Headache Disorders, 3rd edition (beta version). *Cephalalgia* 2013;33:629-808.
- [22] Pilling S, Anderson I, Goldberg D, Meader N, Taylor C. Depression in adults, including those with a chronic physical health problem: summary of NICE guidance. *BMJ* 2009;339:b4108.
- [23] Sullivan MJ, Bishop S.R., Pivik J. The pain catastrophizing scale: development and validation. *Psychological Assessment* 1995;7:524-532.
- [24] Tolle T, Dukes E, Sadosky A. Patient Burden of Trigeminal Neuralgia: Results from a Cross-Sectional Survey of Health State Impairment and Treatment Patterns in Six European Countries. *Pain Practice* 2006;6:153-160.
- [25] VanderPluym J, Richer L. Tic versus TAC: differentiating the neuralgias (trigeminal neuralgia) from the cephalalgias (SUNCT and SUNA). *Curr Pain Headache Rep* 2015;19:473.
- [26] Von Korff M, Dworkin SF, Le Resche L. Graded chronic pain status: an epidemiologic evaluation. *Pain* 1990;40:279-291.

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
- [27] Wu TH, Hu LY, Lu T, Chen PM, Chen HJ, Shen CC, Wen CH. Risk of psychiatric disorders following trigeminal neuralgia: a nationwide population-based retrospective cohort study. *J Headache Pain* 2015;16:64.
- [28] Younis S, Maarbjerg S, Reimer M, Wolfram F, Olesen J, Baron R, Bendtsen L. Quantitative sensory testing in classical trigeminal neuralgia-a blinded study in patients with and without concomitant persistent pain. *Pain* 2016;157:1407-1414.
- [29] Zakrzewska JM. Diagnosis and differential diagnosis of trigeminal neuralgia. *Clin J Pain* 2002;18:14-21.
- [30] Zakrzewska JM, Linskey ME. Trigeminal neuralgia. *Clin Evid (Online)* 2014;2014.
- [31] Zakrzewska JM, Padfield D. The patient's journey through trigeminal neuralgia. *Pain Clinical Update* 2014;22:1-7.
- [32] Zigmond AS, Snaith RP. The hospital anxiety and depression scale. *Acta Psychiatr Scand* 1983;361-437.

Legends for Figures

Figure 1. Medication type prescribed before and after referral. Legend: sky blue-Antibiotics, red-Anticonvulsant, grey-Antidepressants, orange-Analgesics, blue-OTC Analgesics, black-Opioids, dark blue-Others

Figure 2. Association between Pain Severity and Interference on Health Status Domains and general daily activities. Legend: sky blue-Mild (1-3), orange-Moderate (4-6), grey-Severe (7-10). * means $p < 0.001$.

The impact of Trigeminal Neuralgia


Figure 3 Association between Pain Severity and Interference on Health Status Domains and general daily activities including facial status. Legend: sky blue-Mild (1-3), orange-Moderate (4-6), grey-Severe (7-10). * means $p < 0.001$.


1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Summary: Evaluating the impact of trigeminal neuralgia.

This cohort of 225 patients with trigeminal neuralgia describes the demographics and clinical features and provides evidence for the significant psychosocial disability of the condition.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65








Table 1 Surgery type and effectiveness for 23 patients

Surgery type	Effectiveness (n)		
	Complete	Partial	No effect
Gamma Knife	3	0	0
Glycerol	4	5	2
MVD	5	5	0
Peripheral	1	0	1
RFT	3	0	2

Note: some patients had multiple surgeries

Table 2 **Characteristics for 225 TN patients.**

Characteristic	Value (n = 225)
Age in years, mean (SD)	60.9 (12.5)
Age at first attack, median [IQR]	57.0 [46.0, 65.0]
Duration of TN in years, median [IQR]	4.0 [2.0, 7.0]
Female	143 (63.6)
Ethnicity	
Asian	25 (11.0)
Caucasian	189 (83.5)
Others	12 (5.5)
Index of multiple deprivation	
1 (least deprived)	64 (28.4)
2	47 (20.9)
3	47 (20.9)
4	42 (18.7)
5 (most deprived)	25 (11.1)
Profession classification	
Higher managerial, administrative and professional occupations	28 (12.6)
Intermediate occupations	52 (23.3)
Lower managerial, administrative and professional occupations	62 (27.8)
Lower supervisory and technical	10 (4.5)

occupations

Routine occupations	18 (8.1)
Semi-routine occupations	34 (15.2)
Small employers and own account workers	7 (3.1)
Unemployed	12 (5.4)

Employment status

Employed full time	82 (36.6)
Employed part time	21 (9.4)
Full time homemaker	13 (5.8)
Retired	96 (42.9)
Unemployed	12 (5.4)

Referrer to specialist clinic

Dentist	46 (20.6)
GP	102 (45.7)
Specialist	75 (33.6)
Family history of TN	13 (5.8)

Previous services used

GP	180 (80.0)
Dentist	161 (71.6)
Dental Service	95 (42.2)
Dental Specialist	29 (12.9)
Oral Surgeon	74 (32.9)

Dental Procedures	17 (7.6)
Medical Service	132 (58.7)
ENT surgeon	13 (5.8)
Neurosurgeon	43 (19.1)
Neurologist	83 (36.9)
Physician	23 (10.2)
Psychiatrist	1 (0.4)
Psychologist	2 (0.9)
Pain Specialist	14 (6.2)
Other medical procedures	34 (15.1)

No. of secondary dental or medical services

0	40 (17.8)
1	112 (49.8)
2	53 (23.6)
3+	20 (8.9)

Note: values are presented as frequency (%) unless specified.

Table 3 Associated factors and medical history stratified by the type of TN

Associate factors and medical history	Group 1	Group 2	Group 3	P-trend
	TN	TNC	TNA	
	n = 155	n = 32	n = 38	
Altered sensation or numbness	39 (25.3)	12 (40.0)	11 (28.9)	0.258
Any autonomies unilateral	86 (55.5)	17 (53.1)	38 (100.0)	<0.001
Swelling face	19 (12.3)	3 (10.0)	12 (31.6)	0.009
Redness of the face	14 (9.1)	2 (6.7)	13 (34.2)	<0.001
Nasal stuffiness/runny	17 (11.0)	4 (13.3)	23 (60.5)	<0.001
Eye redness	6 (3.9)	1 (3.3)	12 (31.6)	<0.001
Eye tearing	18 (11.7)	7 (23.3)	22 (57.9)	<0.001
Oedema eyelid	4 (2.6)	1 (3.3)	7 (18.4)	<0.001
Earache	13 (8.4)	1 (3.3)	3 (7.9)	0.628
Fullness ears	8 (5.2)	3 (10.0)	6 (15.8)	0.080
Headaches	37 (24.0)	6 (20.0)	10 (26.3)	0.829
Migraines	17 (11.0)	3 (10.0)	7 (18.4)	0.426
Migraines + TTH	12 (7.7)	1 (3.1)	3 (7.9)	0.638
Bruxism	36 (23.7)	8 (26.7)	10 (26.3)	0.906
Medical history				
Hypertension	57 (36.8)	14 (43.8)	13 (34.2)	0.690
CVS	19 (12.3)	4 (12.5)	5 (13.2)	0.989
Diabetes	10 (6.5)	2 (6.2)	0 (0.0)	0.276
Deafness	17 (11.0)	1 (3.3)	2 (5.3)	0.272
Ringing ears	18 (11.7)	0 (0.0)	2 (5.3)	0.083
Other chronic pain	33 (21.3)	9 (28.1)	8 (21.1)	0.686
Neck pain	15 (9.9)	4 (13.3)	4 (10.5)	0.857

Back pain	23 (15.1)	5 (17.2)	5 (13.2)	0.898
Previous surgery TN	15 (9.7)	3 (9.4)	5 (13.2)	0.806

Note: values are presented as frequency (%); P-trend represents comparison across three groups.

Table 4 Pain characteristic stratified by the type of TN

Pain characteristic	Group 1 TN n = 155	Group 2 TNC n = 32	Group 3 TNA n = 38	P- trend
V1	3 (1.9)	0 (0)	0 (0)	0.496
V2	35 (22.7)	6 (18.8)	13 (33.3)	0.289
V3	52 (33.8)	14 (43.8)	4 (10.3)	0.005
V1 + V2	8 (5.2)	4 (12.5)	9 (23.1)	0.002
V2 + V3	46 (29.9)	7 (21.9)	9 (23.1)	0.516
V1 + V2 + V3	9 (5.8)	1 (3.1)	4 (10.3)	0.438
Right	101 (65.6)	18 (56.2)	30 (76.9)	0.720
Left	50 (32.5)	13 (40.6)	9 (23.1)	0.106
Bilateral	3 (1.9)	1 (3.1)	0 (0.0)	0.587
Intra oral	123 (79.9)	26 (81.2)	34 (87.2)	0.578
Extra oral	83 (68.6)	17 (65.4)	22 (62.9)	0.802
Predominant type of attack				0.116
Single stab	71 (46.1)	11 (34.4)	18 (46.2)	
Series of stabs	56 (36.4)	10 (31.3)	9 (23.1)	

Saw tooth	8 (5.2)	6 (18.8)	4 (10.3)	
Single stab + Series of stabs	19 (12.3)	5 (15.6)	8 (20.5)	
Pain severity, median [IQR]				
Worse	10.0 [8.0, 10.0]	9.0 [7.0, 10.0]	10.0 [8.0, 10.0]	0.715
Average	5.0 [3.0, 7.0]	6.0 [4.0, 8.0]	5.0 [3.0, 7.0]	0.298
Least	0.0 [0.0, 3.0]	2.0 [0.0, 6.0]	2.0 [0.0, 5.0]	0.043
Circumstances				0.001
Acute	48 (34.0)	10 (34.5)	11 (30.6)	
Memorable	60 (42.6)	4 (13.8)	20 (55.6)	
Slow to develop	27 (19.1)	9 (31.0)	4 (11.1)	
Cannot remember	6 (4.3)	6 (20.7)	1 (2.8)	
Frequency of pain attack				0.865
Hourly	11 (8.7)	1 (4.0)	2 (6.2)	
Daily	115 (91.3)	24 (96.0)	30 (93.8)	
Duration of attacks				0.872
Seconds	105 (69.5)	20 (66.7)	24 (61.5)	
Minutes	42 (27.8)	9 (30.0)	13 (33.3)	

1-4 hours	4 (2.6)	1 (3.3)	2 (5.1)	
Pain after main attack	50 (36.0)	20 (62.5)	17 (45.9)	0.020
Length of remission				0.385
None	12 (9.0)	2 (6.7)	6 (17.1)	
Days	16 (11.9)	1 (3.3)	3 (8.6)	
Weeks	70 (52.2)	14 (46.7)	16 (45.7)	
Months	23 (17.2)	6 (20.0)	5 (14.3)	
Years	13 (9.7)	7 (23.3)	5 (14.3)	
Remission period change				0.470
No change	42 (31.6)	5 (19.2)	14 (37.8)	
Shorter	84 (63.2)	18 (69.2)	21 (56.8)	
Longer	7 (5.3)	3 (11.5)	2 (5.4)	
Provoking factors				
Spontaneous pain	151 (98.1)	32 (100.0)	38 (97.4)	0.690
Provoked by light touch	143 (92.9)	29 (90.6)	39 (100.0)	0.187
Provoked by other factors				
Cold wind/weather	80 (51.9)	10 (31.2)	23 (59.0)	0.050

Bodily movement	65 (42.2)	12 (37.5)	24 (61.5)	0.063
Noise or light	3 (1.9)	0 (0.0)	1 (2.6)	0.690
Alcohol	1 (0.6)	0 (0.0)	0 (0.0)	0.793

McGill pain questionnaire

Number of words chosen (mean ± sd)	10.3 ± 3.8	12.2 ± 3.9	12.3 ± 2.8	0.003
Pain rating index (mean ± sd)	27.7 ± 12.1	32.8 ± 14.4	31.9 ± 8.4	0.044
Sensory groups	Shooting (110)	Shooting (24)	Shooting (29)	
	Sharp (93)	Sharp (21)	Sharp (26)	
	Stabbing (73)	Stabbing (14)	Stabbing (20)	
Affective	Vicious (43)	Terrifying (8)	Wretched (13)	
	Terrifying (39)	Fearful (6)	Fearful (12)	
Evaluative	Unbearable (58)	Unbearable (16)	Unbearable (13)	
Miscellaneous	Piercing (57)	Piercing (14)	Piercing (16)	

HAD-Anxiety

Nil	72 (53.3)	16 (59.3)	8 (22.2)	
Mild	29 (21.5)	3 (11.1)	18 (50.0)	0.001
Severe	34 (25.2)	8 (29.6)	10 (27.8)	

HAD-Depression

Nil	95 (69.3)	19 (70.4)	16 (44.4)	
Mild	21 (15.3)	5 (18.5)	8 (22.2)	0.047
Severe	21 (15.3)	3 (11.1)	12 (33.3)	

Brief pain inventory, median [IQR]

Pain severity index	3.50 [2.00, 5.25]	3.50 [2.25, 6.25]	5.00 [3.25, 6.31]	0.090
Pain interference-general daily life	2.71 [1.14, 5.29]	3.14 [0.57, 4.57]	3.29 [1.71, 6.00]	0.380
Pain interference-facial status	4.57 [2.07, 7.14]	7.00 [2.43, 7.86]	5.93 [2.21, 7.71]	0.349

Note: values are presented as frequency (%) unless specified; P-trend represents comparison across three group

Table 5 Association between HAD-anxiety and depression and graded chronic pain scale in patients with trigeminal neuralgia and its variants.

HAD-Anxiety	Graded chronic pain scale			
	Grade 1	Grade 2	Grade 3	Grade 4
Nil	19	30	13	19
Mild	2	17	5	17
Severe	1	5	7	25
HAD- Depression				
Nil	18	41	19	26
Mild	3	5	3	16
Severe	1	6	5	19

Pearson's Chi-square test: p-value < 0.001

Table 6 Association between HAD-anxiety and depression and Brief Pain Inventory (BPI) pain severity in patients with trigeminal neuralgia and its variants.

HAD-Anxiety*	BPI pain severity			
	No pain/ not reported	Mild	Moderate	Severe
Nil	12	29	33	11
Mild	3	12	21	10
Severe	1	14	20	12
HAD- Depression**				
Nil	12	42	41	17
Mild	3	6	22	5
Severe	2	8	11	11

*Pearson's Chi-square test: p-value = 0.163 **Pearson's Chi-square test: p-value = 0.022



Click here to access/download


Copyright Transfer Agreement
PAIN_Copyright_Transfer_Form.pdf





Click here to access/download

Copyright Transfer Agreement
coi_disclosure zakrzewska.pdf





[Click here to access/download](#)

Copyright Transfer Agreement

PAIN_Copyright_Transfer_Form Jianhua Wu.pdf





Click here to access/download

Copyright Transfer Agreement

PAIN_Copyright_Transfer_Form Phillips.pdf





Click here to access/download

Copyright Transfer Agreement
PAIN_Copyright_Transfer_Form_mmw.pdf





Click here to access/download

Copyright Transfer Agreement

PAIN_Copyright_Transfer_Form_PAVITT.pdf



Please wait...

If this message is not eventually replaced by the proper contents of the document, your PDF viewer may not be able to display this type of document.

You can upgrade to the latest version of Adobe Reader for Windows®, Mac, or Linux® by visiting http://www.adobe.com/go/reader_download.

For more assistance with Adobe Reader visit <http://www.adobe.com/go/acrreader>.

Windows is either a registered trademark or a trademark of Microsoft Corporation in the United States and/or other countries. Mac is a trademark of Apple Inc., registered in the United States and other countries. Linux is the registered trademark of Linus Torvalds in the U.S. and other countries.

Please wait...

If this message is not eventually replaced by the proper contents of the document, your PDF viewer may not be able to display this type of document.

You can upgrade to the latest version of Adobe Reader for Windows®, Mac, or Linux® by visiting http://www.adobe.com/go/reader_download.

For more assistance with Adobe Reader visit <http://www.adobe.com/go/acrreader>.

Windows is either a registered trademark or a trademark of Microsoft Corporation in the United States and/or other countries. Mac is a trademark of Apple Inc., registered in the United States and other countries. Linux is the registered trademark of Linus Torvalds in the U.S. and other countries.

Please wait...

If this message is not eventually replaced by the proper contents of the document, your PDF viewer may not be able to display this type of document.

You can upgrade to the latest version of Adobe Reader for Windows®, Mac, or Linux® by visiting http://www.adobe.com/go/reader_download.

For more assistance with Adobe Reader visit <http://www.adobe.com/go/acrreader>.

Windows is either a registered trademark or a trademark of Microsoft Corporation in the United States and/or other countries. Mac is a trademark of Apple Inc., registered in the United States and other countries. Linux is the registered trademark of Linus Torvalds in the U.S. and other countries.

Please wait...

If this message is not eventually replaced by the proper contents of the document, your PDF viewer may not be able to display this type of document.

You can upgrade to the latest version of Adobe Reader for Windows®, Mac, or Linux® by visiting <http://www.adobe.com/products/acrobat/readstep2.html>.

For more assistance with Adobe Reader visit <http://www.adobe.com/support/products/acrreader.html>.

Windows is either a registered trademark or a trademark of Microsoft Corporation in the United States and/or other countries. Mac is a trademark of Apple Inc., registered in the United States and other countries. Linux is the registered trademark of Linus Torvalds in the U.S. and other countries.