



UNIVERSITY OF LEEDS

This is a repository copy of *Effects of a 4-week intervention with ready-made bioactive enriched pancakes on biomarkers of the metabolic syndrome*.

White Rose Research Online URL for this paper:  
<http://eprints.whiterose.ac.uk/110989/>

Version: Published Version

---

**Proceedings Paper:**

Sutulic, S, Bosch, C, Marshall, L et al. (4 more authors) (2016) Effects of a 4-week intervention with ready-made bioactive enriched pancakes on biomarkers of the metabolic syndrome. In: Proceedings of the Nutrition Society. Phytochemicals and health: new perspectives on plant based nutrition, 21-22 Mar 2016, Edinburgh. Cambridge University Press .

<https://doi.org/10.1017/S0029665116000513>

---

(c) 2016, the Authors. This version is free to view and download for private research and study only. Not for re-distribution, re-sale or use in derivative works. Published in the Proceedings of the Nutrition Society and available at <https://doi.org/10.1017/S0029665116000513>

**Reuse**

Unless indicated otherwise, fulltext items are protected by copyright with all rights reserved. The copyright exception in section 29 of the Copyright, Designs and Patents Act 1988 allows the making of a single copy solely for the purpose of non-commercial research or private study within the limits of fair dealing. The publisher or other rights-holder may allow further reproduction and re-use of this version - refer to the White Rose Research Online record for this item. Where records identify the publisher as the copyright holder, users can verify any specific terms of use on the publisher's website.

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.



[eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk)  
<https://eprints.whiterose.ac.uk/>

## Effects of a 4-week intervention with ready-made bioactive enriched pancakes on biomarkers of the metabolic syndrome

S. Sutulic<sup>1</sup>, C. Bösch<sup>1</sup>, L. Marshall<sup>1</sup>, A. Bordoni<sup>2</sup>, H. Hingyi<sup>3</sup>, M. Muller<sup>4</sup>, C. Orfila<sup>1</sup> and on behalf of the PATHWAY-27 consortium Grant agreement 311876

<sup>1</sup>School of Food Science and Nutrition, University of Leeds, Leeds LS2 9JT, <sup>2</sup>Dipartimento di Scienze e Tecnologie Agro-Alimentari, Università di Bologna, Italy, <sup>3</sup>Adexgo Kft., Hungary and <sup>4</sup>AdWare Research Kft., Hungary

In the UK it is estimated that 1 in 4 adults are at greater risk of heart disease and stroke due to presence of the metabolic syndrome (MetS)<sup>(1)</sup>. The MetS is a term used to describe any combination of at least three of the following conditions: high blood pressure, increased waist circumference, impaired fasting glucose, increased fasting triglycerides, and low HDL cholesterol.

Components of foods that may improve conditions associated with MetS include the omega-3 fatty acid docosahexaenoic acid (DHA) and the soluble plant fibre beta-glucan (BG), which are considered beneficial for maintaining normal blood triglyceride and cholesterol concentrations<sup>(2,3)</sup>. There is also clinical evidence that anthocyanins (AC) may improve HDL cholesterol in adults with dyslipidaemia<sup>(4)</sup>.

In a double-blind, parallel-arm, randomised pilot study, 35 participants (M = 40 %), mean age 45 y, with or at risk of MetS were recruited to consume ready-made pancakes enriched with DHA, BG, or AC alone or in combinations of DHA + BG or DHA + AC every day for 4 weeks. At baseline and endpoint markers of MetS were measured. Mean change, from baseline to endpoint, in biomarkers of the MetS are summarised in Table 1.

**Table 1:** Mean change in biomarkers of the MetS. HDL-C, HDL cholesterol; SBP, systolic blood pressure; DBP, diastolic blood pressure; SD, standard deviation. Results were not statistically significant.

	DHA		BG		AC		DHA + BG		DHA + AC	
	Δ	SD	Δ	SD	Δ	SD	Δ	SD	Δ	SD
Glucose (mg/dL)	4.5	6.5	0.0	7.5	-4.2	7.9	-2.4	4.1	-2.2	15.1
Triglycerides (mg/dL)	-6.3	46.0	21.0	89.3	-13.2	28.7	0.0	36.7	-11.8	37.8
HDL-C (mg/dL)	5.0	5.3	0.5	5.6	-0.5	6.0	-1.9	5.3	5.8	4.0
SBP (mmHg)	3.1	10.5	0.4	8.8	5.6	9.1	-2.5	11.2	-11.7	10.5
DBP (mmHg)	0.6	6.7	1.8	10.1	5.6	1.9	0.2	12.7	4.5	8.0

The objective of this pilot study was to select the enrichment type most effective at improving fasting blood triglycerides and/or HDL cholesterol, to be consumed in the subsequent PATHWAY-27 12-week placebo-controlled, multi-centre trial. Pancakes enriched with DHA and AC alone and in combination appear to be the most effective. DHA + AC enrichment was selected for further study.

1. Department of Health (2014) NHS Choices: Metabolic syndrome, <http://www.nhs.uk/conditions/metabolic-syndrome/Pages/Introduction.aspx>, (accessed January 2016).
2. EFSA Panel on Dietetic Products, Nutrition and Allergies (2010) *EFSA Journal* 8, (10), 1734.
3. EFSA Panel on Dietetic Products, Nutrition and Allergies (2009) *EFSA Journal* 7, (9), 1254.
4. Qin *et al.* (2009) *Am J Clin Nutr* 90, 485–492.