

Degrees of freedom in social bonds of crested macaque females.

Julie Duboscq<sup>1,2,3,4\*†</sup>, Christof Neumann<sup>1,2,5,6\*\*</sup>, Muhammad Agil<sup>7</sup>, Dyah Perwitasari-Farajallah<sup>8,9</sup>, Bernard Thierry<sup>3,4</sup>, Antje Engelhardt<sup>1,2\*\*\*</sup>

<sup>1</sup> Junior Research Group of Primate Sexual Selection, German Primate Center, Göttingen, Germany, <sup>2</sup> Courant Research Centre for the Evolution of Social Behaviour, Göttingen, Germany, <sup>3</sup> Université de Strasbourg, Institut Pluridisciplinaire Hubert Curien, Strasbourg, France, <sup>4</sup> Centre National de la Recherche Scientifique, Département Ecologie, Physiologie et Ethologie, Strasbourg, France, <sup>5</sup> Junior Research Group of Primate Kin Selection, Max-Planck-Institute for Evolutionary Anthropology, Leipzig, Germany, <sup>6</sup> Institute of Biology, Faculty of Bioscience, Pharmacy and Psychology, University of Leipzig, Leipzig, Germany, <sup>7</sup> Faculty of Veterinary Sciences, Bogor Agricultural University, Bogor, Indonesia, <sup>8</sup> Primate Research Center, Bogor Agricultural University, Bogor, Indonesia, <sup>9</sup> Faculty of Mathematics and Natural Sciences, Bogor Agricultural University, Bogor, Indonesia

\* corresponding author: [julie.a.m.duboscq@gmail.com](mailto:julie.a.m.duboscq@gmail.com)

† present affiliation of Julie Duboscq: Kyoto University Wildlife Research Centre, 2-24 Tanaka-Sekiden-cho, Sakyo, Kyoto, Japan 606-8203.

\*\* present affiliation of Christof Neumann: Université de Neuchâtel, Institute of Biology, Department of Comparative Cognition, Neuchâtel, Switzerland

\*\*\*present affiliation of Antje Engelhardt: Liverpool John Moores University, School of Natural Sciences and Psychology, Liverpool, United Kingdom

Word count main text: 6351



**Abstract**

Social bonds between group members can affect individual fitness and well-being. While the impact of bond strength is well studied, the consequences of bond predictability and equitability are often overlooked. Similarly, whether bonds reflect short-term contingencies and/or long-term social strategies remains understudied. We investigated these questions in female crested macaques (*Macaca nigra*), which display a tolerant social style within a nepotistic hierarchical social structure. We analysed the structure of dyadic social bonds by testing whether similarity within dyads – in kinship, dominance and age – predicted the strength, predictability and equitability of bonds. We then tested the value of social bonds by analysing the effect of their characteristics on three fitness-related behaviours: coalitionary support, feeding-in-proximity and aggression. We found that the bond characteristics of females differed substantially from those of other species with comparable data: bonds were of average strength, of moderate endurance and relatively balanced. Stronger bonds were more equitable but less predictable than weaker bonds. Closely-ranked females, but not kin or age peers, had stronger, more predictable and more equitable bonds than others. Coalitionary support was not related to any of the bond characteristics, feeding-in-proximity was positively associated with strength and predictability and aggression was positively linked to strength and equitability. These results highlight the complex picture of the benefits of social bonds in this species. They reflect the degrees of freedom tolerant macaque females can express in their social relationships within their stable social structure, a pattern that may not be given enough consideration in stable nepotistic hierarchical societies. Comparative research is necessary to establish whether these patterns are more general than previously thought or a specific feature of tolerant macaques. Investigating various characteristics of bonds together is paramount in order to appreciate the dynamics of social relationships and to better understand the social components of fitness.

1 Degrees of freedom in social bonds of crested macaque females.

2

3 Julie Duboscq<sup>1,2,3,4\*†</sup>, Christof Neumann<sup>1,2,5,6\*\*</sup>, Muhammad Agil<sup>7</sup>, Dyah Perwitasari-Farajallah<sup>8,9</sup>,

4 Bernard Thierry<sup>3,4</sup>, Antje Engelhardt<sup>1,2\*\*\*</sup>

5

6 <sup>1</sup> Junior Research Group of Primate Sexual Selection, German Primate Center, Göttingen, Germany, <sup>2</sup>

7 Courant Research Centre for the Evolution of Social Behaviour, Göttingen, Germany, <sup>3</sup> Université de

8 Strasbourg, Institut Pluridisciplinaire Hubert Curien, Strasbourg, France, <sup>4</sup> Centre National de la

9 Recherche Scientifique, Département Ecologie, Physiologie et Ethologie, Strasbourg, France, <sup>5</sup> Junior

10 Research Group of Primate Kin Selection, Max-Planck-Institute for Evolutionary Anthropology,

11 Leipzig, Germany, <sup>6</sup> Institute of Biology, Faculty of Bioscience, Pharmacy and Psychology,

12 University of Leipzig, Leipzig, Germany, <sup>7</sup> Faculty of Veterinary Sciences, Bogor Agricultural

13 University, Bogor, Indonesia, <sup>8</sup> Primate Research Center, Bogor Agricultural University, Bogor,

14 Indonesia, <sup>9</sup> Faculty of Mathematics and Natural Sciences, Bogor Agricultural University, Bogor,

15 Indonesia

16

17 \* corresponding author: [julie.a.m.duboscq@gmail.com](mailto:julie.a.m.duboscq@gmail.com)

18 † present affiliation of Julie Duboscq: Kyoto University Wildlife Research Centre, 2-24 Tanaka-

19 Sekiden-cho, Sakyo, Kyoto, Japan 606-8203.

20 \*\* present affiliation of Christof Neumann: Université de Neuchâtel, Institute of Biology, Department

21 of Comparative Cognition, Neuchâtel, Switzerland

22 \*\*\*present affiliation of Antje Engelhardt: Liverpool John Moores University, School of Natural

23 Sciences and Psychology, Liverpool, United Kingdom

24

25 Word count main text: 6351

26

27

28

29

30 **Abstract**

31

32 Social bonds between group members can affect individual fitness and well-being. While the impact  
33 of bond strength is well studied, the consequences of bond predictability and equitability are often  
34 overlooked. Similarly, whether bonds reflect short-term contingencies and/or long-term social  
35 strategies remains understudied. We investigated these questions in female crested macaques (*Macaca*  
36 *nigra*), which display a tolerant social style within a nepotistic hierarchical social structure. We  
37 analysed the structure of dyadic social bonds by testing whether similarity within dyads – in kinship,  
38 dominance and age – predicted the strength, predictability and equitability of bonds. We then tested  
39 the value of social bonds by analysing the effect of their characteristics on three fitness-related  
40 behaviours: coalitionary support, feeding-in-proximity and aggression. We found that the bond  
41 characteristics of females differed substantially from those of other species with comparable data:  
42 bonds were of average strength, of moderate endurance and relatively balanced. Stronger bonds were  
43 more equitable but less predictable than weaker bonds. Closely-ranked females, but not kin or age  
44 peers, had stronger, more predictable and more equitable bonds than others. Coalitionary support was  
45 not related to any of the bond characteristics, feeding-in-proximity was positively associated with  
46 strength and predictability and aggression was positively linked to strength and equitability. These  
47 results highlight the complex picture of the benefits of social bonds in this species. They reflect the  
48 degrees of freedom tolerant macaque females can express in their social relationships within their  
49 stable social structure, a pattern that may not be given enough consideration in stable nepotistic  
50 hierarchical societies. Comparative research is necessary to establish whether these patterns are more  
51 general than previously thought or a specific feature of tolerant macaques. Investigating various  
52 characteristics of bonds together is paramount in order to appreciate the dynamics of social  
53 relationships and to better understand the social components of fitness.

54

55 **Keywords**

56 social bonds, strength, predictability, equitability, social dynamics, behavioural strategy, primates

57

## 58 **Introduction**

59

60 Social bonds are positive social relationships amongst pairs of individuals of the same group  
61 (Silk, Cheney, & Seyfarth, 2013; Silk 2007a). They are defined in a multidimensional space of  
62 relationship qualities such as relative strength, predictability (or magnitude of change over time) and  
63 equitability (the balance of social exchanges within a dyad) (Silk et al., 2013; Whitehead, 2008).  
64 Variation in these components can affect individual fitness inasmuch as individuals with more  
65 numerous, stable or stronger bonds experience enhanced survival, greater reproductive success or  
66 improved general well-being compared to others (feral horses (*Equus caballus*), Cameron, Setsaas, &  
67 Linklater, 2009; bottlenose dolphins (*Tursiops sp.*) Frère et al., 2010; talapoin monkeys (*Miopethicus*  
68 *talapoin*), Keverne, Martensz, & Tuite, 1989; Barbary macaques (*Macaca sylvanus*), McFarland &  
69 Majolo, 2013; Young, Majolo, Heistermann, Schülke, & Ostner, 2014; chacma baboons (*Papio*  
70 *cynocephalus ursinus*), Silk et al., 2009, 2010; humans (*Homo sapiens*), Uchino 2006). For  
71 individuals, the value of social bonds is also related to the direct or indirect benefits they may obtain  
72 from daily social exchanges (primate males: Ostner & Schülke, 2014; mammalian females: Silk,  
73 2007a), e.g. reciprocation of social grooming, reduction of aggression, increased coalitionary support  
74 during conflicts, or better access to food resources (ravens (*Corvus corax*), Fraser & Bugnyar, 2011;  
75 baboons (*P. ursinus*), King, Clark, & Cowlshaw, 2011; spotted hyenas (*Crocuta crocuta*), Smith,  
76 Memenis, & Holekamp, 2007), which ultimately may impact their fitness and well-being.

77 In order to understand the function and value of social bonds, i.e. which benefits can be  
78 obtained by forming and maintaining them, it is also crucial to investigate their underlying structure,  
79 i.e. the characteristics of the dyads forming particular bonds. In many animal societies, individuals  
80 that are similar in terms of relatedness, dominance status, personality, reproductive state or energetic  
81 needs are more likely to form strong and enduring social bonds than others (yellow-bellied marmots  
82 (*Marmota flaviventris*): Armitage & Schwartz, 2000; chacma baboons: Barrett & Henzi, 2001;  
83 giraffes (*Giraffa camelopardalis*), Carter, Seddon, Frère, Carter, & Goldizen, 2013; mountain goats  
84 (*Oreamnos americanus*), Godde, Côté, & Réale, 2015; rhesus macaques (*M. mulatta*), de Waal &

85 Lutrell, 1986; ring-tailed coatis (*Nasua nasua*), Hirsch, Staton, & Maldonado, 2012). Each of these  
86 characteristics can be uniquely important in influencing the formation and maintenance of a bond. For  
87 instance, although close kin are obvious coalition partners, kin-based coalitionary support may not be  
88 advantageous if such kin are low-ranking (Chapais, 2006), in which case establishing a bond with a  
89 higher-ranking non-relative may be more valuable (primates: Schino, 2007; hyenas: Smith et al.,  
90 2010).

91 Research on the benefits of social bonds amongst same-sex adult group members has so far  
92 mostly considered how bond characteristics at the extreme positive end of the spectrum affect  
93 measures of fitness, health or well-being, e.g. preferred associates (Frère et al., 2011) or top three  
94 partners (Silk et al., 2006a). However, animals may have a variety of options for regulating the  
95 consequences of bonds. First, the predictability and equitability of an individual's social relationships  
96 may be as important as their strength (e.g. the sheer amount of research on cooperation and  
97 reciprocity: Nowak, 2006; Trivers, 1971, 2006). In addition, "weak" bonds, as opposed to "strong"  
98 bonds, within a social network may also be important. For example, weak bonds contribute to  
99 stabilising the overall network in *Escherichia coli* (Csermely, 2004). In humans, although a few  
100 strong bonds are individually more influential, more numerous weak bonds enhance the propagation  
101 of novel information (Bakshy, Rosenn, Marlow, & Adamic, 2012). Finally, it has been shown that  
102 variance in bond strength is actually more predictive of fitness than strength itself (longevity in rock  
103 hyraxes (*Procavia capensis*), Barocas, Ilany, Koren, Kam, & Geffen, 2011; pup production in degus  
104 (*Octodon degus*), Wey, Burger, Ebensperger, & Hayes, 2013). Studies integrating the different  
105 dimensions of social bonds simultaneously and on a continuous scale are therefore indispensable for  
106 deepening our understanding of the link between sociality and fitness.

107 It has also been argued that social bonds are likely to be formed and maintained based on  
108 contingencies (short-term, opportunistic tactics) rather than, or in addition to, long-term, fixed  
109 strategies (Barrett & Henzi, 2001, 2006). For instance, female chacma baboons did not sustain  
110 constant differentiated relationships with other females across time but changed cyclically between  
111 "brief associations", "casual acquaintances" and "constant companionships" according to food  
112 availability (i.e. between food-abundant and food-scarce periods) (Henzi, Lusseau, Weingrill, van

113 Schaik & Barrett, 2009). In addition, pregnant or early lactating female chacma baboons were less  
114 likely to become involved in coalitions and, thus, were not reliable cooperation partners for both kin  
115 and non-kin (Barrett & Henzi, 2001). Therefore, individuals may use varying social strategies  
116 reflecting certain degrees of social freedom according to social context, the spatial or temporal  
117 availability of partners, or environmental conditions, even when living in stable organised societies.  
118 Although this seems rather straightforward in animal societies that display flexibility in their social  
119 structure (i.e. group membership and group size regularly changing, such as in fission-fusion  
120 societies), variation in social strategies only recently began to be considered in species with a stable  
121 nepotistic hierarchical social structure such as those of many primates, of hyenas or African elephants  
122 (Barrett & Henzi, 2001; Ilani, Booms, & Holekamp, 2015; Sick et al., 2014).

123         Macaques (genus *Macaca*) are an ideal candidate for the investigation of variation in social  
124 strategies. Although the different macaque species share the same social organisation (philopatric  
125 females organised in stable matrilineal dominance hierarchies), macaque species are described as  
126 more or less socially tolerant depending on the degree of nepotism, power asymmetries, conciliatory  
127 tendencies and counter-aggression in social relationships (Thierry, 2007; Thierry, 2013). Such social  
128 variation can be expected to influence the structure and function of social bonds (Thierry, 1990;  
129 Butovskaya, 2004). Specifically, when power asymmetries are pronounced and the degree of  
130 nepotism is high, as in less tolerant species, individuals' options regarding who to interact with and  
131 how are limited. These constraints may lead to individuals having a relatively low number of strong,  
132 predictable and equitable partnerships within a social group. In contrast, when social rules are more  
133 relaxed, i.e. when power asymmetries are moderate and the degree of nepotism is weak, as in more  
134 tolerant macaques, individuals are able to interact with diverse partners and develop a greater  
135 diversity and number of social bonds (Thierry, 1990; Butovskaya, 2004; Cooper & Bernstein 2008;  
136 Duboscq et al., 2013). Thus, the degrees of freedom that individuals have in their relationships within  
137 their group can be assessed through the size and diversity of their social network in relation to the  
138 influence of dominance and kinship on an individual's social options – or lack thereof (Thierry, 1990;  
139 Butovskaya, 2004).



140 In this study, we aim to investigate these degrees of freedom and the interplay between the  
141 structure and the value of social bonds in wild female crested macaques (*M. nigra*), which express a  
142 tolerant social style (Petit, Abegg, & Thierry, 1997; Duboscq et al., 2013). The study population of  
143 crested macaques lives in a relatively predictable and safe ecological environment (low predation risk  
144 and abundant food year round; O'Brien & Kinnaird, 1997) while facing dynamic social conditions,  
145 e.g. male migration and hierarchical changes, which are a potential source of social instability in the  
146 group (Neumann, 2013; Marty, Hodges, Agil, & Engelhardt, in press). Females reproduce year round  
147 (Kerhoas et al., 2014), which is another potential source of fluctuation in the amount of time and  
148 attention females can devote to their female social partners (D'Amato, Troisi, Scucchi, & Fuccillo,  
149 1982; Bardi, Shimizu, Fujita, Borgognini Tarli, & Huffman, 2001; Barrett & Henzi, 2001; Brent,  
150 MacLarnon, Platt, & Semple, 2013). Previous studies on the same population showed that female  
151 crested macaques form highly diverse affiliative social networks (Duboscq et al., 2013). On the one  
152 hand, the strength of female-female social bonds was linked to predator deterrence, suggesting that  
153 strong bonds play a role in enhancing survival (Micheletta et al., 2012). On the other hand, bond  
154 strength did not affect the occurrence and frequency of reconciliation, an important conflict  
155 management strategy (Duboscq, Agil, Engelhardt, & Thierry, 2014). Nonetheless, other relationship  
156 qualities, such as equitability and predictability, increased the likelihood of reconciliation (Duboscq et  
157 al., 2014). As such, it seems that social bond characteristics have different values depending on the  
158 context of the social benefits to be gained in this species and we would expect that females express  
159 great degrees of social freedom.

160 Specifically, since macaques form stable matrilineal hierarchical societies (Cords, 2012), kin  
161 and adjacently-ranked dyads are expected to form the strongest, most predictable and equitable bonds  
162 (Silk, 2007b). However, given the tolerant social style of crested macaques and their expected great  
163 degrees of social freedom, we made the hypothesis that these dyad characteristics would not predict  
164 social bond strength, equitability and predictability. To test this prediction and to establish the  
165 structure of bonds, we analysed the relationship between three measures of dyadic similarities (degree  
166 of relatedness, proximity in age and dominance rank) and three social bond characteristics, namely  
167 strength, predictability and equitability. Furthermore, under the hypothesis that social bond

168 characteristics are linked to fitness in a positive predictable way (Silk, 2007a, b), variation in these  
169 characteristics is expected to explain a substantial amount of variation in the occurrence or frequency  
170 of behaviours directly or indirectly linked to fitness benefits, such as a reduction in aggression,  
171 increased coalitionary support during conflicts or better access to food resources. However, again  
172 given the tolerant social style of crested macaques and their expected great degrees of social freedom,  
173 we made the hypothesis that the characteristics of social bonds will not predict the occurrence of these  
174 fitness-related behaviours. To test this prediction and to establish the value of bonds, we tested the  
175 extent to which each social bond characteristic influenced coalitionary support, feeding-in-proximity  
176 and aggression. By taking a more integrated perspective of social bonds in a species with a tolerant  
177 social style, we address the concept of individuals' degrees of social freedom within their stable  
178 network of social relationships (Thierry, 1990; Butovskaya, 2004).

179

## 180 **Methods**

181

182 Behavioural data collection and analysis

183

### 184 *Field site, study animals and data collection*

185 Crested macaques are critically endangered and endemic to the island of Sulawesi, Indonesia  
186 (Sugardjito et al., 1989). The study population inhabits the Tangkoko Reserve, North Sulawesi  
187 (1°33'N, 125°10'E; e.g. Duboscq, Neumann, Perwitasari-Farajallah, & Engelhardt, 2008), broadly  
188 classified as a lowland rainforest with seasonal variation in rainfall and fruit abundance (O'Brien &  
189 Kinnaird, 1997). The study was part of the Macaca Nigra Project, a long-term field project on the  
190 biology of crested macaques that started in 2006. We studied two groups, "PB" and "R1", comprised  
191 of ca. 60 and 80 individuals respectively. The monkeys were fully habituated to human observers and  
192 all adults could be individually identified based on physical characteristics.

193 JD and two field assistants collected behavioural data between October 2008 and May 2010  
194 on all adult females (15 – 18 in PB, 21 – 24 in R1) using focal animal sampling (Martin & Bateson,  
195 1993) (interobserver reliability: Cohen's kappa = 0.69–0.90, correlation coefficients between

196 behavioural variables = 0.79 – 0.98). We collected 30 minute-point-sample observations for activity  
197 (foraging, feeding, socialising, resting, and travelling). Every second minute, we also noted the  
198 identity of neighbours in three proximity categories: in body contact, within one body-length, and  
199 within five body-lengths. We recorded focal social events continuously, including the start and end  
200 time of interactions, the sequence of all behaviours, as well as the identity and behaviours of all social  
201 partners. This study included a total of 2,480 hours of focal data focusing on 35 females that were  
202 present during the entire study period (median<sub>PB</sub> = 68 hours per female, range<sub>PB</sub>: 65 – 78, N<sub>PB</sub> = 14;  
203 median<sub>RI</sub> = 66 hours per female, range<sub>RI</sub>: 59 – 71, N<sub>RI</sub> = 21). Behavioural interactions were expressed  
204 as duration (e.g. social grooming) or frequency (e.g. approach) per focal and per dyadic (sum of two  
205 focals') observation time over the whole study period (i.e. 19 months).

206         Additionally, we collected at least three faecal samples from all females opportunistically (N  
207 = 140, median per female = 4, range = 3 - 4). We followed a two-step alcohol-silica storage protocol  
208 (Nsubuga, et al., 2004), after which the samples were stored at room temperature until DNA  
209 extraction.

210

### 211 *Dyad characteristics*

212         Dominance difference: To account for power asymmetries between females, we used Elo-  
213 rating (R package EloRating, Neumann & Kulik, 2013), which reflects an individual's success in  
214 agonistic interactions and is based on temporal sequences of decided (clear winner and loser)  
215 agonistic interactions (Albers & de Vries, 2001; Neumann et al., 2011). We made use of direct  
216 aggressive interactions (i.e. threats, hits, chases, bites) and displacements or supplantations (i.e. one  
217 individual approaches another one without any menacing behaviour and the other leaves without  
218 protesting; for further definitions and more details, see Duboscq et al., 2013, Thierry et al., 2000)  
219 taken from all agonistic data collected during focal observations and *ad libitum* data. At the beginning  
220 of the observation period, each individual in a group starts with a rating of 1000, which is updated, i.e.  
221 increased or decreased, after each agonistic interaction based on the outcome of the interaction (won  
222 or lost), the previous ratings of both opponents and a determined factor, k (here k = 100, following  
223 Neumann et al., 2011). As we aggregated all other behavioural data over the entire study period, we

224 used the female Elo-rating at the end of the study period. We then computed the absolute difference of  
225 the Elo ratings (hereafter termed dominance difference) between the two members of a dyad.

226 Kinship: DNA was extracted from 100–150 mg of faeces with the GEN-IAL® All-tissue  
227 DNA extraction kit following the manufacturer’s instructions. We amplified 12 short-tandem repeats  
228 (or microsatellites - 10 tetranucleotide loci and 2 dinucleotide loci), proven to be informative in  
229 humans and other primates (see Appendix). We used a two-step multiplex chain polymerase reaction  
230 (PCR) approach (Arandjelovic et al., 2009). In the first step, all loci were amplified in a single  
231 reaction in an Eppendorf® Master Gradient machine following cycles of denaturation, annealing and  
232 elongation (see Appendix). We followed multiplex PCRs with singleplex PCRs, using the same  
233 protocol but for each primer separately (see Appendix). Singleplex PCR products were then  
234 sequenced in an ABI 3130xL sequencer. Allele sizes were finally read into PeakScanner (Applied  
235 Biosystems®). Given that we had several samples per individual, allele sizes were considered  
236 definitive when at least two different samples of the same individual produced the same results in at  
237 least four amplifications for heterozygotes and six for homozygotes (multi-tubes approach, Taberlet et  
238 al., 1996). Consensus genotypes were found for a median of 12 loci (range = 6 – 12) and processed  
239 using COANCESTRY® software, which provides two likelihood methods and five moment  
240 estimators of relatedness (Wang, 2011). We chose the dyadic maximum likelihood (DML) estimator  
241 of Milligan (2003) because it proved to be the most reliable estimator of the mother-infant’s  
242 theoretical degree of relatedness 0.5 (mean  $\pm$  SD = 0.51  $\pm$  0.12, N = 60 mother-infant pairs). DML  
243 between adult females ranged between 0 and 0.72 with a median of 0.05 (median<sub>PB</sub> = 0.05, range<sub>PB</sub> =  
244 0 – 0.53; median<sub>RI</sub> = 0.05, range<sub>RI</sub> = 0 – 0.72).

245 Age difference: We assessed the age category (young, middle-aged or old) females belonged  
246 to based on their reproductive history (e.g. number of dependent infants or cycling status) known  
247 since 2006, the shape of their nipples (e.g. short or long), the presence of physical injuries and their  
248 general appearance. Based on these categories, we then scored dyads as belonging to the same or to  
249 different age classes.

250

251 *Bond characteristics*

252 The strength of dyadic social bonds was quantified with the Composite Sociality Index or CSI  
 253 (Silk et al., 2006b). It is built from matrices of dyadic social interactions and was calculated as  
 254 follows:

$$255 \quad CSI_{ij} = \left[ \left( \frac{G_{ij}}{G} \right) + \left( \frac{P_{ij}}{P} \right) + \left( \frac{P_{posij}}{P_{pos}} \right) \right] / 3,$$

256 where  $G_{ij}$  is the grooming rate (duration of grooming given and received in minutes per hour of  
 257 dyadic observation time) between individual  $i$  and  $j$  and  $G$  is the mean grooming rate across all dyads  
 258 in the group;  $P_{ij}$  is the rate of close proximity (number of instances females were within one body-  
 259 length of each other per hour of dyadic observation time) between individual  $i$  and  $j$  and  $P$  the mean  
 260 proximity rate for all dyads in the group and  $P_{posij}$  is the rate of positive outcome upon approach  
 261 (number of close proximity approaches followed by affiliation per hour of dyadic observation time)  
 262 between individual  $i$  and  $j$  and  $P_{pos}$  the mean rate of positive outcome upon approach for all dyads in  
 263 the group. A CSI above the average of all dyads in the group (equal to 1) indicates a strong bond (Silk  
 264 et al., 2006b).

265 Bond temporal variation (hereafter predictability) was assessed over three periods of six  
 266 months each. We calculated the CSI again for each dyad for each period, then computed the  
 267 coefficient of variation (CV, standard deviation divided by the mean) over the three CSIs for each  
 268 dyad (Majolo, Ventura, & Schino, 2010). The higher the CV, the lower the predictability of CSI  
 269 values across the three periods, i.e. the more CSI values varied across the three periods. Note that, in  
 270 this way, even weak but stable bonds will be considered predictable.

271 Bond equitability represents how balanced social exchanges are within a dyad (Silk et al.,  
 272 2013). The Equitability Index (EI) was calculated as a composite symmetry index (Silk et al., 2013),  
 273 computed from symmetry indices of the behaviours composing the CSI and was calculated as:

$$274 \quad EI_{ij} = \left[ \left( 1 - \left| \frac{G_{ij} - G_{ji}}{G_{ij} + G_{ji}} \right| \right) + \left( 1 - \left| \frac{P_{ij} - P_{ji}}{P_{ij} + P_{ji}} \right| \right) + \left( 1 - \left| \frac{P_{posij} - P_{posji}}{P_{posij} + P_{posji}} \right| \right) \right] / 3,$$

275 where  $G$  is grooming duration,  $P$  the rate of being in close proximity,  $P_{pos}$  the rate of positive outcome  
 276 upon approach, and  $i$  and  $j$  the individuals in the dyad. An index of 1 indicates perfect equitability  
 277 between the two individuals in the dyad, while 0 indicates that one individual alone was responsible

278 for all grooming and proximity interactions. Note that this index takes into account the directionality  
279 of interactions.

280

### 281 *Fitness-related behaviours*

282 We defined coalitionary support as a focal female intervening aggressively or peacefully in  
283 support of another female or receiving such an intervention herself during an aggressive interaction  
284 with another individual (Petit & Thierry, 1994; Duboscq et al., 2014). We calculated the frequency of  
285 support as the number of support instances over the total number of aggressive interactions each  
286 member of the dyad was separately involved in (Duboscq et al., 2014). Due to the low frequency of  
287 occurrences, for subsequent analyses we transformed this variable into a binary variable, i.e. the  
288 behaviour did or did not occur within the dyad (Duboscq et al., 2014). We calculated the frequency of  
289 feeding in proximity as the number of point samples spent feeding while other females were within 5-  
290 body-length proximity, controlling for overall dyadic proximity and observation time. Hourly  
291 frequencies of aggression were taken from Duboscq and colleagues (2013).

292

### 293 Statistical analyses

294

### 295 *Structure of social bonds*

296 We first tested for correlations between the three bond characteristics to assess their  
297 relationships with each other and to test the prediction that stronger bonds would be more predictable  
298 and equitable than weaker bonds. We built symmetric matrices of the CSI scores, the CVs and the EIs  
299 before running a Quadratic Assignment Permutation (QAP) procedure with 1,000 permutations  
300 between those matrices two-by-two (function qaptest in the sna package; Butts, 2008). We then built  
301 three (generalised) linear mixed models (GLMM, Bolker et al., 2008), one for each of the social bond  
302 components as response variables, including relatedness (DML), absolute Elo-rating difference (Elo#)  
303 and age difference (age#, as a categorical variable, close/distant) as predictors and member 1 and  
304 member 2 of the dyad and group as random effects.

305

306 *Function of social bonds*

307 We built three models to investigate the value of social bonds, with the occurrence of  
308 coalitionary support, feeding-in-proximity rate and aggression rate as response variables and bond  
309 strength (CSI), predictability (CV), equitability (EI), as well as relatedness (DML), absolute Elo-  
310 rating difference (Elo#) and age difference (age#) as predictors and member 1 and member 2 of the  
311 dyad and group as random effects.

312 All analyses were done in R version 3.2.1 (R Development Core Team, 2015). We  
313 implemented GLMMs with a Gaussian (and Maximum Likelihood) or binomial error structure using  
314 the function “lmer” from the package “lme4” (v. 1.1-11, Bates, Maechler, Bolker, & Walker, 2015).  
315 We transformed numerical variables whenever necessary (log, square-root or fourth root) and  
316 standardised all numeric variables to a mean of 0 and a standard deviation of 1 for optimal fitting and  
317 the subsequent interpretation or comparison of model estimates. For all models, we checked a variety  
318 of assumptions and diagnostics (normally distributed and homogeneous residuals, variance inflation  
319 factors  $< 2$ , Cooks’ distance, dfbetas; Field, Miles, & Field, 2012). No obvious violation of  
320 assumptions was detected. We tested the final full model (including all fixed and random effects)  
321 against an informed null model, i.e. including only the intercept, control factors (absolute Elo  
322 difference, DML, and age difference) and random factors, using likelihood ratio tests (LRT). We used  
323 95% confidence intervals to assess whether a predictor significantly contributed to explaining the  
324 response variable (interval excluding 0). For clarity, detailed results from the models are given in  
325 Tables A1 to A3 in the Appendix.

326

327 *Testing the effect of kinship measured by a microsatellite-based estimator*

328 Microsatellite-based relatedness estimators have been deemed unreliable for accurately  
329 measuring genetic relatedness in populations without pedigree information (Csilléry et al., 2006; van  
330 Horn, Altmann, & Alberts, 2008). We tackled this issue by using the approach suggested by Tinsley  
331 Johnson and colleagues (2014) of controlling for measurement error in relatedness estimates by  
332 running models repeatedly with a random amount of error added to the observed relatedness value of  
333 a given dyad. In our data, the maximum observed difference between the estimated relatedness

334 (DML) and the true theoretical relatedness ( $r = 0.5$ ) of all 60 known mother-infant pairs was 0.41. We  
335 therefore introduced an error taken from a random uniform distribution of numbers between -0.41 and  
336 +0.41, which we feel is conservative as 95% of the DML values for known mother-infant pairs were  
337 within 0.25 of the pairs' true relatedness ( $r = 0.5$ ). Our costumed simulation proceeded in four steps:  
338 1/ add an error between -0.41 and +0.41 to the DML index of all female-female dyads in the dataset,  
339 2/ run the models again with the modified DML index, 3/ perform a likelihood ratio test between the  
340 full model with modified DML and a reduced model excluding modified DML and 4/ determine the  
341 number of simulations in which the significance of the effect of the modified DML index on the  
342 response variable was different from the tests with the original models with the original data. The  
343 DML index was not a significant predictor of the response variable in 96% to 100% of the 10,000  
344 simulations, depending on the response variable, which indicates relatively robust results. Detailed  
345 results of these simulations are given in Table A3 of the Appendix.

346

## 347 **Results**

348

349 The structure of social bonds

350

351 CSI scores ranged from 0.05 to 3.54 in the R1 group and from 0.16 to 4.99 in the PB group,  
352 with a median of 0.89 and 0.88 respectively (Figure 1), indicating that in both groups, most bonds had  
353 average strength (mean equal to one in each group). The distribution of CSI scores, giving an  
354 assessment of how skewed dyadic affiliative behaviours are, was not right-skewed (Figure 1) as is  
355 typical in other species, indicating that females established and maintained affiliative relationships  
356 with numerous female partners. 45.7% (96/210 in R1) and 40% (42/105 in PB) of female dyads had a  
357 CSI score above the average of the group (i.e., above 1) and the CSI score of the top 10% of dyads  
358 was 2.34 in both groups (Figure 1). Females had a median of 9 (range 2–13) above-average (CSI > 1)  
359 relationships in R1 and 6 (range 2–11) in PB.

360 The mean coefficient of variation (R1 =  $0.66 \pm 0.01$  SD, PB =  $0.70 \pm 0.14$  SD) indicated  
361 moderate to low bond strength predictability across the three 6-months periods (Figure 2). 94% of all



362 females had at least one recurring partner over at least two periods amongst their three top partners  
363 and 47% of all females had at least one recurring partner over all three periods amongst their top three  
364 partners but no female had the same three recurring top partners across all three periods (Figure 2).

365 Bond equitability was overall relatively moderate ( $\text{mean}_{R1} = 0.22 \pm 0.47 \text{ SD}$ ,  $\text{mean}_{PB} = 0.27 \pm$   
366  $0.13 \text{ SD}$ ), indicating relatively balanced social exchanges amongst the two members of a dyad.

367 All three characteristics were positively correlated in both groups (QAP tests, PB:  $N = 105$ :  
368 strength-predictability:  $\rho = 0.59$ ,  $p = 0.001$ ; strength-equitability:  $\rho = 0.91$ ,  $p < 0.001$ ;  
369 predictability-equitability:  $\rho = 0.95$ ,  $p < 0.001$ ; R1:  $N = 210$ : strength-predictability:  $\rho = 0.76$ ,  $p =$   
370  $0.001$ ; strength-equitability:  $\rho = 0.85$ ,  $p < 0.001$ ; predictability-equitability:  $\rho = 0.65$ ,  $p < 0.001$ ),  
371 indicating that stronger bonds were more equitable but less predictable over time than weaker bonds  
372 and that more predictable bonds were less equitable than less predictable ones.

373 Female dyads with smaller dominance differences had significantly stronger, more  
374 predictable and more equitable social bonds than those with greater dominance differences (strength:  
375  $\text{LRT}_{\text{full-null}}: \chi^2 = 39.45$ ,  $\text{d.f.} = 3$ ,  $p < 0.001$ ;  $\beta \pm \text{SE} [95\% \text{ CI}] = -0.33 \pm 0.06 [-0.45 - -0.20]$ ;  
376 predictability:  $\text{LRT}_{\text{full-null}}: \chi^2 = 8.53$ ,  $\text{d.f.} = 3$ ,  $p = 0.036$ ;  $\beta \pm \text{SE} = 0.11 \pm 0.06 [0.00 - 0.19]$ ;  
377 equitability:  $\text{LRT}_{\text{full-null}}: \chi^2 = 21.32$ ,  $\text{d.f.} = 3$ ,  $p < 0.001$ ;  $\beta \pm \text{SE} = -0.25 \pm 0.06 [-0.37 - -0.13]$ ; Table  
378 A1; Figure 2). None of the tested bond characteristics were related to the relatedness and age  
379 difference of the dyads (Table A1).

380

381 The value of social bonds

382

383 Rates of support between adult females were low; we recorded a total of 206 coalitionary  
384 events (48 peaceful interventions, 158 aggressive ones) in the two groups over a total of 3,208  
385 aggressive interactions. A female was found to be feeding in the proximity of another female around  
386 once in every ten times she was found in proximity with that female (median = 0.11, range = 0.00 –  
387 0.39, feeding scan per proximity scan). Females engaged in aggressive interactions with each other  
388 about once every two hours (for details see Duboscq and colleagues (2013)).

389 Coalitionary support was not explained by any of the tested bond components ( $LRT_{full-null}: \chi^2$   
390  $= 1.86$ , d.f. = 3,  $p = 0.603$ ; Table A2; Figure 3). In contrast, dyads that had stronger and less  
391 predictable bonds fed more often in proximity than others ( $LRT_{full-null}: \chi^2 = 11.15$ , d.f. = 3,  $p = 0.011$ ;  
392 strength:  $\beta \pm SE$  [95% CI] =  $0.16 \pm 0.06$  [0.03 - 0.28]; predictability:  $\beta \pm SE$  [95% CI] =  $0.20 \pm 0.07$   
393 [0.04 - 0.35]; Table A2; Figure 3). Finally, dyads that had stronger bonds were more frequently  
394 aggressive towards each other, while those with more equitable bonds fought less often than others  
395 ( $LRT_{full-null}: \chi^2 = 23.54$ , d.f. = 3,  $p < 0.001$ ; strength:  $\beta \pm SE$  [95% CI] =  $0.25 \pm 0.06$  [0.12 - 0.37];  
396 equitability:  $\beta \pm SE$  [95% CI] =  $-0.20 \pm 0.05$  [-0.33 - -0.10]; Table A2; Figure 3).

397

## 398 Discussion

399

400 Social bonds in the studied female crested macaques showed contrasting patterns in their  
401 structure and value compared to other species with nepotistic hierarchical societies for which  
402 comparable data are available. Bonds were mostly of average strength (i.e. median strength close to 1  
403 and moderately left-skewed distribution), relatively equitable but only moderately enduring over the  
404 19 months of the study. Social bonds were stronger, more predictable and more equitable amongst  
405 females close in dominance status than others, but not amongst kin or age peers. The three  
406 components of social bonds were positively related but had differential effects on the three fitness-  
407 related behaviours investigated. Thus, our predictions that dyad characteristics have little influence on  
408 social bond components and that these components have relatively weak effects on fitness-related  
409 behaviours were mostly fulfilled, showing the extent to which the studied females can express great  
410 degrees of freedom within their established network of relationships.

411 We observed a relatively weak influence of kinship on female social relationships; compared  
412 with less related females, more related females did not form significantly stronger, more predictable  
413 or equitable bonds, revealing a weakly nepotistic society. Several factors may contribute to the  
414 observed weak nepotism. First and foremost, the fact that we could not distinguish matriline, due to  
415 the lack of a pedigree, and could not differentiate maternal from paternal relatives may have hidden  
416 kinship effects on behaviour. In several mammals, paternal relatives interact with each other

417 substantially more often than with non-kin but also substantially less often than with maternal kin  
418 (baboons: Smith, Albers, & Altmann, 2003; hyenas: Wahaj et al., 2004; rhesus macaques: Wenzel et  
419 al. 2013). Male reproductive skew and group tenure influence the proportion of paternal relatives in a  
420 group (Widdig 2013); high male reproductive skew and short male tenure, which is a characteristic of  
421 this population of crested macaques (Higham et al., 2012; Marty et al., in press), can lead to a  
422 relatively high proportion of paternal relatives. These conditions have been hypothesised to be a  
423 strong driver for high social tolerance amongst female macaques (Schülke & Ostner, 2008). Indeed, a  
424 weak kin bias amongst numerous paternal relatives may blur a strong kin bias between fewer maternal  
425 relatives. A hint to that effect is the relatively consistent positive influence of small dominance  
426 differences (most likely maternal rather than paternal relatives) on social behaviour. Another related  
427 factor is the use of a continuous measure of kinship which may have obscured any distinction between  
428 kin categories (e.g. mother-daughter, sister-sister) that may be of greater significance for individuals  
429 (Kapsalis & Berman, 1996).

430         Notwithstanding these limitations, these results indicate that an overall weak kin bias in social  
431 relationships constitutes a shared characteristic of tolerant societies, such as Barbary (*M. sylvanus*)  
432 and Tonkean macaques (*M. tonkeana*), which contrasts with more despotic species of macaques and  
433 other primates such as baboons (Paul, 2006; Thierry, 2007; Cords, 2012). This finding is also  
434 consistent with predictions from the primate socioecological model (Sterck, Watts, & van Schaik,  
435 1997); a weakly nepotistic hierarchy may indeed stem from the low profitability of kin support when  
436 rank-related fitness benefits are not pronounced because direct competition for food is relatively low  
437 (Chapais, 2004). The study population indeed lives in a felid-predator-free, food-abundant  
438 environment (O'Brien & Kinnaird, 1997). Higher-ranking females appear to reproduce better than  
439 low-ranking females, but this pattern varies with demographic and ecological conditions (Kerhoas et  
440 al., 2014). The observed weak nepotistic pattern is common in other animal societies with high  
441 cooperation levels and even more flexible social structures (e.g., chimpanzees: Langergraber, Mitani,  
442 & Vigilant, 2007; meerkats: Clutton-Brock, 2009; hyenas: East & Hofer, 2010; raccoons: Hirsch,  
443 Prange, Hauver, & Gehrt, 2013), which suggests great potential for relatively high degrees of freedom  
444 in social relationships in the crested macaques too.

445 Overall, female-female bond strength did not show the typical left-skewed distribution of  
446 more despotic primate species (at least those for which similar data are available: Assamese macaque  
447 (*M. assamensis*) males: Kalbitz, Ostner, & Schülke, 2016; chacma baboon females: Silk et al., 2006b;  
448 Barbary macaque males: Young, Majolo, Schülke, & Ostner, 2014), indicating that female crested  
449 macaques formed mostly average-strength bonds and very few strong bonds. This pattern is strikingly  
450 different from the typical few strong-many weak bonds pattern found in the above mentioned species  
451 and confirms the propensity of female crested macaques to form diverse and large affiliative networks  
452 (Duboscq et al., 2013). Furthermore, bond strength and equitability were negatively related to  
453 predictability, such that stronger and more equitable bonds were less predictable than weaker and less  
454 equitable bonds. This result indicates the limited endurance of strong bonds compared to average or  
455 weak bonds, which also contrasts with findings in other species in which strong bonds appeared very  
456 stable over time (e.g., Kalbitz et al., 2016; Mitani, 2009; Silk et al., 2006a; Young et al., 2014). The  
457 temporal variation found in bond strength suggests that preferred associations of female crested  
458 macaques are of an opportunistic nature. This is likely to be related to varying social contexts that we  
459 did not address in our current study, such as the presence or absence of dependent infants (variable  
460 throughout the year), social instability due to demographic changes (e.g. frequent male migration) or  
461 changes in environmental conditions (e.g. massive seasonal fruiting of fig trees), all of which have  
462 been shown to modulate relationships between group members (birds and mammals: Wrangham &  
463 Rubenstein, 1986; baboons: Barrett & Henzi, 2001; Henzi et al., 2009; hyenas: East & Hofer, 2010).

464 The most consistent finding in our study was the effect of dominance rank differences, i.e.  
465 strong, predictable and equitable bonds were more likely to be formed by dyads with small  
466 differences in dominance rank. In female primates and hyenas, the maintenance of these bonds has  
467 been linked to competition for social partners, as females struggle for access to the highest-ranking  
468 females and end up socialising most with affiliates of adjacent ranks due to competitive exclusion  
469 (Seyfarth, 1977). However, because in the study population power asymmetries between females were  
470 relatively moderate, counter-aggression frequent and affiliative and proximity networks quite diverse  
471 (Duboscq et al., 2013), we argue that social competition was low and competitive exclusion was  
472 ineffective so this hypothesis does not provide a satisfying explanation for our results. Social bond

473 formation and maintenance may instead involve the reciprocal exchange of social commodities if  
474 adjacently-ranked partners were generally more similar – in terms of personality, energetic needs or  
475 reproductive state – or competent partners in cooperation (Chapais, 2006; Schino & Aureli, 2009).  
476 These reciprocal exchanges could be highly dynamic and opportunistic in a biological market  
477 susceptible to environmental and social conditions (Noë & Hammerstein, 1994; Barrett & Henzi,  
478 2006), thereby generating a potential source of variation in the endurance of social bonds too.

479         The potential opportunistic nature of these social bonds seems also apparent in the analyses of  
480 their potential adaptive value. Previous studies on the same population showed somewhat contrasting  
481 results. The strength of social bonds was related to anti-predator responses, indicating their  
482 importance in threatening situations (Micheletta et al., 2012). However, the symmetry and  
483 predictability, rather than strength, of social bonds influenced the occurrence of conflict management  
484 behaviour. Furthermore, the occurrence of reconciliation – an important mechanism of social  
485 cohesion – appears to function as appeasement, a short-term tactic, rather than to repair relationships,  
486 a more long-term strategy (Duboscq et al., 2014). In the current study, variance in bond characteristics  
487 helped only to a certain extent to explain variation in three additional fitness-related behaviours. First,  
488 more strongly bonded females fed more often in proximity but also fought more often with each other  
489 than females with weaker bonds. This indicates that more strongly bonded dyads may be more  
490 resilient to disruptions of their bond (by aggression) over food than less strongly bonded dyads  
491 (Aureli, Fraser, Schaffner, & Schino, 2012). Second, this is consistent with the finding that partners  
492 with less predictable bonds also fed more often in proximity, as stronger bonds tended to be less  
493 predictable. The link between bond predictability and co-feeding frequency suggests either that  
494 partners with enduring relationships avoid endangering the stability of their relationship over feeding  
495 competition, or that the endurance of bonds is affected by another factor that we have not considered,  
496 for example female energetic needs (perhaps in relation with lactation or oestrous). Third, the  
497 negative relationship between equitability and aggression rate could indicate that less equitable dyads  
498 often need to negotiate their relationship through engaging in mild aggression. This could be the case  
499 if one partner gets frustrated to be at the lesser end of the social exchange, especially since less  
500 equitable bonds also tended to be weaker in strength. These results stress the need to consider more

501 than one dimension of social bonds simultaneously in order to get a more integrative picture of how  
502 animals balance the costs and benefits of social bonds. Experiments involving cooperative tasks could  
503 help disentangle the respective weight of bond characteristics in social decision-making in terms of  
504 partner choice, coalition formation, trust or punishment.

505 Overall, it appears that the female crested macaques under study generally form a dynamic  
506 number and large diversity of good average partners, rather than a tight network of enduring strong  
507 ones, perhaps similar to what has been suggested for males of the same species (Neumann, 2013;  
508 Neumann, Agil, Widdig, & Engelhardt, 2013). As such, females seem able to express great degrees of  
509 social freedom with regards to their dominance and kin relationships (Thierry, 1990; Butovskaya,  
510 2004). Nevertheless, females also seem to specifically rely on certain partners, with whom they have  
511 strong, predictable or equitable bonds, in specific contexts or in especially challenging situations  
512 (Duboscq et al., 2014; Micheletta et al., 2012). This “many-good-friends” strategy can be costly  
513 temporally and energetically, but it can also bring a wide range of benefits, including enhanced  
514 negotiation skills, improved collective decision-making, and facilitated cooperation in joint-action  
515 problems (Petit, Desportes, & Thierry, 1992; McComb & Semple, 2005; Hare, Melis, Woods,  
516 Hasting, & Wrangham, 2007; Sueur & Petit, 2008; Thierry et al., 2008).

517 The contrast between the stability of the social structure of macaques, in general, and the  
518 degrees of freedom shown by female crested macaques, in particular, in establishing and maintaining  
519 relationships leads to questions about the temporal dynamics of social bonds and the short- and long-  
520 term reciprocity of social exchanges within stable societies. It highlights the need to consider more  
521 carefully the whole network of bonds, weak and strong, in a more integrated way. More importantly,  
522 fitness-related behaviours, like coalitionary support, are presumably based on long-term alliances.  
523 Thus, to what extent patterns in so-called strategic behaviours can actually resist the magnitude of  
524 changes in social bonds is currently not clear. Investigating this question could help to determine how  
525 dynamic societies actually are and whether species or population differences in dynamics exist.  
526 Fluctuations in social networks are indeed pervasive in species with flexible (Schradin, 2013) or  
527 seasonal sociality (Blumstein, 2013; Brent et al., 2013; Prange, Gehrt, & Hauver, 2011). Whether our  
528 findings reflect a pattern more common than previously thought or are typical for this study

529 population remains to be investigated and requires comparative studies. This is of tremendous  
530 importance to better understand the social components of fitness and the mechanisms linking sociality  
531 to fitness.

532

### 533 **Acknowledgments**

534

535 This article is dedicated to Ugiek and Antri. We gratefully acknowledge the permission of the  
536 Indonesian State Ministry of Research and Technology (RISTEK), the Directorate General of Forest  
537 Protection and Nature Conservation (PHKA) in Jakarta and the Department for the Conservation of  
538 Natural Resources (BKSDA) in Manado, particularly Pak Untung and Pak Yunus, to conduct this  
539 research in the Tangkoko-Batuangus Nature Reserve. We thank all the members of the Macaca Nigra  
540 Project for their support in the field, and especially Dwi Yandhi Febryanti and Jérôme Micheletta for  
541 their help with data collection.

542

### 543 **Compliance with ethical standards**

544

545 Funding: This study was funded by the Volkswagen Foundation (funding initiative Evolutionary  
546 Biology, grant I/84 200) and Primate Conservation Inc. (grant PCI 757) to JD and by a grant from the  
547 German Research Council (EN 916/2) to AE.

548 Conflict of Interest: All authors declare they have no conflict of interest of any kind.

549 Ethical approval: This research adheres to all legal requirements and guidelines of the German and  
550 Indonesian governments and institutions, and to the ASAB/ASB guidelines for the treatment of  
551 animals in behavioural research and teaching.

552

### 553 **References**

554 Albers, P. C. H., & de Vries, H. (2001). Elo-rating as a tool in the sequential estimation of dominance  
555 strengths. *Animal Behaviour*, *61*, 489–495. doi: 10.1006/anbe.2000.1571

556 Arandjelovic, M., Guschanski, K., Schubert, G., Harris, T. R., Thalmann, O., Siedel, H., & Vigilant,  
557 L. (2009) Two-step multiplex polymerase chain reaction improves the speed and accuracy of

558 genotyping using DNA from noninvasive and museum samples. *Molecular Ecology Resources*, 9, 28–  
559 36. doi: 10.1111/j.1755-0998.2008.02387.x

560 Armitage, K. B., & Schwartz, O. A. (2000). Social enhancement of fitness in yellow-bellied marmots.  
561 *Proceedings of the National Academy of Science*, 97, 12149–12152. doi:10.1073/pnas.200196097

562 Aureli, F., Fraser, O. N., Schaffner, C. M., & Schino, G. (2012). The regulation of social  
563 relationships. In: J. C. Mitani, J. Call, P. M. Kappeler, R. Palombit, & J. Silk (Eds.) *The evolution of*  
564 *primate societies* (pp. 531–551). Chicago, U.S.A.: The University of Chicago Press.

565 Barocas, A., Ilany, A., Koren, L., Kam, M., & Geffen, E. (2011). Variance in centrality within rock  
566 hyrax social networks predicts adult longevity. *PLoS ONE*, 6, 1–8. doi:10.1371/journal.pone.0022375

567 Bakshy, E., Rosenn, I., Marlow, C., & Adamic, L. (2012). The role of social networks in information  
568 diffusion. In *Proceedings of the 21st international conference on World Wide Web* (pp. 519–528).  
569 New York: ACM. doi:10.1145/2187836.2187907

570 Bardi, M., Shimizu, K., Fujita, S., Borgognini Tarli, S. M., & Huffman, M. (2001). Social behavior  
571 and hormonal correlates during the perinatal period in Japanese macaques. *Hormones & Behavior*, 39,  
572 239–246. doi: 10.1006/hbeh.2001.1651

573 Barrett, L., & Henzi, P. (2001). Constraints on relationship formation among female primates.  
574 *Behaviour*, 139, 263–289. doi:10.1163/156853902760102672.

575 Barrett, L., & Henzi, P. (2006). Monkeys, market and minds: biological markets and primate sociality.  
576 In: P. M. Kappeler & C. P. van Schaik (Eds.) *Cooperation in Primates and Humans: Mechanisms and*  
577 *Evolution* (pp. 209–232). Berlin, Germany: Springer.

578 Bates, D. M., Maechler, M., Bolker, B., & Walker, S. (2015). Fitting linear mixed-effects models  
579 using lme4. *Journal of Statistical Software*, 67, 1–48. doi:10.18637/jss.v067.i01.

580 Blumstein, D. T. (2013). Yellow-bellied marmots: insights from an emergent view of sociality.  
581 *Philosophical Transactions of the Royal Society B: Biological Sciences*, 368, 20120349.  
582 doi:10.1098/rstb.2012.0349

583 Bolker, B., Brooks, M. E., Clark, C. J., Geange, S. W., Poulsen, J. R., Stevens, M. H. H., & White, J.-  
584 S. S. (2008) Generalized linear mixed models: a practical guide for ecology and evolution. *Trends in*  
585 *Ecology and Evolution*, 24, 127–135. doi: 10.1016/j.tree.2008.10.008

586 Brent, L. J. N., MacLarnon, A. M., Platt, M. L., & Semple, S. (2013). Seasonal changes in the  
587 structure of rhesus macaque social networks. *Behavioral Ecology and Sociobiology*, 67, 349–359. doi:  
588 10.1007/s00265-012-1455-8

589 Butovskaya, M. L. (2004). Social space and degrees of freedom. In: B. Thierry, M. Singh, & W.  
590 Kaumanns (Eds.) *Macaque societies, a model for the study of social organisation* (pp. 182–185).  
591 Cambridge, U.K.: Cambridge University Press.

592 Butts, C. T. (2008). Social Network Analysis with sna. *Journal of Statistical Software*, 24, 1–51.

593 Cameron, E. Z., Setsaas, T. H., & Linklater, W. L. (2009). Social bonds between unrelated females  
594 increase reproductive success in feral horses. *Proceedings of the National Academy of Sciences*, 106,  
595 13850–13853. doi:10.1073/pnas.0900639106

596 Carter, K. D., Seddon, J., Frère, C., Carter, J. K., & Goldizen, A. W. (2013). Fission-fusion dynamics  
597 in wild giraffes may be driven by kinship, spatial overlap and individual social preferences. *Animal*  
598 *Behaviour*, 85, 385–394. doi:10.1016/j.anbehav.2012.11.011

599 Chapais, B. (2006). Kinship, competence and cooperation in primates. In: P. M. Kappeler, & C. P.  
600 van Schaik (Eds.) *Cooperation in primates and humans: mechanisms and evolution* (pp. 47–64).  
601 Berlin: Springer.



602 Chapais, B. (2004). How kinship generates dominance structures: a comparative perspective. In B.  
603 Thierry, M. Singh, & W. Kaumanns (Eds.), *Macaque Societies* (pp. 186–204). Cambridge, U.K.:  
604 Cambridge University Press.

605 Clutton-Brock, T. H. (2009). Cooperation between non-kin in animal societies. *Nature*, *462*, 51–57.  
606 doi:10.1038/nature08366

607 Cooper, M. A., & Bernstein, I. S. (2008) Evaluating dominance styles in Assamese and rhesus  
608 macaques. *International Journal of Primatology*, *29*, 225–243. doi:10.1007/s10764-008-9236-y

609 Cords, M. (2012). The behavior, ecology and social evolution of cercopithecine monkeys. In: J. C.  
610 Mitani, J. Call, P. M. Kappeler, R. Palombit, & J. Silk (Eds.) *The evolution of primate societies* (pp.  
611 91–112). Chicago, U.S.A.: The University of Chicago Press.

612 Csilléry, K., Johnson, T., Beraldi, D., Clutton-Brock, T. H., Coltman, D., Hansson, B., Spong, G., &  
613 Pemberton, J. M. (2006) Performance of marker-based relatedness estimators in natural populations of  
614 outbred vertebrates. *Genetics*, *173*, 2091–2101.

615 Csermely, P. (2004). Strong links are important, but weak links stabilize them. *Trends in Biochemical*  
616 *Sciences*, *29*, 331–334. doi:10.1016/j.tibs.2004.05.004

617 D'Amato, F. R., Troisi, A., Scucchi, S., & Fuccillo, R. (1982) Mating season influence on  
618 allogrooming in a confined group of Japanese macaques: a quantitative analysis. *Primates*, *23*, 220–  
619 232. doi: 10.1093/beheco/aru099

620 Duboscq, J., Agil, M., Engelhardt, A., & Thierry, B. (2014). The function of post-conflict  
621 interactions: new prospects from the study of a tolerant species of primate. *Animal Behaviour*, *87*,  
622 107–120. doi:10.1016/j.anbehav/2013-10-1018

623 Duboscq, J., Micheletta, J., Agil, M., Hodges, K., Thierry, B., & Engelhardt, A. (2013). Social  
624 tolerance in wild female crested macaques, *Macaca nigra*, in Tangkoko-Batuangus Nature Reserve,  
625 Sulawesi, Indonesia. *American Journal of Primatology*, *75*, 361–375. doi:10.1002/ajp.22114

626 Duboscq, J., Neumann, C., Perwitasari-Farajallah, D., & Engelhardt, A. 2008. Daytime birth of a baby  
627 crested black macaque (*Macaca nigra*) in the wild." *Behavioural processes*, *79*, 81-84.  
628 doi:10.1016/j.beproc.2008.04.010

629 East, M. L., & Hofer, H. (2010) Social environments, social tactics and their fitness consequences in  
630 complex mammalian societies. In: T. Székely T, A. J. Moore, J. Komdeur (Eds) *Social Behaviour:*  
631 *Genes, Ecology and Evolution* (pp. 360-391). Cambridge, U.K.: Cambridge University Press.

632 Field, A., Miles, J., & Field, Z. (2012). *Discovering statistics using R*. London, U.K.: SAGE  
633 Publications Ltd.

634 Fraser, O. N., & Bugnyar, T. (2011) Ravens reconcile after aggressive conflicts with valuable  
635 partners. *PLoS ONE*, *6*, e18118–e18118. doi:10.1371/journal.pone.0018118

636 Frère, C. H., Krützen, M., Mann, J., Connor, R. C., Bejder, L., & Sherwin, W. B. (2010). Social and  
637 genetic interactions drive fitness variation in a free-living dolphin population. *Proceedings of the*  
638 *National Academy of Sciences*, *107*, 19949–19954. doi:10.1073/pnas.1007997107

639 Godde, S., Côté, S. D., & Réale, D. (2015). Female mountain goats, *Oreamnos americanus*, associate  
640 according to kinship and reproductive status. *Animal Behaviour*, *108*, 101–107.  
641 doi:10.1016/j.anbehav.2015.07.005

642 Hare, B., Melis, A. P., Woods, V., Hasting, S., & Wrangham, R. W. (2007). Tolerance allows  
643 bonobos to outperform chimpanzees on a cooperative task. *Current Biology*, *17*, 619–623.  
644 doi:10.1016/j.cub.2007.02.040

645 Hemelrijk, C. K. (1999). An individual-orientated model of the emergence of despotic and egalitarian  
646 societies. *Proceedings of the Royal Society B: Biological Sciences*, *266*, 361–369.  
647 doi:10.1098/rspb.1999.0646

- 648 Henzi, S. P., Lusseau, D., Weingrill, T., van Schaik, C. P., & Barrett, L. (2009). Cyclicality in the  
649 structure of female baboon social networks. *Behavioral Ecology and Sociobiology*, 63, 1015–1021.  
650 doi:10.1007/s00265-009-0720-y
- 651 Hirsch, B. T., Prange, S., Hauver, S. A., & Gehrt, S. D. (2013) Genetic relatedness does not predict  
652 racoon social network structure. *Animal Behaviour*, 85, 463–470. doi:10.1016/j.anbehav.2012.12.011
- 653 Hirsch, B. T., Stanton, M. A., Maldonado, J. E. (2012). Kinship shapes affiliative social networks but  
654 not aggression in ring-tailed coatis. *PLoS ONE*, 7, e37301. doi:10.1371/journal.pone.0037301
- 655 Ilany, A., Booms, A. S., & Holekamp, K. E. (2015). Topological effects of network structure on long-  
656 term social network dynamics in a wild mammal. *Ecology Letters*, 687–695. doi:10.1111/ele.12447
- 657 Kalbitz, J., Ostner, J., & Schülke, O. (2016). Strong, equitable and long-term social bonds in the  
658 dispersing sex in Assamese macaques. *Animal Behaviour*, 113, 13-22.  
659 doi:10.1016/j.anbehav.2015.11.005
- 660 Kapsalis, E., & Berman, C. M. (1996). Models of affiliative relationships among free-ranging rhesus  
661 monkeys (*Macaca mulatta*) I. Criteria for kinship. *Behaviour*, 133, 1209–1234.  
662 doi:10.1163/156853996X00387
- 663 Kerhoas, D., Mundry, R., Perwitasari-Farajallah, D., Agil, M., Widdig, A. & Engelhardt, A. (2014).  
664 Social and ecological factors influencing offspring survival in wild macaques. *Behavioral Ecology*,  
665 25, 1164–1172. doi:10.1093/beheco/aru099
- 666 Keverne, E. B., Martensz, N. D., & Tuite, B. (1989) Beta-endorphin concentrations in cerebrospinal  
667 fluid of monkeys are influenced by grooming relationships. *Psychoneuroendocrinology*, 14, 155–161.  
668 doi:10.1016/0306-4530(89)90065-6
- 669 King, A. J., Clark, F. E., & Cowlishaw, G. (2011) The dining etiquette of desert baboons: the roles of  
670 social bonds, kinship, and dominance in co-feeding networks. *American Journal of Primatology*, 73,  
671 1–7. doi:10.1002/ajp.20918
- 672 Langergraber, K. E., Mitani, J. C., & Vigilant, L. (2007). The limited impact of kinship on  
673 cooperation in wild chimpanzees. *Proceedings of the National Academy of Science*, 104, 7786–7790.  
674 doi:10.1073/pnas.0611449104
- 675 Majolo, B., Ventura, R., & Schino, G. (2010). Assymetry and dimensions of relationships quality in  
676 the Japanese macaque (*Macaca fuscata yakui*). *International Journal of Primatology*, 31, 736–750.  
677 doi:10.1007/s10764-010-9424-4
- 678 Martin, P., & Bateson, P. (1993). *Measuring Behaviour - An Introductory Guide* (2nd ed). Cambridge,  
679 U.K.: Cambridge University Press.
- 680 Marty, P. R., Hodges, K., Agil, M., & Engelhardt, A. (in press). Alpha male replacements and delayed  
681 dispersal in crested macaques (*Macaca nigra*). *American Journal of Primatology*. doi:  
682 10.1002/ajp.22448
- 683 McComb, K., & Semple, S. (2005). Coevolution of vocal communication and sociality in primates.  
684 *Biology Letters*, 1, 381–385. doi: 10.1098/rsbl.2005.0366
- 685 Micheletta, J., Waller, B. M., Panggur, M. R., Neumann, C., Duboscq, J., Agil, M., & Engelhardt, A.  
686 (2012). Social bonds affect anti-predator behaviour in a tolerant species of macaques, *Macaca nigra*.  
687 *Proceedings of the Royal Society B: Biological Sciences*, 279, 4042–4050. doi:  
688 10.1098/rspb.2012.1470
- 689 Milligan, B. (2003). Maximum-likelihood estimation of relatedness. *Genetics*, 163, 1153–1167.
- 690 Mitani, J. C. (2009). Male chimpanzees form enduring and equitable social bonds. *Animal Behaviour*,  
691 77, 633–640. doi:10.1016/j.anbehav.2008.11.021
- 692 Neumann, C., & Kulik, L. (2013). EloRating: Animal dominance hierarchies by Elo Rating.  
693 <http://CRAN.R-project.org/package=EloRating>

694 Neumann, C. (2013). Achievement and maintenance of dominance in male crested macaques  
695 (*Macaca nigra*) (Doctoral thesis). Leipzig, Germany: University of Leipzig.

696 Neumann, C., Agil, M., Widdig, A., & Engelhardt, A. (2013). Personality of wild male crested  
697 macaques (*Macaca nigra*). *PLoS ONE*, 8, e69383. doi:10.1371/. doi: 10.1371/journal.pone.0069383

698 Neumann, C., Duboscq, J., Dubuc, C., Ginting, A., Irwan, A. M., Agil, M., Widdig, A., & Engelhardt,  
699 A. (2011) Assessing dominance hierarchies: validation and advantages of progressive evaluation with  
700 Elo-rating. *Animal Behaviour*, 82, 911–921. doi: 10.1016/j.anbehav.2011.07.016

701 Newberry, R., & Swanson, J (2001) Breaking Social Bonds. In L. J. Keeling, & H. W. Gonyou (Eds.),  
702 *Social Bonds in Farm Animals* (pp. 307–331). Oxon, U.K.: CABI Publishing.

703 Noë R., & Hammerstein, P. (1994). Biological markets: supply and demand determine the effect of  
704 partner choice in cooperation, mutualism and mating. *Behavioral Ecology and Sociobiology*, 35, 1–  
705 11. doi:10.1007/BF00167053

706 Nowak, M. A. (2006). Five Rules for the Evolution of Cooperation. *Science*, 314, 1560–1563.  
707 doi:10.1126/science.1133755

708 Nsubuga, A. M., Robbins, M. M., Roeder, A. D., Morin, P., Boesch, C., & Vigilant, L. (2004) Factors  
709 affecting the amount of genomic DNA extracted from ape faeces and the identification of an  
710 improved sample storage method. *Molecular Ecology*, 13, 2089–2094. doi: 10.1111/j.1365-  
711 294X.2004.02207.x

712 O'Brien, T. G., & Kinnaird, M. F. (1997). Behavior, diet, and movements of the Sulawesi crested  
713 black macaque (*Macaca nigra*). *International Journal of Primatology*, 18, 321–351. doi:  
714 10.1023/A:1026330332061

715 Ostner, J., & Schülke, O. (2014). The evolution of social bonds in primate males. *Behaviour*, 151,  
716 871–906. doi: 10.1163/1568539X-00003191

717 Paul, A. (2006). Kinship and behaviour in Barbary macaques. In: J.K. Hodges, & J. Cortes (Eds) *The*  
718 *Barbary macaques: Biology, Management and Conservation* (pp. 47 - 61). Nottingham, U.K.:  
719 Nottingham University Press.

720 Petit, O., Abegg, C., & Thierry, B. (1997). A comparative study of aggression and conciliation in  
721 three cercopithecine monkeys (*Macaca fuscata*, *Macaca nigra*, *Papio papio*). *Behaviour*, 134, 415–  
722 432. doi: 10.1163/156853997X00610

723 Petit, O., Desportes, C., & Thierry, B (1992). Differential probability of “coproduction” in two  
724 species of macaques (*Macaca tonkeana*, *M. mulatta*). *Ethology*, 90, 107–120. doi: 0.1111/j.1439-  
725 0310.1992.tb00825.x

726 Petit, O., & Thierry, B. (1994). Aggressive and peaceful interventions in conflicts in Tonkean  
727 macaques. *Animal Behaviour*, 48, 1427–1436. doi:10.1006/anbe.1994.1378

728 Prange, S., Gehrt, S. D., & Hauver, S. (2011). Frequency and duration of contacts between free-  
729 ranging raccoons: uncovering a hidden social system. *Journal of Mammalogy*, 92, 1331–1342.  
730 doi:10.1644/10-MAMM-A-416.1

731 R Development Core Team (2015) R: A language and environment for statistical computing, version  
732 3.2.1.

733 Schino, G. (2007). Grooming and agonistic support: a meta-analysis of primate reciprocal altruism.  
734 *Behavioral Ecology*, 18, 115–120. doi: 10.1093/beheco/arl045

735 Schino, G., & Aureli, F. (2009). Reciprocal altruism in primates: partner choice, cognition, and  
736 emotions. *Advances in the Study of Behavior*, 39, 45–69. doi: 10.1016/S0065-3454(09)39002-6

737 Schradin, C. (2013) Intraspecific variation in social organization by genetic variation, developmental  
738 plasticity, social flexibility or entirely extrinsic factors. *Philosophical Transactions of the Royal*  
739 *Society B: Biological Science*, 368, 20120346. doi: 10.1098/rstb.2012.0346

- 740 Schülke, O., & Ostner, J. (2008). Male reproductive skew, paternal relatedness, and female social  
741 relationships. *American Journal of Primatology*, *70*, 695–698. doi: 10.1002/ajp.20546
- 742 Seyfarth, R. M. (1977). A model of social grooming among adult female monkeys. *Journal of*  
743 *Theoretical Biology*, *65*, 671–698. doi:10.1016/0022-5193(77)90015-7
- 744 Seyfarth, R. M., Silk, J. B., & Cheney, D. L. (2014). Social bonds in female baboons: the interaction  
745 between personality, kinship and rank. *Animal Behaviour*, *87*, 23–29.  
746 doi:10.1016/j.anbehav.2013.10.008
- 747 Sick, C., Carter, A. J., Marshall, H. H., Knapp, L. A., Dabelsteen, T., & Cowlshaw, G. (2014).  
748 Evidence for varying social strategies across the day in chacma baboons. *Biology Letters*, *10*,  
749 20140249. doi:10.1098/rsbl.2014.0249
- 750 Silk, J. B., Cheney, D. L., & Seyfarth, R. M. (2013). A practical guide to the study of social  
751 relationships. *Evolutionary Anthropology*, *22*, 213–225. doi:10.1002/evan.21367
- 752 Silk, J. B., Beehner, J. C., Bergman, T. J., Crockford, C., Engh, A. L., Moscovice, L. R., et al. (2010).  
753 Strong and Consistent Social Bonds Enhance the Longevity of Female Baboons. *Current Biology*, *20*,  
754 1359–1361. doi:10.1016/j.cub.2010.05.067
- 755 Silk, J. B., Beehner, J. C., Bergman, T. J., Crockford, C., Engh, A. L., Moscovice, L. R., et al. (2009).  
756 The benefits of social capital: close social bonds among female baboons enhance offspring survival.  
757 *Proceedings of the Royal Society B: Biological Sciences*, *276*, 3099–3104.  
758 doi:10.1098/rspb.2009.0681
- 759 Silk, J. B. (2007a). The adaptive value of sociality in mammalian groups. *Philosophical Transaction*  
760 *of the Royal Society B: Biological Sciences*, *362*, 539–559. doi: 10.1098/rstb.2006.1994.
- 761 Silk, J. B. (2007b). Social components of fitness in primate groups. *Science*, *317*, 1347–1351. doi:  
762 10.1126/science.1140734
- 763 Silk, J. B., Altmann, J., & Alberts, S. C. (2006a). Social relationships among adult female baboons  
764 (*Papio cynocephalus*) II. Variation in the quality and stability of social bonds. *Behavioral Ecology*  
765 *and Sociobiology*, *61*, 197–204. doi: 10.1007/s00265-006-0250-9
- 766 Silk, J. B., Altmann, J., & Alberts, S. C. (2006b). Social relationships among adult female baboons (*Papio*  
767 *cynocephalus*) I. Variation in the strength of social bonds. *Behavioral Ecology and Sociobiology*, *61*,  
768 183–195. doi: 10.1007/s00265-006-0249-2
- 769 Smith, J. E., van Horn, R. C., Powning, K. S., Cole, A. R., Graham, K. E., Memenis, S. K.,  
770 Holekamp, K. E. (2010). Evolutionary forces favoring intragroup coalitions among spotted hyenas  
771 and other animals. *Behavioral Ecology*, *21*, 284–303. doi:10.1093/beheco/arp181
- 772 Smith, J. E., Memenis, S. K., Holekamp, K. E. (2007). Rank-related partner choice in the fission-  
773 fusion society of the spotted hyena (*Crocuta crocuta*). *Behavioral Ecology and Sociobiology*, *61*,  
774 753–765. doi: 10.1007/s00265-006-0305-y
- 775 Smith, K. L., Alberts, S. C., & Altmann, J. (2003). Wild female baboons bias their social behaviour  
776 towards paternal half-sisters. *Proceedings of the Royal Society B: Biological Science*, *270*, 503–510.  
777 doi: 10.1098/rspb.2002.2277
- 778 Sterck, E. H. M., Watts, D. P., van Schaik C. P. (1997). The evolution of female social relationships  
779 in nonhuman primates. *Behavioral Ecology and Sociobiology*, *41*, 291–309.  
780 doi:10.1007/s002650050390
- 781 Sueur, C., & Petit, O. (2008). Shared or unshared consensus decision in macaques? *Behavioral*  
782 *Processes*, *78*, 84–92. doi: 10.1016/j.beproc.2008.01.004
- 783 Sugardjito, J., Southwick, C. H., Supriatna, J., Kohlhaas, A., Baker, S. C., Erwin, J., Froehlich, J., &  
784 Lerche, N. (1989). Population survey of macaques in northern Sulawesi. *American Journal of*  
785 *Primatology*, *18*, 285–301. doi: 10.1002/ajp.1350180403

- 786 Taberlet, P., Griffin, S., Goossens, B., Questiau, S., Manceau, V., Escaravage, N., Waits, L. P.,  
787 Bouvet, J. (1996) Reliable genotyping of samples with very low DNA quantities using PCR. *Nucleic*  
788 *Acids Research*, 24, 3189–3194. doi:10.1093/nar/24.16.3189
- 789 Thierry, B. (2013). Identifying constraints in the evolution of primate societies. *Philosophical*  
790 *Transactions of the Royal Society B: Biological Sciences*, 368, 20120342. doi:10.1098/rstb.2012.0342
- 791 Thierry, B., Aureli, F., Nunn, C. L., Petit, O., Abegg, C., & De Waal, F. B. M. (2008). A comparative  
792 study of conflict resolution in macaques: insights into the nature of trait covariation. *Animal*  
793 *Behaviour*, 75, 847–860. doi: 10.1016/j.anbehav.2007.07.006
- 794 Thierry, B. (2007). Unity in diversity: lessons from macaque societies. *Evolutionary Anthropology*,  
795 16, 224–238. doi: 10.1002/evan.20147
- 796 Thierry, B., Bynum, E. L., Baker, S. C., Kinnaird, M. F., Matsumura, S., Muroyama, Y., et al. (2000).  
797 The social repertoire of Sulawesi macaques. *Primate Research*, 16, 203–226. doi:10.2354/psj.16.203
- 798 Thierry, B. (1990). Feedback loop between kinship and dominance: the macaque model. *Journal of*  
799 *Theoretical Biology*. 145, 511–522. doi:10.1016/S0022-5193(05)80485-0
- 800 Tinsley Johnson, E., Snyder-Mackler, N., Beehner, J. C., & Bergman, T. J. (2014) Kinship and  
801 dominance rank influence the strength of social bonds in female geladas (*Theropithecus gelada*).  
802 *International Journal of Primatology*, 35, 288–304. doi: 10.1007/s10764-013-9733-5
- 803 Uchino, B. N. (2006). Social Support and Health: A Review of Physiological Processes Potentially  
804 Underlying Links to Disease Outcomes. *Journal of Behavioral Medicine*, 29, 377–387.  
805 doi:10.1007/s10865-006-9056-5
- 806 van Horn, R. C., Altmann, J., & Alberts, S. C. (2008). Can't get there from here: inferring kinship  
807 from pairwise genetic relatedness. *Animal Behaviour*, 75, 1173–1180.  
808 doi:10.1016/j.anbehav.2007.08.027
- 809 van Schaik, C. P. (1989). The ecology of social relationships amongst female primates. In: V.  
810 Standen, & R. A. Foley (Eds) *Comparative Socioecology: The Behavioural Ecology of Humans and*  
811 *other Mammals* (pp. 195-218). Oxford, U.K.: Blackwell Scientific Publications.
- 812 Ventura R, Majolo B, Koyama NF, et al (2006) Reciprocation and interchange in wild Japanese  
813 macaques: grooming, co-feeding, and agonistic support. *Am J Primatol* 68:1138–1149.  
814 doi:10.1002/ajp.20314
- 815 Wahaj, S. A., van Horn, R. C., van Horn, T. L., Dreyer, R., Hilgris, R., Schwarz, J., Holekamp, K. E.  
816 (2004). Kin discrimination in the spotted hyena (*Crocuta crocuta*): nepotism among siblings.  
817 *Behavioral Ecology and Sociobiology*, 56, 237–247. doi: 10.1007/s00265-004-0783-8
- 818 Wang, J. (2011). COANCESTRY: a program for simulating, estimating and analysing relatedness and  
819 inbreeding coefficients. *Molecular Ecology Resources*, 11, 141–145. doi: 10.1111/j.1755-  
820 0998.2010.02885.x
- 821 Wenzel, S., Ostner, J., & Schülke, O. (2013). Paternal relatedness predicts the strength of social bonds  
822 among female rhesus macaques. *PLoS ONE*, 8, e59789–e59789. doi: 10.1371/journal.pone.0059789
- 823 Wey, T. W., Burger, J. R., Ebensperger, L. A., & Hayes, L. D. (2013). Reproductive correlates of  
824 social network variation in plurally breeding degus (*Octodon degus*). *Animal Behaviour*, 85, 1407–  
825 1414. doi:10.1016/j.anbehav.2013.03.035
- 826 Whitehead, H. (2008). *Analysing animal societies: quantitative methods for vertebrates social*  
827 *analysis*. Chicago, U.S.A.: The University of Chicago Press.
- 828 Widdig, A. (2013) The impact of male reproductive skew on kin structure and sociality in multi-male  
829 groups: Altmann (1979) revisited. *Evolutionary Anthropology*, 22, 239–250. doi: 10.1002/evan.21366

830 Wrangham, R. W., & Rubenstein, D. I. (1986) Social evolution in birds and mammals. In: D. I.  
831 Rubenstein, & Wrangham R. W. (Eds) *Ecological Aspects of Social Evolution - Birds and Mammals*  
832 (pp. 452–470). Princeton, USA: Princeton University Press.

833 Young, C., Majolo, B., Heistermann, M., Schülke, O., & Ostner, J. (2014). Responses to social and  
834 environmental stress are attenuated by strong male bonds in wild macaques. *Proceedings of the*  
835 *National Academy of Science*, *111*, 195–200. doi:10.1073/pnas.1411450111

836 Young, C., Majolo, B., Schülke, O., & Ostner, J. (2014). Male social bonds and rank predict supporter  
837 selection in cooperative aggression in wild Barbary macaques. *Animal Behaviour*, *95*, 23–32.  
838 doi:10.1016/j.anbehav.2014.06.007

839

## 840 **Figure legends**

841

842 **Figure 1:** Distribution of overall CSI scores of female-female dyads in the two study groups, up PB  
843 and down R1. The blue arrow indicates the median and the red arrow the limit of 10% of the strongest  
844 CSI scores. The fact that the distribution is only moderately skewed to the left (i.e. towards 0)  
845 indicates how average most of the bonds between females are contrary to what is known for other  
846 species where similar data are available.

847

848 **Figure 2:** Variation in CSI scores of female-female dyads in the two study groups, left PB and right  
849 R1, across three six-months-periods. Row and column labels represent female identities. A circle  
850 denotes that the female in the column was amongst the top three partners of the female in the row at  
851 least once (small grey), twice (medium blue) or three times (big red) across the three periods. The  
852 presence of few red circles but many grey dots illustrate how bond strength changes across periods  
853 and lacks of predictability.

854

855 **Figure 3:** Effect of dominance difference (left panels), relatedness (middle panels) and age difference  
856 (right panels) on bond strength (top panels), bond predictability (middle panels), and bond equitability  
857 (bottom panels). The straight full line represents the estimate variation as predicted by the model, the  
858 dotted lines are the associated lower and upper 95% confidence intervals of the estimate, the grey  
859 points are the original data points transformed and scaled as in the statistical model.

860

861 **Figure 4:** Effect of bond strength (left panels), bond predictability (middle panels), and bond  
862 equitability (right panels) on coalitionary support (top panels), feeding in proximity frequency (middle  
863 panels), and aggression frequency (bottom panels). The straight full line represents the estimate  
864 variation as predicted by the model, the dotted lines are the associated lower and upper 95%  
865 confidence intervals of the estimate, the grey points are the original data points transformed and  
866 scaled as in the statistical model.

Degrees of freedom in social bonds of crested macaque females.

Julie Duboscq<sup>1,2,3,4\*†</sup>, Christof Neumann<sup>1,2,5,6\*\*</sup>, Muhammad Agil<sup>7</sup>, Dyah Perwitasari-Farajallah<sup>8,9</sup>, Bernard Thierry<sup>3,4</sup>, Antje Engelhardt<sup>1,2\*\*\*</sup>

<sup>1</sup> Junior Research Group of Primate Sexual Selection, German Primate Center, Göttingen, Germany, <sup>2</sup> Courant Research Centre for the Evolution of Social Behaviour, Göttingen, Germany, <sup>3</sup> Université de Strasbourg, Institut Pluridisciplinaire Hubert Curien, Strasbourg, France, <sup>4</sup> Centre National de la Recherche Scientifique, Département Ecologie, Physiologie et Ethologie, Strasbourg, France, <sup>5</sup> Junior Research Group of Primate Kin Selection, Max-Planck-Institute for Evolutionary Anthropology, Leipzig, Germany, <sup>6</sup> Institute of Biology, Faculty of Bioscience, Pharmacy and Psychology, University of Leipzig, Leipzig, Germany, <sup>7</sup> Faculty of Veterinary Sciences, Bogor Agricultural University, Bogor, Indonesia, <sup>8</sup> Primate Research Center, Bogor Agricultural University, Bogor, Indonesia, <sup>9</sup> Faculty of Mathematics and Natural Sciences, Bogor Agricultural University, Bogor, Indonesia

\* corresponding author: [julie.a.m.duboscq@gmail.com](mailto:julie.a.m.duboscq@gmail.com)

## Highlights

Most bonds were of average strength, of moderate endurance and balanced

Close-ranked females had stronger, more predictable and equitable bonds than others

Bond characteristics and fitness-related behaviours were linked in complex ways

Tolerant females can express great degrees of freedom in their social relationships



These patterns are under-appreciated in stable nepotistic hierarchical societies



### **Acknowledgments**

This article is dedicated to Ugiek and Antri. We gratefully acknowledge the permission of the Indonesian State Ministry of Research and Technology (RISTEK), the Directorate General of Forest Protection and Nature Conservation (PHKA) in Jakarta and the Department for the Conservation of Natural Resources (BKSDA) in Manado, particularly Pak Untung and Pak Yunus, to conduct this research in the Tangkoko-Batuangus Nature Reserve. We thank all the members of the Macaca Nigra Project for their support in the field, and especially Dwi Yandhi Febryanti and Jérôme Micheletta for their help with data collection.


Figure



Figure


PB

ap bp cp dp ep fp gp hp ip jp lp np rp sp yp




R1

as bs cs ds es gs hs is js ks ms ns os ps qs rs ss ts us xs ys




Figure


dominance difference






relatedness index






age difference



bond predictability




bond equitability




Figure


**bond strength**






**bond predictability**






**bond equitability**



feeding in proximity



aggression frequency



**Supplementary material for on-line publication only**

[Click here to download Supplementary material for on-line publication only: social bonds\\_Duboscq et al\\_appendix\\_20160628.doc](#)