

Match physical performance of elite  
female soccer players during international  
competition

**1 ABSTRACT**

2 The purpose of the present study was to provide a detailed  
3 analysis of the physical demands of competitive international  
4 female soccer match-play. A total of 148 individual match  
5 observations were undertaken on 107 outfield players  
6 competing in competitive international matches during the  
7 2011-2012 and 2012-2013 seasons, using a computerized  
8 tracking system (Prozone Sports Ltd., Leeds, England). Total  
9 distance (TD) and total high-speed running distances (THSR)  
10 were influenced by playing position, with central midfielders  
11 (CM) completing the highest ( $10985\pm706$  m and  $2882\pm500$  m)  
12 and central defenders (CD) the lowest ( $9489\pm562$  m and  
13  $1901\pm268$  m) distances, respectively. Greater total very high-  
14 speed running (TVHSR) distances were completed when a  
15 team was without ( $399\pm143$  m) compared to with ( $313\pm210$  m)  
16 possession of the ball. The majority of sprints were over short  
17 distances with 76 % and 95 % being less than 5 m and 10 m,  
18 respectively. Between half reductions in physical performance  
19 were present for all variables, independent of playing position.  
20 The current study provides novel findings regarding the  
21 physical demands of different playing positions in competitive  
22 international female match-play and provides important  
23 insights for physical coaches preparing elite female players for  
24 competition.

25

26 **Key Words:** football; match analysis; tracking system; playing

27 position; high-speed running

28

29

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

**51 INTRODUCTION**

52 A comprehensive understanding of the physical demands of  
53 match-play is necessary in order to apply a systematic approach  
54 to training and testing protocols.<sup>1</sup> As a consequence, global  
55 positioning system (GPS) technology and semi-automated  
56 camera systems have been extensively used to provide a  
57 detailed analysis of specific elements of a player's physical  
58 performance in men's soccer.<sup>2-4</sup> Despite advancements in the  
59 understanding of the physical demands of match-play in elite  
60 male players, limited research currently exists on elite female  
61 players. This predominantly reflects the fact that female  
62 matches are rarely played in stadiums equipped with semi  
63 automated camera systems. Furthermore, the high financial  
64 costs that are associated with other contemporary technologies,  
65 often prohibit their use in female soccer.<sup>5,6</sup> Consequently, a  
66 large proportion of the research undertaken to date has been  
67 derived from relatively small samples using traditional video-  
68 based technology.<sup>7-10</sup> Collectively, these factors limit the depth  
69 of analysis possible; therefore, it is important that further  
70 information relating to female match-play is derived to better  
71 inform female-specific training prescription and testing  
72 protocols.

73

74 Available data on female match-play indicates that the standard  
75 of competition influences physical performance with greater

76 total distances observed in European club football<sup>11</sup> compared  
77 to friendly international competition.<sup>6</sup> Furthermore, greater  
78 high-speed running (HSR) and sprinting have also been  
79 observed during friendly international matches compared to  
80 domestic club matches.<sup>12</sup> However, to date, no information  
81 utilizing contemporary techniques exists on the demands of  
82 competitive international match-play, which represents the  
83 highest standard within the female game. Furthermore, due to  
84 the limited sample sizes available, the majority of studies  
85 examining the influence of playing position on match physical  
86 performance have been restricted to more generic assessments  
87 (e.g. defenders, midfielders and attackers) with only one  
88 study<sup>11</sup> further differentiating between central and wide  
89 positions. Bradley and colleagues<sup>11</sup> presented activity profiles  
90 for female match-play across five playing positions; however,  
91 the primary focus of their research was to compare male and  
92 female match-play and as such detailed female positional  
93 comparisons were lacking. Consequently, a comprehensive  
94 positional analysis of the physical demands of elite female  
95 match-play is necessary in order to provide applied  
96 practitioners working with elite players, pertinent information  
97 to better inform position-specific training prescription.  
98 Therefore, the aim of the current investigation was to provide a  
99 detailed analysis of the physical demands of different playing  
100 positions during competitive international female match-play.

101

102 **METHODS**

103

104 **EXPERIMENTAL APPROACH TO THE PROBLEM**

105 To quantify the demands of competitive international female  
106 match-play, physical performance data were collected during  
107 the 2011-2012 and 2012-2013 seasons. Data were derived from  
108 ten matches, featuring thirteen teams playing in different  
109 stadiums across Europe.

110

111 **SUBJECTS**

112 A total of 148 individual match observations were undertaken  
113 on 107 outfield players (goalkeepers were excluded) with a  
114 median of two matches per player (range = 1-4). Data were  
115 only included for those players completing entire matches (i.e.  
116 90 minutes). Data were collected as a condition of employment  
117 in which player performance is routinely measured during  
118 match-play.<sup>13</sup> Therefore, usual appropriate ethics committee  
119 clearance was not required. Nevertheless, to ensure team and  
120 player confidentiality, all physical performance data were  
121 anonymised before analysis. Permission to publish this data  
122 was granted by Prozone (Prozone Sports Ltd., Leeds, UK).

123

124 **PROCEDURES**

125 Match physical performance data were collected using a  
126 computerized semi-automated multi-camera image recognition  
127 system (Prozone Sports Ltd., Leeds, UK). This system  
128 provides valid<sup>14</sup> and reliable<sup>15</sup> estimations of a variety of match  
129 performance indices. Players were categorized by playing  
130 position; central defenders (CD) (n = 25; 35 match  
131 observations), wide defenders (WD) (n = 28; 34 match  
132 observations), central midfielders (CM) (n = 31; 40 match  
133 observations), wide midfielders (WM) (n = 17; 20 match  
134 observations) and attackers (A) (n = 16; 19 match observations)  
135 to determine the influence of playing position on match  
136 physical performance. The influence of playing position on the  
137 difference in activity between the first and second half periods  
138 was undertaken. Within half changes in physical performance  
139 were also assessed by examining 15 and 5-minute time periods.

140

141 The following activity classifications were used: total distance  
142 (TD), walking (0.7-7.1 km.h<sup>-1</sup>), jogging (7.2-14.3 km.h<sup>-1</sup>),  
143 running (14.4-19.7 km.h<sup>-1</sup>), HSR (19.8-25.1 km.h<sup>-1</sup>) and  
144 sprinting (>25.1 km.h<sup>-1</sup>) distance. Total high-speed running  
145 (THSR) (>14.4 km.h<sup>-1</sup>) and total very high-speed running  
146 (TVHSR) (>19.8 km.h<sup>-1</sup>) were also computed.<sup>16</sup> The above  
147 velocity thresholds for each activity have been extensively  
148 employed to quantify the physical demands of male match-  
149 play.<sup>2-4</sup> Recent commentary<sup>17</sup> has suggested that transposing

150 these thresholds to the performances of female players will  
151 underestimate match-play demands by reducing the amount of  
152 high-speed activities completed by individuals. While the  
153 present authors support this view in general, there has been a  
154 reluctance to adopt such thresholds in the current data as a  
155 consequence of the confidence that can be associated with  
156 current recommendations that exist regarding female specific  
157 velocity thresholds.<sup>17</sup> For example, female specific HSR and  
158 sprint thresholds derived from small samples (n = 5-14) of non-  
159 elite players (domestic level players).<sup>9,18</sup> have been proposed  
160 without consideration for the key methodological  
161 considerations required when determining velocity  
162 thresholds.<sup>19</sup> This includes the use of match activity zones that  
163 are expressed relative to individual players physical  
164 capabilities.<sup>20</sup> Furthermore, if physiological thresholds are used  
165 to demarcate individualized match activity zones they should  
166 be ascertained from activity patterns that replicate the  
167 movement demands of soccer in order to account for the  
168 increased energy cost associated with unorthodox modes of  
169 motion (e.g. backwards and sideways running) experienced  
170 during match-play.<sup>21</sup> Consequently, the authors feel that the  
171 suggested velocities<sup>17</sup> will not be representative of the abilities  
172 of either elite female players (as used in the present study) or  
173 female soccer players more generally. As such it may be that  
174 activity classifications derived from these thresholds may not



175 be any more valid than the arbitrary male thresholds presently  
176 used.

177

178 Total very high-speed running ( $>19.8 \text{ km}\cdot\text{h}^{-1}$ ) was expressed as  
179 both TVHSR distance completed when the respective player's  
180 team were in possession (VHSRP) or were without possession  
181 (VHSRWP) of the ball. Further analysis of sprinting activity  
182 ( $>25.1 \text{ km}\cdot\text{h}^{-1}$ ) was also considered, with the distance covered  
183 and the type of sprint classified. Sprints were classed as either  
184 explosive or leading sprints. An explosive sprint was defined as  
185 the attainment of sprint speed from standing, walking, jogging  
186 or running with time spent in the HSR category less than 0.5 s.  
187 Conversely, a leading sprint was defined as the attainment of  
188 sprint speed from standing, walking, jogging or running whilst  
189 entering the HSR category for a minimum of 0.5 s.<sup>15</sup>

190

## 191 **STATISTICAL ANALYSIS**

192 Data are presented as mean $\pm$ SD, with significance set at  $p <$   
193 0.05. Data were analyzed using factorial linear mixed modeling  
194 using the Statistical Package for Social Sciences (Version 21).  
195 Linear mixed modeling can be applied to repeated measures  
196 data from unbalanced designs, which was the case in our study  
197 since players differed in terms of the number of repeated  
198 matches they participated in. Linear mixed modeling can also  
199 cope with the mixture of random and fixed level effects that

200 occur with performance analysis data<sup>22</sup> as well as with missing  
201 and ‘nested’ data (hierarchical models). Significant main  
202 effects of each factor were followed up with Bonferroni-  
203 corrected multiple contrasts. Effect size (ES), estimated from  
204 the ratio of the mean difference to the pooled standard  
205 deviation, were also calculated. The ES magnitude was  
206 classified as trivial (<0.2), small (>0.2-0.6), moderate (>0.6-  
207 1.2), large (>1.2-2.0) and very large (>2.0-4.0).<sup>23</sup>

208

## 209 **RESULTS**

210

### 211 **TOTAL MATCH PERFORMANCE**

212 The average ‘ball in play time’ was  $62.0 \pm 7.7$  % of the total  
213 match duration. The distance covered in all speed classification  
214 zones was influenced by playing position ( $p < 0.001$ ) (Table 1).  
215 Total distance was greater in CM compared to all other playing  
216 positions (ES 1.0-2.3;  $p < 0.05$ ) except WM (ES 0.5); conversely  
217 CD completed less total distance compared to all other  
218 positions (ES 1.1-2.3;  $p < 0.05$ ). Total high-speed running  
219 distance was similar between all positions (ES 0.1-0.6) with the  
220 exception of CD who completed the least distance (ES 1.6-2.4;  
221  $p < 0.001$ ) and between CM and WD (ES 0.7,  $p < 0.05$ ).  
222 Positional differences for running, HSR and sprinting were also  
223 apparent. Physical performance was generally similar between  
224 wide players (WD and WM) and A, with no differences

225 observed in TD, jogging, running, HSR or sprinting distances  
226 (Table 1).

227

228 Both VHSRP and VHSRWP also differed between positions  
229 ( $p < 0.001$ ) (Table 1). The VHSRP was greater in A and WM  
230 compared to defenders (CD and WD) and CM (ES 0.9-4.4;  
231  $p < 0.05$ ). The VHSRP was similar in WD and CM (ES 0.0),  
232 however, CD completed less VHSRP than all other playing  
233 positions (ES 1.5-4.4;  $p < 0.001$ ). The VHSRWP was greater in  
234 CM (ES 0.8-1.5;  $p < 0.05$ ) compared to all other playing positions  
235 except WD (ES 0.5). Attackers completed less VHSRWP than  
236 all other playing positions with moderate to large differences  
237 observed (ES 0.8-1.5) (Table 1).

238

239 **\*\*\*\*Table 1 near here\*\*\*\***

240

241 There were no significant differences between playing  
242 positions for either the percentage of explosive (ES 0.0-0.7) or  
243 leading (ES 0.0-0.7) sprints. However, CM generally  
244 completed a greater percentage of explosive sprints compared  
245 to WD and A (ES 0.6-0.7). Central midfielders completed a  
246 greater proportion of sprints that were explosive compared to  
247 leading in nature (ES 0.8;  $p < 0.05$ ) (Table 1). The total number  
248 of sprints was influenced by playing position ( $p < 0.001$ ) (Figure  
249 1). Attackers completed more sprints than defenders (ES 0.8-

250 2.5;  $p < 0.05$ ) but a similar number to WM (ES 0.1). Similar  
251 numbers of sprints (ES 0.2) were also observed between WD  
252 and CM. Central defenders completed less sprints than all other  
253 playing positions (ES 0.9-2.5;  $p < 0.05$ ).

254

255 A similar number of very short sprints ( $< 5$  m) were completed  
256 by A, WM and CM (ES 0.1-0.3), with trends for WD to  
257 complete less than A (ES 0.7). Central defenders completed  
258 fewer very short sprints (ES 1.0-2.1;  $p < 0.05$ ) compared to all  
259 positions. Wide midfielders completed more 5.1-10.0 m sprints  
260 than CD (ES 1.2;  $p < 0.05$ ) and A completed more than both CD  
261 and CM (ES 0.9-2.0;  $p < 0.05$ ). Attackers also completed more  
262 10.1-15.0 m sprints than CD (ES 0.8;  $p < 0.05$ ), with no other  
263 significant positional differences found between 5.1-10.0 m  
264 (ES 0.1-0.7) and 10.1-15.0 m sprints (ES 0.0-0.6). There was a  
265 trend (ES 0.6-0.7) for A to complete more mid-range sprints  
266 (5.1-15.0 m) than WD. All players completed a similar number  
267 of 15.1-20.0 m sprints (ES 0.0-0.4), but WM produced  
268 marginally more  $> 20$  m sprints than defenders and CM (ES 0.6;  
269  $p < 0.05$ ) (Figure 1).

270

271 **\*\*\*\*Figure 1 near here\*\*\*\***

272

273 **BETWEEN HALF MATCH PERFORMANCE:**

274 **INFLUENCE OF PLAYING POSITION**

275 There was a reduction in the average 'ball in play time' in the  
276 second ( $59.9\pm 7.8$  %) compared to the first ( $64.1\pm 7.3$  %) half  
277 (ES 0.6). When considering the sample as a whole there was a  
278 reduction in TD ( $365\pm 270$  m (ES 0.8;  $p<0.001$ )), THSR  
279 ( $141\pm 169$  m (ES 0.5;  $p<0.001$ )) and TVHSR ( $47\pm 100$  m (ES  
280 0.4;  $p<0.001$ )) during the second half compared to first. These  
281 differences were mainly attributed to a reduction in jogging  
282 ( $217\pm 188$  m (ES 0.8;  $p<0.001$ )), running ( $93\pm 108$  m (ES 0.5;  
283  $p<0.001$ )) and HSR ( $38\pm 71$  m (ES 0.4;  $p<0.001$ )) and to a  
284 lesser extent sprinting ( $10\pm 41$  m (ES 0.2;  $p<0.05$ )). Trivial to  
285 small reductions in VHSRP ( $16\pm 66$  m (ES 0.1;  $p<0.05$ )) and  
286 VHSRWP ( $24\pm 65$  (ES 0.3;  $p<0.001$ )) were also observed  
287 during the second half compared to the first half. The  
288 magnitude of the reduction in physical performance between  
289 the first and second half was independent of playing position.  
290 There were no differences in the percentage of explosive or  
291 leading sprints between halves for any playing position (ES  
292 0.0-0.4).

293

#### 294 **WITHIN HALF MATCH PERFORMANCE (15 MINUTE** 295 **INTERVALS)**

296 Total high-speed running distance during the final 15-min  
297 period of the match was lower (12-35 %) compared to all other  
298 15-min blocks (ES 0.4-1.1;  $p<0.001$ ) (Figure 3). In both halves,  
299 THSR was lower in the final 15 minutes compared to the first

300 and second 15-minute interval (1<sup>st</sup> half, ES 0.2-0.5;  $p < 0.05$ ; 2<sup>nd</sup>  
301 half, ES 0.4-0.7;  $p < 0.001$ ) (Figure 2).

302

303 **\*\*\*\*Figure 2 near here\*\*\*\***

304

305 **WITHIN HALF MATCH PERFORMANCE (5 MINUTE**  
306 **INTERVALS)**

307 The peak THSR distance in a 5-minute period was  $223 \pm 47$  m.  
308 In the following 5-minute period, the amount of THSR was 39  
309 % lower ( $p < 0.001$ ) ( $135 \pm 47$  m, ES 1.9;  $p < 0.001$ ) but was not  
310 different to the mean distance covered during all 5-minute  
311 intervals not including the peak distance ( $135 \pm 32$  m) (ES 0.0).

312

313 **DISCUSSION**

314 The present study represents the largest single analysis of elite  
315 female match-play data to date and provides novel insights into  
316 the physical demands of different playing positions during  
317 competitive international match-play using contemporary  
318 techniques. The present data highlights large differences in the  
319 physical demands of match-play between playing positions and  
320 the number of high-speed efforts is lower across the duration of  
321 the match in all positions. Collectively, the current data  
322 provides physical coaches with new insights into the position-  
323 specific physical demands of competitive international match-

324 play which will inform the design and implementation of  
325 training drills for elite female players.

326

327 The TD covered in this current investigation ( $10321 \pm 859$  m) is  
328 similar to values previously observed in European club football  
329 ( $10754$  m)<sup>11</sup> and college soccer ( $9496$ - $10297$  m)<sup>24</sup> but appear  
330 greater than the TD reported during a small sample of  
331 international friendlies ( $9292$ - $9631$  m).<sup>6</sup> This increase in TD  
332 covered during competitive international matches relative to  
333 international friendlies<sup>6</sup> appears consistent across playing  
334 positions (defenders =  $9864$  vs.  $8759$  m, midfielders =  $10864$   
335 vs.  $10150$  m, attackers =  $10262$  vs.  $9442$  m). Whilst some  
336 caution should be exercised when comparing data between  
337 studies that have utilized different data capture methods<sup>25-27</sup>  
338 and small sample sizes, the moderate to large effect size  
339 suggests an increased overall physical demand of competitive  
340 versus friendly international match-play. This to some extent  
341 may simply reflect the greater importance associated with  
342 competitive matches.

343

344 Low-speed activity (walking and jogging) accounts for the  
345 majority (~85 %) of total distance covered in elite females,  
346 during domestic-level matches.<sup>7,10,12</sup> However, it is high-speed  
347 activity that is widely regarded as an important component of  
348 match physical performance as these activities are often critical

349 to the outcome of matches by directly impacting goal scoring  
350 opportunities.<sup>15,28</sup> Interestingly, in the current study a distance  
351 of ~2520 m was covered at high-speed, accounting for 24 % of  
352 the total distance. These observations suggest that a greater  
353 proportion of high-speed activity may be undertaken during  
354 competitive international football relative to domestic-level  
355 matches.<sup>7,10,12</sup> As noted previously, there remains no consensus  
356 in the literature regarding female specific velocity thresholds.<sup>17</sup>  
357 The female specific thresholds that have recently been  
358 proposed<sup>17</sup> are not representative of this elite population and  
359 therefore may not be any more valid than the arbitrary male  
360 thresholds that frequent the literature. The findings from the  
361 current study indicate similar proportions (23 % in males and  
362 24 % in females) of high-speed activity relative to total  
363 distance when compared to male players.<sup>16</sup> As a consequence, a  
364 focus on high-intensity soccer-specific conditioning<sup>29,30</sup> should  
365 represent an integral component of the training methodology  
366 applied to the development of elite female players.

367

368 Previous investigations examining sprint activity in women's  
369 soccer are largely limited to the analysis of total sprint  
370 distance.<sup>6-8,10,24</sup> The sprint distance covered in the current  
371 investigation (168±82 m) was less (ES 1.2-4.9) than values  
372 previously observed (221-380 m) in elite players during  
373 domestic level matches.<sup>7,10</sup> Since greater THSR was observed



374 in the present study relative to domestic level matches,<sup>7,10,12</sup> it  
375 is possible this increase largely reflects an increase in HSR  
376 activity rather than any changes in sprint activity. The present  
377 study is the first to provide a comprehensive analysis of both  
378 the range of sprint distances and types of sprints undertaken by  
379 elite female players. Sprint distances between 0-5 m and 0-10  
380 m accounted for 76 % and 95 % of all sprints, respectively.  
381 Whilst female sprint data has not previously been presented in  
382 this format, average sprint distances of  $15.1 \pm 9.4$  m have been  
383 observed in players from a professional league in the United  
384 States.<sup>31</sup> It is likely that this distance is greater than the average  
385 sprint distance in the current sample of players since 95 % of  
386 all sprints were shorter than 10 m. Alongside a high proportion  
387 of shorter sprints, the present data demonstrates an even  
388 distribution of explosive and leading sprints ( $51 \pm 10\%$  vs.  
389  $49 \pm 10\%$ ). Interestingly, these findings suggest that women  
390 adopt a greater proportion of explosive sprints compared to  
391 males (77 % leading vs. 23 % explosive).<sup>32</sup> This observation  
392 could reflect differences in how the game is played with  
393 females being more reactive to match-play events relative to  
394 males, or that males obtain the sprint threshold at a lower  
395 proportion of their maximum sprint velocity, however, further  
396 work is needed in order to confirm this. Collectively, the  
397 present findings indicate that sprint training in elite female  
398 players should include a particular focus on sprinting over short

399 distances (<10 m) with a combination of sprinting from a  
400 stationary and rolling start. This emphasis on short sprints and  
401 accelerations is necessary due to the explosive nature of  
402 activity reported in the current findings. However, it should be  
403 noted that sprint training drills over longer distances (>20 m)  
404 are required in order to condition players for the longer sprint  
405 distances that arise in match-play, albeit infrequently, and also  
406 to develop maximum sprinting speed.<sup>33</sup> It should be  
407 acknowledged that although the present study provides novel  
408 data concerning the [locomotor](#) demands of elite female match-  
409 play it fails to quantify the true physical demands. For example,  
410 a limitation of camera based tracking systems, such as the one  
411 used in the present study, is their inability to provide a valid  
412 assessment of acceleration and deceleration activity. Similarly,  
413 camera based systems, unlike GPS that are equipped with  
414 triaxial accelerometers, cannot provide information pertaining  
415 to mechanical loading. [Consequently, it is not possible from the](#)  
416 [current dataset to gain a full understanding of the physical](#)  
417 [demands of match-play due to the inability to quantify](#)  
418 [variables such as the number of tackles, jumps or the instances](#)  
419 [that a player goes to ground.](#) As the use of GPS monitors in  
420 competitive match-play has now been sanctioned, a more  
421 comprehensive analysis of the overall physical demands of  
422 match-play should be more permissible. This detailed

423 understanding will aid practitioners in developing complete  
424 physical training regimes.

425

426 Understanding the physical demands of specific playing  
427 positions represents an integral component of training  
428 prescription. Due to the limited sample sizes employed in  
429 previous studies, the examination of playing position has  
430 largely been restricted to basic positional comparisons (e.g.  
431 defenders, midfielders and attackers) with only one study<sup>11</sup>  
432 further differentiating between central and wide positions. The  
433 present findings support previous research which has  
434 highlighted that midfielders cover greater TD<sup>6,7,24</sup> and THSR<sup>6,7</sup>  
435 than defenders. Large differences (ES 1.4) in TD were  
436 observed between defenders and midfielders in the present  
437 study. These positional differences are similar (ES 1.6) to  
438 those previously noted in international match-play<sup>7</sup> using  
439 video-based technology. However, larger differences (ES 2.7)  
440 have been noted between defenders and midfielders during  
441 domestic match-play,<sup>7</sup> which may be a consequence of reduced  
442 tactical and physical demands of domestic relative to  
443 international match-play.

444

445 To the authors knowledge the current study is the first to  
446 examine the physical demands of specific defensive and  
447 midfield positions in competitive international female match-

448 play. Numerous differences in the physical activity profiles  
449 between CD and WD and also CM and WM were noted.  
450 Specifically, CM covered more TD and THSR than WD and  
451 CD (and A for TD only). Central defenders completed less TD  
452 and THSR than all other playing positions. The activity profile  
453 of CD is in contrast to WD, as they complete more TD, THSR  
454 and TVHSR than their central defensive counterparts. This  
455 confirms the need to analyze physical match performance  
456 across five playing positions. The findings from the current  
457 study which highlight that CM cover the greatest TD and CD  
458 the least are in accordance with previous data on European club  
459 football.<sup>11</sup> The positional differences observed in the current  
460 study are similar to those reported in male match-play<sup>2,15</sup> and  
461 are likely to be a direct consequence of the tactical role of each  
462 playing position within the team. The high requirement of  
463 midfielders to cover distance to support attacking and defensive  
464 movements is accepted and thus their greater values of TD and  
465 THSR are to be expected.

466

467 It has previously been shown that attackers complete a greater  
468 sprint distance during match-play than defenders and  
469 midfielders.<sup>8,10</sup> This finding was in part corroborated in the  
470 present study with moderate to large effect sizes shown for  
471 differences in sprinting distance between CD and other playing  
472 positions (CM (ES 1.0), WM (1.2) and A (ES 2.3)). There was

473 a trend for WM and A to complete a greater number of short  
474 sprints (<15 m) than other positions with WM undertaking a  
475 greater number of longer sprints (>15 m). Differences in the  
476 percentage of sprint type were only highlighted in CM who  
477 completed a higher proportion of explosive relative to leading  
478 sprints. The differences in sprinting profile between playing  
479 positions is again likely to be related to positional requirements  
480 in match-play. The tendency for a higher percentage of CM  
481 sprints to be explosive and shorter in nature may reflect the  
482 tighter spaces within which they operate and the tactical role of  
483 these individuals as they attempt to counteract the movement of  
484 the opposition.<sup>15</sup> Conversely, the fact that attacking players  
485 (WM and A) complete more longer sprints may be a function  
486 of their need to complete fast movements away from defending  
487 players to generate space or to capitalize on goal scoring  
488 opportunities.<sup>15</sup> The majority of differences between positions  
489 were related to CD completing less actions and distances than  
490 other playing positions across a number of the measured  
491 indices, which is most likely due to their predominant  
492 involvement being limited to defensive actions. This finding  
493 highlights the importance of analyzing positional subsets, i.e.  
494 CD versus WD not only for an understanding of match-play but  
495 also for the direct impact on training regimes.  
496

497 A unique element of the current investigation was to  
498 differentiate high-speed activity with and without the ball,  
499 which enabled the effectiveness of high-speed efforts in  
500 relation to crucial match actions to be evaluated.<sup>15</sup> A small  
501 increase in the amount of TVHSR completed when a team was  
502 without possession of the ball was observed (399±143 m vs.  
503 313±210 m, ES 0.5) as previously reported in male match-  
504 play.<sup>2,15</sup> A link between TVHSR when out of possession and  
505 team success has been demonstrated in male match-play with  
506 less successful teams completing more VHSRWP,<sup>15</sup> this  
507 analysis was beyond the scope of our study but is a  
508 recommendation for future work. Despite, an overall increase  
509 in TVHSR by the team when out of possession, the amount of  
510 TVHSR undertaken with or without possession was dependent  
511 upon playing position. Attacking positions (A, WM and CM)  
512 completed more TVHSR when the team was in possession with  
513 defensive players (CD and WD) completing more TVHSR  
514 when the team was without possession. These trends are  
515 similar to those previously reported in male match-play.<sup>2,15</sup> The  
516 observed differences in high-speed activity when a team is with  
517 and without possession, particularly between different playing  
518 positions, provides important insights for both technical and  
519 physical coaches regarding the influence of styles of play and  
520 tactical formations on the physical demands of match-play.  
521 Practitioners should consider the implementation of position-

522 specific training drills that reflect the nature of TVHSR, for  
523 example, attacking players may benefit from undertaking a  
524 greater proportion of their high-speed training with the ball  
525 compared to more defensive players, as activity that  
526 incorporates the ball has an increased energetic cost, rating of  
527 perceived exertion and blood lactate response.<sup>34</sup> However, it  
528 should be noted that the analysis of team metrics, as in the  
529 current study, limit the level of specificity that can be applied  
530 to individual players.

531

532 Previous research has used changes in physical performance  
533 both between halves and within each half as possible indicators  
534 of fatigue.<sup>35</sup> Reductions in physical performance in the second  
535 half have frequently been observed with specific reference to  
536 TD, THSR<sup>7,10</sup> and sprint distance.<sup>10</sup> In the present study, TD,  
537 THSR and sprint distances were reduced during the second  
538 half. The moderate reduction in TD (361 m; ES 0.8) between  
539 halves was greater than those reported in other studies,  
540 however, the small reduction in THSR (ES 0.5) and sprinting  
541 (ES 0.2) respectively were similar to previous reports.<sup>6,7,10</sup>  
542 Within half decreases in THSR were also currently observed,  
543 with less THSR completed during the final 15-minutes of each  
544 half compared to the previous 15-minutes. There was also a 35  
545 % reduction in THSR in the last 15-minutes of match-play  
546 compared to the first 15-minute interval. This finding was

547 similar to the 26 % reduction shown by Hewitt et al.<sup>6</sup> but less  
548 than the 57 % reduction demonstrated by Mohr et al.<sup>10</sup> These  
549 findings suggest that in some instances elite female players  
550 may be unable to perform at the required speed for the duration  
551 of the match. A second half reduction in physical performance  
552 by females has previously been attributed in part to fatigue  
553 development and an insufficient training capacity of  
554 players.<sup>7,9,10</sup> However, due to a lack of data on the match  
555 outcome, tactics, fitness status of players or biochemical  
556 markers of fatigue it is difficult to provide a clear explanation  
557 for the transient changes in high-speed activity presently  
558 observed. Furthermore, little information is currently available  
559 regarding the variability of within-game physical performance,  
560 measures. However, it is likely that differences in activity may  
561 be mediated to some extent by the inherent variation in a  
562 player's match physical performance that is associated with  
563 changes in the tactical and technical requirements of the game  
564 as opposed to fatigue.<sup>36</sup>

565

566 The current investigation reported a 39 % reduction in THSR  
567 from the most intense 5-minute period to the next 5-minutes,  
568 which was in agreement but less substantial than previous  
569 studies (48-58 %).<sup>7,10</sup> In contrast to earlier reports, the current  
570 study failed to demonstrate transient fatigue immediately after  
571 the most intense period of the match which is in agreement



572 with other more recent findings.<sup>11</sup> In the current study the  
573 reductions in THSR both toward the end of the match and  
574 following intense activity, were not as pronounced as studies  
575 that were conducted over 5 years ago. This smaller decrease in  
576 THSR may be a consequence of increased levels of  
577 professionalism and training status of female players in recent  
578 years; however, the issues of methodological differences and  
579 within game variability must also be considered. There were  
580 very few differences between positions for the changes in  
581 physical performance shown between halves, which is  
582 consistent with previous findings in females.<sup>10</sup>

583

#### 584 **PRACTICAL APPLICATIONS**

585 The present study provides an overview of the position-specific  
586 [locomotor](#) demands of competitive international female match-  
587 play. These findings are of relevance to applied practitioners  
588 responsible for the physical development of elite female  
589 players. [In order to elicit a comprehensive analysis of the](#)  
590 [overall physical demands of match-play, practitioners should](#)  
591 [combine the current dataset with information derived from GPS](#)  
592 [technology, which provide data on acceleration and](#)  
593 [deceleration profiles as well as mechanical loading. As the use](#)  
594 [of GPS monitors has now been sanctioned for use in match-](#)  
595 [play, such data will become readily available in the future.](#) A  
596 number of differences were highlighted [in the current study](#)

597 between wide and central defensive playing positions which  
598 suggest that it may be necessary for WD to complete more  
599 high-intensity soccer-specific conditioning, relative to CD, in  
600 order to cope with the increased [locomotor](#) of their playing  
601 position. During match-play the majority of sprints are less  
602 than 10 m in distance and are both explosive and leading in  
603 nature. Consequently, soccer-specific sprint drills should focus  
604 on short acceleration based activities from both a stationary and  
605 rolling start. Sprint training over longer distances (>20 m) is  
606 also required in order to condition players for longer sprint  
607 distances that may be required during match-play and to  
608 develop maximum sprinting speed. The finding that attacking-  
609 based players complete more high-speed activity when a team  
610 is in possession whilst defensive players complete more high-  
611 speed activity when a team is out of possession provides an  
612 important link between tactical and physical decision-making.  
613 Specifically, this information may be used by the coach to  
614 affect decision-making on substitutions or by the physical  
615 trainer to direct post-match training and recovery routines.  
616 Reductions in physical performance are apparent between and  
617 within halves and although these may not be entirely attributed  
618 to fatigue it emphasizes the importance of appropriate  
619 conditioning levels in order to maintain work rate.

620

621 **REFERENCES**

622

623 1 Reilly T. An ergonomics model of the soccer training  
624 process. *J Sports Sci.* 2005;23(6):561–572.

625 2 Bradley PS, Carling C, Gomez Diaz A, Hood P, Barnes  
626 C, Ade J, Boddy M, Krustup P, Mohr M. Match  
627 performance and physical capacity of players in the top  
628 three competitive standards of English professional  
629 soccer. *Hum Mov Sci.* 2013;32:808-821.

630 3 Bush M, Barnes C, Archer DT, Hogg B, Bradley PS.  
631 Evolution of match performance parameters for various  
632 playing positions in the English Premier League. *Hum*  
633 *Mov Sci.* 2015;39:1-11.

634 4 Di Salvo V, Pigozzi F, González-Haro C, Laughlin M,  
635 De Witt J. Match performance comparison in top  
636 English soccer leagues. *Int J Sports Med.*  
637 2013;34(06):526–532.

638 5 Datson N, Hulton A, Andersson H, Lewis T, Weston M,  
639 Drust B, Gregson W. Applied physiology of female  
640 soccer: an update. *Sports Med.* 2014;44(9):1225–1240.

641 6 Hewitt A, Norton K, Lyons K. Movement profiles of  
642 elite women soccer players during international matches  
643 and the effect of opposition's team ranking. *J Sports*  
644 *Sci.* 2014;32(20):1874-1880.

645 7 Andersson HÅ, Randers MB, Heiner-Møller A,  
646 Krustup P, Mohr M. Elite female soccer players

- 647 perform more high-intensity running when playing in  
648 international games compared with domestic league  
649 games. *J Strength Cond Res.* 2010;24(4):912-919.
- 650 8 Gabbett TJ, Mulvey MJ. Time-motion analysis of  
651 small-sided training games and competition in elite  
652 women soccer players. *J Strength Cond Res.*  
653 2008;22(2):543-552.
- 654 9 Krstrup P, Mohr M, Ellingsgaard H, Bangsbo J.  
655 Physical demands during an elite female soccer game:  
656 importance of training status. *Med Sci Sports Exerc.*  
657 2005;37(7):1242–1248.
- 658 10 Mohr M, Krstrup P, Andersson H, Kirkendal D,  
659 Bangsbo J. Match activities of elite women soccer  
660 players at different performance levels. *J Strength Cond*  
661 *Res.* 2008;22(2):341–349.
- 662 11 Bradley PS, Dellal A, Mohr M, Castellano J, Wilkie A.  
663 Gender differences in match performance  
664 characteristics of soccer players competing in the UEFA  
665 Champions League. *Hum Mov Sci.* 2014;33:159-171.
- 666 12 Krstrup P, Andersson H, Mohr M, Randers MB,  
667 Jensen M, Zebis M, Kirkendal D, Bangsbo J. Match  
668 activities and fatigue development of elite female soccer  
669 players at different levels of competition. In Reilly T,  
670 Korkusuz F, ed. *Science and Football VI.* Abingdon:  
671 Routledge. 2008:205-211.

- 672 13 Winter EM and Maughan RJ. Requirements for ethics  
673 approval. *J Sports Sci.* 2009;27:985
- 674 14 Di Salvo V, Collins A, McNeill B, Cardinale M.  
675 Validation of Prozone®: A new video-based  
676 performance analysis system. *Int J Perf Anal Sport.*  
677 2006;6(1):108–119.
- 678 15 Di Salvo V, Gregson W, Atkinson G, Tordoff P, Drust  
679 B. Analysis of high intensity activity in Premier League  
680 soccer. *Int J Sports Med.* 2009;30(03):205-212.
- 681 16 Bradley PS, Sheldon W, Wooster B, Olsen P, Boanas P,  
682 Krusturup P. High-intensity running in English FA  
683 Premier League soccer matches. *J Sports Sci.*  
684 2009;27(2):159-168.
- 685 17 Bradley PS, Vescovi J. Velocity thresholds for women's  
686 soccer matches: sex specificity dictates high-speed-  
687 running and sprinting thresholds - female athletes in  
688 motion (FAiM). *Int J Sports Physiol Perf.* 2015;10:112-  
689 116.
- 690 18 Dwyer DB, Gabbett T. Global positioning system data  
691 analysis: velocity ranges and a new definition of  
692 sprinting for field sport athletes. *J Strength Cond Res.*  
693 2012;26(3):818-824.
- 694 19 Weston M, Castagna C, Impellizzeri FM, Bizzini M,  
695 Williams AM, Gregson W. Science and medicine  
696 applied to soccer refereeing. *Sports Med.*

- 697 2012;42(7):615-631.
- 698 20 Weston M, Castagna C, Impellizzeri, FM, Rampinini,  
699 E, Abt, G. Analysis of physical match performance in  
700 English Premier League soccer referees with particular  
701 reference to first half and player work rates. *J Sci Med*  
702 *Sport*. 2007;10(6):390-397.
- 703 21 Reilly T, Bowen T. Exertional costs of unorthodox  
704 modes of motion. *Percept Motor Skills*. 1984;58:49-50/
- 705 22 Cnaan A, Laird NM, Slasor P. Using the general linear  
706 mixed model to analyse unbalanced repeated measures  
707 and longitudinal data. *Stat Med*. 1997;16: 2349–2380.
- 708 23 Batterham AM, Hopkins WG. Making meaningful  
709 inferences about magnitudes. *Int J Sports Physiol Perf*.  
710 2006;1:50-57.
- 711 24 Vescovi JD, Favero TG. Motion characteristics of  
712 women's college soccer matches: female athletes in  
713 motion (FAiM) study. *Int J Sports Physiol Perf*.  
714 2014;9:405-414.
- 715 25 Buchheit M, Allen A, Poon TK, Modonutti M, Gregson  
716 W, Di Salvo V. Integrating different tracking systems in  
717 football: multiple camera semi-automatic system, local  
718 position measurement and GPS technologies. *J Sports*  
719 *Sci*. 2014;32(20):1844-1857.
- 720 26 Harley JA, Lovell RJ, Barnes CA, Portas MD, Weston  
721 M. The interchangeability of global positioning system

- 722 and semiautomated video-based performance data  
723 during elite soccer match play. *J Strength Cond Res.*  
724 2011;25(8):2334–2336.
- 725 27 Randers MB, Mujika I, Hewitt A, Santisteban J,  
726 Bischoff R, Solano R, Zubillaga A, Peltola E, Krustup  
727 P, Mohr M. Application of four different football match  
728 analysis systems: A comparative study. *J Sports Sci.*  
729 2010;28(2):171–182.
- 730 28 Mohr M, Krustup P, Bangsbo J. Match performance of  
731 high-standard soccer players with special reference to  
732 development of fatigue. *J Sports Sci.* 2003;21(7): 519–  
733 528.
- 734 29 Hoff J, Wisløff U, Engen LC, Kemi OJ, Helgerud J.  
735 Soccer specific aerobic endurance training. *Br J Sports*  
736 *Med.* 2002;36:218-221
- 737 30 Kelly DM, Gregson W, Reilly T, Drust B. The  
738 development of a soccer-specific training drill for elite-  
739 level players. *J Strength Cond Res.* 2013;27(4):938–  
740 943.
- 741 31 Vescovi JD. Sprint profile of professional female soccer  
742 players during competitive matches: Female Athletes in  
743 Motion (FAiM) study. *J Sports Sci.* 2012;30(12):1259–  
744 1265.
- 745 32 Di Salvo V, Baron R, González-Haro C, Gormasz C,  
746 Pigozzi F, Bachl N. Sprinting analysis of elite soccer

- 747 players during European Champions League and UEFA  
 748 Cup matches. *J Sports Sci.* 2010;28(14):1489–1494.
- 749 33 Little T, Williams AG. Specificity of acceleration,  
 750 maximum speed, and agility in professional soccer  
 751 players. *J Strength Cond Res.* 2005;19(1):76-78.
- 752 34 Reilly T, Ball D. The net physiological cost of dribbling  
 753 a soccer ball. *Research Quarterly for Exercise and*  
 754 *Sport.* 1984;55:267-271.
- 755 35 Mohr M, Krustup P, Bangsbo J. Fatigue in soccer: a  
 756 brief review. *J Sports Sci.* 2007;23(6):593–599.
- 757 36 Gregson W, Drust B, Atkinson G, Di Salvo V. Match-to  
 758 match variability of high-speed activities in Premier  
 759 league soccer. *Int J Sports Med.* 2010;31(4):237-242.

## 760 TABLE AND FIGURE CAPTIONS

761 **Table 1** Influence of playing position on match  
 762 physical activity profile. TD = total distance; HSR = high-  
 763 speed running; THSR = total high-speed running; TVHSR =  
 764 total very high-speed running; VHSRP = total very high-speed  
 765 running with team in possession of the ball; VHSRWP = total  
 766 very high-speed running without team in possession of the ball  
 767 (mean±SD). Significant difference (p<0.05): +different from all  
 768 other playing positions, \*different from CD, ^different from A,  
 769 #different from CM, †different from WD, ‡different from WM,  
 770 \$different from percentage of leading sprints, §different from  
 771 percentage of explosive sprints. Numbers denote magnitude of



772 Effect Size for significant differences: 3 = moderate ES (>0.6-  
773 1.2), 4 = large ES (>1.2–2.0) and 5 = very large ES (>2.0).

774

775

776 **Figure 1.** Influence of playing position on the total  
777 number of sprints and the number of sprints completed over  
778 different distances (mean±SD). Significant difference  
779 (p<0.05): +different from all other playing positions, \*different  
780 from CD, ^different from A, #different from CM, †different  
781 from WD, ‡different from WM. Numbers denote magnitude of  
782 Effect Size for significant differences: 2 = small (ES>0.2-0.6),  
783 3 = moderate ES (>0.6-1.2), 4 = large ES (>1.2–2.0) and 5 =  
784 very large ES (>2.0).

785

786 **Figure 2** Influence of time (15-minute periods) on total  
787 high speed running (THSR) distance (mean±SD). Significant  
788 difference (p<0.05): +different from all other time points,  
789 #different from all time points except 16-30 mins, \*different  
790 from all time points except 46–60 mins, ^different from all time  
791 points except 61-75 mins. Numbers denote magnitude of Effect  
792 Size for significant differences: 1 = trivial (ES<0.2), 2 = small  
793 (ES>0.2-0.6), 3 = moderate ES (>0.6-1.2).