



The reinfection threshold

M. Gabriela M. Gomes^{a,*}, Lisa J. White^b, Graham F. Medley^b^a*Instituto Gulbenkian de Ciência, Apartado 14, 2781-901 Oeiras, Portugal*^b*Ecology and Epidemiology Group, Department of Biological Sciences, University of Warwick, Coventry, CV4 7AL, UK*

Received 6 January 2005; received in revised form 23 February 2005

Abstract

© 2005 Published by Elsevier Ltd.

Keywords: ■; ■; ■

Breban and Blower (2005) challenge our use of the word “threshold” to name the “Reinfection Threshold” (Gomes et al., 2004a) on the basis that this is not a bifurcation point. Assuming that a misnomer of the concept would invalidate its implications, they deduce that our results have no epidemiological consequences. Here we explain the terminology but, more importantly, we emphasize that the choice of name does not affect the implications of the concept. We assert that the epidemiological behaviour associated with $R_0 = 1/\sigma$ remains unchanged.

The recognition that partial immunity divides the transmissibility axis into two distinct modes of transmission emerged almost simultaneously from two studies: a systematic analysis of microparasite transmission under suboptimal immunity (Gomes et al., 2004a); and a representation of the transmission dynamics of tuberculosis leading to a hypothesis for the widely debated variable efficacy of the bacille Calmette-Guérin (BCG) vaccine (Gomes et al., 2004b). The two studies overlap through a modified *SIR* model

$$\begin{aligned}\frac{dS}{dt} &= (1-v)e - R_0IS - eS, \\ \frac{dI}{dt} &= R_0I(S + \sigma R) - I,\end{aligned}\quad (1)$$

where S and I are the proportions of fully susceptible and infected individuals, respectively, and $R = 1 - S - I$ is the proportion recovered with partial immunity.

In units of average duration of infection, births and deaths occur at a rate e , and a proportion v of the population is vaccinated at birth to enter compartment R . Individuals in S are infected at a rate, R_0I , and individuals in R are infected at a reduced rate, σR_0I , where $0 \leq \sigma \leq 1$. The existence of these two distinct rates of infection is the key to our findings.

The endemic equilibria of model (1) are represented in Fig. 1 as a function of the basic reproduction number, R_0 . The proportion of infectious individuals is represented in linear (a) and logarithmic (b) scales, and the four curves correspond to different vaccination coverage: $v = 0, 0.67, 0.89, 1$, from left to right. For the purpose of illustration we chose $\sigma = 0.25$, but the conclusions that follow remain valid for any value between 0 and 1. We set $e = 0.0012$, which is attained under a life expectancy of 70 years and an average duration of infection of 1 month. Shorter infectious periods would enhance the effects that follow.

Breban and Blower (2005) carry out a bifurcation analysis of the $v = 0$ model, and correctly conclude that

DOI of original article: 10.1016/j.jtbi.2004.12.026

*Corresponding author. Fax: +351 214 407 970.

E-mail address: ggomes@igc.gulbenkian.pt (M.G.M. Gomes).




Fig. 1. Endemic equilibria, and potential impact of vaccination as a function of transmissibility illustrated for $\sigma = 0.25$. (a, b) proportion infectious in linear and logarithmic scale, respectively, under four levels of vaccination coverage: $v = 0, 0.67, 0.89, 1$, from left to right. The non-vaccination line marks the epidemic threshold ($R_0 = 1$), and the full-vaccination line (dashed) marks the reinfection threshold ($R_0 = 1/\sigma$). (c) vaccination coverage required to eliminate infection in the controllable (light) zone. Mass vaccination is generally ineffective in the uncontrollable (dark) zone.

the only bifurcation point is the epidemic threshold, $R_0 = 1$ (Kermack and McKendrick, 1927). However, we note that all curves show a marked increase in the prevalence of infection as R_0 increases across $R_0 = 1/\sigma$. In Fig. 1(a) this effect is more evident, and in Fig. 1(b) we observe that vaccination can eliminate infection for $1 < R_0 < 1/\sigma$, while this is ineffective for $R_0 > 1/\sigma$. Fig. 1(c) shows the vaccination coverage required to eliminate infection in the controllable (light) zone ($1 < R_0 < 1/\sigma$) while this is impossible in the uncontrollable (dark) zone ($R_0 > 1/\sigma$). We attribute this behaviour to a bifurcation in the reinfection submodel that takes place at $R_0 = 1/\sigma$. The reinfection submodel is obtained

by setting $v = 1$, and the associated equilibria are thus represented by the dashed curve in Fig. 1(b). Consequently, we term $R_0 = 1/\sigma$ as the “reinfection threshold”. When $v < 1$, the reinfection threshold is no longer a bifurcation point, but remains as a useful value to interpret levels of infection and to determine the potential impact of vaccination programmes.

The reinfection threshold is determined by the degree of protection conferred by a previous infection or vaccine, whichever is more protective. Therefore, its position can be manipulated by more potent vaccines widening the controllable zone. The reinfection threshold sets the targets for vaccine development and use, and

1 its quantitative characterization cannot be overempha-
 2 sized. This is perhaps the most important message for
 3 public health.

5 **Final remarks**

7 Our final remarks refer to the last paragraph of
 9 Breban and Blower (2005). Their final sentence is
 10 misleading.¹ The authors restate that no epidemiological
 11 consequences can be assigned to the reinfection thresh-
 12 old because $R_0 = 1/\sigma$ is not a bifurcation point. They
 13 claim that our findings concerning reinfection reduce to
 14 a nonlinear increase in the prevalence of infection as R_0
 15 increases, as previously reported by Blower et al. (1998)
 16 and Feng et al. (2000). This is not the case. As described
 17 above we have analysed the form of the nonlinear
 18 increase and the variability in the effectiveness of
 19 vaccination. We have detected a bifurcation in the
 20 reinfection submodel at $R_0 = 1/\sigma$, we have shown that
 21 levels of infection of the full epidemic model increase by
 22 two orders of magnitude as R_0 increases across a vicinity
 23 of $1/\sigma$, we have demonstrated that the same vaccine can
 24 eliminate infection if $R_0 < 1/\sigma$ and appears impotent if

$R_0 > 1/\sigma$. We named this critical value the reinfection
 27 threshold and discussed its implications for vaccine
 28 development and public health policy.

31 **References**

- 33 Blower, S.M., Porco, T.C., Lietman, T., 1998. Tuberculosis: the
 34 evolution of antibiotic resistance and the design of epidemic
 35 control strategies. In: Horn, M.A., Siminett, G., Webb, G.F.
 36 (Eds.), *Mathematical Models in Medical and Health Sciences*.
 37 Vanderbilt University Press, Nashville & London.
- 38 Breban, R., Blower, S., 2005. The reinfection threshold does not exist.
 39 *J. Theor. Biol.* (this issue).
- 40 Feng, Z., Castillo-Chavez, C., Capurro, A.F., 2000. A model for
 41 tuberculosis with exogenous reinfection. *Theor. Popul. Biol.* 57,
 42 235–247.
- 43 Gomes, M.G.M., White, L.J., Medley, G.F., 2004a. Infection,
 44 reinfection, and vaccination under suboptimal immune protection:
 45 epidemiological perspectives. *J. Theor. Biol.* 228, 539–549.
- 46 Gomes, M.G.M., Franco, A.O., Gomes, M.C., Medley, G.F., 2004b.
 47 The reinfection threshold promotes variability in tuberculosis
 48 epidemiology and vaccine efficacy. *Proc. R. Soc. London B* 271,
 49 617–623.
- Kermack, W.O., McKendrick, A.G., 1927. A contribution to the
 mathematical theory of epidemics. *Proc. R. Soc. A* 115, 700–721.

¹Breban and Blower (2005) refer to tuberculosis and quote from a
 previous article: “the higher the susceptibility to reinfection the easier
 it will be to achieve eradication” (Blower et al., 1998). The full sentence
 in the original article is: “For any particular (endemic equilibrium)
 incidence rate, the higher the susceptibility to reinfection the easier it
 will be to achieve eradication” (Blower et al., 1998). The two
 statements have different meanings. The first could imply that
 increasing susceptibility to reinfection with *M. tuberculosis* (e.g. by
 spreading AIDS) would bring us closer to eradicating tuberculosis.
 This does not make sense. The second applies solely to the
 interpretation of a given endemic equilibrium: the higher the
 reinfection, the lower the R_0 required for a given endemic equilibrium
 and, in principle, lower R_0 implies easier eradication.