

# Transcriptional Profiling of Arabidopsis Tissues Reveals the Unique Characteristics of the Pollen Transcriptome<sup>1[w]</sup>

Jörg D. Becker, Leonor C. Boavida, Jorge Carneiro, Matthias Haury, and José A. Feijó\*

Instituto Gulbenkian de Ciência, PT-2780-156 Oeiras, Portugal (J.D.B., L.C.B., J.C., M.H., J.A.F.); and Centro de Biotecnologia Vegetal, Faculdade de Ciências, Universidade de Lisboa, Campo Grande, Ed.C2. PT-1749-016 Lisboa, Portugal (L.C.B., J.A.F.)

**AQ: A** Pollen tubes are a good model for the study of cell growth and morphogenesis because of their extreme elongation without cell division. Yet, knowledge about the genetic basis of pollen germination and tube growth is still lagging behind advances in pollen physiology and biochemistry. In an effort to reduce this gap, we have developed a new method to obtain highly purified, hydrated pollen grains of Arabidopsis through flowcytometric sorting, and we used GeneChips (Affymetrix, Santa Clara, CA; representing approximately 8,200 genes) to compare the transcriptional profile of sorted pollen with those of four vegetative tissues (seedlings, leaves, roots, and siliques). We present a new graphical tool allowing genomic scale visualization of the unique transcriptional profile of pollen. The 1,587 genes expressed in pollen showed a 90% overlap with genes expressed in these vegetative tissues, whereas one-third of the genes constitutively expressed in the vegetative tissues were not expressed in pollen. Among the 469 genes enriched in pollen, 162 were selectively expressed, and most of these had not been associated previously with pollen. Their functional classification reveals several new candidate genes, mainly in the categories of signal transduction and cell wall biosynthesis and regulation. Thus, the results presented improve our knowledge of the molecular mechanisms underlying pollen germination and tube growth and provide new directions for deciphering their genetic basis. Because pollen expresses about one-third of the number of genes expressed on average in other organs, it may constitute an ideal system to study fundamental mechanisms of cell biology and, by omission, of cell division.

**Fn1** Pollen has been the subject of intense studies not only for its importance as the male partner in plant reproduction, but also as a model for the study of cell growth and morphogenesis in a broader sense (Feijó et al., 2001). Whereas early studies focused on pollen physiology and biochemistry (for review, see Mascarenhas, 1975), the last 20 years have been marked by increasing efforts to decipher the genetic basis of pollen development and functions (for reviews, see Scott et al., 1991; McCormick, 1993; Preuss, 1995; Taylor and Hepler, 1997; Franklin-Tong, 1999; Hepler et al., 2001; Lord and Russell, 2002). These efforts led to the identification of more than 150 pollen-expressed genes from more than 28 species (Twell, 2002). These genes encode proteins thought or known to be involved in pollen development, pollen germination, and pollen tube growth, as well as in interactions with the stigma/transmitting tissue or

the female gametophyte. Many of these studies were conducted in *Lilium* sp. or in Solanaceae species, because their pollen is easily collected in sufficient quantities, and methods for in vitro germination are robust. However, the available genetic information for these species is limited. The availability of the genome sequence of Arabidopsis (The Arabidopsis Genome Initiative, 2000) and the concomitant increase in available genomic tools (Wixon, 2001) make Arabidopsis a preferable model for large scale genetic studies of pollen germination and tube growth. The shift toward Arabidopsis is reflected in recent studies of the cellular organization and ultrastructure of in vivo- and in vitro-grown pollen tubes of Arabidopsis (Lennon and Lord, 2000; Derksen et al., 2002) and by efforts to improve in vitro germination of Arabidopsis pollen (Fan et al., 2001).

Studies in several plant species have indicated that the bulk of mRNAs needed for pollen germination and tube growth accumulates in pollen grains well before germination (Mascarenhas, 1989; Guyon et al., 2000). Thus, we expected that identifying transcripts that were enriched (up-regulated gene expression) or even selectively expressed in hydrated pollen grains on a genomic scale would increase significantly our knowledge of the genetic basis of pollen germination and tube growth. We chose an approach using pollen of Arabidopsis and overcame the limitations of small-scale transcriptional analysis applying

<sup>1</sup> This work was supported by the Fundação para a Ciência e a Tecnologia (FCT; project nos. POCTI/BCI/41725/2001, POCTI/BIA/34772/1999, and POCTI/BCI/46453/2002). J.D.B. and L.C.B. were supported by FCT fellowships SFRH/BPD/3619/2000 and SFRH/BD/1128/2000.

<sup>[w]</sup> The online version of this article contains Web-only data.

\* Corresponding author; e-mail [jose.feijo@fc.ul.pt](mailto:jose.feijo@fc.ul.pt); fax XXX-XXX-XXXX.

**AQ: M** Article, publication date, and citation information can be found at [www.plantphysiol.org/cgi/doi/10.1104/pp.103.028241](http://www.plantphysiol.org/cgi/doi/10.1104/pp.103.028241).


Becker et al.

oligonucleotide-based microarray technology. The Arabidopsis Genome GeneChip array (Affymetrix, Santa Clara, CA) has been used to examine the transcriptional regulation of genes in Arabidopsis by the circadian clock (Harmer et al., 2000) and to identify gene expression changes during shoot development (Che et al., 2002). As a prerequisite for high-quality transcriptional profiles of pollen grains, we developed a protocol using fluorescence-activated cell sorting (FACS) to obtain highly purified hydrated pollen grains of Arabidopsis. We obtained transcriptional profiles for hydrated pollen grains and for four types of vegetative tissues (leaves, roots, seedlings, and siliques) of Arabidopsis. By comparison of these data sets we show that the transcriptional profile of Arabidopsis pollen is clearly distinguishable from those of the vegetative tissues. About 1,500 genes (approximately 90%) expressed in pollen grains are also expressed in at least one of the vegetative tissues, whereas the remaining 10% (162 genes) are selectively expressed in pollen. Most of these 162 genes have not been described as selectively expressed in Arabidopsis pollen before, and their functional classification yields new insights into several aspects of the genetic program underlying germination and tube growth of Arabidopsis pollen.

## RESULTS AND DISCUSSION

### FACS Yields Highly Purified Arabidopsis Pollen Grains

An essential requirement for obtaining high-quality DNA array data is purity of the tissue or cellular source for RNA extraction, because any kind of impurity could result in an inaccurate transcriptional profile. Because the reported sensitivity of the Arabidopsis Genome GeneChip arrays is one transcript in 100,000 to one transcript in 300,000 (Zhu and Wang, 2000), we consider this requirement especially important for pollen. Therefore, we have developed a new protocol to obtain highly purified, hydrated pollen grains. Prehydrated pollen was washed out from flowers in buffer and subjected to filtering steps. The resulting solution contained fully hydrated pollen grains (oval shape), non-hydrated and/or destroyed pollen grains (round shape) as well as smaller impurities (Fig. 1A). For a final purification step, we used FACS using the size and autofluorescence properties of pollen. To separate hydrated from non-hydrated pollen, a two-dimensional histogram using the forward scatter versus pulse width parameters was used, and an additional gate was then applied on the 670/40 and 580/20 nm detection channels to further purify the pollen grains based on their characteristic autofluorescence properties (see Fig. 2). The characteristic oval shape of fully hydrated pollen grains caused a longer pulse when passing the laser, thus allowing a separation of the non-hydrated pollen grains and smaller impurities from the fully hydrated



**Figure 1.** Purification steps and viability of Arabidopsis pollen. Pollen and impurities before sorting (A) were separated into non-hydrated (B) and hydrated pollen (C) by flowcytometric sorting. D, A viability stain of the sorted, hydrated pollen grains.


pollen grains (Fig. 1, B and C). The purity of the hydrated pollen grain fraction was routinely >99%. The viability of the hydrated pollen grains was assessed by fluorescein diacetate staining (Fig. 1D). The remaining impurities (<1%) consisted of very small debris, which most likely did not include organelles. Hydrated pollen grains (150,000) yielded 1  $\mu$ g of total RNA. More than 500,000 non-hydrated pollen grains resulted in the same yield, supporting our assumption that the majority of the non-hydrated pollen grains were not intact.

### Pollen Grains Have a Unique Transcriptional Profile

Gene expression patterns of approximately 8,200 genes, representing roughly one-third of the Arabidopsis genome, were obtained for Arabidopsis pollen grains and for several vegetative tissues: seedlings, leaves, root, and siliques. The results were highly reproducible as underlined by the high correlation coefficients of the replicates, which ranged from 0.977 to 0.992. For each vegetative tissue, a similar percentage of genes were called Present by the MAS 5 algorithm, with a mean of 59% in seedlings, 56% in leaves, and 64% in roots and siliques. In contrast, only 21% of the genes represented on the arrays were called Present in the pollen samples (1584 unique genes). Normalization reduces variation of non-biological origin and is therefore a prerequisite for the direct comparison of expression profiles from different arrays. The large differences in the transcriptional profiles in this study, i.e. the

AQ: C  
F1

F2



**Figure 2.** Flowcytometric sorting of Arabidopsis pollen. Arabidopsis pollen was identified through its size (A) and autofluorescence properties (B). A, Hydrated pollen was located in region R1 of the pulse width versus FSC display, whereas non-hydrated pollen was mostly contained in R2. B, For sorting, a logical gate combination of regions R1 and R3 was used.


sample-specific differences in Presence calls, precluded the use of global scaling/normalization methods. Instead, we employed a sample-wise normalization to the median median probe cell intensity of all arrays, implemented into version 1.3 of the dChip software (<http://www.dchip.org>; Wong laboratory, Harvard, Boston). This method works independently from the overall intensity of an array. After normalization and model-based computation of expression values, we excluded genes called Absent in all arrays and genes with inconsistent expression levels within the replicate arrays. Thus, for further analysis, our data set contained 6,459 genes.

We developed a new graphical tool to visualize the striking differences between the transcriptional profiles of the vegetative tissues and pollen (Fig. 3). This tool, "Snail View", compares and displays the changes in expression levels of thousands of genes simultaneously, but still allows meaningful interpretations of the overview obtained (the software can be downloaded at <http://eao.igc.gulbenkian.pt/ti/Soft/SnailView/>). The average expression value in the seedling samples were chosen as reference, considering that seedlings probably contain cell types found in roots, leaves and, most likely, in siliques. The high correlation of data derived from the replicates is exemplified by the comparison of expression values obtained for the single-seedling replicates. This high correlation is visualized by the small deviations from the line representing the average value of the seedling replicates. The expression values of

leaves and of seedlings are strongly correlated, especially for those genes with the highest expression level in seedlings. The cotyledons contribute the largest part to the biomass of 4-d-old seedlings, so this similarity was anticipated. Expression values obtained for siliques and root samples show high deviations from the seedling reference. Genes highly expressed in seedlings are down-regulated (most considerably in roots), and those with low expression levels in seedlings are up-regulated (most considerably in siliques). The most dramatic differences are seen in the pollen to seedling comparison. Most genes with high or medium expression values in seedlings show low expression values in pollen. This trend is reversed for genes with low expression values in seedlings, because a high proportion of these genes are highly expressed in pollen and reach expression values comparable with those of genes with the highest expression in seedlings. The correlation coefficients (0.032, 0.029, 0.040, and 0.067) of the expression values of pollen relative to expression values of seedlings, leaves, siliques, and roots, show that pollen has a transcriptional profile that is clearly distinguished from that of vegetative tissues.

#### Ten Percent of the Genes Expressed in Pollen Are Not Expressed in the Sporophyte

We assumed that gene products of transcripts that were highly enriched or even selectively expressed in pollen grains might be of major importance, if not



**Figure 3.** "Snail view" representation of tissue-dependent gene expression patterns. The expression for 5,999 genes is represented in angular coordinates, in which angle encodes gene rank (clockwise from top) and radius encodes the logarithm of gene expression (values  $<1$  were set to 1 for better visualization). Genes were ranked according to increasing mean expression in seedlings. For each tissue, the mean pattern in seedlings (black line) is coplotted with the mean pattern of specific tissue (gray dots), except for seedling where one replicate is coplotted. The values are the correlation coefficient between the gene expression patterns in the two tissues (for seedlings between the replicates).



Becker et al.


crucial for successful pollen germination and tube growth. To identify such genes, we followed two complementing approaches. First, we measured enrichment in gene expression as the lower limit (lower confidence bound) of a 90% confidence interval for the -fold change in gene expression. Gene expression values in pollen were compared with those in each of the vegetative tissues, and a score above 1.2 was used as the criterion to select genes enriched in pollen; 469 genes met this criterion in all four comparisons (supplemental Table I). Second, we used Affymetrix MAS 5 Present calls to sort genes; 5,775 genes were reproducibly called Present in at least one of the four vegetative tissues. Of the 1,584 genes called Present in pollen (for hierarchical clustering, see Fig. 4A), 1,422 (90%) showed an overlap with the genes detected in vegetative tissues (Fig. 4B). The remaining 10% (162 genes) were called Present only in pollen and are referred to as selectively expressed from hereon. The two methods yielded an overlap of 150 genes that we characterized in more detail.

Thus, our expression analysis of roughly one-third of the annotated genes of Arabidopsis shows that 10% of the genes expressed in pollen are selectively

expressed in pollen, whereas the other 90% are also expressed in one or more vegetative tissues. This substantial overlap of genes active in the sporophyte and in the male gametophyte had been predicted earlier based on isoenzyme studies in pollen and vegetative tissues in barley (*Hordeum vulgare*; Pederson et al., 1987), heterologous hybridizations of pollen cDNA with shoot poly(A) RNA and shoot cDNA to pollen poly(A) RNA in maize (*Zea mays*; Willing et al., 1988), as well as colony hybridizations of maize pollen cDNA libraries with cDNAs from pollen and vegetative tissues (Stinson et al., 1987).

### Functional Classification of Selectively Expressed Genes

We sorted the 150 selectively expressed genes into functional categories (Table I; Fig. 4C), taking into consideration several aspects of current knowledge about the pollen grain and pollen tube physiology. This categorization is based on known functions of the gene products as well as gene ontology annotations derived from homologies. Our analysis confirmed the expression of some genes already known



**Figure 4.** Analysis of genes enriched or selectively expressed in pollen. A, Hierarchical cluster analysis of the 1,584 genes called Present in pollen. Default parameters in dChip were used (standardization and clustering methods follow Golub et al. [1999] and Eisen et al. [1998]). Each column represents a single gene, and each row a single tissue type. Genes mainly enriched in pollen are found in the right cluster, whereas those expressed in pollen but with higher expression values in at least one of the vegetative tissues are found in the left cluster. P, Pollen; Se, seedling; L, leaf; Si, silique; R, root. B, Venn diagram depicting total numbers of genes expressed in vegetative tissues and in pollen. Genes were scored as Present in the vegetative tissues, if they received at least one Present call (MAS 5) in the seedling, leaf, root, or silique sample (5,775 genes). In pollen, 1,422 genes of these were also called Present. A set of 162 genes was exclusively called Present in pollen. Genes recognized by more than one probe set in the Affymetrix array were only counted once. C, Overview of the functional classification of 150 pollen selectively expressed genes listed in Table I. Total number of genes in each category is given in brackets. Functional classification is based on established as well as putative functions.

**Table 1.** Functional classification of genes selectively expressed in pollen

Annotation	Signal	FC	Probe Set	Arabidopsis Genome Initiative Identification
<b>Signal transduction (37)</b>				
Putative protein kinase	4,728	76.8	17803_at	At1g61860
Putative receptor-like protein kinase	4,061	78.4	18205_s_at	At4g39110
Putative Pro extensin-like receptor kinase PERK7	4,010	20.3	17812_at	At1g49270
Putative calcium-dependent protein kinase CPK24	3,538	160.8	16806_s_at	At2g31500
Putative 4,5 PIP kinase	3,425	45.9	19788_at	At1g01460
Putative calmodulin	3,326	214.9	13415_at	At4g03290
Putative Leu-rich repeat transmembrane protein kinase AtPRKc	3,051	80.7	19047_at	At2g07040
Putative phosphatidylinositol/phosphatidylcholine transfer protein	1,887	15.0	19052_at	At2g18180
Putative calcium-dependent protein kinase CPK14	1,511	250.6	20236_s_at	At2g41860
Putative protein kinase	1,083	16.9	18152_at	At2g24370
Putative Leu-rich repeat transmembrane protein kinase AtPRKd	1,026	17.2	19103_at	At5g35390
Calcium-dependent protein kinase-like protein CPK26	854	26.4	20241_at	At4g38230
Putative Ser/Thr protein kinase	725	21.7	14364_at	At1g04700
Putative protein kinase (cdc2 kinase homolog cdc2MaC, <i>M. sativa</i> )	670	48.9	16824_at	At4g10010
Putative GTP-binding protein	652	11.6	18107_at	At2g33870
Putative inositol polyphosphate 5'-phosphatase	636	8.8	19795_at	At2g43900
Putative proline extensin-like receptor kinase PERK5	611	29.0	16876_at	At4g34440
Putative protein (GTPase-activating protein, <i>Yarrowia lipolytica</i> )	574	4.9	16758_at	At4g27100
Putative inositol polyphosphate 5'-phosphatase	563	17.1	19582_at	At2g31830
Putative protein (various predicted protein kinases, Arabidopsis)	532	11.0	18193_at	At5g26150
AK20 gene protein kinase catalytic domain (fragment)	519	6.1	19111_at	
Putative receptor-like Ser/Thr kinase RKF2	483	10.4	17496_at	At1g19090
Putative Pro extensin-like receptor kinase PERK12	335	14.7	17359_at	At1g23540
Hypothetical protein (contains similarity to wall-associated kinase 1, Arabidopsis)	324	10.8	18186_at	At1g78940
Putative Pt $\alpha$ kinase interactor (similar to Pt $\alpha$ kinase interactor 1, tomato)	300	3.8	18825_at	At4g13190
Putative GTP-binding protein	292	16.3	12489_at	At2g22290
Protein phosphatase 2C (PP2C)	275	4.8	19345_at	At2g40180
Putative calcium-dependent protein kinase CPK20	241	8.8	12311_at	At2g38910
Putative calmodulin (calmodulin PIR1:MCKM, <i>Chlamydomonas reinhardtii</i> )	229	4.7	19274_i_at	At4g12860
Protein kinase-like protein	214	5.7	19556_at	At4g31220
Putative protein (various predicted protein kinases, Arabidopsis)	159	2.2	19071_at	At5g35380
Putative Ser/Thr protein kinase	110	4.0	16866_at	At1g79250
Putative STE20/PAK-like protein kinase (putative MAP4K similar to yeast STE20)	74	4.7	18999_at	At1g70430
Putative receptor-like kinase	65	3.4	14275_at	At4g31230
Putative protein kinase	62	1.8	16848_at	At2g20470
Putative phosphatidylinositol-4-phosphate 5-kinase	55	4.3	19310_at	At2g41210
Calcium-dependent Ser/Thr protein kinase CPK18	44	1.8	16849_at	At4g36070
<b>Cell wall biosynthesis and regulation (16)</b>				
Pectinesterase family (nearly identical to pollen-specific BP10 protein, canola)	7,180	110.3	17753_at	At1g55570
Exopolysaccharuronase	7,059	51.9	18923_g_at	At3g07850
Extensin-like protein	6,221	71.5	15765_at	At4g33970
Putative polygalacturonase	5,660	91.2	15716_at	At2g23900
Putative cellulose synthase (AtCalD1—homolog of NaCalD1)	4,911	58.9	20243_at	At2g33100
Glycosyl hydrolase family 10 (xylan endohydrolase isoenzyme X-I, barley)	4,536	40.1	13393_at	At4g33860
Polysaccharide lyase family 1 (similar to pectate lyase P59, tomato)	4,226	184.1	16837_at	At2g02720
Polysaccharide lyase family 1 (At59)	3,914	83.0	16857_at	At1g14420
Putative cellulose synthase (AtCalD4—homolog of NaCalD1)	3,489	28.5	16782_at	At4g38190
Putative protein (bp4A protein, Brassica napus)	1,196	51.4	15360_at	At4g11760
COBRA-like protein COBL-11	929	13.1	15331_at	At4g27110
Cellose synthase (1,3- $\beta$ -glucan synthase) family (AtGal2, homolog of NaGal1)	925	2.7	18515_at	At2g13675
Pectinesterase family	662	19.2	18519_at	At4g33230
Glycosyl hydrolase family 17 (BETAG5, $\beta$ -1,3-glucanase bg5)	438	5.9	18539_at	At5g20340
Putative xyloglucan endotransglycosylase	214	5.8	19530_at	At4g18990
Glycosyl hydrolase family 17 (BETAG4, $\beta$ -1,3-glucanase bg4)	214	2.3	18120_i_at	At5g20330

(Table continues on following page.)

Becker et al.

**Table I.** (Continued from previous page.)

Annotation	Signal	FC	Probe Set	Arabidopsis Genome Initiative Identification
<b>Stress response (11)</b>				
Putative plant defensin protein PDF2.6	9,739	53.2	19928_at	At2g02140
Putative peroxiredoxin (similar to type 2 peroxiredoxin, <i>Brassica rapa</i> )	3,640	7.3	15966_i_at	At1g60740
17.6-kD heat shock protein (AA 1–156)	1,637	26.5	13276_at	At1g53540
Heat shock protein 17.6A (At-HSP17.6A)	731	17.2	13278_f_at	At5g12030
Heat shock protein 22.0 (At-HSP22.0)	431	8.6	17076_s_at	At4g10250
Putative protein (similar to Mlo proteins, barley)	402	4.8	14639_at	At2g33670
Heat shock protein 18	299	3.1	13280_at	At5g59720
Putative stress protein	130	6.1	13378_at	At2g01330
Putative peroxidase	108	4.4	17828_at	At4g17690
Mitochondrion-localized small heat shock protein (AtHSP23.6-mito)	69	2.0	13282_s_at	At4g25200
Putative disease resistance protein (TIR-NBS-LRR class)	61	2.8	12258_at	At4g14370
<b>Metabolism (9)</b>				
Glycosyl hydrolase family 35 ( $\beta$ -galactosidase) BGAL11	6,152	194.9	16874_at	At4g35010
Glycosyl hydrolase family 32 (invertase)	3,759	54.4	16369_s_at	At3g52600
Glycosyl hydrolase family 35 ( $\beta$ -galactosidase) BGAL13	3,074	69.7	17316_at	At2g16730
FAD-linked oxidoreductase family	2,481	126.4	15302_at	At1g11770
Putative thioredoxin	2,236	21.2	19937_at	At2g33270
Putative acyl-CoA:1-acylglycerol-3-phosphate acyltransferase	656	10.4	18423_at	At1g51260
3-hydroxy-3-methylglutaryl CoA reductase HMG1 isoform HMGR1L	550	3.0	20160_at	
Glycosyl hydrolase family 1	101	18.6	19601_at	At2g25630
Putative CDP-diacylglycerol synthetase	53	1.7	19016_at	At4g26770
<b>Ion transport (8)</b>				
Putative cation/H <sup>+</sup> antiporter AtCHX8	3,259	17.1	17246_at	At2g28180
Putative protein (anion exchange protein 2, <i>Homo sapiens</i> )	3,208	67.1	18837_at	At4g32510
Shaker potassium channel SPIK	1,489	51.8	16783_at	At2g25600
Putative cation/H <sup>+</sup> antiporter AtCHX15	618	4.3	14334_at	At2g13620
Ion transporter-like protein AtNRAMP5	404	30.2	18118_at	At4g18790
Putative cyclic nucleotide-regulated ion channel AtCNGC16	205	8.3	14291_at	At3g48010
G subunit of vacuolar-type H <sup>+</sup> -ATPase (vag1)	78	5.3	16226_at	At4g25950
Putative cation/H <sup>+</sup> antiporter AtCHX13	60	2.4	15704_at	At2g30240
<b>Cytoskeleton (8)</b>				
Actin 4	3,262	110.7	14356_at	At5g59370
Actin depolymerizing factor-like protein	3,092	37.4	12529_at	At4g25590
Actin 12	1,815	35.8	15697_at	At3g46520
Prolilin 4	2,691	25.8	12411_s_at	At2g19770
Myosin heavy chain-like protein AtVIII B	341	19.7	12257_at	At4g27370
Putative myosin heavy chain AtXID	224	2.7	16327_at	At2g33240
Putative myosin II heavy chain ( <i>Naegleria fowleri</i> )	179	6.2	15329_at	At4g33390
Kinesin-related protein	97	3.3	16794_at	At1g09170
<b>Transcriptional regulation (5)</b>				
Putative protein (NorM, <i>Vibrio parahaemolyticus</i> )	1,471	33.5	14992_at	At4g21900
Transcription factor MYB97 (similar to anther-specific myb-related protein 2, tobacco)	309	10.4	17979_at	At4g26930
MADS-box protein AGL30 (homolog of MADS1;11 tobacco)	167	5.2	18443_at	At2g03060
MADS-box protein AGL29	154	15.8	19309_at	At2g34440
CONSTANS B-box zinc finger family protein	23	1.9	15698_at	At4g15250
<b>Proteolysis (5)</b>				
Putative Ser carboxypeptidase II	7,644	55.0	17765_at	At2g05850
Ser-type carboxypeptidase-like protein	4,000	117.4	18137_at	At3g52010
Subtilisin-like Ser protease	1,348	26.8	16349_at	At4g21326
Ser-type carboxypeptidase-like protein	721	12.1	17814_at	At3g52000
Ser-type carboxypeptidase-like protein	64	6.4	18461_at	At3g52020
<b>Vesicle trafficking (5)</b>				
Syntaxin AtSYP124	1,243	10.7	15275_at	At1g61290
ARF GTPase-activating domain-containing protein (similar to ARFGAP1, <i>H. sapiens</i> )	829	15.2	12639_at	At2g35210
Clathrin protein family	631	8.9	13499_at	At1g05020
ARF GTPase-activating domain-containing protein	598	4.4	17446_at	At2g14490
Unknown protein (similar to yeast Sec7p protein)	280	7.9	18381_at	At2g30690

(Table continues on following page.)

Table 1. (Continued from previous page.)

Annotation	Signal	FC	Probe Set	Arabidopsis Genome Initiative Identification
Sugar and protein transport (3)				
Monosaccharide transporter	4,441	129.6	19270_at	At5g23270
Putative ABC transporter AtNAP12	437	4.6	13049_at	At2g37010
Putative vacuolar sorting receptor	182	2.8	18509_at	At2g30290
Transposition (3)				
Ac-like transposase (related to Ac/Ds transposon family of maize)	198	4.8	16730_at	At2g16040
Putative Mutator-like transposase	118	2.1	12987_s_at	At2g30640
Putative TNP2-like transposon protein (related to the En/Spm transposon family of maize)	60	2.6	19339_i_at	At2g10140
Water channel (2)				
Aquaporin-like protein TIP5;1 (aquaporin, <i>Vernicia fordii</i> )	1,497	13.6	18856_at	At3g47440
Putative water channel protein—major intrinsic protein (MIP) TIP1;3	544	6.1	17798_at	At4g01470
Translational regulation (2)				
Putative protein (similar to polyadenylate-binding protein PABP)	429	10.9	16836_at	At3g16380
Putative translation initiation factor	65	5.2	18609_at	At2g39820
Cell cycle regulation (1)				
Cyclin A2;1	89	1.8	16851_s_at	At5g25380
Unknown and hypothetical (35)				
Arabinogalactan-protein AGP6	16,351	50.8	12003_at	At5g14380
Nectarin-like protein (nectarin I precursor, <i>Nicotiana plumbaginifolia</i> )	11,576	81.8	19014_at	At5g26696
Hypothetical protein	9,322	89.6	16201_at	At1g49490
Unknown protein (homolog of N;t; auxin-repressed/dormancy-associated protein ARPL 1;1)	9,259	181.8	19385_at	At1g54070
Putative LIM-domain protein (AtPLIM2)	8,733	82.0	20478_at	At2g45800
Unknown protein	8,161	92.0	19301_at	At4g36490
Unknown protein	7,359	35.4	12568_at	At2g33420
Putative pistil-specific protein (similar to potato pistil-specific protein STS15)	7,018	149.3	14936_at	At4g02250
Hypothetical protein	4,933	47.1	13912_at	At1g04540
Unknown protein	3,744	101.5	12638_at	At1g04470
Hypothetical protein	3,186	61.3	18426_at	At2g13350
Unknown protein	1,225	42.7	17230_at	At2g40990
Unknown protein	962	96.2	20549_i_at	At2g26850
Hypothetical protein	818	33.4	17687_at	At1g60240
Putative GDP-dissociation inhibitor	753	28.9	19790_at	At1g12070
Hypothetical protein	466	5.4	15925_at	At1g43630
Unknown protein	420	13.4	13836_at	At1g06750
Unknown protein	379	27.4	15838_at	At2g29790
Hypothetical protein	377	2.7	13304_at	At1g24330
Putative protein (MSP1, <i>Saccharomyces cerevisiae</i> )	348	21.5	19604_at	At4g28000
Unknown protein (related to MO25, early mouse development protein family)	329	28.7	17711_at	At2g03410
RALF-LIKE 10	313	7.0	18385_at	At2g19020
Hypothetical protein	258	6.9	13814_at	At2g29040
Unknown protein	254	5.8	15277_at	At2g41440
Glu-rich protein (mature-parasite-infected erythrocyte surface antigen MESA, <i>P. falciparum</i> )	238	22.1	16656_at	At4g20160
Putative protein	227	18.3	15322_at	At4g24580
Unknown protein	221	7.6	19849_at	At2g18080
Unknown protein	219	4.3	14496_at	At2g45610
Hypothetical protein	155	4.7	14379_at	At2g33320
Unknown protein	128	2.6	18454_g_at	At4g08605
Unknown protein	75	1.9	15016_at	At1g04090
Putative protein	51	2.1	12076_at	At4g36600
Hypothetical protein	49	2.3	15266_at	At4g08630
Hypothetical protein	45	3.5	15262_at	At2g33190
Putative protein (fragment), (BRCA1-associated RING domain protein, <i>H. sapiens</i> )	32	1.4	17790_g_at	At4g21070



Becker et al.

to be expressed in pollen, but it also led to the identification of several genes not known to be expressed in the male gametophyte so far.

### Cytoskeleton

A fundamental aspect of tip growth of pollen tubes is the continuous deposition of new cell wall and plasma membrane at the tube apex. Vesicles delivering this material are transported by the actin cytoskeleton. In addition to components of the cytoskeleton in pollen tubes, which have been described as such, i.e. the actin genes ACT4/12 (Huang et al., 1996) and profilin PRF4 (Kandasamy et al., 2002), we have identified a previously uncategorized actin-depolymerizing factor-like protein (At4g25590). Furthermore, three potential motor proteins are selectively expressed. The myosins AtVIIIID and AtXID might function in cargo movement along actin filaments (Reddy and Day, 2001), whereas the kinesin-related protein At1g09170 might be involved in movement along microtubules. The already described members of the myosin family, MYA1 to -3 (Kinkema et al., 1994), are not pollen selectively expressed but are pollen-enriched (supplemental Table I). We confirmed that profilin PRF3 (At4g29340) is highly expressed in pollen grains but is otherwise only called Present in roots, which is in disagreement with its reported constitutive, strong expression in all vegetative tissues (Kandasamy et al., 2002).

### Vesicle Trafficking

Exo- and endocytosis are required to release the contents of the transported vesicles and to reincorporate excess membrane material. The syntaxin AtSYP124 (At1g61290) and a homolog of the yeast Sec7p protein (At2g30690) fall into the large group of genes presumably involved in vesicle trafficking. Other potential SNAREs that might be required for vesicle fusion in pollen are encoded by the pollen-enriched genes AtBET12 (At4g14450) and AtVAMP725 (At2g32670). The clathrin family protein At1g05020 is selectively expressed, and a clathrin assembly protein (At1g03050), which is highly enriched in pollen, might have endocytic functions. Furthermore, two putative ARF GTPases (At2g35210 and At2g14490) are selectively expressed.

### Cell Wall Biosynthesis and Regulation

The pollen tube wall is a bipartite structure with an inner sheath of 1,3- $\beta$ -glucan (callose) covered by an outer fibrillar layer of 1,4- $\beta$ -glucan (cellulose) and  $\alpha$ -linked pectic polysaccharides. Recently, NaCslD1 and NaGsl1 were identified as pollen-expressed genes, potentially encoding the major  $\beta$ -glucan polysaccharide synthases in *Nicotiana glauca* pollen tubes (Doblin et al., 2001). Here, we show that AtCslD4 (putative cellulose synthase; At4g38190) and AtGsl2 (callose synthase; At2g13675) are selectively expressed homologs in Arabidopsis pollen, as assumed

in that study. However, AtCslD1 (At2g33100) is also selectively expressed in pollen, and it shows a higher expression than AtCslD4.

Arabidopsis pollen grains express a whole range of cell wall hydrolytic and cell wall-loosening enzymes such as polygalacturonases, pectate lyases, pectin esterases, glycosyl hydrolases, and expansins. The genes encoding these proteins are among those with the highest expression levels in pollen grains in our study, e.g. the putative polygalacturonase At3g07820. Besides their putative roles in modifications of the pollen tube wall, they may be important for the penetration of the stigmatic tissue.

Glycosylphosphatidylinositol (GPI)-anchored proteins are targeted to the cell surface and presumably are involved in remodeling of the extracellular matrix and/or in signaling (Borner et al., 2002). The putative GPI-anchored COBL11 (At4g27110), a member of the recently described COBRA family (Roudier et al., 2002), is selectively expressed in pollen in our study. COBRA is implicated in the regulation of oriented cell expansion in the root (Schindelman et al., 2001), and we propose that COBL11 might fulfill a similar function in pollen tubes. Two other putatively GPI-anchored proteins identified in this study are the arabinogalactan proteins AGP6 (At5g14380) and AGP23 (At3g57690), with the highest expression values of the selectively expressed and the pollen-enriched genes, respectively. Although the exact function of AGPs in pollen tubes and styles remains to be determined, it is interesting that AGP23 encodes an AG-peptide (Schultz et al., 2002) of only 61 amino acids length. If cleaved and thus released from its lipid anchor, it might serve as a diffusible signal molecule.

### Ion Dynamics

Besides calcium fluxes, several studies indicate the involvement of polarized internal gradients and/or external fluxes of protons, potassium, and chloride in pollen tube growth (for review, see Hepler et al., 2001). However, channels and transporters accounting for the observed ion fluxes across the plasma membrane in pollen tubes remain largely unknown. Thus, the identification of ion transporters that are selectively expressed or enriched in pollen could open new avenues (Feijó et al., 2001). The important role of the selectively expressed, inwardly rectifying K<sup>+</sup> channel SPIK (At2g25600) for pollen tube development and competitive ability has been shown by Mouline et al. (2002) and the pollen enriched, outwardly rectifying K<sup>+</sup> channel SKOR (At3g02850) has been characterized extensively in root stelar tissues (Gaymard et al., 1998; Lacombe et al., 2000). Interestingly, SKOR is permeable to both monovalent cations and Ca<sup>2+</sup> and decreasing cytoplasmic pH reduces SKOR-mediated currents. The selectively expressed, putative cyclic-nucleotide-gated ion channel AtCNGC16 (At3g48010) could be involved in the control of [Ca<sup>2+</sup>]<sub>cyt</sub> too, because it has been shown



that an increase in cytoplasmic cAMP or cGMP elicits both a  $\text{Ca}^{2+}$  influx and a rise in  $[\text{Ca}^{2+}]_{\text{cyt}}$  in plant cells (Kurosaki et al., 1994; Volotovski et al., 1998). Furthermore, an overlap of the cyclic nucleotide-binding domain and a calmodulin-binding site in AtCNGC1 and -2 (Kohler and Neuhaus, 2000) suggests a regulation of the channels by cyclic nucleotides and calmodulin. Because it has been demonstrated that cAMP can modulate pollen tube growth and reorientation (Moutinho et al., 2001), AtCNGC16 could constitute a possible link between increases in cAMP concentrations and increases in  $[\text{Ca}^{2+}]_{\text{cyt}}$  in pollen tubes.

The putative cation/ $\text{H}^+$  antiporters AtCHX8, -13, and -16 (At2g28180, At2g13620, and At2g30240) are possible regulators of proton fluxes observed during pollen tube growth (Feijó et al., 1999), although their specificity and membrane localization are still to be determined (Maser et al., 2001). In addition, the identification of a pollen selectively expressed G subunit of the vacuolar type  $\text{H}^+$ -ATPase underlines the possible importance of V-ATPases for proton homeostasis in growing pollen tubes. Recently,  $\text{Cl}^-$  fluxes have been shown to play a role in growth and cell volume regulation in pollen tubes (Zonia et al., 2002). The chloride channel CLC-c (At5g49890) is pollen enriched, and its function in cation homeostasis has been demonstrated by suppression of the Gef1 mutant phenotype in yeast (Gaxiola et al., 1998). Thus, CLC-c might fulfill a similar function in pollen. Furthermore, we identified two tonoplast intrinsic proteins (TIPs), TIP5;1 (At3g47440) and TIP1;3 (At4g01470), as selectively expressed in pollen.

### Signal Transduction

Inhibitor studies with pollen of several plant species (for review, see Mascarenhas, 1975) indicate that pollen germination and early tube growth are largely independent of transcription, but strictly dependent on translation. Thus, signal transduction and translational control might play a more important role than transcriptional control. In fact, 25% of the selectively expressed genes fall under the signal transduction category. Most of these genes (26 of 37) encode putative protein kinases. In addition to the Arabidopsis receptor-like kinase RKF1 (At1g29750; Takahashi et al., 1998), the Leu-rich repeat receptor-like kinases LePRK1-3 and ZmPRK1 were characterized as examples of pollen-specific receptor kinases possibly involved in these signaling events (Muschiatti et al., 1998; Kim et al., 2002). Here, we show that the Arabidopsis homologs AtPRKc (At2g07040) and AtPRKd (At5g35390) are not only expressed in pollen, but that their expression is restricted to pollen. For RKF2 (At1g19090), which was reportedly expressed at low levels in several organs (Takahashi et al., 1998), we found that it was selectively expressed in pollen. Moreover, we identify the putative Pro-rich, extensin-like receptor kinases AtPERK5, -7, -12, and -4 (At4g34440, At1g49270, At1g23540, and

At2g18470) as selectively expressed and pollen enriched, respectively. PERK1 is a canola (*Brassica napus*) homolog of this novel family of plant receptor-like kinases and is rapidly induced by wounding (Silva and Goring, 2002). Sequence similarities of its extracellular domain to extensins might indicate an involvement of the Arabidopsis pollen homologs in a signal transduction pathway, signaling changes in the mechanical properties of the pollen tube cell wall to target proteins in the pollen cytoplasm.

Potential ligands to pollen receptor kinases might be expressed by the pistil or by the pollen itself as shown for LePRK2 and its pollen-expressed ligand LAT52 (Tang et al., 2002). The potential signaling peptide RALF-LIKE 10 (At2g19020; Olsen et al., 2002) is selectively expressed in pollen. Its tobacco (*Nicotiana tabacum*) homolog rapid alkalization factor (RALF) encodes a ubiquitous 115-amino acid prepro-peptide, which is processed into a 5-kD signaling peptide (Pearce et al., 2001). RALF causes a rapid alkalization of tobacco cell cultures and an arrest of root growth and development when supplied to germinating tomato (*Lycopersicon esculentum*) and Arabidopsis seeds. RALF-LIKE 10, encoding a 73-amino acid protein with a potential N-terminal signal peptide for export, might be a putative ligand, e.g. for Leu-rich receptor-like kinases in the plasma membrane of pollen or the pistil. The importance of intracellular signaling for pollen tube growth has been demonstrated by studies covering several types of molecular switches, e.g. Rop/Rac GTPases (Gu et al., 2003) and the MAP kinase kinase kinase AtMAP3K $\gamma$  (Jouannic et al., 1999). We confirm the pollen-enriched expression of the Rho-related GTPases AtRac1 (At2g17800) and AtRac6 (At4g35950) and of AtMAP3K $\gamma$  (At5g66850), as well as the pollen-enriched expression of the G-protein AtRAB2 (At4g17170; Moore et al., 1997). The Rho-related GTPase Rop1At (At3g51300), reported to be specifically expressed in anthers (Li et al., 1998), is highly enriched in pollen but was called Present in roots and siliques. Moreover, our study enlarges this list significantly, including several putative protein kinases, a putative STE20/PAK-like protein kinase (MAP4K-At1g70430), and two putative G-proteins (At2g33870 and At2g22290). Considering the established link between elevation of cytosolic  $\text{Ca}^{2+}$  at the pollen tube tip and its growth (for review, see Franklin-Tong, 1999) our identification of two selectively expressed putative calmodulins (At4g03290 and At4g12860) and the five calcium-dependent protein kinases CPK14, -18, -20, -24, and -26 (At2g41860, At4g36070, At2g38910, At2g31500, and At2g31500) points out six new potential players in  $\text{Ca}^{2+}$ -mediated signaling in Arabidopsis pollen. In addition, several genes presumably involved in phosphoinositide signaling are selectively expressed in pollen (At2g18180, At2g43900, At2g31830, and At2g41210).

## Translational and Transcriptional Regulation

Although only two genes encoding potential regulators of translation were selectively expressed (At3g16380 and At2g39820), there are several more potential regulators of translation showing an enriched expression in pollen (supplemental Table I). Protein turnover in pollen might also contribute to regulation, and therefore the five genes involved in proteolysis are notable. Three percent of the genes that are selectively expressed encode proteins potentially involved in transcriptional regulation. Among these are transcription factor MYB97 (At4g26930) and the MADS-box proteins AGL29 and AGL30 (At2g34440 and At2g03060). AGL30 is a homolog of MADS1;11 of tobacco, which is thought to be a regulator of gene expression during early pollen tube growth (Steiner et al., 2003). Several more potential transcription factors are pollen enriched, including the response regulator ARR2 (At4g16110), as described previously (Lohrmann et al., 2001). The expression of several genes encoding transcription factors in hydrated pollen grains is surprising, because it might indicate that more *de novo* synthesis of RNA takes place in germinating pollen and during tube growth than had been anticipated by early inhibitor studies (for review, see Mascarenhas, 1975).

## Cell Cycle

The cell cycle in the vegetative cell of pollen is believed to be arrested. However the sperm cells continue through the S phase of the cell cycle after pollination and are deposited into the embryo sac with a 2C content of DNA in G2 (Friedman, 1999). According to our results, cyclin A2;1 (At5g25380) is selectively expressed, and cyclin B3;1 (At1g16330) is enriched in pollen with an additional Present call only in roots. Both genes were described recently as showing a peak of expression in M phase of the cell cycle in synchronized cell suspensions of *Arabidopsis* (Menges et al., 2002). Interestingly, the mitotic cyclin gene *CycA1;1* of maize was found to be expressed in isolated sperm and zygotes, although sperm of maize remain in G1 until fertilization (Sauter et al., 1998). We cannot exclude that the cyclin transcripts we have detected are derived from expression in the sperm cells, but whatever their origin, their expression in pollen might be important for fertilization or post-fertilization events rather than for pollen tube growth.

## Stress Response

Stress response-related genes, such as the small heat shock protein gene At-HSP17.6A (At5g12030), represent 7% of the pollen selectively expressed genes. The expression of At-HSP17.6A was shown to be induced by heat and osmotic stress (Sun et al., 2001). Thus, the expression of this gene could be

explained by osmotic stress during desiccation and rehydration of the pollen grain, or the flow-cytometric sorting might have caused the induction of stress response-related genes.

Hypothetical proteins and proteins with unknown function account for 23% of the 150 selectively expressed genes. The high expression levels for some of them indicate their potential importance for the male gametophyte. Furthermore, our data confirm the expression of 10 genes that had hypothetical status so far (i.e. no representation in EST databases existing), which makes them interesting candidates for a functional characterization of their encoded gene products.

## One-Third of Constitutively Expressed Genes in Vegetative Tissues Are Not Expressed in Pollen

Our main goal was to identify enriched or selectively expressed genes in pollen to gain insight into the genetic basis of pollen germination and tube growth, but identifying genes that are specifically down-regulated in pollen is also informative. For this purpose, we used a list of constitutively expressed genes from samples from 5-week-old *Arabidopsis* (leaves, roots, inflorescence stems, and flowers; Zhu et al., 2001). Comparing the analysis of Zhu et al. (2001) with our data yielded 283 genes that showed constitutive expression in vegetative tissues in our study as well. We used the lower confidence bound 1.2 criterion and MAS 5 calls to identify those of the 283 genes that were down-regulated or called Absent in pollen with high confidence, resulting in a list of 104 genes, which were functionally characterized (Table II; supplemental Table II). Thus, 37% of 283 genes constitutively expressed in vegetative tissues are not expressed in pollen grains. The largest set (27%) of transcripts is functionally related to protein biosynthesis; all of the 29 genes in this category encode putative or known ribosomal proteins. Labeling experiments of lily and *Tradescantia* sp. pollen tubes indicated transcriptional inactivation of rRNA genes in immature pollen grains and during pollen tube growth (for review, see Mascarenhas, 1975), and our data is in accordance with these early observations. This supports the view that in many species, the majority of rRNAs and ribosomal proteins as well as tRNAs, mRNAs, and other proteins are already stored in the mature pollen grain to ensure rapid germination and initial tube growth on the stigma (Mascarenhas, 1989). Thus, in most cases, the transcript abundance that we have measured in *Arabidopsis* pollen grains most likely reflects the accumulation and storage of transcripts during earlier stages of pollen development.

The next largest set (26%) of transcripts encodes proteins involved in diverse functions related to metabolism. Surprisingly, the third largest group (8%) consists of genes encoding membrane intrinsic proteins. Besides two TIPs (TIP1;1 and TIP1;2), six of these encode plasma membrane intrinsic proteins

**Table II.** Functional classification of constitutively expressed genes not expressed in pollen

Protein biosynthesis (29)
At2g34480, At1g43170, At4g11050, At2g09990, At4g27090, At1g34030, At1g67430, At1g09690, At1g70600, At2g39460, At4g34670, At2g19730, At4g16720, At3g49010, At2g25210, At3g52580, At1g27400, At4g14320, At2g18020, At3g47370, At4g00100, At2g36160, At1g72370, At2g19750, At4g26230, At2g33370, At2g40510, At2g31610, At4g18730
Metabolism (28)
At2g25450, At1g04410, At3g61440, At2g26670, At4g21960, At5g35630, At2g28000, At5g43940, At2g30860, At2g31570, At3g58610, At4g23860, At5g04590, At2g30490, At1g70310, (12881_s_at), At1g53240, At1g50480, At2g13360, At4g09320, At5g39950, At2g30870, At2g43090, At5g60920, (20709_s_at), At1g65960, At1g75330, At2g36580
Water channel (8)
At2g37170, At3g53420, At3g26520, At3g61430, At4g35100, At2g45960, At4g23400, At2g36830
Signal transduction (6)
At5g10450, At2g42590, At5g55190, At5g42080, At1g02130, At5g38480
Stress response (5)
At1g47128, At4g24190, At1g20620, At1g20440, At2g02130
Transcriptional regulation (5)
At3g16770, At2g32080, At4g40060, At2g33340, (18032_i_at)
Translational regulation (4)
At3g04730, At1g30230, At1g09640, At2g22670
Others
At2g27020, At1g13060, At4g26840 (proteolysis); At5g49720, At4g19410 (cell wall biosynthesis and regulation); At3g06720, At2g14720 (protein transport); At4g30270 (senescence); At5g12250 (cytoskeleton); At1g04400 (phototropism); At5g40890 (ion transport)
Unknown (8)
At2g41430, At3g15353, (12847_at), At1g13930, At4g36040, At2g35810, At4g16520, At1g14910

(PIPs; PIP1;1, PIP1;2, PIP1;5, PIP2;1, PIP2;2, and PIP2;7). This observation prompted us to study the expression levels and calls of the remaining seven annotated Arabidopsis PIPs (Johanson et al., 2001). Probe sets for five of these genes (PIP1;3, PIP1;4, PIP2;5, PIP2;6, and PIP2;8) were found on the version of the Arabidopsis array used in this study. Taking all our available data about PIPs together, we identified PIP1;3 (At1g01620) as the only PIP expressed in pollen. It has been assumed that water fluxes might be linked to the flux of Cl<sup>-</sup> (Zonia et al., 2001). Plant major intrinsic proteins have been reported to be enriched in zones of fast cell division and expansion or in areas where water flow or solute flux density would be expected to be high (for review, see Johanson et al., 2001; Tyerman et al., 2002). Our data indicate the sole expression of PIP1;3 (old name TMP-B) in pollen, although 13 PIPs are annotated in the Arabidopsis genome. Interestingly, it has been shown that PIP1;3 expression is turgor responsive (Shagan et al., 1993). The selectively expressed TIPs, TIP1;3 and TIP5;1, might be involved in cytosolic osmoregulation, too. The subcellular localizations indicated by the TIP and PIP labels should be taken as putative (Barkla et al., 1999) because they are solely based on sequence data in most cases.

## CONCLUSIONS

We have identified transcripts of 1,587 genes in Arabidopsis pollen, of which 30% are pollen enriched and 10% pollen selectively expressed. Thus, our study significantly increases the current knowledge of genes expressed in the male gametophyte of Arabidopsis. The specific down-regulation of otherwise

constitutively expressed genes emphasizes that a particular genetic program underlies the unique growth of pollen tubes. T-DNA insertion lines are available for the majority of the 150 genes selectively expressed in pollen, and their characterization might support the ongoing efforts to combine genetic and physiological evidence into a model for pollen germination and pollen tube growth. We show here that pollen possesses a significantly lower amount of expressed genes compared with other vegetative tissue and yet retains remarkable self-organized regulatory mechanisms of growth. This makes pollen an excellent model for the study of cell growth and morphogenesis on apical growing cells because it seems to be using a “minimal” set of genes encoding a mechanism with obvious evolutionary success.

## MATERIALS AND METHODS

### Plant Material and Growth Conditions

Arabidopsis ecotype Col-0 was used in this study. To minimize interplant variability, tissues from a minimum of 12 plants were pooled for each RNA extraction. For seedling and root, seeds were surface-sterilized and then spread on petri dishes containing B-5 medium (Duchefa, Haarlem, The Netherlands) solidified with 0.8% (w/v) phytagar (Duchefa). The seeds were cold-treated for 3 d at 4°C to ensure uniform germination. The plates were transferred to short-day conditions (8 h of light at 21°C–23°C) and grown in a horizontal (for harvest of seedlings) or vertical position (for harvest of roots). For the seedling samples more than 25 seedlings from five petri dishes were collected after 4 d of growth. For the root samples more than 25 roots from 10 petri dishes were harvested after 13 d.

Plants for the leaf, silique, and pollen samples were grown on soil for 12 weeks in short-day conditions (8 h of light at 21°C–23°C) and then changed to long-day conditions (16 h of light) to induce flowering. After bolting, more than 12 rosette leaves from different plants per leaf sample were collected. Young siliques were harvested 2 weeks later. Old parts of the flower, especially stamens, were removed from the siliques to ensure pollen-

AQ: E



Becker et al.

free silique samples, and more than 25 siliques from different plants were pooled per sample.

## Isolation and FACS Sorting of Pollen Grains

A detailed protocol of the purification of hydrated pollen grains can be found in the supplemental "Materials and Methods". In brief, flower heads were cut and placed in a humid chamber for 2 h. Then the flower heads were agitated three times in 500 mL of pollen-sorting buffer (10 mM CaCl<sub>2</sub>, 1 mM KCl, 2 mM MES, and 5% [w/v] Suc, pH 6.5, with NaOH, in double-distilled water). After consecutive filtration and centrifugation steps, the resulting pellet, highly enriched in pollen, was re-suspended in 10 mL of pollen-sorting buffer. Hydrated pollen grains were separated from non-hydrated and/or destroyed pollen grains and other impurities in a final purification step using FACS based on size and autofluorescence criteria of pollen. Pollen viability was assessed by enzymatically induced fluorescence using fluorescein diacetate according to Heslop-Harrison and Heslop-Harrison (1970).

## RNA Isolation, Target Synthesis, and Hybridization to Affymetrix GeneChips

Total RNA was extracted from the tissue and cell samples, respectively, using the RNeasy Mini Plant Kit (Qiagen, Hilden, Germany). RNA quality was assessed by agarose gel electrophoresis and spectrophotometry. RNA was processed for use on Affymetrix Arabidopsis Genome GeneChip arrays, according to the manufacturer's protocol. In brief, 7 µg of total RNA was used in a reverse transcription reaction (SuperScript II, Invitrogen, Paisley, UK) to generate first-strand cDNA. After second-strand synthesis, double-strand cDNA was used in an in vitro transcription reaction to generate biotinylated cRNA. After purification and fragmentation, 15 µg of cRNA was used in a 300-µL hybridization containing added in vitro transcription controls. Two hundred microliters of mixture was hybridized on arrays for 16 h at 45°C. Standard post hybridization wash and double-stain protocols were used on an Affymetrix GeneChip Fluidics Station 400. Arrays were scanned by using a Gene Array scanner (Hewlett Packard, Palo Alto, CA).

## Data Analysis

Scanned arrays were analyzed first with Affymetrix MAS 5.0 software to obtain Absent/Present calls and to assure that all quality parameters were in the recommended range. For subsequent analysis, dChip 1.3 (<http://www.dchip.org>; Wong laboratory, Harvard) was used. The following conditions were applied to ensure reliability of the analyses (for details, see supplemental "Materials and Methods"): First, each GeneChip experiment was performed with biological replicates and triplicates in the case of pollen, respectively. Second, we used a sample-wise normalization to the median median probe cell intensity of all arrays. Third, normalized median median probe cell intensities of the 11 arrays were used to obtain model-based gene expression indices based on a PM-only model (Li and Hung Wong, 2001). Finally, all genes compared were considered to be differentially expressed if they were called Present in at least one of the arrays and if the 90% lower confidence bound of the -fold change between experiment and baseline was above 1.2.

To achieve a higher stringency for the identification of constitutively expressed genes in the vegetative tissues, we combined our data with a set identified by Zhu et al. (2001). Because a proprietary pre-version of the Arabidopsis Genome GeneChip array was used by these authors, 81 of the 346 probe sets they identified as constitutively expressed were not included in our study, resulting in an overlap of 283 probe sets.

## Gene Annotation

For gene annotation, we used the updated The Arabidopsis Information Resource annotation (October 2002 release) for the Arabidopsis Genome GeneChip array (<http://www.arabidopsis.org>). Genes were classified into functional categories using the Gene Ontology information available from The Arabidopsis Information Resource as of October 2002. Genes represented by two or more probe sets on the array were analyzed manually, and only the most significant probe set for this gene was included in the final tables.

## ACKNOWLEDGMENTS

We thank Philip Benfey (New York University [now at Duke University, Durham, NC]) for initial support and critical reading of the manuscript, Pedro Coutinho (Instituto Gulbenkian de Ciência) for his help to update the gene annotation, and Sheila McCormick (Plant Gene Expression Center, XXX, XX) for critical reading of the manuscript.

Received June 7, 2003; returned for revision July 7, 2003; accepted July 9, 2003.

## LITERATURE CITED

- Arabidopsis Genome Initiative (2000) Analysis of the genome sequence of the flowering plant *Arabidopsis thaliana*. *Nature* **408**: 796–815
- Barkla BJ, Vera-Estrella R, Pantoja O, Kirch HH, Bohnert HJ (1999) Aquaporin localization: how valid are the TIP and PIP labels? *Trends Plant Sci* **4**: 86–88
- Borner GH, Sherrier DJ, Stevens TJ, Arkin IT, Dupree P (2002) Prediction of glycosylphosphatidylinositol-anchored proteins in Arabidopsis: a genomic analysis. *Plant Physiol* **129**: 486–499
- Che P, Gingerich DJ, Lall S, Howell SH (2002) Global and hormone-induced gene expression changes during shoot development in *Arabidopsis*. *Plant Cell* **14**: 2771–2785
- Derksen J, Knuiman B, Hoedemaekers K, Guyon A, Bonhomme S, Pierson ES (2002) Growth and cellular organization of *Arabidopsis* pollen tubes *in vitro*. *Sex Plant Reprod* **15**: 133–139
- Doblin MS, De Melis L, Newbigin E, Bacic A, Read SM (2001) Pollen tubes of *Nicotiana glauca* express two genes from different  $\beta$ -glucan synthase families. *Plant Physiol* **125**: 2040–2052
- Eisen MB, Spellman PT, Brown PO, Botstein D (1998) Cluster analysis and display of genome-wide expression patterns. *Proc Natl Acad Sci USA* **95**: 14863–14868
- Fan LM, Wang YF, Wang H, Wu WH (2001) In vitro *Arabidopsis* pollen germination and characterization of the inward potassium currents in Arabidopsis pollen grain protoplasts. *J Exp Bot* **52**: 1603–1614
- Feijó JA, Sainhas J, Hackett GR, Kunkel JG, Hepler PK (1999) Growing pollen tubes possess a constitutive alkaline band in the clear zone and a growth-dependent acidic tip. *J Cell Biol* **144**: 483–496
- Feijó JA, Sainhas J, Holdaway-Clarke T, Cordeiro MS, Kunkel JG, Hepler PK (2001) Cellular oscillations and the regulation of growth: the pollen tube paradigm. *Bioessays* **23**: 86–94
- Franklin-Tong VE (1999) Signaling and modulation of pollen tube growth. *Plant Cell* **11**: 727–738
- Friedman WE (1999) Expression of the cell cycle in sperm of *Arabidopsis*: implications for understanding patterns of gametogenesis and fertilization in plants and other eukaryotes. *Development* **126**: 1065–1075
- Gaxiola RA, Yuan DS, Klausner RD, Fink GR (1998) The yeast CLC chloride channel functions in cation homeostasis. *Proc Natl Acad Sci USA* **95**: 4046–4050
- Gaymard F, Pilot G, Lacombe B, Bouchez D, Bruneau D, Bouchez J, Michaux-Ferriere N, Thibaud JB, Sentenac H (1998) Identification and disruption of a plant shaker-like outward channel involved in K<sup>+</sup> release into the xylem sap. *Cell* **94**: 647–655
- Golub TR, Slonim DK, Tamayo P, Huard C, Gaasenbeek M, Mesirov JP, Coller H, Loh ML, Downing JR, Caligiuri MA et al. (1999) Molecular classification of cancer: class discovery and class prediction by gene expression monitoring. *Science* **286**: 531–537
- Gu Y, Vernoud V, Fu Y, Yang Z (2003) ROP GTPase regulation of pollen tube growth through the dynamics of tip-localized F-actin. *J Exp Bot* **54**: 93–101
- Guyon VN, Astwood JD, Garner EC, Dunker AK, Taylor LP (2000) Isolation and characterization of cDNAs expressed in the early stages of flavonol-induced pollen germination in petunia. *Plant Physiol* **123**: 699–710
- Harmer SL, Hogenesch JB, Straume M, Chang HS, Han B, Zhu T, Wang X, Kreps JA, Kay SA (2000) Orchestrated transcription of key pathways in *Arabidopsis* by the circadian clock. *Science* **290**: 2110–2113
- Hepler PK, Vidali L, Cheung AY (2001) Polarized cell growth in higher plants. *Annu Rev Cell Dev Biol* **17**: 159–187
- Heslop-Harrison J, Heslop-Harrison Y (1970) Evaluation of pollen viability by enzymatically induced fluorescence; intracellular hydrolysis of fluorescein diacetate. *Stain Technol* **45**: 115–120

AQ: J

AQ: F

AQ: G  
AQ: H  
AQ: I



- Huang SR, An YQ, McDowell JM, McKinney EC, Meagher RB (1996) The *Arabidopsis thaliana* ACT4/ACT12 actin gene subclass is strongly expressed throughout pollen development. *Plant J* 10: 189–202
- Johanson U, Karlsson M, Johansson I, Gustavsson S, Sjovald S, Frayse L, Weig AR, Kjellbom P (2001) The complete set of genes encoding major intrinsic proteins in *Arabidopsis* provides a framework for a new nomenclature for major intrinsic proteins in plants. *Plant Physiol* 126: 1358–1369
- Jouannic S, Hamal A, Leprince AS, Tregear JW, Kreis M, Henry Y (1999) Characterisation of novel plant genes encoding MEKK/STE11 and RAF-related protein kinases. *Gene* 229: 171–181
- Kandasamy MK, McKinney EC, Meagher RB (2002) Plant profilin isoforms are distinctly regulated in vegetative and reproductive tissues. *Cell Motil Cytoskeleton* 52: 22–32
- Kim HU, Cotter R, Johnson S, Senda M, Dodds P, Kulikauska R, Tang W, Ezcurea I, Herzmark P, McCormick S (2002) New pollen-specific receptor kinases identified in tomato, maize and *Arabidopsis*: The tomato kinases show overlapping but distinct localization patterns on pollen tubes. *Plant Mol Biol* 50: 1–16
- Kinkema M, Wang HY, Schiefelbein J (1994) Molecular analysis of the myosin gene family in *Arabidopsis thaliana*. *Plant Mol Biol* 26: 1139–1153
- Kohler C, Neuhaus G (2000) Characterisation of calmodulin binding to cyclic nucleotide-gated ion channels from *Arabidopsis thaliana*. *FEBS Lett* 471: 133–136
- Kurosaki F, Kaburaki H, Nishi A (1994) Involvement of plasma membrane-located calmodulin in the response decay of cyclic nucleotide-gated cation channel of cultured carrot cells. *FEBS Lett* 340: 193–196
- Lacombe B, Pilot G, Gaymard F, Sentenac H, Thibaud JB (2000) pH control of the plant outwardly-rectifying potassium channel SKOR. *FEBS Lett* 466: 351–354
- Lennon KA, Lord EM (2000) In vivo pollen tube cell of *Arabidopsis thaliana*: I. Tube cell cytoplasm and wall. *Protoplasma* 214: 45–56
- Li C, Hung Wong W (2001) Model-based analysis of oligonucleotide arrays: model validation, design issues and standard error application. *Genome Biol* 2: RESEARCH0032
- Li H, Wu G, Ware D, Davis KR, Yang Z (1998) *Arabidopsis* Rho-related GTPases: differential gene expression in pollen and polar localization in fission yeast. *Plant Physiol* 118: 407–417
- Lohrmann J, Sweere U, Zabaleta E, Baurle I, Keitel C, Kozma-Bognar L, Brennicke A, Schafer E, Kudla J, Harter K (2001) The response regulator ARR2: a pollen-specific transcription factor involved in the expression of nuclear genes for components of mitochondrial complex I in *Arabidopsis*. *Mol Genet Genomics* 265: 2–13
- Lord EM, Russell SD (2002) The mechanisms of pollination and fertilization in plants. *Annu Rev Cell Dev Biol* 18: 81–105
- Mascarenhas JP (1975) The biochemistry of angiosperm pollen development. *Bot Rev* 41: 259–314
- Mascarenhas JP (1989) The male gametophyte of flowering plants. *Plant Cell* 1: 657–664
- Maser P, Thomine S, Schroeder JI, Ward JM, Hirschi K, Sze H, Talke IN, Amtmann A, Maathuis FJ, Sanders D et al. (2001) Phylogenetic relationships within cation transporter families of *Arabidopsis*. *Plant Physiol* 126: 1646–1667
- McCormick S (1993) Male gametophyte development. *Plant Cell* 5: 1265–1275
- Menges M, Hennig L, Gruißem W, Murray JA (2002) Cell cycle regulated gene expression in *Arabidopsis*. *J Biol Chem* 277: 41987–42002
- Moore I, Diefenthal T, Zarsky V, Schell J, Palme K (1997) A homolog of the mammalian GTPase Rab2 is present in *Arabidopsis* and is expressed predominantly in pollen grains and seedlings. *Proc Natl Acad Sci USA* 94: 762–767
- Mouline K, Very AA, Gaymard F, Boucherez J, Pilot G, Devic M, Bouchez D, Thibaud JB, Sentenac H (2002) Pollen tube development and competitive ability are impaired by disruption of a Shaker K(+) channel in *Arabidopsis*. *Genes Dev* 16: 339–350
- Moutinho A, Hussey PJ, Trewavas AJ, Malho R (2001) cAMP acts as a second messenger in pollen tube growth and reorientation. *Proc Natl Acad Sci USA* 98: 10481–10486
- Muschietti J, Eyal Y, McCormick S (1998) Pollen tube localization implies a role in pollen-pistil interactions for the tomato receptor-like protein kinases LePRK1 and LePRK2. *Plant Cell* 10: 319–330
- Olsen AN, Mundy J, Skriver K (2002) Peptomics, identification of novel cationic *Arabidopsis* peptides with conserved sequence motifs. *In Silico Biol* 2: 0039
- Pearce G, Moura DS, Stratmann J, Ryan CA Jr (2001) RALF, a 5-kDa ubiquitous polypeptide in plants, arrests root growth and development. *Proc Natl Acad Sci USA* 98: 12843–12847
- Pederson S, Simonsen V, Loeschcke V (1987) Overlap of gametophytic and sporophytic gene expression in barley. *Theor Appl Genet* 75: 200–206
- Preuss D (1995) Being fruitful: genetics of reproduction in *Arabidopsis*. *Trends Genet* 11: 147–153
- Reddy AS, Day IS (2001) Analysis of the myosins encoded in the recently completed *Arabidopsis thaliana* genome sequence. *Genome Biol* 2: RESEARCH0024
- Roudier F, Schindelman G, DeSalle R, Benfey PN (2002) The COBRA family of putative GPI-anchored proteins in *Arabidopsis*: a new fellowship in expansion. *Plant Physiol* 130: 538–548
- Sauter M, von Wieggen P, Lorz H, Kranz E (1998) Cell cycle regulatory genes from maize are differentially controlled during fertilization and first embryonic cell division. *Sex Plant Reprod* 11: 41–48
- Schindelman G, Morikami A, Jung J, Baskin TI, Carpita NC, Derbyshire P, McCann MC, Benfey PN (2001) COBRA encodes a putative GPI-anchored protein, which is polarly localized and necessary for oriented cell expansion in *Arabidopsis*. *Genes Dev* 15: 1115–1127
- Schultz CJ, Rumsewicz MP, Johnson KL, Jones BJ, Gaspar YM, Bacic A (2002) Using genomic resources to guide research directions: the arabinogalactan protein gene family as a test case. *Plant Physiol* 129: 1448–1463
- Scott R, Hodge R, Paul W, Draper J (1991) The molecular biology of anther differentiation. *Plant Sci* 80: 167–191
- Shagan T, Meraro D, Bar-Zvi D (1993) Nucleotide sequence of an *Arabidopsis thaliana* turgor-responsive TMP-B cDNA clone encoding transmembrane protein with a major intrinsic protein motif. *Plant Physiol* 102: 689–690
- Silva NF, Goring DR (2002) The proline-rich, extensin-like receptor kinase-1 (PERK1) gene is rapidly induced by wounding. *Plant Mol Biol* 50: 667–685
- Steiner C, Bauer J, Amrhein N, Bucher M (2003) Two novel genes are differentially expressed during early germination of the male gametophyte of *Nicotiana tabacum*. *Biochim Biophys Acta* 1625: 123–133
- Stinson JR, Eisenberg AJ, Willing RP, Pe ME, Hanson DD, Mascarenhas JP (1987) Genes expressed in the male gametophyte of flowering plants and their isolation. *Plant Physiol* 83: 442–447
- Sun W, Bernard C, Van de Cotte B, Van Montagu M, Verbruggen N (2001) At-HSP17.6A, encoding a small heat-shock protein in *Arabidopsis*, can enhance osmotolerance upon overexpression. *Plant J* 27: 407–415
- Takahashi T, Mu JH, Gasch A, Chua NH (1998) Identification by PCR of receptor-like protein kinases from *Arabidopsis* flowers. *Plant Mol Biol* 37: 587–596
- Tang W, Ezcurea I, Muschietti J, McCormick S (2002) A cysteine-rich extracellular protein, LAT52, interacts with the extracellular domain of the pollen receptor kinase LePRK2. *Plant Cell* 14: 2277–2287
- Taylor LP, Hepler PK (1997) Pollen germination and tube growth. *Annu Rev Plant Physiol Plant Mol Biol* 48: 461–491
- Twell D (2002) The developmental biology of pollen. In JA Roberts, ed, *Plant Reproduction*, Vol 6. Sheffield Academic Press, Sheffield, UK, pp 86–153
- Tyerman SD, Niemietz CM, Bramley H (2002) Plant aquaporins: multifunctional water and solute channels with expanding roles. *Plant Cell Environ* 25: 173–194
- Volotovskii ID, Sokolovsky SG, Molchan OV, Knight MR (1998) Second messengers mediate increases in cytosolic calcium in tobacco protoplasts. *Plant Physiol* 117: 1023–1030
- Willing RP, Bashe D, Mascarenhas JP (1988) An analysis of the quantity and diversity of messenger RNAs from pollen and shoots of *Zea mays*. *Theor Appl Genet* 75: 751–753
- Wixon J (2001) *Arabidopsis thaliana*. *Comp Funct Genom* 2: 91–98
- Zhu T, Budworth P, Han B, Brown D, Chang HS, Zou GZ, Wang X (2001) Toward elucidating the global gene expression patterns of developing *Arabidopsis*: parallel analysis of 8,300 genes by a high-density oligonucleotide probe array. *Plant Physiol Biochem* 39: 221–242
- Zhu T, Wang X (2000) Large-scale profiling of the *Arabidopsis* transcriptome. *Plant Physiol* 124: 1472–1476
- Zonia L, Cordeiro S, Feijó JA (2001) Ion dynamics and hydrodynamics in the regulation of pollen tube growth. *Sex Plant Reprod* 14: 111–116
- Zonia L, Cordeiro S, Tupy J, Feijó JA (2002) Oscillatory chloride efflux at the pollen tube apex has a role in growth and cell volume regulation and is targeted by inositol 3,4,5,6-tetrakisphosphate. *Plant Cell* 14: 2233–2249

AQ: L

AQ: K