

1 **STR-based genetic structure of the Berber population of Bejaia (Northern Algeria) and**
2 **its relationships to various ethnic groups**

3

4

5 Nadir AMIR ^a, Mohamed SAHNOUNE ^b, Lounes CHIKHI ^{cde} and Djebbar ATMANI ^{a*}

6

7

8

9 ^aLaboratoire de Biochimie Appliquée, Faculté des Sciences de la Nature et de la Vie,
10 Université de Bejaia, 06000 Algeria.

11 ^bLaboratoire d'Ecologie et Environnement, Faculté des Sciences de la Nature et de la Vie,
12 Université de Bejaia, 06000 Algeria

13 ^cCNRS, Université Paul Sabatier, ENFA; UMR5174 EDB (Laboratoire Évolution & Diversité
14 Biologique) ; 118, Route de Narbonne, F-31062 Toulouse, France.

15 ^dUniversité Toulouse 3 Paul Sabatier, CNRS ; UMR5174 EDB, F-31062 Toulouse, France.

16 ^eInstituto Gulbenkian de Ciência, Rua da Quinta Grande, n°6, 2780-56, Oeiras, Portugal.

17

18 *Corresponding author: DjebbarAtmani

19 Laboratoire de Biochimie Appliquée, Faculté des Sciences de la Nature et de la Vie,
20 Université de Bejaia, 06000 Algeria.

21 E-mail: djatmani@yahoo.com

22 Tel/Fax : +21334214762

23

24

25

26

27

28

29

30

31

32

33

34

35Abstract

36Patterns of genetic variation in human populations have been described for decades. However,
37North Africa has received little attention and Algeria, in particular, is poorly studied, Here we
38genotyped a Berber-speaking population from Algeria using 15 short tandem repeat (STR)
39loci D8S1179, D21S11, D7S820, CSF1PO, D3S1358, TH01, D13S317, D16S539, D2S1338,
40D19S433, vWA, TPOX, D18S51, D5S818 and FGA from the commercially available
41AmpF/STR Identifiler kit. Altogether 150 unrelated North Algerian individuals were sampled
42across 10 administrative regions or towns from the Bejaia Wilaya (administrative district). We
43found that all of the STR loci met Hardy–Weinberg equilibrium expectations, after Bonferroni
44correction and that the Berber-speaking population of Bejaia presented a high level of
45observed heterozygosity for the 15 STR system (>0.7). Genetic parameters of forensic interest
46such as combined power of discrimination (PD) and combined probability of exclusion (PE)
47showed values higher than 0.999, suggesting that this set of STRs can be used for forensic
48studies. Our results were also compared to those published for 42 other human populations
49analyzed with the same set. We found that the Bejaia sample clustered with several North
50African populations but that some geographically close populations, including the Berber-
51speaking Mozabite from Algeria were closer to Near-Eastern populations. While we were able
52to detect some genetic structure among samples, we found that it was not correlated to
53language (Berber-speaking versus Arab-speaking) or to geography (east versus west). In other
54words, no significant genetic differences were found between the Berber-speaking and the
55Arab-speaking populations of North Africa. The genetic closeness of European, North African
56and Near-Eastern populations suggest that North Africa should be integrated in models aiming
57at reconstructing the demographic history of Europe. Similarly, the genetic proximity with
58sub-Saharan Africa is a reminder of the links that connect all African regions.

59

60**Keywords:** STR diversity, Forensics, Berber/Arab-speaking populations, North Africa,
61Continuity.

631. Introduction

64Global patterns of genetic diversity are becoming increasingly important to reconstruct the
65demographic history of human populations. While some regions have received significant
66attention, others, like North Africa, have been generally less sampled and less studied. This is
67the case for Algeria despite its geographical position linking the Mediterranean area and Sub-
68Saharan Africa. Today the Algerian population is composed of two main linguistic groups, the
69Berber- and the Arab-speaking populations, and it is usually considered that the majority of
70the Algerians descend from Berbers and Arabs (Taïeb, 2004). However, the history of Algeria
71and North Africa is rather complex. For instance, the Berber-speaking region of Bejaia has
72witnessed many successive invasions and conquests that caused important cultural, linguistic
73and religious reshuffles among which the most important is probably the Arab conquest that
74started in the seventh century. Chronologically, the region was submitted to the influence of
75the Romans (33 BC), the Vandals (429 AC), the Byzantines (533 AC), the Arabs (647 AC),
76the Spanish (1510 AC), the Ottomans (1555 AC) and the French (1832 AC) (Cote, 1991;
77Laporte, 2004). In addition to these migrations, there have been internal reshuffles, with the
78introduction of Jewish and sub-Saharan African populations. At the fall of Andalusia (1610
79AC), many of its expelled citizens came to establish settlements in Bejaia (see Gaid, 2008).
80Thus, while Berbers are likely to be the most ancient inhabitants of the region, gene flow,
81immigration and language switching may have obscured the relationships between
82neighboring or distant populations. Genetic data could therefore be useful to identify
83connections between populations speaking different languages today within Algeria or at a
84wider geographical scale. For instance, Henn et al. (2012), using genomic data, estimated that
85the North African populations are likely of Berber origin with substantial shared ancestry with
86the Near East and, to a lesser extent, eastern and western sub-Saharan Africa and Europe.

87 Though the number of studies on North Africa is relatively limited, there have been important
88 studies using various markers that have contributed to the anthropogenetic characterization of
89 North African Berber populations. These studies have focused on the GM immunoglobulin
90 allotypic system (Dugoujon et al., 2004; Coudray et al., 2004; Coudray et al., 2006), others on
91 mitochondrial DNA (Fadhlaoui-Zid et al., 2004; Ennafaa et al., 2009, Coudray et al., 2009),
92 the Y chromosome (Arredi et al., 2004), autosomal microsatellites (STR) (Bosch et al., 2000;
93 Bosch et al., 2001; Coudray et al., 2006; Coudray et al., 2007a; khodjet-el-khil et al., 2008, El
94 Ossmani, 2010, Khodjet-El-Khil et al., 2012, Gaibar et al., 2012), SNP (Henn et al., 2012),
95 and Alu Sequences (Gonzalez-Pérez et al., 2003). Very few studies have been carried out on
96 Algerian Berber populations (Bosch et al., 2001; Achilli et al., 2005; Lefevre-Witier et al.,
97 2006; Coudray et al., 2009; Pereira et al., 2010, Bekada et al., 2013).

98 The present study is part of a wider project on the anthropogenetic characterization of
99 Algerian populations. In this paper we used 15 independent autosomal STR loci to genotype
100 a sample of 150 individuals from the Berber-speaking population of the Bejaia wilaya to
101 provide data on allele frequencies distribution and forensic parameters. The allele frequencies
102 were exploited, using multidimensional scaling (MDS) and **tree analysis (UPGMA)**, to assess
103 the relationships between the Bejaia population and 42 other populations from North Africa,
104 Sub-Saharan Africa, the Middle-East, Europe, Asia and South America. **Analysis of molecular**
105 **variance (AMOVA) was performed to assess the genetic structure of 17 populations**
106 **(including Bejaia). A STRUCTURE analysis was also conducted.**

1072. **Materials and methods**

1082.1. **Population**

109 Buccal swab samples were collected from unrelated healthy Berber-speaking donors (n=150
110 individuals, 300 gametes) from the Bejaia area in North Algeria (Fig.1), after written
111 informed consent was obtained. Donors provided genealogical information for at least three

112previous generations. Samples were collected in accordance with the ethical guidelines
113specified by the institutions involved in this study.

114

Figure 1

1152.2. DNA extraction and amplification

116Genomic DNA extraction was performed on the saliva samples with the QIAamp DNA Mini
117Kit (Qiagen, Hilden, Germany) according to the manufacturer's recommendations. Fifteen
118independent autosomal tetranucleotide STR loci (see Butler, 2006), namely D13S317,
119D16S539, D2S1338, vWA, TPOX, D18S51, D5S818, FGA, D8S1179, D21S11, D7S820,
120D19S433, CSF1PO, TH01 and D3S1358, were coamplified in a multiplex PCR amplification
121reaction. Amplification was performed in a GeneAmp PCR system 9700 (Applied
122Biosystems, Foster City, CA) using the AmpF/STR Identifiler PCR Amplification Kit
123(Applied Biosystems, Foster City, CA) according to the manufacturer's specifications
124(Applied Biosystems, 1998).

1252.3. Electrophoresis and genotyping

126DNA fragments were separated by multi-capillary electrophoresis on an ABI Prism 3130xl
127Genetic Analyzer using the ABI GeneScan 500 LIZ internal size standard as a basis for
128comparison. Fragment sizes were obtained using the software GeneMapper® v3.2 (Applied
129Biosystems, Foster City, CA) and alleles were identified by comparison to an allelic ladder
130supplied by the manufacturer (Applied Biosystems, Foster City, CA).

1312.4. RelPair analysis

132 To detect intra-population pairs of close relatives, we used the program RelPair Version
1332.01 (Epstein et al., 2000). Each population was separately analyzed following the suggested
134settings of Pemberton et al. (2013), namely with a critical value set to 100 and a genotyping
135error rate of 0.008. When related individuals were identified, one of them was discarded from
136the analysis. In order to minimize the number of individuals removed, we preferentially
137omitted the individuals present in two or more related pairs while favoring those with less

138missing data. We applied this analysis to all the populations for which we managed to obtain
139genotype data (See table 1 for populations' codes). The number of individuals retained out of
140the initial number for each population is 116/150 (BJ), 40/44 (MB), 46/48 (SM), 94/105 (AN),
14190/104 (BH), 86/98 (SW), 87/99 (MA), 86/100 (CA), 73/80 (AM), 57/63 (BM), 57/59 (SH).

1422.5. **Statistical and phylogenetic analysis**

143Allele frequencies, expected (H_e) and observed (H_o) heterozygosity (Nei, 1987) and the exact
144test of Hardy-Weinberg equilibrium (Levene, 1949; Guo and Thompson, 1992) were
145computed using the Arlequin Software Version 3.5.1.2 (Excoffier and Lischer, 2010). The
146forensic parameters (matching probability (MP), power of discrimination (PD), polymorphism
147information content (PIC), probability of exclusion (PE) and typical paternity index (TPI))
148were calculated using Powerstats Version 1.2
149(<http://www.promega.com/geneticidtools/powerstats/>).

150The expected number of genotypes was computed as $N_g = \Pi(k_j^2 + k_j)/2$ and the number of
151pairwise haplotype allele associations as $N_a = [(\sum k_j)^2 - \sum k_j^2]/2$ (where k is the number of
152alleles at a considered locus and j the allele index). Bonferroni correction (Weir, 1996) was
153applied to adjust P values in Hardy-Weinberg assumptions ($P = 0.05/15 = 0.0033$ where 15 is
154the number of loci).

155In order to determine the genetic relationship of our sample with other ethnic groups, we
156compared it to 42 populations from Europe, Asia, America and Africa using homologous
157microsatellite loci (Table 1). Pairwise uncorrected F_{st} distances between the 43 populations
158were used to perform a standard non-metric MDS using Statistica 8.0 (StatSoft, 2008) and
159infer a UPGMA tree using POPTREE2 (Takezaki et al., 2010) available
160at:<http://www.med.kagawa-u.ac.jp/~genomelb/takezaki/poptree2/index.html>. Tree robustness
161was evaluated using Bootstrap tests on 1000 permutations (Felsenstein, 1985). UPGMA rather
162than NJ method was used because it was more bootstrap-supported than the NJ one. Note that
163the trees were simply used as a graphical representation of the genetic distances computed.

164 They cannot be seen as a reliable representation of the relationships between populations due
165 to the fact that such trees ignore the existence of gene flow, which is a crucial feature of
166 human populations (Barbujani and Chikhi, 2007).

167 The MDS and Tree analyses were performed on all the 15 loci (including those with missing
168 data) as well as after removing those with missing data (i.e. D16S539, D2S1338 and
169 D19S433).

170 The significance of discriminance between groups in the MDS plot was determined using
171 one-way ANOVA followed by unequal HSD (Honestly Significant Difference) test as
172 implemented in Statistica 8.0 (StatSoft, 2008). The homogeneity of variances was checked
173 using Levene's and Cochran's tests. When required, equality of variances was achieved by
174 dividing data by the standard deviation values and comparing the standardized data.

175 Locus-by-locus allele frequency based AMOVA was performed using Arlequin v.3.5.1.2:
176 Three plans of grouping were tested: (1) Grouping in relation to spoken language (Group 1 =
177 Arab-speaking populations (RB, DM, MA, CA, AM, SH) ; Group 2 = Berber-speaking
178 populations (BJ, MB, SM, AN, BH, AZ, KM, TN, LY, SW, BM)); (2) Grouping in
179 accordance to geographical distribution (Group 1 = Western North African populations (SM,
180 AN, BH, RB, AZ, KM, DM, AM, BM, SH); Group 2 = Central North African populations
181 (BJ, MB, TN, LY); Group 3 = Eastern North African populations (SW, MA, CA); (3)
182 Grouping in relation to UPGMA clustering (Group 1 = BJ, AZ, RB, LY, AN, BH, KM; Group
183 2 = MA, DM, SW, CA, Group 3 = MB, SM, SH, AM, BM; Group 4 = TN) (see Tab. 1 for
184 population codes). **An analysis of population structure was also carried out using the**
185 **STRUCTURE software (Pritchard et al., 2000) but since our data were uninformative and did**
186 **not lead to any clearly identifiable genetic clusters, the results are presented as supplementary**
187 **material.**

188The correlations between genetic (uncorrected F_{st} s) and geographical distances and between
189initial and final MDS distances were evaluated using Mantel test (Mantel, 1967; Smouse et
190al., 1992), and the fixation indices were tested using the permutation procedure (1000
191iterations), as implemented in Arlequin 3.5.1.2.

192

Table 1

1933. Results

194Observed heterozygosity (H_o), expected heterozygosity (H_e) and Hardy-Weinberg
195equilibrium tests (P_h) estimated on the 116 individuals of the Bejaia population, are given in
196Table 2 (See Supplementary Table 1 for allele frequencies). Altogether, for the 15 loci, there
197were a total of 140 alleles among which 56 (40%) were rare (frequency < 0.05). The highest
198observed frequencies were 43.1% for TPOX (allele 8), 37.93% for D5S818 (allele 12),
19934.05% for D7S820 (allele 10), 32.75% and 31.462% for D13S317 (alleles 11 and 12,
200respectively), 30.17% for both CSF1PO (alleles 11) and D19S433 (allele 13). Each of the 15
201loci showed a high level of polymorphism as expressed by the numbers and frequencies (<
2020.95) of alleles per locus. The number of alleles per locus varied from 6 to 15 alleles with a
203mean of 9.33 ± 2.58 . The observed heterozygosities varied from 0.84 (D2S1338 and FGA) to
2040.66 (TPOX) with a mean value of 0.77 ± 0.05 . No significant departure from Hardy-Weinberg
205equilibrium after Bonferroni correction ($p < 0.0033$) was detected for all loci.

206The power of discrimination (PD), the probability of excluding paternity (PE) and the
207polymorphic information content (PIC) are displayed in Table 2. PD ranged from 0.911
208(D19S433) to 0.975 (D2S1338), PE from 0.388 (TPOX) to 0.624 (D2S1338 and FGA) and
209 PIC from 0.740 (D19S433) to 0.900 (D2S1338 and FGA). All the 15 STR loci were highly
210polymorphic ($PIC > 0.7$). The combined power of discrimination and combined probability of
211exclusion showed values higher than 0.999. With a $PIC > 0.8$, seven of the fifteen loci
212(D21S11, D7S820, D3S1358, D2S1338, VWA, D18S51 and FGA) can be considered as very

213informative for genetic personal identification (with a combined PE=0,987). For the 15 loci (6
214to 15 alleles per locus) the computed number of possible genotypes was 7.11×10^{24} .

215

Table 2

216The standard non-metric multidimensional scaling (MDS) based on *Fst* distances (15 loci)
217split the 17 North African populations including Bejaia (Table 1) into two main groups
218significantly discriminated (Fig. 2 A), one (*North Africa 1*, of 11 populations including
219Bejaia) close to the European populations, and the other (*North Africa 2*, of 5 populations),
220close to the Arabian Peninsula populations. As to Tunisia (TN), it behaved as outgroup to all
221other groups (Fig. 2 A). The population of Bejaia (BJ) went with the *North Africa 1*
222populations. Its closest neighbors are Lybia (LY), Rabat (RB) and Azrou (AZ) of Morocco, all
223geographically close to Algeria (Fig. 2 A and B). The North African populations appeared to
224be the most heterogeneous in comparison with other regions groups included in this analysis
225(Fig. 2 A and B). Another MDS (Not shown) based on 12 loci (after removing those
226containing missing data, namely D16S539, D2S1338 and D19S433) gave roughly same
227results as above except that only MB (Mozabites) and CA (Copt Adaima) went with the
228Arabian peninsula populations.

229

Fig. 2

230The UPGMA tree inferred using the *Fst* distances between the 43 populations (Fig. 3) was
231congruent with the MDS results (Fig. 2) and exhibited higher bootstrap values than the
232neighbor-joining (NJ) tree (not shown,). However, most of the deepest nodes showed very
233low bootstrap values connecting reasonably well supported clusters. This tree emphasized the
234heterogeneity revealed by the MDS for the North African populations but the low bootstrap
235values suggest that caution is required in interpretation. The group containing BJ, AZ, RB, LY,
236AN, BH and KM appears as a sister cluster of that regrouping European and the three Middle
237East populations (LB, IR and IQ) but many topologies could explain the data and this mostly
238suggests that connections exist between all these populations. Four other populations, MA,

239DM, SW and CA, exhibit longer branch but no clear clustering. As to MB, SM, SH, AM and
240BM, they formed a sister cluster to that constituted by the Arabian Peninsula populations (DB,
241OM, SA and YE), but again the low bootstrap values suggest caution. TN still behaved as an
242out-group to all other North African populations, hence confirming its isolated position in the
243MDS plot.

244

Fig. 3

245Locus-by-locus AMOVA revealed no significant difference between the Berber- and Arab-
246speaking groups, with *F_{ct}* values that varied from -0.00246 to 0.00104 and percentages of
247variation from 0.02 to 0.25%. Similarly, the three geographical groups did not exhibit strong
248signals of differentiation with *F_{ct}* values that varied from -0.00097 to 0.00454 and
249percentages of variation from 0.01 to 0.45%. Significant differences between these
250geographical groups were revealed only for D8S1197 and TH01 and no significant correlation
251($R = -0.089$, $P = 0.68$) between genetic distances (*F_{st}*) and geographical distances was
252detected by the Mantel test. Groupings defined according to the UPGMA tree (Fig. 3) showed
253similarly low *F_{ct}* values and percentages of variation of -0.00062-0.00689 and 0.01-0.69%
254respectively. For this grouping, significant differences were observed for 8 out of 15 loci
255(Table 3): D8S1197, D7S820, CSF1PO, D3S1358, TH01, D16S539, TPOX and FGA.
256However, for all the three plans of grouping (spoken language, geographical location and
257cluster affiliation), AMOVA revealed highly significant differences between populations
258within each the groups and for all loci (Supplementary Table 2).

259

Table 3

260The *F_{st}* distances between the 17 North African populations (supplementary table 3) suggest
261that the closest populations to Bejaia were AZ and AN (Berber- speaking populations from
262Morocco) with $F_{st} = 0.005$ and $F_{st} = 0.006$, respectively; whereas the most distant
263populations from Bejaia were MB (Berber-speaking from Central South Algeria) and TN

264(Berber-speaking from Tunisia) both with $F_{st} = 0.029$. The closest populations to each other
265were BJ (Bejaia), AZ (Azrou) and AN (Asni) with $F_{st} = 0.005-0.006$, and the most distant
266populations were CA (Copt Adaima) and TN (Tunisia) with $F_{st} = 0.052$. Altogether these
267values were relatively low.

2684. Discussion

269These results constitute the first data reported on genetic diversity of the Bejaia population.
270The 15 STR loci were highly polymorphic with a significant proportion (40%) of rare alleles
271(Tab. 2 and ST. 2). The power of discrimination (0.911-0.975), the probability of exclusion
272(0.388-0.624) and the polymorphic information content (0.74-0.90) (Tab. 2) were high with
273combined PD and PE values higher than 0.999. As expected, the most polymorphic loci
274($PIC > 0.8$) were also the most discriminating. Altogether these results strongly support the use
275of this set of genetic markers for forensic personal identification and paternity testing in the
276Bejaia region.

277Regarding population structure, the different analyses provided concordant results but
278exhibited also slightly different levels of discrimination. The STRUCTURE and AMOVA
279analyses identified little genetic structure across North Africa and between major linguistic or
280geographical groupings. The MDS and Tree analyses (Figs. 2 and 3) confirmed this but
281identified possible subgroups. For instance, some populations, such as MB (Mozabites), SM
282(Berbers from South Morocco), SH (Sahrawis), AM (Arabs from Morocco) and BM
283(Northern Morocco Berbers), appeared closer to the Arabian peninsula populations, while
284others, including Bejaia (BJ), Lybia (LY), Asni (AN), Bouhria (BH) (Figs. 2 and 3), appeared
285closer to European and other Middle East (Lebanon, Iran and Iraq) populations. However, the
286main result is that the different populations are slightly differentiated from each other but the
287differentiation exists both among geographically close or distant populations and among
288populations speaking the same or a different language.

289The genetic heterogeneity of North African populations with more or less affinities with
290Middle East, Europe and Sub-Saharan Africa has been suggested by authors using mtDNA
291(Plaza et al., 2003; Coudray et al., 2009), Y-chromosome DNA (Arredi et al., 2004; Capelli et
292al., 2006), STR markers (Capelli et al., 2006; El Ossmani et al., 2010) and SNPs (Botigué et
293al., 2013; Henn et al., 2012). In some studies, a West-to-East gradient, ranging from West
294Sahara to the Middle East has been described, which we could not detect in the present study,
295probably due to the limited number of markers or populations used. As suggested by the
296results of Henn et al. (2012), if larger numbers of populations and markers were sampled from
297North Africa and the neighboring regions (Europe, Middle-East and Sub-Saharan), one can
298expect a continuous complex with multi-polar gradients between the various ancestries
299admitted to North African populations (Maghrebi = Berber, Sub-Saharan, Middle-Eastern and
300European), as suggested by Serre and Pääbo (2004).

301While we did not find a clear geographical pattern, we can note that some geographical
302structure appear when the two dimensions of Figure 2 are considered separately. Indeed, the
303North African populations (including Tunisia) are distributed as a gradient between Sub-
304Saharan and European groups according to dimension 1 and as another gradient between Sub-
305Saharan and Arabian Peninsula groups. In relation to dimension 1, Sub-Saharan, North Africa
306 (including Mozabites) and Arabian Peninsula populations are confounded. This suggests
307that a finer geographical sampling and a larger number of markers would be necessary to
308identify the regions through which gene flow connected all these regions.

309At a smaller geographical scale and based on the MDS and tree analyses (Figs. 2-3 and Tab.
3104), the closest neighbors of the Bejaia population (BJ) were Azrou (AZ), Rabat (RB), Lybia
311(LY), Asni (AN), Bouhria (BH) with genetic distances (*Fst*) lower than 0.01. The Mozabite
312(MB) and the Tunisian populations (TN) were more distant from the Bejaia population (both
313with *Fst*=0.029) despite their geographical closeness and the shared Berber language. The out

314layer position of the Tunisian Chenini population (Figs. 2 and 3) can be attributed to genetic
315drift due to the small size of the sampled population as proposed by Khodjet-El-Khil et al.
316(2008) and Bentayebi et al. (2014).

317These observations illustrate the absence of correlation of genetic distances with both
318geographical distances and the spoken language demonstrated by the AMOVA and Mantel
319tests (Tab. 2). The absence of correlation between genetic and geographical distances at this
320geographical scale may be due to population relocations, isolations and genetic drift. Indeed,
321most studies that have shown that genetic distances are correlated with geographical distances
322(Ramachandran et al., 2005; Lao et al., 2008) were performed at large geographic scale.
323Studies carried out at smaller scale (within regions such as North Africa) are likely more
324influenced by population relocations and isolations (Ramachandran et al., 2005).

325No significant genetic differences were found in this study between the Berber- and the Arab-
326speaking populations (Tab. 3). This lack of differentiations between these two groups of
327populations have also been found by several studies using classical markers (Bosh et al.,
3281997), Alu insertion polymorphism (Coma et al., 2000), Y chromosome (Bosh et al., 2001),
329mtDNA (Fadhlaoui-Zid, 2004) and autosomal STRs (Bosh et al., 2000; Khodjet el khil et al.,
3302008 and 2012). This suggests that either the presence of Arab-speaking groups in north
331Africa was mostly a cultural process, with limited gene flow between Arabs and Berbers
332(Bosh et al, 2000), or that these populations were genetically very similar when they met.

333Our results show that language boundaries are not correlated with genetic distances for North
334African populations, probably due to the fact that the Arabisation is recent in the region.
335However, this is not necessarily a general rule since several authors found correlation between
336language boundaries and genetic differentiation (Barbujani and Sokal, 1990; Chen et al.,
3371995).

338 SNP-based STRUCTURE analysis (730,000 sites) of 7 North African populations in
339 comparison with populations from Middle East (Qatar), Europe (Tuscan and Basque) and
340 Sub-Saharan Africa (6 populations) has revealed a putative autochthonous North African
341 ancestry (referred to as Maghrebi = Berber ancestry component) decreasing in frequency from
342 Western Sahara eastward to Egypt (interrupted only by the isolated Tunisian Berber Chenini
343 population) with a parallel and equal increase of the Middle Eastern and European ancestry
344 components, with lesser and irregular Sub-Saharan influence (Henn et al., 2012). **Our STR-**
345 **based STRUCTURE analysis did not retrieve this east-to-west ancestry gradient in North**
346 **African populations (SF. 1)**, and did not detect the genetic heterogeneity suggested by MDS,
347 phylogenetic and AMOVA (Figs. 2-3 and Table 3) analyses. This may be due to the low
348 number of markers used in our study (15 STR) and/or the sample size of the populations
349 analyzed. As demonstrated by Pritchard et al., 2000, the accuracy of inferences improves with
350 sample size, number of loci, and degree of divergence between populations. **Our results are in**
351 **agreement with previously reported observation (Bosch et al., 2000, Khodjet-El-Khil et al.,**
352 **2008 and 2012).**

353 Altogether our results show that the language spoken today may not reflect the history of the
354 populations, with several Arab-speaking populations being Berbers who shifted their language
355 after the Arab conquest. Another possibility is that genetic drift in some of them has led to
356 significant differences in allele frequencies which blurred the historical relationships. Also,
357 admixture and gene flow between Arab-speaking and Berber-speaking population may have
358 contributed to the present-day situation where linguistic and genetic distances are less
359 correlated than they perhaps were in the past.

360 In this study, we do not wish to make strong statements and draw conclusions on these issues.
361 Our aim was to identify useful markers for forensic studies and quantify genetic diversity in
362 the Bejaia area compared to other previously analyzed populations. In order to reconstruct the

363history of the Bejaia population we would need a better geographical sampling of Algeria and
364North Africa. We would also need to apply more complex and advanced statistical methods
365that those used here. In particular, it would be interesting to better understand the relationships
366between north Africa and the Andalusians of Moroccan origin who came to settle around
367Bejaia after the fall of Andalusia in 1610 (see Gaid, 2008). Similarly it would be interesting to
368quantify the impact of the various invaders of the Bejaia region during history. Historical texts
369and the genetic closeness of the Bejaia population to its neighbours found here suggests that
370these contributions were probably limited but it would still be interesting to quantify them
371using genomic approaches and inferential methods such as Approximate Bayesian
372Computation (ABC, Beaumont, 2010). One could for instance test whether it is true that
373Berbers were little impacted by external gene flow as a consequence of their taking refuge in
374difficultly accessible mountains. More populations and more different markers must be used
375before drawing decisive conclusions.

376While it is not new to state that spoken languages do not constitute a reliable criterion of
377ethnic origin, our results show that it is also true in North Africa between Berber-speaking and
378Arab-speaking populations. This suggests that genomic studies using Mozabites as
379representative of Berber-speaking populations should perhaps be regarded as very
380approximate. Interestingly the genetic heterogeneity of the North African Berber populations
381together with their relative closeness to the European and Middle Eastern populations
382revealed here suggest that these populations should probably be more integrated in models
383aiming at understanding the recent demographic history of Europe, including both historical
384and prehistoric events such as the Neolithic transition.

385**Acknowledgements**

386

387We thank all the donors who made this study possible and Isa Pais for technical help in the
388laboratory (Instituto Gulbenkian de Ciência). We thank Elena Bosch, Maria Esther Esteban,
389Jean-Michel Dugoujon, and Ana Fernandez-Santander for providing STR data for North

390African populations, Naoko Takezaki for help with tree software. LC is partly funded by the
391Laboratoire d'Excellence (LABEX) entitled TULIP (ANR-10-LABX-41).

392

393**Conflict of interest:** the authors declare they have no conflict of interest.

394

395References

396

397Achilli, A., Rengo, C., Battaglia, V., Pala, M., Olivieri, A., et al. 2005. Saami and Berbers—an
398unexpected mitochondrial DNA link. *Am. J. Hum. Genet.* 76, 883–886.

399Alshamali, F., Alkhayata, A.Q., Budowl, B., Watson, N.D., 2005. STR population diversity in
400nine ethnic populations living in Dubai. *Forensic Sci. Int.* 152, 267–279.

401Alves, C., Gusmao, L., Lopez-Parra, A.M., Mesa, M. S., Amorim, A., Arroyo-Pardo, E.,
4022005. STR allelic frequencies for an African population sample (Equatorial Guinea) using
403AmpFI STR Identifiler and Powerplex 16 kits. *Forensic Sci. Int.* 148, 239–242.

404Alves, C., Gusmao, L., Damasceno, A., Soares, B., Amorim, A., 2004. Contribution for an
405African autosomic STR database (AmpF/STR Identifiler and Powerplex 16 system) and a
406report on genotypic variations. *Forensic Sci. Int.* 139, 201–205.

407Applied Biosystems, 1998. AmpF STR Profiler™ PCR Amplification Kit User's Manual. PE
408Applied Biosystems, San Jose CA.

409Arredi, B., Poloni, E.S., Paracchini, S., Zerjal, T., Fathallah, D.M., Makrelouf, M., Pascali,
410V.L., Novelletto, A., Tyler-Smith, C., 2004. A Predominantly Neolithic Origin for Y-
411Chromosomal DNA Variation in North Africa. *Am. J. Hum. Genet.* 75, 338–345.

412Barbujani, G., Chikhi, L., 2007. Human genetic diversity and its history. In *Handbook of*
413*Statistical Genetics*, eds D. Balding, M. Bishop, and C. Cannings, 3rd edition.

414Barbujani, G., Sokal, R.R., 1990. Zones of sharp genetic change in Europe are also linguistic
415boundaries. *Proc. Natl. Acad. Sci. USA* 87, 1816–19.

416Barni, F., Berti, A., Pianese, A., Boccellino, A., Miller, M. P., Caperna, A., Lago, G., 2007.
417Allele frequencies of 15 autosomal STR loci in the Iraq population with comparisons to other
418populations from the middle-eastern region. *Forensic Sci. Int.* 167, 87–92.

419Beaumont M.A., 2010. Approximate Bayesian Computation in Evolution and Ecology. *Annu.*
420*Rev. Ecol. Evol. Syst.* 41, 379–406.

421Bekada, A., Fregel, R., Cabrera, V.M., Larruga, J.M., Pestano, J., et al. 2013. Introducing the
422Algerian Mitochondrial DNA and Y-Chromosome Profiles into the North African Landscape.
423*PLoS ONE* 8, e56775.

- 424Beleza, S., Alves, C., Reis, F., Amorim, A., Carracedo, A., Gusmao, L., 2004. 17 STR data
425(AmpF/STR Identifiler and Powerplex 16 System) from Cabinda (Angola). *Forensic Sci. Int.*
426141, 193–196.
- 427Bentayebi, K., Abada, F., Ihzmad, H., Amzazi, S., 2014. Genetic ancestry of a Moroccan
428population as inferred from autosomal STRs. *Meta Gene* 2, 427–438
- 429Bernal, L.P., Borjas, L., Zabala, W., Portillo, M.G., Fernandez, E., Delgado, W., Tovar, F.,
430Lander, N., Chiurillo, M.A., Ramirez, J.L., Garcia, O., 2006. Genetic variation of 15 STR
431autosomal loci in the Maracaibo population from Venezuela. *Forensic Sci. Int.* 161, 60–63.
- 432Bosch, E., Calafell, F., Pérez-Lezaun, A., Clarimon, J., Comas, D., Mateu, E., Martinez-Arias,
433R., Morera, B., Brakez, Z., Akhayat, O., Sefiani, A., Hariti, G., Cambon-Thomsen, A.,
434Bertranpetit, J., 2000. Genetic structure of north-west Africa revealed by STR analysis. *Eur. J.*
435*Hum. Genet.* 8, 360–366.
- 436Bosch, E., Calafell, F., Perez-Lezaun, A., Coma, D., Mateu, E., and Bertranpetit, J., 1997.
437Population history of North Africa: evidence from classical genetic markers. *Hum. Biol.* 69,
438295 – 311.
- 439Bosch, E., Clarimon, J., Perez-Lezaun, A., Calafell, F., 2001. STR data for 21 loci in
440northwestern Africa. *Forensic Sci. Int.* 116, 41–51.
- 441Botigué, L.R., Henn, B.M., Gravel, S., Maples, B.K., Gignoux, C.R., Corona, E., Atzmon, G.,
442Burns, E., Ostrer, H., Flores, C., Bertranpetit, J., Comas, D., Bustamante, C.D., 2013. Gene
443flow from North Africa contributes to differential human genetic diversity in southern Europe.
444*Proc. Natl. Acad. Sci.* 110 (29), 11791–11796.
- 445Butler, J. M., 2006. Genetics and Genomics of Core Short Tandem Repeat Loci Used in
446Human Identity Testing. *J. Forensic Sci.* 51, 253–265.
- 447
- 448Capelli, C., Redhead, N., Romano, V., Cali, F., Lefranc, G., Delague, V., Megarbane, A.,
449Felice, A. E., Pascali, V. L., Neophytou, P. I., Poulli, Z., Novelletto, A., Malaspina, P.,
450Terrenato, L., Berebbi, A., Fellous, M., Thomas, M. G. & Goldstein, D.B. (2006) Population
451structure in the Mediterranean basin: a Y chromosome perspective. *Ann. Hum. Genet.* **70**,
452207–25.
- 453
- 454Cerny, V., Mulligan, C.J., Ridl, J., Zaloudkova, M., Edens, C.M., Hajek, M., Pereira, L., 2008.
455Regional differences in the distribution of the sub-Saharan, West Eurasian, and South Asian
456mtDNA lineages in Yemen. *Am. J. Phys. Anthropol.* 136,128– 137.
- 457
- 458Chen, J., Sokal, R. R., Ruhlen, M., 1995. Worldwide analysis of genetic and linguistic
459relationships of human populations. *Hum. Biol.* 67, 595–612.
- 460Chouery, E., Coble, M.D., Strouss, K.M., Saunier, J.L., Jalkh, N., Medlej-Hashim, M.,
461Ayoub, F., Mégarbané, A., 2010. Population genetic data for 17 STR markers from Lebanon.
462*Leg. Med.* 12, 324–326.

- 463Comas, D., Calafell, F., Benchemsi, N., Helal, A., Lefranc, G., Stoneking, M., Batzer, M.A.,
464Bertranpetit, J. and Sajantila, A., 2000. Alu insertion polymorphisms in NW Africa and the
465Iberian Peninsula: evidence of a strong genetic boundary through the Gibraltar straits. *Hum.*
466*Genet.* 107, 312 – 319.
- 467Cote, M., 1991. « Béjaïa ». *Encyclopédie berbère*, 9 | *Baal – Ben Yasla*, Aix-en-Provence,
468Edisud, pp. 1408–1415.
- 469Coudray, C., Calderon, R., Guitard, E., Ambrosio, B., Gonzalez-Martin, A., Dugoujon, J.M.,
4702007c. Allele frequencies of 15 tetrameric short tandem repeats (STRs) in Andalusians from
471Huelva (Spain). *Forensic Sci. Int.* 168, e21–e24.
- 472Coudray, C., Guitard, E., El-Chennawi, F., Larrouy, G., Dugoujon, J.M., 2007b. Allele
473frequencies of 15 short tandem repeats (STRs) in three Egyptian populations of different
474ethnic groups. *Forensic Sci. Int.* 169, 260–265.
- 475Coudray, C., Guitard, E., Kandil, M., Harich, N., Melhaoui, M., Baali, A., Sevin, A., Moral,
476P., Dugoujon, J.M., 2006. Study of GM immunoglobulin allotypic system in Berbers and
477Arabs from Morocco. *Am. J. Hum. Biol.* 18, 23–34.
- 478Coudray, C., Guitard, E., Keyser-Tracqui, C., Melhaoui, M., Cherkaoui, M., Larrouy, G.,
479Dugoujon, J.M., 2007a. Population genetic data of 15 tetrameric short tandem repeats (STRs)
480in Berbers from Morocco. *Forensic Sci. Int.* 167, 81–86.
- 481Coudray, C., Guitard, E., Lemaire, O., Cherkaoui, M., Baali, A., Hilali, K., Sevin, A., Kandil,
482M., Harich, N., Melhaoui, M., Larrouy, G., Moral, P., Dugoujon, J.M., 2004. Les allotypes Gm
483des immunoglobulines chez les Berbères du Maroc. *Antropo.* 6, 63–69.
- 484Coudray, C., Torroni, A., Achilli, A., Pala, M., Olivieri, A., Larrouy, G., Dugoujon, J.M. 2009.
485Les lignées mitochondriales et l’histoire génétique des populations berbérophones du nord de
486l’Afrique. *Antropo.* 18, 63–72.
- 487Decorte, R., Engelen, M., Larno, L., Nelissen, K., Gilissen, A., Cassiman, J.J., 2004. Belgian
488population data for 15 STR loci (AmpFISTR1 SGM Plus and AmpFISTR1M profiler PCR
489amplification kit). *Forensic Sci. Int.* 139, 211–213.
- 490Dugoujon, J.M., Hazout, S., Loirat, F., Mourrieras, B., Crouau-Roy, B., Sanchez-Mazas, A.,
4912004. GM haplotype diversity of 82 populations over the world suggests a centrifugal model
492of human migrations. *Am. J. Phys. Anthropol.* 125, 175–192.
- 493Earl, D.A. and vonHoldt, B.M., 2012. STRUCTURE HARVESTER: a website and program
494for visualizing STRUCTURE output and implementing the Evanno method. *Conservation*
495*Genetics Resources* vol. 4 (2), 359-361
- 496El Ossmani, H., Bouchrif, B., Talbi, J., El Amri, H., Chafik, A., 2007. La diversité génétique
497de 15 STR chez la population arabophone de Rabat-Salé-Zemmour-Zaer. *Antropo.* 15, 55–62.

498El Ossmani, H., Bouchrif, B., Aboukhalid, R., Bouabdillah, M., Gazzaz, B., Zaoui, D.,
499Chafik, A., Talbi, J., 2010. Assessment of phylogenetic structure of Berber-speaking
500population of Azrou using 15 STRs of Identifiler kit. *Leg. Med.* 12(1), 52-56.

501Ennafaa, H., Cabrera, V.M., Abu-Amero, K.K., González, A.M., Amor, M.B., Bouhaha, R.,
502Dzimiri, N., Elgaaïed A.B., Larruga J.M., 2009. Mitochondrial DNA haplogroup H structure in
503North Africa. *BMC Genetics* 10:8.

504Epstein, M.P., Duren, M.L. and Boehnke M., 2000. Improved Inference of Relationship for
505Pairs of Individuals. *Am. J. Hum. Genet.* 67,1219–1231.

506Evanno, G., Regnaut, S., Goudet, J., (2005). Detecting the number of clusters of individuals
507using the software structure: a simulation study. *Molecular Ecology* 14 (8), 2611-2620.

508Excoffier, L. and Lischer, H.E. L., 2010. Arlequin suite ver 3.5: A new series of programs to
509perform population genetics analyses under Linux and Windows. *Mol. Ecol. Resour.* 10, 564-
510567.

511Fadhlaoui-Zid, K., Plaza, S., Calafell, F., Ben Amor, M., Comas, D., Elgaaïed, A.B., 2004.
512Mitochondrial DNA heterogeneity in Tunisian Berbers. *Ann. Hum. Genet.* 68, 222–233.

513Felsenstein, J., 1985. Confidence limits on phylogenies: an approach using the bootstrap.
514*Evolution* 39,783–791.

515Gaïbar, M., Esteban, M. E., Via, M., Harich, N., Kandil, M., Fernández-Santander, A., 2012
516Usefulness of autosomal STR polymorphisms beyond forensic purposes: data on Arabic- and
517Berber-speaking populations from central Morocco. *Ann. Hum. Biol.* 39(4), 297-304.

518Gaïd M., 2008. Histoire de Béjaïa et de sa région, depuis l'antiquité jusqu'à 1954. Editions
519Mimouni, Alger.

520Gonzalez-Pérez, E., Via, M., Esteban, E., Lopez-Alomar, A., Mazieres, S., Harich, N.,
521Kandil, M., Dugoujon, J.M., Moral, P., 2003. Alu insertions in the Iberian Peninsula and north
522west Africa - genetic boundaries or melting pot?. *Coll. Antropol.* 27, 491–500.

523Gorostiza, A, Gonzalez-Martin, A., Ramirez, C. L., Sanchez, C., Barrot, C., Ortega, M.,
524Huguet, E., Corbella, J., Gené, M., 2007. Allele frequencies of the 15 AmpF/Str Identifiler
525loci in the population of Metztlán (Estado de Hidalgo), México. *Forensic Sci. Int.* 166, 230–
526232.

527Guo, S. and Thompson, E. 1992 Performing the exact test of Hardy-Weinberg proportion for
528multiple alleles. *Biometrics* 48,361-372.

529Havas, D., Jeran, N., Efremovska, L., Dorpevic, D., Rudan, P., 2007. Population genetics of
53015 AmpfI STR Identifiler loci in Macedonians and Macedonian Romani (Gypsy). *Forensic Sci.*
531*Int.* 173, 220–224.

532Henn, B. M., Botigue, L. R., Gravel, S., Wang, W., Brisbin, A., Byrnes, J. K., Fadhlou-Zid,
533K., Zalloua, P. A., Moreno-Estrada, A., Bertranpetit, J., Bustamante, C. D., Comas, D., 2012.
534Genomic Ancestry of North Africans Supports Back-to-Africa Migrations. *PLoS Genet.* 8(1),
535e1002397.

536khodjet-el-khil, H., fadhlaoui-zid, K., Gusmao, L., Alves, C., Benammar-elgaaied, A.,
537Amorim, A., 2008. Substructure of a Tunisian Berber Population as Inferred from 15
538Autosomal Short Tandem Repeat Loci. *Human Biology* 80, 435–448.

539Khodjet-el-Khil, H., fadhlaoui-zid, K., Gusmao, L., Alves, C., Benammar-elgaaied, A.,
540Amorim, A., 2012. Allele frequencies for 15 autosomal STR markers in the Libyan population.
541*Ann. Hum. Biol.* 39, 80–83.

542Kim, Y.L. Hwang, J.Y., Kim, Y.J., Lee, S., Chung, N.G., Goh, H.G., Kim, C.C., Kim, D.W.,
5432003. Allele frequencies of 15 STR loci using AmpF/STR Identifiler kit in a Korean
544population. *Forensic Sci. Int.* 136, 92–95.

545Lao, O., Lu, T.T., Nothnagel, M., Junge, O., Freitag-Wolf, S., et al., 2008, Correlation
546between Genetic and Geographic Structure in Europe. *Current Biology* 18, 1241–1248.

547Laporte, J.P., 2004. « Kabylie : La Kabylie antique ». *Encyclopédie berbère*, 26 | Judaïsme –
548Kabylie, Aix-en-Provence, Edisud, pp. 4000–4015.

549Lefevre-Witier P., Aireche H., Benabadji M., Darlu P., Melvin K., Andre Sevin A., Crawford
550M.H. 2006. Genetic Structure of Algerian Populations. *Am. J. Hum. Bio.* 18, 492–501.

551Levene, H., 1949. On a matching problem arising in genetics. *Annals of Mathematical*
552*Statistics* 20, 91-94.

553Lopes, V., Serra, A., Gamero, J., Sampaio, L., Balsa, F., Oliveira, C., Batista, L., Corte-Real,
554F., Vieira, D.N., Vide, M.C., Anjos, M.J., Carvalho, M., 2009. Allelic frequency distribution
555of 17 STRs from Identifiler and PowerPlex-16 in Central Portugal area and the Azores
556archipelago. *Genetics* 4, e1–e7.

557Mantel, N. 1967. The detection of disease clustering and a generalized regression approach.
558*Cancer Res.* 27, 209-220.

559Matamoros, M., Pinto, Y., Inda, F.J., Garcia, O., 2008. Population genetic data for 15 STR loci
560(Identifiler™ kit) in Honduras. *Leg. Med.* 10, 281–283.

561Montelius, K., Karlsson, A.O., Holmlund, G., 2008. STR data for the AmpF™STRIdentifiler
562loci from Swedish population in comparison to European, as well as with non-European
563population. *Forensic Sci. Int. Genetics* 2, e49–e52.

564Nei, M., 1987. *Molecular Evolutionary Genetics*. Columbia University Press, New York, NY,
565USA.

566Pemberton, P., Laurent T. J., Kemp, R., Gonzalez-Oliver, B. M., et al. 2014. Patterns of
567Admixture and Population Structure in Native Populations of Northwest North America. *PLoS*
568*Genetics* 10, 1-17.

569Pemberton, T.J., DeGiorgio, M. and Rosenberg, N.A., 2013. Population Structure in a
570Comprehensive Genomic Data Set on Human Microsatellite Variation.
571*Genes/Genomes/Genetics* 3, 891-916.

572Pereira, L., Silva, N.M., Franco-Duarte, R., Fernandes, V., Pereira, J.B., Costa, M.D., Martins,
573H., Soares, P., Behar, D.M., Richards, M.B., Macaulay V., 2010. Population expansion in the
574North African Late Pleistocene signalled by mitochondrial DNA haplogroup U6. *BMC Evol.*
575*Biol.* 10:390.

576
577Plaza S., F. Calafell, A. Helal, N. Bouzerna, G. Lefranc, J. Bertranpetit and D. Comas, 2003.
578Joining the Pillars of Hercules: mtDNA Sequences Show Multidirectional Gene Flow in the
579Western Mediterranean. *Annals of Human Genetics* 67: 312–328.

580Pritchard, J.K., Stephens, M., Donnelly, P., 2000. Inference of population structure using
581multilocus genotype data. *Genetics* 155, 945 – 959.

582Ramachandran, S., Deshpande, O., Roseman, C. C., Rosenberg, N.A., Feldman, M.W., and
583Cavalli-Sforza, L.L., 2005. Support from the relationship of genetic and geographic distance
584in human populations for a serial founder effect originating in Africa. *PNAS* 102 (44), 15942–
58515947.

586Rebala, K., Wysocka, J., Kapinska, E., Cybulska, L., Mikulich, A.I., Tsybovsky, I.S.,
587Szczerkowska, Z., 2007. Belarusian population genetic database for 15 autosomal STR loci.
588*Forensic Sci. Int.* 173, 235–237.

589Regueiro, M., Carril, J.C., Pontes, M.L., Pinheiro, M.F., Luis, J.R., Caeiro, B., 2004. Allele
590distribution of 15 PCR-based loci in the Rwanda Tutsi population by multiplex amplification
591and capillary electrophoresis. *Forensic Sci. Int.* 143, 61–63.

592Rerkamnuaychoke, B., Rinthachai, T., Shotivaranon, J., Jomsawat, U., Siriboonpiputtana, T.,
593Chaiatchanarat, K., Pasomsub, E., Chantratita, W., 2006. Thai population data on 15 tetrameric
594STR loci—D8S1179, D21S11, D7S820, CSF1PO, D3S1358, TH01, D13S317, D16S539,
595D2S1338, D19S433, vWA, TPOX, D18S51, D5S818 and FGA. *Forensic Sci. Int.* 158, 234–
596237.

597Rocabado, O., Taboada, P., Inda, F.J., Yurrebaso, I., Garcia, O., 2009. Population genetic data
598for 15 STR loci (Identifiler™ kit) in Bolivia. *Leg. Med.* 11, 302–304.

599Rosenberg N.A., Mahajan, S., Ramachandran, S., Zhao, C., Pritchard, J. K., Feldman, M. W.
6002005. Clines, Clusters, and the Effect of Study Design on the Inference of Human Population
601Structure. *PLoS Genetics* 1, 660-671.

602Rosenberg, N. A, Pritchard, J. K., Weber, J. L., Cann, H. M., Kidd, K.K., Zhivotovsky, L. A.,
603Feldmanl, M. W. 2002. Genetic structure of human populations. *Science* 298, 2381–2385.

604Serre, D., Pääbo, S. 2004. Evidence for gradients of human genetic diversity within and
605among continents. *GenomeResearch*, 14, 1679–1685.

606Shepard, E.M., Herrera, R.J., 2006. Iranian STR variation at the fringes of biogeographical
607demarcation. *Forensic Sci. Int.* 158, 140–148.

608Smouse, P. E., Long, J. C., 1992. Matrix correlation analysis in Anthropology and Genetics. *Y.*
609*Phys. Anthop.* 35, 187-213.

610StatSoft Inc., 2008. STATISTICA (Data Analysis Software System), Version 8.0.
611<http://www.statsoft.com>.

612Taïeb, J., 2004. « Juifs du Maghreb : onomastique et langue, une composante berbère ? ». *Y.*
613*Encyclopédie berbère*, 26 | Judaïsme – Kabylie, Aix-en-Provence, Edisud, p. 3969–3975.

614Takezaki, N., Nei, M., Tamura, K., 2010. POPTREE2: Software for Constructing Population
615Trees from Allele Frequency Data and Computing Other Population Statistics with Windows
616Interface. *Mol. Biol. Evol.* 27, 747–752.

617Wang, C.W., Chen, D.P., Chen, C.Y., Lu, S.C., Sun, C.F., 2003. STR data for the AmpF1STR
618SGM Plus and Profiler loci from Taiwan. *Forensic Sci. Int.* 138, 119–122.

619Yan, J., Shen, C., Li, Y., Yu, X., Xiong, X., Mue, H., Huang, Y., Deng, Y., Yu, J., 2007.
620Genetic analysis of 15 STR loci on Chinese Tibetan in Qinghai Province.*Forensic Sci. Int.*
621169, e3–e6.

622Yang, B., Wang, G., Liu, Y., Yang, W., 2005.Population data for the AmpF‘STRIdentifiler™
623PCR Amplification Kit in China Han in Jilin Province, China.*Forensic Sci. Int.* 151, 293–297.