

Page containing authors

Hemodynamic Profiles of Functional and Dysfunctional forms of Repetitive Thinking

Cristina Ottaviani¹, PhD, Jos F. Brosschot², PhD, Antonia Lonigro^{1,3}, PhD, Barbara Medea³, PhD, Ilse Van Diest⁴, PhD, Julian F. Thayer⁵, PhD

¹ Santa Lucia Foundation, Rome, Italy

² Leiden University, Leiden, The Netherlands

³ Sapienza University of Rome, Italy

⁴ University of Leuven, Leuven, Belgium

⁵ The Ohio State University, Columbus, Ohio

Correspondence concerning this article should be addressed to Cristina Ottaviani, Neuroimaging Laboratory, Santa Lucia Foundation, via Ardeatina 306, 00142 Rome, Italy. E-mail: cristina.ottaviani@uniroma1.it.

Compliance with Ethical Standards

Funding: Support for the last author was received from the Italian Ministry of Health Young Researcher Grant (2011-02348232).

Conflict of Interest: The authors declare that they have no conflict of interest.

Ethical approval: All procedures performed in studies involving human participants were in accordance with the ethical standards of the institutional research committee and with the 1964 Helsinki declaration and its later amendments or comparable ethical standards.

Informed consent: Informed consent was obtained from all individual participants included in the study.

Hemodynamic Profiles of Functional and Dysfunctional forms of Repetitive Thinking

Abstract

Background The ability of the human brain to escape the here and now (mind wandering) can take functional (problem solving) and dysfunctional (perseverative cognition) routes. Although it has been proposed that only the latter may act as a mediator of the relationship between stress and cardiovascular disease, both functional and dysfunctional forms of repetitive thinking have been associated with blood pressure (BP) reactivity of the same magnitude. However, a similar BP reactivity may be caused by different physiological determinants, which may differ in their risk for cardiovascular pathology. **Purpose** To examine the way (hemodynamic profile) and the extent (compensation deficit) to which total peripheral resistance and cardiac output compensate for each other in determining BP reactivity during functional and dysfunctional types of repetitive thinking. **Methods** Fifty-six healthy participants randomly underwent a perseverative cognition, a mind wandering, and a problem solving induction, each followed by a 5-min recovery period while their cardiovascular parameters were continuously monitored. **Results** Perseverative cognition and problem solving (but not mind wandering) elicited BP increases of similar magnitude. However, perseverative cognition was characterized by a more vascular (versus myocardial) profile compared to mind wandering and problem solving. As a consequence, BP recovery was impaired after perseverative cognition compared to the other two conditions. **Conclusions** Given that high vascular resistance and delayed recovery are the hallmarks of hypertension the results suggest a potential mechanism through which perseverative cognition may act as a mediator in the relationship between stress and risk for developing precursors to cardiovascular disease.

Keywords Perseverative Cognition; Mind Wandering; Problem Solving; Hemodynamic Profile; Cardiac Output; Total Peripheral Resistance

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

People spend a large part of their time engaging in mind wandering, which often leads to going over the same thoughts again and again, called repetitive thinking. Such an extremely common cognitive process can take functional (problem solving) and dysfunctional (rumination, worry) routes [1]. The Perseverative Cognition Hypothesis specifically suggests that rumination about the past and worrisome thoughts about the future (i.e., perseverative cognition) cause a “fight-or-flight” action tendency, followed by a cascade of biological events such as increases in cardiovascular activity and this persistent physiological activation may have an impact on an individual’s health ultimately leading to somatic disease [2, 3]. A recent meta-analysis supported this view, providing evidence of increased cardiovascular, autonomic, and endocrine nervous system activity associated with rumination and worry [4].

However, previous studies repeatedly showed that even ostensibly more functional forms of repetitive thinking, such as mind wandering and problem solving could be associated with increased physiological activity. For example, Verkuil and colleagues [5] found cardiac effects of the same magnitude during worrying and problem solving, leading the authors to conclude that “mere mental load may be responsible for at least a part of the physiological effects of worry” (page 448). Similarly, several studies linked mind wandering per se with increased physiological activation [6-8].

If both functional and dysfunctional forms of repetitive thinking were associated with physiological activity, why would only the latter be associated with increased risk for somatic disease? In terms of cardiovascular activity, it has to be noted that elevations in blood pressure (BP) of the same magnitude can be elicited by different patterns of compensatory changes in cardiac output and total peripheral resistance [9]. Therefore, looking at the physiological determinants of BP becomes more informative than focusing on BP responses per se [10]. The term “hemodynamic profile” describes the relationship between cardiac output and total peripheral resistance in the homeostatic regulation of BP [11]. The first aim of the present study was to examine the hemodynamic profiles of functional and dysfunctional forms of repetitive thinking.

1 We hypothesized that perseverative cognition would be characterized by a predominantly vascular
2 hemodynamic profile whereas problem solving and mind wandering would have a more myocardial
3 or mixed profile. If this were true, then the cardiovascular reactivity of the same magnitude that has
4 been shown to characterize perseverative cognition and problem solving would be associated with
5 distinctive hemodynamic patterns, with different implications for health. In fact elevated BP driven
6 by total peripheral resistance, compared to cardiac output, has been linked to increased risk of
7 cardiac events and mortality [9, 12, 13].
8
9
10
11
12
13
14
15
16

17 The second aim of the present study, closely connected to the first, was to show that another
18 crucial difference between functional and dysfunctional forms of repetitive thinking concerns the
19 duration of the concomitant physiological activation. Indeed only prolonged or chronic activation
20 can lead to the pathogenic state that eventually leads to organic disease [2]. A major consequence
21 of the dominance of reactivity-based theories has been the failure to examine the duration of
22 activation. This is an important limitation considering that a recent meta-analysis showed that poor
23 recovery from laboratory challenges provided incremental value for predicting adverse
24 cardiovascular outcomes beyond reactivity per se [14]. We hypothesized that functional and
25 dysfunctional forms of repetitive thinking would be characterized by an equivalent BP reactivity but
26 only dysfunctional forms would be associated with delayed BP recovery.
27
28
29
30
31
32
33
34
35
36
37
38
39
40

41 The two hypothesis of the present study are closely interconnected. In terms of hemodynamics,
42 delayed recovery has been primarily associated with vascular responding [15]. For example,
43 Steptoe and Marmot [16] found that an increase in BP over a 3-year period was predicted by
44 impaired post-stress recovery and that the elevation in BP recorded during the recovery period was
45 determined by vascular rather than cardiac responses. Consistent with this idea, extended mental
46 stress seems to be characterized by transient increases in cardiac output but prolonged changes in
47 total peripheral resistance [17]. Thus, if both our hypothesis are confirmed, it is likely that the
48 delayed recovery that characterizes perseverative cognition would actually be due to its “vascular
49 nature”.
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

In light of the role played by anxiety, depression, and anger-in in augmenting cardiovascular risk [18], the present study also examined which of these personality factors, as well as the dispositional tendency to engage in rumination and worry and state levels of sadness and anxiety, better predicted the hemodynamic profile that characterizes perseverative cognition.

To the best of our knowledge, this is the first study to examine the physiological mechanisms underlying the cardiovascular consequences of functional and dysfunctional forms of repetitive thinking and to examine the hemodynamic mechanisms through how this may ultimately lead to cardiovascular disease.

Method

Participants

The sample was composed of university students and employees. Of the 65 subjects who agreed to participate in the study, 9 were excluded due to Portapres device (see 'cardiovascular monitoring' below) malfunction. The final sample was composed of 26 women and 30 men with a mean age 24.5 (3.9) years. All subjects were Caucasian. Exclusionary criteria, assessed during a pre-screening questionnaire, were: diagnosis of psychiatric disorders (current and/or past), diagnosis of hypertension or heart disease, any other disease or use of drugs/medications that might affect cardiovascular function, obesity (body mass index $> 32 \text{ kg/m}^2$), menopause, use of oral contraceptives during the previous 6 months, and pregnancy or childbirth within the last 12 months.

Participants were compensated for their time. The protocol was approved by the Bioethical Committee of S. Lucia Foundation, Rome, Italy.

Procedure

Participants were informed of the following restrictions: no caffeine, alcohol, nicotine, or strenuous exercise for 2 h prior to the appointment. After reading and signing the informed consent form, the continuous BP cuff was attached on the middle finger of participants' right hand. After

1
2 calibration, the experimental protocol started with a ‘vanilla’ baseline period [19]. Then, all
3 participants took part in three experimental conditions: a perseverative cognition induction, a mind
4 wandering induction, and a cognitive problem-solving task.
5
6

7 The experimental conditions were presented in counterbalanced order. Each condition lasted 5
8 minutes and was followed by a 5-minute recovery period. After baseline, each experimental
9 condition, and each recovery period, participants rated their mood and thoughts by a series of visual
10 analogue scales (VAS). Cardiovascular parameters were continuously monitored by the Portapres
11 device throughout the experimental session.
12
13
14
15
16
17
18
19
20
21

22 Experimental manipulation instructions

23
24 Perseverative cognition induction: *“Now I would like you to recall an episode that happened in*
25 *the past year that made you feel sad, anxious, or stressed, or something that may happen in the*
26 *future that worries you. Then, I would like you to think about this episode in detail, for example*
27 *about its possible causes, consequences, and your feelings about it. Please take as much time as you*
28 *need to identify the event and press the button whenever you are ready”*.
29
30
31
32
33
34
35

36 Mind wandering induction: *“Now I would like you to let your mind wander without getting*
37 *stuck on any particular thought”*.
38
39
40

41 Problem solving induction: *“Now I would like you to solve a series of syllogisms. Please select*
42 *“Yes” if you think that the presented syllogism is valid, select “No” otherwise”*. Example: No A are
43 B. Some C are B. Therefore, some C are not A.
44
45
46
47

48 Recovery periods: *“The task is terminated. Now I would like you to rest until the instructions*
49 *for the following task appear on the screen”*.
50
51
52
53
54
55

56 Measures

57 Cardiovascular Monitoring

58
59 Noninvasive continuous measurement of beat-to-beat BP was obtained throughout the study
60
61
62
63
64
65

1 with the Portapres II (FMS; The Netherlands) device, which has been shown to reliably compare
2 with intra-aortic pressure measurement [20]. The arterial pressure signal was analyzed using
3 BeatScope® software to obtain systolic (SBP) and diastolic BP (DBP), cardiac output, and total
4 peripheral resistance. SBP, DBP, cardiac output, and total peripheral resistance reactivity values
5 were computed by subtracting the initial baseline from task values. SBP, DBP, cardiac output, and
6 total peripheral resistance recovery values were computed by subtracting task values from post-task
7 values. Hemodynamic Profile and Compensation Deficit were computed as detailed in the
8 following section.
9
10
11
12
13
14
15
16
17
18
19
20
21

22 *Hemodynamic Profile and Compensation Deficit*

23
24 Hemodynamic profile (HP) and compensation deficit were assessed following the orthogonal,
25 physiologically grounded model proposed by Gregg and colleagues [21]. The model is derived
26 from the multiplicative relationship between cardiac output and total peripheral resistance in
27 determining mean arterial pressure [22] and the computation is based on the following equation:
28
29
30
31
32

$$33 \log(\text{cardiac output})_r + \log(\text{total peripheral resistance})_r = \log(\text{mean arterial pressure})_r$$

34 where r indicates:

- 35
36
37
38 a) a ratio of task to baseline values for reactivity periods [10, 21];
39
40
41 b) a ratio of recovery to task values for recovery periods [15].
42

43
44 This approach has the advantage of not being based on any artificial taxonomy that would
45 ignore the continuous nature of cardiovascular measurements. The outcome is a continuous
46 variable by which participants are described as more vascular (greater HP values) when the
47 algebraic increase in $\log(\text{total peripheral resistance})_r$ exceeds that in $\log(\text{cardiac output})_r$, and
48 more myocardial when the algebraic increase in $\log(\text{cardiac output})_r$ exceeds that in $\log(\text{total}$
49 $\text{peripheral resistance})_r$. Compensation deficit (CD) increases as the algebraic sum of the
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

cardiac output.

Visual Analog Scales (VAS)

After baseline, each task and each recovery period, participants were asked to rate their current levels of feeling Anxious, Angry, Happy, Tired, and Sad on separate visual analogue 100-point scales. The same scale was used to inquire about participants' ongoing cognitive activity: "How much for the duration of the task were you": 1) mind wandering, 2) ruminating, 3) worrying. For each VAS, change scores were computed by subtracting the initial baseline from task values for reactivity periods and by subtracting the task from post-task values for recovery periods.

Questionnaires

Ruminative Response Scale is a measure of depressive rumination tendencies assessed by how often people engage in responses to depressed mood that are self-focused (e.g., I think "Why do I react this way?"), symptom-focused (e.g., I think about how hard it is to concentrate), and focused on the possible consequences and causes of one's mood (e.g., I think "I won't be able to do my job if I don't snap out of this") [RRS; 23]. Reliability and validity of the RRS were supported through several longitudinal studies with Cronbach's α ranging from 0.75 to 0.80 [24, 25].

Penn State Worry Questionnaire is a 16-item self-report questionnaire mainly focused on future outcomes (e.g., As soon as I finish one task, I start to worry about everything else I have to do) and commonly used to assess pathological worry in both clinical and non-clinical populations [PSWQ; 26]. The PSWQ has demonstrated good discriminant validity [27-29] and high internal consistency and test-retest reliability with Cronbach's alphas ranging between .86 and .95 [28].

State Trait Anxiety Inventory includes a measure of trait dispositional anxiety that targets how respondents "generally feel" (e.g., I am a steady person) [STAI; 30]. High validity and reliability (Cronbach's alpha from .86 to .95) have been documented [31, 32].

Center for Epidemiologic Studies Depression Scale is a 20-item self-report scale designed to

1 measure depressive symptomatology (e.g., I felt that everything I did was an effort) over the
2 previous week in the general population [CES-D; 33]. The validity of the CES-D has been
3 repeatedly confirmed, although some specific items are currently a matter of debate [34].
4
5 Cronbach's alphas are above .85 in the general population and .90 in depressed patients confirming
6 high reliability [33].
7

8
9
10
11
12 *Anger-In subscale* of the *Spielberger Trait Anger Expression Inventory* is a measure of the
13 tendency of individuals to hold in or suppress responses to anger provocation (e.g., I control my
14 urge to express my angry feelings) [STAXI; 35]. The STAXI has consistently demonstrated
15 evidence to support its validity and reliability as an instrument to assess anger (overall Cronbach's
16 alpha above .90) [36].
17
18
19
20
21
22

23 24 25 26 Data Analysis

27
28 All data are expressed as means (SD). Differences at $p \leq .05$ were regarded as significant.
29
30 Data processing was performed with the software modules of SPSS 23 (IBM). SBP, DBP, cardiac
31 output, and total peripheral resistance reactivity and recovery, hemodynamic profile, compensation
32 deficit, and scores on personality questionnaires were treated as continuous variables.
33
34
35
36

37
38 Sex differences were analyzed by t -tests and χ^2 tests.

39
40
41 To test for differences in reactivity and recovery levels for the three experimental conditions, a
42 series of 3 (Induction: Perseverative Cognition, Mind Wandering, Problem Solving) x 2 (Time:
43 Reactivity, Recovery) x 6 (Order) General Linear Models (GLMs) were performed on SBP, DBP,
44 cardiac output, total peripheral resistance, and each VAS. Reactivity and recovery change scores
45 were used in these GLMs.
46
47
48
49
50
51

52
53 Consistent with the approach adopted by James and Gregg [37], one-sample t tests were used to
54 test the difference from zero of hemodynamic profile and compensation deficit scores for each
55 condition and subsequent recovery periods. A significant t -test result for hemodynamic profile was
56 taken to indicate either a vascular (positive t value) or a myocardial profile (negative t value). A
57
58
59
60
61
62
63
64
65

1 nonsignificant hemodynamic profile result coupled with a significant compensation deficit result
2 means that the response was mixed (i.e., neither vascular nor myocardial). No hemodynamic
3
4 response at all was deemed to have occurred when both hemodynamic profile and compensation
5
6 deficit were not significant.
7

8
9 To examine the predictive power of both state moods and dispositional traits in determining the
10 hemodynamic profile that characterizes functional and dysfunctional forms of repetitive thinking
11 above and beyond traditional predictors, three hierarchical multiple regression analyses were
12 conducted with sex entered in the first stage (see below); mood and thought changes (anxious, sad,
13 ruminating, worrying) during the corresponding induction entered in the second stage, and the
14 questionnaires scores entered in the third stage as the independent variables. Hemodynamic profile
15 during the perseverative cognition, mind wandering, and problem solving inductions served as
16 dependent variables.
17
18
19
20
21
22
23
24
25
26
27

28 **Results**

29 **Descriptive statistics**

30
31 Significant sex differences emerged for baseline DBP ($t(54) = 2.18; p = .03$), and scores on the
32 CESD ($t(54) = 2.91; p = .01$), STAI ($t(54) = 3.12; p = .003$), PSWQ ($t(54) = 2.65; p = .01$), and
33 RRS ($t(54) = 2.17; p = .04$). As shown in Table 1, men were characterized by higher levels of DBP
34 and trait worry, whereas women had higher tendencies toward anxiety, depression, and depressive
35 rumination, therefore sex was included as a covariate in all the analyses concerning these variables.
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50

51 **Blood Pressure, Total Peripheral Resistance, and Cardiac Output Reactivity and Recovery**

52
53 The GLM having SBP as the dependent variable yielded significant effects of Time, $F(1,50) =$
54 $10.19, p = .002; \eta^2 = .17$, Induction $F(2,100) = 7.82, p = .001; \eta^2 = .14$, and Time x Induction
55 interaction, $F(2,100) = 6.20, p = .003; \eta^2 = .11$. As depicted in Figure 1, post hoc comparisons
56 showed that SBP increases were significantly larger during perseverative cognition and problem
57
58
59
60
61
62
63
64
65

1 solving compared to mind wandering ($ps < .002$). Moreover, SBP significantly decreased from
2 problem solving to recovery from the same task ($p < .0001$), but not for mind wandering, and
3 marginally increased from perseverative cognition to the subsequent recovery period ($p = .07$).
4 Within the recovery period, SBP was higher during perseverative cognition compared to mind
5 wandering and problem solving ($ps < .002$), with no significant differences between the last two
6 conditions.
7

8
9
10 As for DBP, significant effects of Time, $F(1,50) = 7.37, p = .01; \eta^2 = .13$, Induction $F(2,100) =$
11 $3.83, p = .02; \eta^2 = .07$, and Time x Induction interaction, $F(2,100) = 5.59, p = .005; \eta^2 = .10$
12 emerged. As illustrated in Figure 1, during both perseverative cognition and problem solving there
13 were significantly larger DBP increases compared to mind wandering ($ps < .01$). Moreover, DBP
14 significantly decreased from problem solving to recovery from the same task ($p = .01$), but this was
15 not the case for mind wandering and perseverative cognition. Within the recovery period, DBP was
16 higher during perseverative cognition compared to mind wandering and problem solving ($ps < .01$),
17 with no significant differences between the last two conditions.
18

19
20
21 The GLM having cardiac output as the dependent variable yielded a marginally significant
22 effect of Induction, $F(2,100) = 2.68, p = .07; \eta^2 = .05$ with larger increases in cardiac output during
23 problem solving compared to perseverative cognition and mind wandering ($ps < .02$) and during
24 mind wandering compared to perseverative cognition ($p = .08$).
25

26
27
28 As for total peripheral resistance, significant effects of Time, $F(1,50) = 5.97, p = .02; \eta^2 = .11$,
29 Induction $F(2,100) = 26.53, p < .0001; \eta^2 = .35$, and Time x Induction interaction, $F(2,100) = 3.93,$
30 $p = .02; \eta^2 = .07$ emerged. Post hoc comparisons showed that total peripheral resistance increased
31 more during perseverative cognition compared to mind wandering and problem solving ($ps < .0001$)
32 with no differences between the last two conditions. A marginally significant result emerged for
33 the recovery phase, with higher total peripheral resistance after perseverative cognition compared to
34 the other two inductions ($p = .068$).
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Hemodynamic Profile and Compensation Deficit

Figure 2 shows the relationship between hemodynamic profile and compensation deficit scores for the different tasks. No significant sex differences emerged in the hemodynamic profile induced by the different experimental conditions. One-sample *t*-tests for hemodynamic profile and compensation deficit indicated that a vascular profile was produced by the perseverative cognition induction, $t(55) = 3.85, p < .0001$, whereas the problem-solving condition evoked a myocardial profile, $t(55) = -3.35, p < .0001$, and no hemodynamic response occurred during the mind wandering induction, $t(55) = 0.02, p = .98$. A significant compensation deficit emerged during both the perseverative cognition ($t(55) = 6.64, p < .0001$), and the problem solving ($t(55) = 5.86, p < .0001$) -but not during the mind wandering- inductions providing a potential explanation for the increase in BP that characterized these two tasks.

When recovery after the perseverative cognition induction was examined, no hemodynamic response seemed to occur in the transition from reactivity to recovery ($t(55) = -0.42, p = .68$ and $t(55) = 0.57, p = .57$ for hemodynamic profile and compensation deficit, respectively), indicating that the vascular profile provoked by the induction did not change during the subsequent recovery period. A mixed profile characterized the transition from reactivity to recovery periods for both mind wandering ($t(55) = 0.06, p = .96$ and $t(55) = 2.90, p = .005$ for hemodynamic profile and compensation deficit, respectively) and problem solving ($t(55) = 1.03, p = .31$ and $t(55) = 3.14, p = .003$ for hemodynamic profile and compensation deficit, respectively).

VAS

To control for violations of sphericity degrees of freedom were corrected using Greenhouse-Geisser estimates of epsilon.

As shown in Figure 3 and Table 2, self-reported levels of mind wandering, rumination, and worry confirmed effectiveness of the experimental manipulations. For the GLM having Mind wandering as the dependent variable, main effects of Time, $F(1,50) = 29.08, p < .0001, \eta^2 = .37$,

1 and Induction $F(2,100) = 7.57, p = .005, \epsilon = .60, \eta^2 = .13$, were qualified by an interaction
2 between Time and Induction, $F(2,100) = 21.71, p < .0001, \epsilon = .82, \eta^2 = .30$. A significant
3 main effect of Task Order, $F(5,50) = 3.19, p = .01, \eta^2 = .24$ and a Task Order x Induction
4 interaction, $F(10,100) = 3.25, p = .001, \eta^2 = .25$ also emerged. As to Rumination, significant main
5 effects of Time, $F(1,50) = 14.65, p < .0001, \eta^2 = .23$, and Induction, $F(2,100) = 89, p < .0001$,
6 $\epsilon = .54, \eta^2 = .64$, and Time x Induction interaction ($F(2,100) = 84.51, p < .0001, \epsilon =$
7 $.74, \eta^2 = .63$) emerged. For Worry, the GLM yielded a main effect of Induction, $F(2,100) = 26.31,$
8 $p < .0001, \epsilon = .55, \eta^2 = .35$ and a significant Time x Induction interaction, $F(2,100) = 46.83, p$
9 $< .0001, \epsilon = .77, \eta^2 = .48$. Figure 3 illustrates significant post hoc comparisons.

10
11
12
13
14
15
16
17
18
19
20
21
22 As to self-reported mood (see Table 2 for means and standard deviations and Figure 3 for
23 significant post-hoc results), the GLM having Angry as a dependent variable revealed main effects
24 of Time, $F(1,50) = 25.32, p < .0001, \eta^2 = .34$, and Induction $F(2,100) = 42.52, p < .0001, \epsilon =$
25 $.54, \eta^2 = .46$, qualified by an interaction between Time and Induction, $F(2,100) = 25.52, p < .0001,$
26 $\epsilon = .76, \eta^2 = .34$.

27
28
29
30
31
32
33
34 For Anxious, a significant effect of Induction, $F(2,100) = 5.46, p = .01, \epsilon = .55, \eta^2 = .10$,
35 and a Time x Induction interaction, $F(2,100) = 14.11, p < .0001, \epsilon = .81, \eta^2 = .22$ emerged,
36 with higher increases during perseverative cognition compared to the other two inductions.
37
38

39
40
41 No significant effects emerged for Tired.

42
43
44 The GLM having Happy as a dependent variable revealed main effects of Time, $F(1,50) =$
45 $12.28, p = .001, \eta^2 = .20$, and Induction $F(2,100) = 14.60, p < .0001, \epsilon = .55, \eta^2 = .23$,
46 qualified by an interaction between Time and Induction, $F(2,100) = 4.80, p = .01, \epsilon = .79, \eta^2$
47 $= .09$.

48
49
50
51
52
53 As to Sad, the GLM revealed main effects of Time, $F(1,50) = 6.98, p = .01, \eta^2 = .12$, Induction
54 $F(2,100) = 26.25, p < .0001, \epsilon = .60, \eta^2 = .34$, and Time x Induction interaction, $F(2,100) =$
55 $21.92, p < .0001, \epsilon = .73, \eta^2 = .31$.

56
57
58
59
60 Overall, participants reported to be angrier, sadder, and more anxious during the perseverative
61
62
63
64
65

cognition induction compared to the mind wandering and problem solving inductions.

Hierarchical Regression

Given the high correlation between RRS and CESD scores ($r = .81; p < .0001$) and between PSWQ and STAI ($r = .77; p < .0001$) scores, only STAI, CESD, and STAXI-In were included as predictors, to prevent multicollinearity. For the same reason, only Sad and Anxious (correlations with Rumination: $r = .31; p = .02$ and $r = .50; p < .0001$, respectively) were included in the models.

Results from the hierarchical regression for the prediction of hemodynamic profile during perseverative cognition did not reveal a significant relationship between sex and the dependent variable ($\beta = .08; p = .65; R^2 = .006$). Momentary mood changes significantly added to the prediction with a higher anxiety response to the perseverative cognition induction being a significant predictor of a more vascular profile during perseverative cognition ($\beta = .41; p = .04; R^2 = .10$). Dispositional traits further added to the prediction of hemodynamic profile during perseverative cognition, particularly with higher Anger-In scores being associated with a more vascular profile during perseverative cognition, $\beta = .35; p = .04; R^2 = .28$.

Neither Sex ($\beta = -.05; p = .77; R^2 = .01$), nor momentary mood changes (Sad, $\beta = -.13; p = .47$; Anxious, $\beta = .08; p = .67; R^2 = .02$) were significant predictors of hemodynamic profile during problem solving. Among dispositional traits, higher Anger-In scores significantly predicted a more vascular profile during problem solving, $\beta = .38; p = .04; R^2 = .16$.

Neither Sex ($\beta = .15; p = .35; R^2 = .02$), nor momentary mood changes (Sad, $\beta = .04; p = .84$; Anxious, $\beta = -.02; p = .93; R^2 = .02$), nor dispositional traits, (STAI, $\beta = .21; p = .50$; CES-D, $\beta = .05; p = .87$; STAXI-In, $\beta = .08; p = .65; R^2 = .06$) were significantly associated with hemodynamic profile during mind wandering.

Results did not change when the analyses were performed replacing STAI and CES-D with PSWQ and RRS, with these two variables not being significant predictors in any of the examined hierarchical regression models. Similar results were also obtained if Anxious and Sad were

1 replaced by Rumination and Worry, with higher levels of Rumination predicting a more vascular
2 profile during the perseverative cognition ($\beta = .45$; $p = .01$; $R^2 = .11$) and the problem solving
3 induction ($\beta = .31$; $p = .03$; $R^2 = .07$). Worry did not significantly add to the predictions (see the
4 Limitation section for a possible explanation).
5
6
7
8

9 The absence of excessive multicollinearity was suggested by variance inflating factors not
10 substantially greater than one and tolerance well above 0.2.
11
12
13
14
15

16 **Discussion**

17
18 The present findings supported our hypotheses showing that perseverative cognition was
19 associated with the same BP reactivity as more functional forms of repetitive thinking, but was
20 uniquely characterized by a more vascular hemodynamic profile and (subsequently) delayed
21 recovery.
22
23
24
25
26
27

28
29 Previous studies similarly showed that problem solving and worry elicited a cardiovascular
30 reactivity of the same magnitude [5]. The present investigation extended such results with the
31 inclusion of mind wandering as a comparison condition. Some authors previously reported an
32 association between episodes of mind wandering and increases in heart rate, skin conductance [6,
33 7], and enhancement of the blink reflex [8]. However, when mind wandering was directly
34 compared to or differentiated from perseverative cognition, its association with physiological
35 reactivity disappeared both in laboratory [38] and in ambulatory studies [39-41]. These findings are
36 underscored by the lower BP responses during the mind wandering induction in the current study.
37
38
39
40
41
42
43
44
45
46
47

48 Furthermore, we examined the hemodynamic correlates of functional and dysfunctional forms
49 of repetitive thinking. Even if that was not the main objective of a previous study, the hemodynamic
50 profile of angry rumination in comparison with a series of stressful tasks including a logical-
51 mathematical task (which can be viewed as a form of problem solving) has been reported [10].
52
53
54
55
56
57
58 These findings are in line with our current results, in the way that the authors report a more vascular
59 profile during rumination and a mixed profile during the logical-mathematical task. Here, we
60
61
62
63
64
65

1 consistently found a vascular profile during perseverative cognition and a more myocardial profile
2 during problem solving, as well as a mixed profile during mind wandering.
3

4 The present findings are in line with the Obrist's distinction of active versus passive coping
5 [42]. In the Obrist view, active coping, which refers to an individual's attempts to exert personal
6 control over environmental events, leads to significant beta-adrenergic influence on myocardial
7 responses. However, such beta-adrenergic reactions become attenuated in situations that offer little
8 opportunity to exercise instrumental control (i.e., passive coping), in which a significant vascular
9 response is instead elicited. Problem solving can be considered as an example of active coping
10 whereas perseverative cognition can be representative of passive coping. In our opinion, due to the
11 nature of our tasks, it is possible to exclude that our results are due to quantitative differences in
12 mental effort [5]. In terms of mental engagement, problem solving may be viewed as the most
13 effortful condition but mind wandering (the default mode of operation of our brain) would be the
14 least effortful or equal to perseverative cognition, making it difficult to interpret the present
15 cardiovascular differences between these two experimental conditions solely in terms of mental
16 effort. As a limit, this argument has not been demonstrated in this study in any empirical manner.
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35

36 Drawing on Obrist's theory [42], Blascovich developed the biopsychosocial model of challenge
37 and threat, according to which perceived challenge versus perceived threat reliably result in distinct
38 patterns of physiological changes [43]. In this model, challenge is characterized by higher cardiac
39 output and lower total peripheral resistance (a pattern similar to that taking place during aerobic
40 exercise), whereas threat is characterized by the opposite pattern (i.e., higher total peripheral
41 resistance and lower cardiac output). This view nicely fits with our finding of higher state anxiety
42 being a significant predictor of a more vascular profile during perseverative cognition. In the
43 Blascovich view, the threat cardiovascular pattern, which is characterized by arterial constriction
44 rather than dilation, can result in strain on the coronary arteries, leading to damage, scarring, plaque
45 deposits, and eventually ischemic heart disease or, if prolonged or repeated over time, hypertension.
46 Indeed persistent or excessive vasoconstriction is a pathognomonic indicator of hypertension [9, 12,
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

13].

1
2 Keeping in mind the limitation that a 5-minute recovery period may not have been adequate in
3
4 length [16], the second core result of the present study suggests that another crucial difference
5
6 between functional and dysfunctional forms of repetitive thinking relies on the duration of the
7
8 concomitant physiological activation. Perseverative cognition was in fact not only characterized by
9
10 an increase in BP during the induction itself but also by a lack of BP recovery at the end of the task.
11
12 Such sustained cardiovascular activation is not surprising as self-reported levels of rumination and
13
14 worry during the recovery period suggest that participants were not able to stop perseverative
15
16 cognition when asked to do so. This is of particular interest in light of evidence that hypertension is
17
18 characterized by both elevated total peripheral resistance and delayed recovery [9, 44, 45]. It has
19
20 further been suggested that recovery impairments may be among the earliest precursors to the
21
22 development of essential hypertension in normotensive subjects at genetic risk of hypertension [46].
23
24
25
26
27

28
29 The effects of perseverative cognition on mood are one of the most well-replicated findings in
30
31 this field [47] and do not need to be further commented.
32

33
34 When we looked at possible associations with dispositional traits, we found that a more
35
36 vascular hemodynamic profile during perseverative cognition was predicted by higher levels of
37
38 dispositional anger-in. Delayed recovery following anger provocation has been previously
39
40 described and specifically linked to rumination [39-41]. More specifically, suppression of anger
41
42 expression (i.e., anger-in) was specifically related to high BP, atherosclerosis, and delayed recovery
43
44 [48-52]. A previous study found an association between anger-in levels and baroreceptor
45
46 sensitivity during anger rumination [53], which is in line with the delayed recovery of BP that was
47
48 seen in the present study after the perseverative cognition induction. Lastly, a vascular
49
50 hemodynamic profile during angry rumination has been reported elsewhere [10], enhancing the
51
52 robustness of the present results.
53
54
55
56
57

58
59 Some limitations need to be mentioned. First, the sample size was relatively small and may not
60
61 have been adequate in some of the comparisons. Second, with a recovery period of adequate
62
63
64
65

1 duration we might have been able to capture the time needed for BP to fully recover after the
2 perseverative cognition induction. Third, we experimentally induced perseverative cognition, mind
3 wandering, and problem solving in the laboratory, whereas in the real world spontaneity is an
4 intrinsic feature of these cognitive processes. This is particularly true for the mind wandering
5 induction, whose investigation is commonly challenged by the lack of direct experimental control
6 and its covert nature [54]. Among the methods generally used to investigate mind wandering, we
7 preferred retrospective report to experience sampling to avoid the risk of altering the natural
8 dynamics of the experience by periodically disrupting it [54]. A growing body of research
9 employing resting state imaging measures and retrospective reports of mind wandering indicate that
10 -in the absence of tasks requiring deliberative processing- the mind tends to wander [55, 56]. The
11 retrospective report of mind wandering employed at the end of the induction has the advantage to
12 preserve the integrity of time-course data and has been proven to be particularly suited to relate
13 mind wandering to its physiological signatures, as suggested by studies using pupillometry [57] and
14 EEG [58]. Directly related to this point, a further limitation of the present study is that we relied on
15 post-task subjective reports (VAS) as the only measure of effectiveness for our inductions.

16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36 Replication studies should include more objective measures of the distinct ongoing mental
37 activities, such as EEG and test the possibility that rumination and worry have their own unique
38 hemodynamic signatures. Our exploratory analysis seems to suggest that state rumination is a
39 better predictor of hemodynamic profile than worry. This result should be, however, interpreted
40 with caution especially considering that the Italian meaning of the terms used to measure state
41 rumination and state worry are not exactly the same as in English. In Italian, the distinction
42 between these two words does not exist in daily language; moreover, the word rumination
43 encompasses threat (as confirmed with the significant correlation with state anxiety), whereas worry
44 has a much milder meaning in terms of the evoked emotion compared to English. Lastly, despite
45 the strength of the within-subject design used, and the counterbalancing of the order of the three
46 inductions, and they may still have influenced each other in a non-natural way, each of them either
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 enhancing or flattening the response to another. Our methodological approach could reduce the
2 strength of the emotion experienced if some emotional episodes are not recalled as reliably and if
3 the emotions are not relived as vividly.
4
5

6
7 Limitations notwithstanding, the present study is clinically relevant in that it provides further
8 insights into the consequences of perseverative cognition for cardiovascular risk, furnishing
9 information on its hemodynamics compared to more functional forms of repetitive thinking.
10
11

12 Obviously, replication with a larger sample size, a wider range of inductions, and a longer recovery
13 period is needed to test the robustness of the present findings.
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Table1 Sex differences at baseline

	Women (<i>n</i> = 26)	Men (<i>n</i> = 30)	<i>t</i> / χ^2
Age (years)	24.2 (3.4)	24.7 (4.5)	0.41
BMI (Kg/m ²)	21.9 (2.7)	22.7 (4.1)	0.80
Education	11 L, 14 M, 1 H	13 L, 15 M, 2 H	0.25
SBP (mmHg)	125.7 (23.4)	133.7 (18.3)	1.44
DBP (mmHg)	74.5 (13.2)	81.8 (11.9)	2.18*
CO (L/min)	5.2 (1)	5.7 (1.2)	1.87
TPR (dyn/cm ² /s)	1192.3 (231)	1152.4 (149.7)	0.78
Nicotine	16 N, 10 Y	24 N, 6 Y	2.33
RRS	51.5 (13.4)	42.7 (11.1)	2.17*
PSWQ	43.3 (10.3)	54.2 (11.9)	2.65*
STAI	49.1 (10.6)	39.3 (8.9)	3.12*
CES-D	43.3 (10.3)	34.5 (8.6)	2.92*

Note. BMI = Body Mass Index; SBP = Systolic Blood Pressure; DBP = Diastolic Blood Pressure;

CO = Cardiac Output; TPR = Total Peripheral Resistance; RRS = Ruminative Response Scale;

PSWQ = Penn State Worry Questionnaire; STAI = State Trait Anxiety Inventory; CES-D = Center for Epidemiologic Studies Depression Scale.

* $p < 0.05$

Table 2 Means and standard deviations of VAS scores during the three experimental conditions (reactivity) and subsequent recovery periods

	Perseverative Cognition		Mind Wandering		Problem Solving	
	Reactivity	Recovery	Reactivity	Recovery	Reactivity	Recovery
Ruminating	44.6 (33.4)	41.2 (35.0)	-85.8 (60.4)	-44.6 (33.4)	-7.9 (24.2)	6.9 (27.2)
Worrying	24.2 (35.1)	29.9 (34.5)	-54.1 (62.9)	-24.2 (35.1)	-19.1 (28.4)	16.0 (29.7)
Wandering	-14.2 (36.7)	26.6 (39.3)	40.8 (70.1)	14.2 (36.7)	-9.5 (26.9)	26.8 (35.8)
Happy	-12.6 (21.4)	3.8 (18.9)	16.4 (34.29)	12.6 (21.4)	-6.7 (21.9)	0.5 (15.7)
Sad	20.2 (27.2)	-12.6 (26.3)	-32.8 (47.1)	-20.2 (27.2)	-2.1 (19.7)	1.8 (11.9)
Tired	3.9 (29.3)	4.5 (17.4)	0.6 (38.8)	-3.9 (29.3)	1.1 (19.6)	3.4 (14.9)
Anxious	9.9 (28.2)	-12.5 (23.3)	-22.4 (45.0)	-9.9 (28.2)	-7.2 (23.9)	1.7 (17.1)
Angry	22.0 (25.4)	-13.3 (23.5)	-35.3 (43.6)	-22.0 (25.4)	6.1 (20.3)	0.4 (16.5)

References

1. Watkins ER. Constructive and unconstructive repetitive thought. *Psychol Bull.* 2008; 134: 163-206.
2. Brosschot JF, Gerin W, Thayer JF. The perseverative cognition hypothesis: A review of worry, prolonged stress-related physiological activation, and health. *J Psychos Res.* 2006; 60: 113-124.
3. Brosschot JF, Verkuil B, Thayer JF. Conscious and unconscious perseverative cognition: Is a large part of prolonged physiological activity due to unconscious stress? *J Psychosom Res.* 2010; 69: 407-416.
4. Ottaviani C, Thayer JF, Verkuil B, et al. Physiological concomitants of perseverative cognition: a systematic review and meta-analysis. *Psychol Bull.* 2016; 142: 231-259.
5. Verkuil B, Brosschot JF, Borkovec TD, Thayer JF. Acute autonomic effects of experimental worry and cognitive problem solving: Why worry about worry? *Int J Clin Health Psychol.* 2009; 9: 439-453.
6. Smallwood J, Davies JB, Heim D, et al. Subjective experience and the attentional lapse: Task engagement and disengagement during sustained attention. *Conscious Cog.* 2004; 13: 657-690.
7. Smallwood J, O'Connor RC, Sudberry MV, Haskell C, Ballantyne C. The consequences of encoding information on the maintenance of internally generated images and thoughts: the role of meaning complexes. *Conscious Cog.* 2004; 13: 789-820.
8. Smilek D, Carriere JS, Cheyne JA. Out of mind, out of sight: Eye blinking as indicator and embodiment of mind wandering. *Psychol. Sci.* 2010; 21: 786-789.
9. Julius S. Transition from high cardiac output to elevated vascular resistance in hypertension. *Am Heart J.* 1988; 116: 600-606.
10. Ottaviani C, Shapiro D, Goldstein IB, James JE, Weiss R. Hemodynamic profile, compensation deficit, and ambulatory blood pressure. *Psychophysiology* 2006; 43: 46-56.
11. Gregg ME, James JE, Matyas TA, Thorsteinsson EB. Hemodynamic profile of stress-induced

anticipation and recovery. *Int J Psychophysiol.* 1999; 34: 147-162.

12. Mayet J, Hughes A. Cardiac and vascular pathophysiology in hypertension. *Heart.* 2003; 89: 1104-1109.
13. Sung BH, Lovallo WR, Pincomb GA, Wilson MF. Cardiac adaptation to elevated blood pressure in borderline hypertensive men. *Am J Cardiol.* 1993; 72: 407-412.
14. Panaite V, Salomon K, Jin A, Rottenberg J. Cardiovascular recovery from psychological and physiological challenge and risk for adverse cardiovascular outcomes and all-cause mortality. *Psychosom Med.* 2015; 77: 215-226.
15. Ottaviani C, Shapiro D, Goldstein IB, Mills PJ. Vascular profile, delayed recovery, inflammatory process, and ambulatory blood pressure: Laboratory-to-life generalizability. *Int J Psychophysiol.* 2007; 66: 56-65.
16. Steptoe A, Marmot M. Impaired cardiovascular recovery following stress predicts 3-year increases in blood pressure. *J Hypertens.* 2005; 23: 529-536.
17. Ring C, Burns VE, Carroll D. Shifting hemodynamics of blood pressure control during prolonged mental stress. *Psychophysiology* 2002; 39: 585-590.
18. Rutledge T, Hogan BE. A quantitative review of prospective evidence linking psychological factors with hypertension development. *Psychosom Med.* 2002; 64: 758-766.
19. Jennings JR, Kamarck T, Stewart C, Eddy M, Johnson P. Alternate cardiovascular baseline assessment techniques: Vanilla or resting baseline. *Psychophysiology.* 1992; 29: 742-750
20. Eckert S, Horstkotte D. Comparison of Portapres non-invasive blood pressure measurement in the finger with intra-aortic pressure measurement during incremental bicycle exercise. *Blood Press Monit.* 2002; 7: 179-183.
21. Gregg ME, Matyas TA, James JE. A new model of individual differences in hemodynamic profile and blood pressure reactivity. *Psychophysiology* 2002; 39: 64-72.
22. Guyton A. *Human physiology and mechanisms of disease* (4th ed). Philadelphia: W.B.

Saunders; 1987.

23. Nolen-Hoeksema S, Morrow J. A prospective study of depression and posttraumatic stress symptoms after a natural disaster: the 1989 Loma Prieta Earthquake. *J Pers Soc Psychol.* 1991; 61:115-121.
24. Nolen-Hoeksema S, Parker LE, Larson J. Ruminative coping with depressed mood following loss. *J Pers Soc Psychol.* 1994; 67:92-104.
25. Nolen-Hoeksema S, Davis CG. "Thanks for sharing that": ruminators and their social support networks. *J Pers Soc Psychol.* 1999; 77:801-14.
26. Meyer TJ, Miller ML, Metzger RL, Borkovec TD. Development and validation of the Penn State Worry Questionnaire. *Behav. Res. Ther.* 1990; 28: 487-495.
27. Stober J. Reliability and validity of two widely-used worry questionnaires: self-report and self-peer convergence. *Pers Individ Dif.* 1998; 24: 887-890.
28. Molina S, Borkovec TD. The Penn State Worry Questionnaire: psychometric properties and associated characteristics. In: Davey GCL, Tallis F, eds. *Worrying: perspectives on theory, assessment and treatment.* New York, NY: Wiley; 1994: 265-283.
29. Brown TA, Marten PA, Barlow DH. 1995. Discriminant validity of the symptoms comprising the DSM-III-R and DSM-IV associated symptom criterion of Generalized Anxiety Disorder. *J Anxiety Disord.* 1995; 9: 317-328.
30. Spielberger CD, Gorsuch RL, Lushene RE. *STAI Manual.* Palo Alto, CA: Consulting Psychologists Press; 1970.
31. Spielberger CD. *State-Trait Anxiety Inventory: Bibliography (2nd ed.).* Palo Alto, CA: Consulting Psychologists Press, 1989.
32. Spielberger CD, Gorsuch RL, Lushene R, Vagg PR, Jacobs GA. *Manual for the State-Trait Anxiety Inventory.* Palo Alto, CA: Consulting Psychologists Press, 1983.
33. Radloff LS. The CES-D scale: a self-report depression scale for research in the general population. *Appl. Psychol. Meas.* 1977; 1: 385-401.

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
34. Carleton RN, Thibodeau MA, Teale MJN, Welch PG, Abrams MP, Robinson T, Asmundson GJ. (2013) The Center for Epidemiologic Studies Depression Scale: A Review with a Theoretical and Empirical Examination of Item Content and Factor Structure. *PLoS One* 2013; 8: e58067.
 35. Spielberger CD, Johnson EH, Russell SF, Crane RJ, Jacobs GA, Worden TJ. The experience and expression of anger: construction and validation of an Anger Expression Scale. In: Chesney MA, Rosenman RH, eds. *Anger and hostility in cardiovascular and behavioral disorders*. Washington DC: Hemisphere; 1985: 5-30.
 36. Spielberger CD, Reheiser EC. Measuring anxiety, anger, depression, and curiosity as emotional states and personality traits with the STAI, STAXI, and STPI. In: Hersen M, Segal DL, Hilsenroth M, eds. *Comprehensive Handbook of Psychological Assessment (Vol. 2): Personality Assessment*. New York: Wiley; 2004: 74-80.
 37. James JE, Gregg ME. Hemodynamic effects of dietary caffeine, sleep restriction, and laboratory stress. *Psychophysiology* 2004; 41: 914-923.
 38. Ottaviani C, Shapiro D, Couyoumdjian A. Flexibility as the key for somatic health: from mind wandering to perseverative cognition. *Biol Psychol.* 2013; 94: 38-43.
 39. Ottaviani C, Couyoumdjian A. Pros and cons of a wandering mind: a prospective study. *Front Psychol.* 2013; 4: 524, DOI: 10.3389/fpsyg.2013.00524.
 40. Ottaviani C, Medea B, Lonigro A, Tarvainen M, Couyoumdjian A. Cognitive rigidity is mirrored by autonomic inflexibility in daily life perseverative cognition. *Biol Psychol.* 2015; 107: 24-30.
 41. Ottaviani C, Shahabi L, Tarvainen M, Cook I, Abrams M, Shapiro D. Cognitive, behavioral, and autonomic correlates of mind wandering and perseverative cognition in major depression. *Front Neurosci.* 2015; 8: 433. doi: 10.3389/fnins.2014.00433
 42. Obrist PA. *Cardiovascular psychophysiology*. New York: Plenum; 1981.
 43. Blascovich J. Challenge, threat, and health. In: Shah JY, Gardner WL, eds. *Handbook of*

Motivation Science. New York: Guilford; 2008: 481-493.

- 1
2 44. Schuler JL, O'Brien WH. Cardiovascular recovery from stress and hypertension risk factors: a
3
4 meta-analytic review. *Psychophysiology*. 1997; 34: 649-659.
5
6
- 7 45. Steptoe A, Donald AE, O'Donnell K, Marmot M, Deanfield JE. Delayed blood pressure
8
9 recovery after psychological stress is associated with carotid intima-media thickness: Whitehall
10
11 psychobiology study. *Arterioscler Thromb Vasc Biol*. 2006; 26: 2547-2551.
12
13
- 14 46. Schneider GM, Jacobs DW, Gevirtz RN, O'Connor DT. Cardiovascular haemodynamic
15
16 response to repeated mental stress in normotensive subjects at genetic risk of hypertension:
17
18 Evidence of enhanced reactivity, blunted adaptation, and delayed recovery. *J Hum Hypertens*.
19
20 2003; 17: 829-840.
21
22
- 23 47. Nolen-Hoeksema S, Wisco BE, Lyubomirsky S. Rethinking rumination. *Perspect Psychol Sci*.
24
25 2008; 3: 400-424.
26
27
- 28 48. Gerin W, Davidson KW, Christenfeld NJ, Goyal T, Schwartz JE. The role of angry rumination
29
30 and distraction in blood pressure recovery from emotional arousal. *Psychosom Med*. 2006; 68:
31
32 64-72.
33
34
- 35 49. Ottaviani C, Shapiro D, Fitzgerald L. Rumination in the laboratory: What happens when you go
36
37 back to everyday life? *Psychophysiology*. 2011; 48: 453-461.
38
39
- 40 50. Suchday S, Carter MM, Ewart CK, Larkin KT, Desiderato O. Anger cognitions and
41
42 cardiovascular recovery following provocation. *J Behav Med*. 2004; 27: 319-341.
43
44
- 45 51. al'Absi M, Bongard S. Neuroendocrine and behavioral mechanisms mediating the relationship
46
47 between anger expression and cardiovascular risk: Assessment considerations and
48
49 improvements. *J Behav Med*. 2006; 29: 573-591.
50
51
- 52 52. Vella EJ, Friedman BH. Hostility and anger in: Cardiovascular reactivity and recovery to mental
53
54 arithmetic stress. *Int J Psychophysiol*. 2009; 72: 253-259.
55
56
- 57 53. Ottaviani C, Shapiro D, Davydov DM, Goldstein IB, Mills PJ. The autonomic phenotype of
58
59 rumination. *Int J Psychophysiol*. 2009; 72: 267-275.
60
61
62
63
64
65

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
54. Smallwood J, Schooler JW. The science of mind wandering: empirically navigating the stream of consciousness. *Annu Rev Psychol.* 2015; 66: 487-518.
55. Fox KC, Spreng RN, Ellamil M, Andrews-Hanna JR, Christoff K. The wandering brain: meta-analysis of functional neuroimaging studies of mind-wandering and related spontaneous thought processes. *Neuroimage.* 2015; 111: 611-621.
56. Mason MF, Norton MI, Van Horn JD, Wegner DM, Grafton ST, Macrae CN. Wandering minds: the default network and stimulus-independent thought. *Science* 2007; 315: 393-395.
57. Smallwood J, Brown KS, Baird B, Mrazek MD, Franklin MS, Schooler JW. Insulation for daydreams: a role for tonic norepinephrine in the facilitation of internally guided thought. *PLoS One* 2012; 7:e33706.
58. Barron E, Riby LM, Greer J, Smallwood J. Absorbed in thought: the effect of mind wandering on the processing of relevant and irrelevant events. *Psychol Sci.* 2011; 22: 596-601.

Figure captions

1
2 Figure 1. Systolic and diastolic blood pressure reactivity and recovery for each experimental
3
4 condition.
5

6
7 Note. Error bars represent standard deviations of the mean. PC = Perseverative Cognition; MW =
8
9 Mind Wandering; PS = Problem Solving; SBP = Systolic Blood Pressure; DBP = Diastolic Blood
10
11 Pressure.
12
13

14
15
16
17 Figure 2. Scatterplots for hemodynamic profile and compensation deficit during each task. A
18
19 “more vascular” profile is associated with more positive values along the hemodynamic profile
20
21 axis and a “more myocardial” profile is associated with more negative values along the
22
23 hemodynamic profile axis. A “higher deficit” in compensating is associated with more positive
24
25 values on the compensation deficit axis and a “lower deficit” in compensating is associated with
26
27 more negative values on the compensation deficit axis.
28
29
30

31
32
33
34 Figure 3. VAS reactivity and recovery change scores for perseverative cognition (black), mind
35
36 wandering (light grey), and problem solving (dark grey).
37

38
39 Note. Error bars represent standard deviations of the mean. * $p < .05$; ** = $p < .0001$. MW =
40
41 Mind Wandering.
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65