

2
3 Date Submitted (Original): 20-January-2014

4
5 Revised Date: 20-May-2014

6
7 Accepted Date: 02-July-2014
8
9

10
11 **Title: Displacement of bedtime by screen time in schoolchildren: the importance**
12 **of area deprivation.**

13
14
15 **Running title:** Children, late bedtime, screen time and deprivation.

16 ***Ayodele A. Ogunleye (PhD)^a, Christine Voss (PhD)^b, Gavin R. Sandercock**
17 **(PhD)^c**

18 ^aObesity Research & Management, Department of Medicine, University of Alberta,
19 87th Avenue and 112th Street, Edmonton, AB, T6G 2E1, Canada.

20 ^bCentre for Hip Health and Mobility, University of British Columbia, Vancouver,
21 Canada.

22 ^cCentre for Sports and Exercise Science, School of Biological Sciences, University of
23 Essex, Colchester, CO4 3SQ, United Kingdom.

24
25 *Correspondence to: Ayodele A. Ogunleye;

26 Email: aogunley@ualberta.ca

27 Tel: +1 780 4926641

28 Fax: +1 780 4929414

29

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as doi: 10.1111/ped.12447

30 **Abstract**

31 **Background:** Sleep duration is an important predictor of obesity and health. We aim
32 to evaluate the association between late bedtime with screen time and the role of
33 geographical deprivation in English schoolchildren.

34 **Methods:** We collected Sleeping & waking times, screen time, socio-demographic
35 data and measured body mass index in a cross-section of 1,332 (45.7% females) 11-
36 15 year old schoolchildren participating in the East of England healthy heart study.
37 Logistic regressions were used to determine the likelihood of late bedtimes in
38 schoolchildren with different screen time and from a different geographic location.
39 Mean differences were assessed either by ANOVA or t-test.

40 **Results:** About 42% of males go to bed late at night compared with 37% females.
41 When compared to those with <2hours of daily screen time, schoolchildren who
42 spend 2-4 hours on screen time were more likely 1.50(1.07 to 2.09) to sleep late at
43 night while those with > 4hours of daily screen time were most likely 1.97(1.34 to
44 2.89) to sleep late at night. Late bedtimes were associated with deprivation in
45 schoolchildren.

46 **Conclusions:** High screen time and deprivation may explain lateness in bedtime in
47 English schoolchildren. This explanation may vary according to area deprivation and
48 geographic location. Family centred interventions and parental support is important to
49 reducing screen time, late bedtimes and sleep duration.

50 **Keywords:** Bedtime; screen time; deprivation; geographic location; children.

51

52

53

54

55 **Introduction**

56 The average sleep duration of schoolchildren has declined greatly ¹, a common
57 behavioural issue brought to the attention of paediatricians ². Recent evidence

58 suggests that adequate sleep is important health behaviour, following the
59 identification of potential mechanistic pathways linking sleep with obesity^{3,4}. There
60 is evidence that insufficient amount of sleep (short sleep duration or sleep
61 deprivation) is an independent risk factor for excessive weight gain⁵, obesity⁴ and
62 cardiometabolic risk⁶. Although causative inferences cannot be made between sleep
63 and health outcomes, the decline in average sleep duration has been concurrent with
64 increases in screen time and obesity pandemic.

65 Screen time, a high prevalent behaviour among schoolchildren, is not encouraged in
66 children under two years of age⁷, and should be limited to not more than two hours
67 per day in older children^{7,8}. Recent study suggests that one in three English
68 schoolchildren may be exposed to over two hours of screen time in a day⁹. In
69 children age 3-5 years, evening (after 7pm) media use is associated with sleep
70 problem¹⁰. A study on schoolchildren from a different population, New Zealand,
71 shows that screen time shortly before bedtime delay onset of sleep¹¹. There is
72 evidence also that screen time in adults¹² and in children¹³ are associated with
73 deprivation. Despite these studies, evidence on differing screen time in English
74 children living in varying location, and that have late bedtimes is lacking. As a result
75 therefore, we aim to assess the association of late bedtime, as opposed to sleep
76 duration, with screen time in schoolchildren and whether geographical location was
77 related to late bedtime. We also assessed the importance of deprivation on the
78 association between screen time and late bedtimes.

79

80 **Methods**

81 The study participants came from the ongoing East of England Healthy Hearts Study.
82 Following approval by the University Ethical Review Committee, data were gathered
83 from 1332 (45.7% females) 11-15 year olds attending three state-run, comprehensive
84 schools, with differing area deprivation levels. One School (school 1 here-in) was
85 from a less deprived location, school 2 was from the less deprived rural location,

86 while school 3 was a deliberate booster sample to include schoolchildren from a
87 highly deprived location. All data collection occurred in the summer months of 2010
88 and 2011. We sent letters to schools in the East of England region inviting them to
89 participate in this study, and then purposefully selected a representative mix of
90 volunteer schools to take part in the study, a detail methodology has been described
91 previously⁹.

92

93 **Assessment of Bedtime**

94 Participants self-reported bedtimes by answering the following question: ‘What time
95 do you usually go to bed on school nights’. These questions were adapted from the
96 general sleep questionnaire and have been validated for use in this age group
97 previously¹⁴. Schools’ 2 and 3 have are the same opening time, while school 1 opens
98 five minutes later. Since school opening times may have an effect on the bedtime the
99 previous night, we assumed that a five minutes difference in the opening time should
100 not have a significant effect on the bedtimes. Participant’s bedtimes were classified as
101 either early- or late-bed, using median splits for age- and sex-adjusted bedtimes on a
102 school day. This method is much preferred and has been used previously¹⁵ than
103 choosing an arbitrary bedtimes. Bedtimes on weekdays (school nights’) were used
104 because it is likely to be more constant than bedtimes on weekends.

105

106 **Screen time**

107 Participants self-reported daily screen time by answering the following question:
108 ‘How much time do you spend on average each day watching television, watching
109 DVDs or videos, using a computer or games console’. Answers were given on a 0–5
110 point scale with the following answers: none, 0–30 min, 30–60 min, 1–2, 2–4 and .4
111 h. Participants were grouped according to whether they reported <2 h screen time as
112 recommended^{7, 8}, 2–4 or >4 h. The latter value is proposed as another important
113 threshold representing heavy use¹⁶.

114

115 **Body composition**

116 Participants' mass and stature were measured, to the nearest 0.1 kg and 0.1 cm,
117 respectively, wearing light clothing (T-shirts and shorts) and without shoes. Body
118 mass index (BMI) was calculated (kg/m^2) and z -scores generated using the UK 1990
119 Growth Reference which adjusts for age, sex and skewness¹⁷. We categorized BMI in
120 two ways to determine the potential effects of our method of categorization.
121 Schoolchildren BMI were categorized according to the International Obesity Task
122 force (IOTF) criteria¹⁸.

123

124 **Area-level Deprivation**

125 We obtained an area-level measure of deprivation for each participant using their
126 home postcode as detailed previously⁹. Briefly, The English Index of Multiple
127 Deprivation 2007 (IMD 2007) is measured based on the small area geographical units
128 known as Lower Super Output Areas (LSOAs); each LSOA contains between 1,000
129 and 3,000 inhabitants with an average population of 1,500 people allowing
130 identification of small pockets of deprivation by area¹⁹. In IMD 2007, there are a total
131 of 38 indicators, distributed across the seven domains of deprivation (income,
132 employment, health and disability, education, skills and training, barriers to housing
133 and services, living environment, crime)¹⁹. A low IMD score indicates affluence, and
134 a high score suggests an area of deprivation.

135

136 **Statistical analyses**

137 Binary logistic regression analysis was used to assess the relationship between
138 bedtime (dichotomous bedtime early-bed versus late-bed) was the outcome variable
139 with categorical screen time (<2h, 2-4h and >4h) as the determinant. A univariate
140 model was initially produced followed by a multivariate model controlling for: sex,
141 age, school, BMI and deprivation. The differences in area deprivation between

142 schools were carried out using analysis of variance (ANOVA) with as the *post hoc*
143 (Bonferroni) tests for multiple comparisons. Statistical analyses were performed using
144 IBM SPSS 19.0 for windows (SPSS Inc.: an IBM Company, Chicago, IL, USA).

145

146 **Results**

147 The sample included 9.7% and 22.1% obese and overweight schoolchildren
148 respectively. The proportion of those who reported 2 - 4 hours screen time daily was
149 19.7%, with 15.3% reporting >4 hours.

150 Table 1 shows the demographic characteristics of schoolchildren according to whether
151 they reported going to bed early or late. Overall, 42.2% of males reported going to
152 bed late compared with 37.3% females. Prevalence of late-bed increased with higher
153 reported screen time; 51.5% schoolchildren who spent >4 hours engaged in screen-
154 time were classed as late-bed, compared with those that spend between 2-4 hours
155 (45.8%) or less than 2hours (35.1%) screen time.

156 There was a significant difference in IMD scores between early-bed and late-bed
157 groups (mean difference in IMD score = -2.89, 95%CI: -4.80 to -0.97, p= 0.003) as
158 shown in table 2. There was no significant difference (p>0.05) in the mean BMI z-
159 score between early-bed and late-bed groups (mean difference= -0.74, 95%CI:-0.22 to
160 0.07, p=0.32).

161 Analysis of variance showed that there was a significant main effects for IMD score
162 among the three schools (F=499.7, p< 0.001). The mean IMD score in the third school
163 (34.9±14.2) was significantly different from the other two schools (p<0.05). Mean
164 IMD score was not different (p>0.05) between the first (13.3±7.59) and second school
165 (12.8±5.89).

166

167 **Adjusted and unadjusted likelihood of late bedtime**

168 Adjusted for age, sex, school, deprivation and weight status, the odds ratios for late
169 night sleeping (i.e., > bedtimes greater than median splits for age- and sex-adjusted
170 bedtimes) were 1.00 for screen time <2 hours (reference category), 1.50(1.07 to 2.09)
171 for 2-4 hours screen time and 1.97(1.34 to 2.89) for over 4 hours of daily screen time.
172 The unadjusted odds ratios for screen time were very similar to these values (table 3).

173 Late bedtime may be common in schoolchildren from a more deprived location
174 according to IMD. Where school 1 was the reference category, the unadjusted odds
175 ratios for late bedtime were 2.30(1.59 to 3.32) in school 3 (in a more deprived
176 location) and 1.31(0.89 to 1.94) in school 2. When we adjusted for age, sex, school,
177 deprivation and weight status, the odds ratios for late-bed were 1.66(0.96 to 2.85) and
178 1.12(0.73 to 1.74) in school 3 and school 1 respectively. Age was associated with late
179 night sleeping, but not in a linear manner. Compared to 11 years old, 12 years old
180 schoolchildren were over 2 times (2.19(1.46 to 3.27)) more likely to go to bed late at
181 night; while the adjusted odds ratio was in 1.58(0.93 to 2.71) in the 15 years old.

182

183 **Influence of deprivation**

184 When accounting for school (already an area-level factor), adjusting for deprivation
185 had very little influence on the association between late sleeping and screen time.

186 Deprivation, using IMD 2007, seemed not to be a significant determinant of sleep
187 time in schoolchildren (table 3). However, school location may be an important
188 determinant of late sleeping in schoolchildren. These schools in our study have
189 different levels of deprivation. Schoolchildren in the most deprived school were more
190 likely (1.64(1.07 to 2.52)) to go to bed late at night than a less deprived reference
191 category school.

192 **Discussion**

193 This study shows that late bedtime habits are associated with shorter total sleep
194 duration in children, especially during schooldays. This is the first study comparing
195 bedtimes in English schoolchildren of different deprivation categories. Screen time
196 displaces physical activity and may also displace bedtime; both factors are important
197 determinants of weight status and obesity as shown in figure 1. Schoolchildren who
198 report >2 hours daily screen time were more likely to go to bed at a time deemed late
199 at night. Those who live in deprived area were twice as likely to report late bedtimes.
200 In order to improve sleep duration, screen time (evening screen time) should be
201 reduced. Bed times also are different in schoolchildren living in different geographic
202 locations and areas with different levels of deprivation. The proportion of children
203 reporting late bedtime may be as high as 45% in more deprived schools, more
204 common than in more affluent ones. Sleep duration has been linked with childhood
205 obesity in previous studies^{20,21}, but the present study found that obese and
206 overweight schoolchildren were no more likely to report late bedtimes than those of
207 normal weight.

208
209 There are multiple reasons for insufficient sleep, including: insomnia (a sleep
210 disorder), stressors such as preparation for examinations as well as excessive screen
211 time. These reasons in adults may differ from that of children. But addressing the
212 behavioural reasons/causes for sleep deprivation, not insomnia, may be important in
213 combating obesity pandemic.

214
215 Few studies have examined the association between screen time and late bedtime in
216 schoolchildren. In fact there is little data on English schoolchildren with high screen
217 time⁹. Those that have examined the association between obesity and screen time
218 have done so either in adults²², indirectly²³ or in populations^{24,25} likely to accumulate
219 lower daily screen time than the present population. The only one of these studies that

220 examined the influence of socioeconomic status or deprivation, did so in a population
221 with high socioeconomic status ²⁵.

222 In 4 to 13 year old Dutch children ²⁵ short sleep duration was associated with being
223 overweight. Short sleep duration was determined by late bedtimes and was strongly
224 associated with higher screen-time. We understand that late bedtime and sleep
225 duration are different constructs; and that late sleeping may be associated with sleep
226 duration especially on a school weekday ²⁵. More recent findings also show that, the
227 bedtimes of schoolchildren may be important in addition to total sleep duration ¹⁵.

228 Though this sample of their study is of a different population with a low study
229 response rate ¹⁵ compared to that of our study, sleeping pattern was associated with
230 physical activity levels, screen time and weight status in schoolchildren.

231 Previously we observed that, Age-specific prevalence for >2 h daily screen time
232 increased at around 13 years of age⁹.

233 The proportion of schoolchildren with daily screen time >2 h rises sharply at 13 years
234 of age, while the duration of sleep start falling during this age. There are other studies
235 that have reported a similar increase in screen time⁹ or late bedtime²⁶ at this age,
236 possibly due to an increase in computer use for educational purposes at this age. Such
237 increases may, however, also be associated with 13 years being the lower age limit for
238 registration on a number of the world's most popular social networking websites
239 including FacebookTM.

240

241 Inequality may be central to the screen time – sleep time relationship. Prior studies
242 have shown that both low sleep duration and socioeconomic status were predictors of
243 obesity in schoolchildren²⁷. Also, screen time is shown to relate to obesity, and
244 previous studies in schoolchildren did not find any significant trend between
245 deprivation categories and screen time ⁹ even though socioeconomic status is related
246 to high screen time in adults, or deprived adults (defined using area deprivation)
247 engage in high screen time ¹². Deprived children are more likely to go to be late at

248 night. Our result suggests that socio economic status may be an important determinant
249 of sleep time. Parents of low socioeconomic status may be indulging in of high
250 screen time in the evening and may lack the control of reducing high or late night
251 screen time in their children. There is need for parents' guidance on the best ways of
252 preventing late bedtime and associated high levels of screen time.

253

254 Increased television viewing is associated with shorter sleep duration^{20, 28, 29}. After
255 adjusting for television viewing, these studies did not find television viewing to be
256 independently associated with either sleep duration or obesity. The obesity-sleep
257 duration relationship may be one thing and screen time-sleep duration is another and
258 may be independent. Based on the current findings, it seems that the relationship
259 between bedtimes and screen time in schoolchildren are independent of weight status.
260 Of note, is that we have measured screen time in our study with television viewing
261 inclusive among other devices and we only studied bedtimes (not sleep duration).

262

263 The present study suggests that area-level deprivation may be associated with late
264 bedtime in schoolchildren. Previous studies in schoolchildren have mainly used
265 family structure indicator such as living with a single parent and the presence of other
266 siblings, low level of parental education, or unemployment^{4, 30}; maternal education,
267 maternal work and family income³⁰. Direct associations between socioeconomic
268 status/deprivation, bedtimes and screen time in adults may be visceral, but the
269 association in schoolchildren may be indirect.

270 Parents face difficulties in making their children go to bed early, and may have to
271 undertake interactive routines³⁰ such as reading, storytelling, singing prayer, and
272 putting off the lights. Difficulty can arise because they are unsure of the appropriate
273 time to send them to bed or late night working by the parents or they sleep earlier than
274 their children. Sleeping in lounge are not uncommon in schoolchildren especially
275 sleeping with television or computer game still on. It may be difficult for parent to

276 identify the right time to send schoolchildren to sleep as some may want to study or
277 be preparing for an examination.

278

279 Previous suggestions^{4,5} favour the development and the testing behavioural
280 interventions that will improve sleeping habits. Interventions to reduce screen time in
281 schoolchildren especially in the evening and before bed are important. Family
282 regulations to reduce television viewing or other screen based devices use at a
283 particular time, to give schoolchildren ample time for sleeping may be beneficial.
284 Paediatric health professionals working with schoolchildren should also consider
285 asking about bedtime in addition to their sleep duration.

286

287 **Study strengths**

288 Our study is an improvement over studies that have used parental-reported bedtime of
289 schoolchildren. In this age group, self-reported may be better than parental-reported
290 bedtime. The relatively large sample size provides a robust support for our findings
291 presented here.

292

293 Deprivation or socioeconomic status is difficult to measure in some parts of the
294 population. An example is schoolchildren. However, area deprivation may be a better
295 indicator for schoolchildren than socioeconomic status. Both area-level deprivations
296 with 37 indicators measured through children postcode were used here and we also
297 compared three schools, which can act as a cluster, in this study that varies in location
298 and built.

299

300 **Study limitations**

301 We have not identified or separated children who may be suffering from insomnia
302 from our study; self-imposed sleep deprivation was our aim.

303 Self-reported sleep/wake and screen time habits was used and we recognised that
304 bedtimes may vary between by days, weeks and seasons. Therefore to minimise the
305 bias this might bring we have not included bedtimes on weekends with is highly
306 varied and irregular. We have used a less costly and a less stressful measure on
307 participants. We understand the possibility of social desirability and satisficing in our
308 study do to the use of questionnaire. Due to the cross sectional nature of our study, no
309 conclusions can be drawn regarding causal links or causality.

• 310 Reverse causality is also possible, may be English schoolchildren are generally late
311 bed goers, and found themselves exposed to screen as a result of that habit/behaviour
312 rather than the other way round. Randomised controlled trials and cohort studies are
313 needed to confirm a temporal relationship between screen time and bedtimes in
314 schoolchildren.

315

316 High screen time may be a factor preventing English schoolchildren from going to
317 bed early or sleeping for an adequate duration. However, some of the participants may
318 sleep late due to other factors (e.g. reading) and not due to screen based activity.

319 Future studies on sleep-obesity relationship should not only evaluate the association
320 between intermediate factors like physical activity levels and sleep duration⁴, but in
321 schoolchildren, studies should consider closely the association between sleep duration
322 and mode of transport to school (figure 1). Objective measures of what

323 schoolchildren do after school and during the time before they go to bed need to be
324 investigated more closely. Randomised controlled trials promoting earlier bed times
325 and increases sleep duration may also be effective in establishing that screen based
326 activities (rather than reading) is what is depriving English schoolchildren of adequate
327 sleeping time. Implication of switching off television and other screen based devices
328 in the home at a particular time, say 9.00pm, and how this would affect sleeping time,
329 sleep duration, late night eating and weight status would be interesting to explore.

330 Future studies should also evaluate the association between built environment and

331 screen time; especially after school hours screen time. Safer places with brilliant
332 outdoor facilities may have different screen time-bedtime pattern compared to other
333 places.

334 In agreement with previous research¹⁵ that the emphasis has been on sleep duration,
335 but that the importance of bed time may have been neglected in relation to child
336 health to date.

337 Schoolchildren's activities in the evening may be important in potential public health
338 interventions as this is the period they are likely to engage in screen based activities or
339 become sedentary. What a high sedentary time would it have been for schoolchildren
340 that sleep at midnight? Interventions that are aimed at reducing sedentary behaviours
341 or late sleeping time in schoolchildren may be tailored to evenings after schools when
342 they are at home engaging in screen/media use and are not in parks or on bed.

343

344 **Conclusion**

345 Previous studies have shown that short sleep duration is associated with physical
346 inactivity³¹ and high caloric intake⁵. The findings from this study suggest that high
347 screen time and deprivation may explain lateness in bedtime in schoolchildren and
348 possibly in turn sleep duration on schooldays. Interventions that support family rules
349 and support for parents may be effective in combating high screen time, late bedtime,
350 short sleep duration and obesity in English schoolchildren.

351 It is intuitive to suggest that sleep deprivation or duration may be improved by
352 reducing evening screen time in schoolchildren. Interventions trying to improve sleep
353 duration in schoolchildren that targets bed times should target screen times as well.

354 Just like previous study²⁵, interventions should also focus on improving parenting
355 skills and encouraging rules to govern the home. Limiting screen time may reduce late
356 bedtimes and in turn improve weight status via increase in sleep duration of school
357 children.

358 There are still many questions that remain unanswered. Future studies methodology
359 should see how geospatial technologies such as GIS (Geographic Information
360 Systems) / GPS (Global Positioning System) could be used. Also, can the association
361 between screen time and BMI be mediated by sleep duration? Since short sleep
362 duration is associated with high screen time.

363

364

365 **Acknowledgements**

366 Authors wish to acknowledge the people who contributed to the development of this
367 project, and those who were involved in the data collection process in the schools. We
368 also wish to thank all of the children and the Physical Education department staff of
369 the schools who participated in this project.

370 **Funding:** This study was funded by the University of Essex Research Promotion
371 Fund (grant code: DB014B5).

372 **Conflict of interest statement:** None declared. Authors have no financial conflicts of
373 interest.

374 **Authors' contributions:**

375 A.A.O., Dr. C.V., and Dr. G.R.S., conceptualized the study, designed the protocol and
376 were involved in data collection in the schools. A.A.O. performed the regression
377 analysis, and wrote parts of the Introduction and Results sections of the manuscript.
378 Dr. G.R.S. proofread and wrote the Discussion section of the manuscript. Dr. CV.
379 edited, proofread and reviewed the manuscript.

380

381

382 **References**

383

384 1 Matricciani L, Olds T, Petkov J. In search of lost sleep: secular trends in the
385 sleep time of school-aged children and adolescents. *Sleep Med Rev.* 2012; **16**: 203-11.

386 2 Owens JA. The practice of pediatric sleep medicine: results of a community
387 survey. *Pediatrics.* 2001; **108**: E51.

388 3 Spiegel K, Leproult R, L'Hermite-Baleriaux M, Copinschi G, Penev PD, Van
389 Cauter E. Leptin levels are dependent on sleep duration: relationships with
390 sympathovagal balance, carbohydrate regulation, cortisol, and thyrotropin. *J Clin*
391 *Endocrinol Metab.* 2004; **89**: 5762-71.

392 4 Patel SR, Hu BF. Sleep deprivation and Obesity. In: Frank H (ed.). *Obesity*
393 *Epidemiology.* Oxford University Press, New York, 2008a; 320-41.

- 394 5 Patel SR, Hu FB. Short sleep duration and weight gain: a systematic review.
395 *Obesity (Silver Spring)*. 2008b; **16**: 643-53.
- 396 6 Knutson KL. Sleep duration and cardiometabolic risk: a review of the
397 epidemiologic evidence. *Best Pract Res Clin Endocrinol Metab*. 2010; **24**: 731-43.
- 398 7 American Academy of Pediatrics. Committee on Public Education. American
399 Academy of Pediatrics: Children, adolescents, and television. *Pediatrics*. 2001; **107**:
400 423-6.
- 401 8 Tremblay MS, Leblanc AG, Janssen I, Kho ME, Hicks A, Murumets K, *et al*.
402 Canadian sedentary behaviour guidelines for children and youth. *Appl Physiol Nutr*
403 *Metab*. 2011; **36**: 59-64; 65-71.
- 404 9 Ogunleye AA, Voss C, Sandercock GR. Prevalence of high screen time in
405 English youth: association with deprivation and physical activity. *J Public Health*
406 *(Oxf)*. 2012; **34**: 46-53.
- 407 10 Garrison MM, Liekweg K, Christakis DA. Media use and child sleep: the
408 impact of content, timing, and environment. *Pediatrics*. 2011; **128**: 29-35.
- 409 11 Foley LS, Maddison R, Jiang Y, Marsh S, Olds T, Ridley K. Presleep
410 activities and time of sleep onset in children. *Pediatrics*. 2013; **131**: 276-82.
- 411 12 Stamatakis E, Hillsdon M, Mishra G, Hamer M, Marmot M. Television
412 viewing and other screen-based entertainment in relation to multiple socioeconomic
413 status indicators and area deprivation: the Scottish Health Survey 2003. *J Epidemiol*
414 *Community Health*. 2009; **63**: 734-40.
- 415 13 Sisson SB, Church TS, Martin CK, Tudor-Locke C, Smith SR, Bouchard C, *et*
416 *al*. Profiles of sedentary behavior in children and adolescents: the US National Health
417 and Nutrition Examination Survey, 2001-2006. *Int J Pediatr Obes*. 2009; **4**: 353-9.
- 418 14 Owens JA, Spirito A, McGuinn M. The Children's Sleep Habits Questionnaire
419 (CSHQ): psychometric properties of a survey instrument for school-aged children.
420 *Sleep*. 2000; **23**: 1043-51.
- 421 15 Olds TS, Maher CA, Matricciani L. Sleep duration or bedtime? Exploring the
422 relationship between sleep habits and weight status and activity patterns. *Sleep*. 2011;
423 **34**: 1299-307.
- 424 16 Tammelin T, Ekelund U, Remes J, Nayha S. Physical activity and sedentary
425 behaviors among finnish youth. *Med Sci Sport Exer*. 2007; **39**: 1067-74.
- 426 17 Cole TJ, Freeman JV, Preece MA. Body mass index reference curves for the
427 UK, 1990. *Arch Dis Child*. 1995; **73**: 25-9.
- 428 18 Cole TJ, Bellizzi MC, Flegal KM, Dietz WH. Establishing a standard
429 definition for child overweight and obesity worldwide: international survey. *Bmj*.
430 2000; **320**: 1240-3.
- 431 19 Department of Communities and Local Governments. English Indices of
432 Deprivation. Assessed 3rd of August, 2012. Vol. 2012. London, 2007.
- 433 20 Reilly JJ, Armstrong J, Dorosty AR, Emmett PM, Ness A, Rogers I, *et al*.
434 Early life risk factors for obesity in childhood: cohort study. *Bmj*. 2005; **330**: 1357.
- 435 21 Locard E, Mamelie N, Munoz F, Miginiac M, Billette A, Rey S. [Life style of
436 children and obesity in a population of 5-year-old children]. *Rev Epidemiol Sante*
437 *Publique*. 1992; **40**: 460-6.
- 438 22 Custers K, Van den Bulck J. Television viewing, internet use, and self-
439 reported bedtime and rise time in adults: implications for sleep hygiene
440 recommendations from an exploratory cross-sectional study. *Behav Sleep Med*. 2012;
441 **10**: 96-105.
- 442 23 Taylor AW, Winefield H, Kettler L, Roberts R, Gill TK. A population study of
443 5 to 15 year olds: full time maternal employment not associated with high BMI. The
444 importance of screen-based activity, reading for pleasure and sleep duration in
445 children's BMI. *Matern Child Health J*. 2012; **16**: 587-99.
- 446 24 Van den Bulck J. Television viewing, computer game playing, and Internet
447 use and self-reported time to bed and time out of bed in secondary-school children.
448 *Sleep*. 2004; **27**: 101-4.

449 25 de Jong E, Stocks T, Visscher TL, Hirasing RA, Seidell JC, Renders CM.
450 Association between sleep duration and overweight: the importance of parenting. *Int J*
451 *Obes (Lond)*. 2012.

452 26 Olds T, Maher C, Blunden S, Matricciani L. Normative data on the sleep
453 habits of Australian children and adolescents. *Sleep*. 2010; **33**: 1381-8.

454 27 O'Dea JA, Dibley MJ, Rankin NM. Low sleep and low socioeconomic status
455 predict high body mass index: a 4-year longitudinal study of Australian
456 schoolchildren. *Pediatr Obes*.

457 28 Locard E, Mamelle N, Billette A, Miginiac M, Munoz F, Rey S. Risk factors
458 of obesity in a five year old population. Parental versus environmental factors. *Int J*
459 *Obes Relat Metab Disord*. 1992a; **16**: 721-9.

460 29 Chaput JP, Brunet M, Tremblay A. Relationship between short sleeping hours
461 and childhood overweight/obesity: results from the 'Quebec en Forme' Project. *Int J*
462 *Obes (Lond)*. 2006; **30**: 1080-5.


463 30 Hale L, Berger LM, LeBourgeois MK, Brooks-Gunn J. Social and
464 demographic predictors of preschoolers' bedtime routines. *J Dev Behav Pediatr*. 2009;
465 **30**: 394-402.

466 31 Taheri S, Lin L, Austin D, Young T, Mignot E. Short sleep duration is
467 associated with reduced leptin, elevated ghrelin, and increased body mass index. *PLoS*
468 *Med*. 2004; **1**: e62.

469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503

504 **Figure 1:** Flowchart showing the potential mechanism in the association between
505 screen time, sleep deprivation, intermediate factors and obesity (Note: arrows on
506 represent associations and not causal links).

507
508
509
510
511
512



514
515
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551

	<i>Early bed: n(%)</i>	<i>Late bed:n(%)</i>
Sex		
Female	382(62.73)	227(37.27)
Male	418(57.81)	305(42.19)
Age (years)		
11	475(67.19)	232(32.81)
12	89(41.59)	125(58.41)
13	117(75.48)	38(24.51)
14	69(48.59)	73(51.41)
15	50(43.86)	64(56.14)
*BMI Z score	0.66(±1.32)	0.73(±1.25)
BMI categories		
Normal	520(59.56)	353(40.44)
Overweight	181(64.07)	106(35.93)
Obese	75(58.14)	54(41.86)
Screen time		
<2hours	556(64.95)	300(35.05)
2-4 hours	141(54.23)	119(45.77)
>4hours	99(48.53)	105(51.47)
*Deprivation	22.82(±15.36)	25.71(±15.82)
‡Schools		
1	298(66.52)	150(33.48)
2	120(72.29)	46(27.71)
3	382(53.20)	336(46.80)

553 Data shown are n(%) or *Mean (±SD)

554 BMI: body mass index

555 ‡School 1 was from a less deprived location; school 2 was from the least deprived rural location, while school 3
556 was from the most deprived location.

557

558

559

560

561 **Table 2:** Bonferroni tests for multiple comparisons of mean differences and 95%
562 Confidence intervals of index of multiple deprivation (IMD) scores.

<i>Multiple comparisons</i>	<i>Mean difference in IMD score (95% CI)</i>	<i>p-values</i>
‡Schools:		
School 1 compared to school 2	0.49(-2.17 to 3.16)	1.00
School 1 compared to school 3	-21.63(-23.42 to -19.85)	<0.001
School 2 compared to school 3	-22.13(-24.71 to -19.55)	<0.001
Bedtimes:		
Early-bed compared to Late- bed	-2.89(-4.80 to -0.97)	0.003
Screen time:		
<2hours compared to 2-4 hours	-1.69(-4.63 to 1.25)	0.505
<2hours compared to 4 hours	-5.73(-9.11 to -2.35)	<0.001
2-4hours compared to 4 hours	-4.04(-8.04 to -0.04)	0.047

563 ‡School 1 was from a less deprived location; school 2 was from the least deprived rural location, while school 3
564 was from the most deprived location.

565

566

567

568

569

570

571

572

573

574

575

576 Table 3: Adjusted and unadjusted odds ratio (95% confidence intervals) of late
 577 bedtimes in English schoolchildren.

	<i>Unadjusted OR (95%CI)</i>	<i>Adjusted OR (95%CI)</i>	<i>Adjusted OR (95%CI) without deprivation</i>
Screen time			
>4hours	1.97(1.44 to 2.68)	1.97(1.34 to 2.89)	1.70(1.22 to 2.36)
2-4 hours	1.56(1.18 to 2.07)	1.50(1.07 to 2.09)	1.43(1.06 to 1.39)
<2hours	1.00	1.00	1.00
Sex			
Males	1.23(0.99 to 1.53)	1.16(0.88 to 1.51)	1.20(0.95 to 1.53)
Females	1.00	1.00	1.00
Age			
15	2.62(1.75 to 3.92)	1.58(0.93 to 2.71)	1.77(1.11 to 2.84)
14	2.17(1.50 to 3.12)	1.65(0.99 to 2.77)	1.50(0.96 to 2.34)
13	0.67(0.45 to 0.99)	0.54(0.33 to 0.87)	0.55(0.36 to 0.84)
12	2.88(2.10 to 3.94)	2.19(1.46 to 3.27)	2.35(1.65 to 3.33)
11	1.00	1.00	1.00
BMI			
Obese	1.06(0.73 to 1.54)	1.01(0.65 to 1.59)	0.93(0.63 to 1.39)
Overweight	0.83(0.63 to 1.09)	0.87(0.63 to 1.20)	0.84(0.63 to 1.11)
Normal weight	1.00	1.00	1.00
Deprivation	1.01(1.00 to 1.02)	1.00(0.99 to 1.01)	-
Schools (different Geographical location)			
3	2.30(1.59 to 3.32)	1.66(0.96 to 2.85)	1.64(1.07 to 2.52)
2	1.31(0.89 to 1.94)	1.12(0.73 to 1.74)	1.28(0.85 to 1.93)
1	1.00	1.00	1.00

578
 579
 580
 581
 582
 583