

This is an accepted manuscript of an article published by Elsevier Ltd. in  
LWT- Food Science and Technology, 25/07/2016 available at: DOI: 10.1016/j.lwt.2016.07.051

## Accepted Manuscript

Microbiological quality of ready-to-eat fresh vegetables and their link to food safety environment and handling practices in restaurants

Dima Faour-Klingbeil, Ewen C.D. Todd, Victor Kuri



PII: S0023-6438(16)30462-5

DOI: [10.1016/j.lwt.2016.07.051](https://doi.org/10.1016/j.lwt.2016.07.051)

Reference: YFSTL 5625

To appear in: *LWT - Food Science and Technology*

Received Date: 4 May 2016

Revised Date: 18 July 2016

Accepted Date: 21 July 2016

Please cite this article as: Faour-Klingbeil, D., Todd, E.C.D., Kuri, V., Microbiological quality of ready-to-eat fresh vegetables and their link to food safety environment and handling practices in restaurants, *LWT - Food Science and Technology* (2016), doi: 10.1016/j.lwt.2016.07.051.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2016. This manuscript version is made available under the CC-BY-NC-ND 4.0 license

<http://creativecommons.org/licenses/by-nc-nd/4.0/>

Contact at University of Plymouth:  
[v.kuri@plymouth.ac.uk](mailto:v.kuri@plymouth.ac.uk)

## Microbiological quality of ready-to-eat fresh vegetables and their link to food safety environment and handling practices in restaurants

Dima Faour-Klingbeil<sup>a\*</sup>, Ewen C. D. Todd<sup>b</sup>, Victor Kuri<sup>a</sup>

<sup>a</sup>School of Biological Sciences, Plymouth University, UK

<sup>b</sup>Visiting Professor, Department of Nutrition and Food Science, American University of Beirut, Lebanon, and Ewen Todd Consulting, Okemos, Michigan, USA

\*Correspondence author: dima.faour@gmail.com

### ABSTRACT

1 The increased consumption of ready-to-eat salads outside homes as a result of a fast  
2 paced lifestyle, awareness on their nutritional attributes and enhanced processing  
3 technology is well documented. This study aimed to determine the microbiological  
4 quality of fresh-cut salads vegetables in small and medium sized foodservice  
5 establishments (SMEs) and to identify risk factors and handling practices through  
6 observational assessment in order to investigate if an association between  
7 microbiological quality and visual assessment (inspection) scores can be established.  
8 A total of 118 samples fresh-cut vegetable salads were collected from 50 inspected  
9 locations and analysed microbiologically, in addition to 49 swabs of knives and cutting  
10 boards. There was no statistically significant correlation between visual assessment  
11 scores and bacteriological counts on vegetables or cutting boards. Nonetheless, the  
12 consistent relationship between inspection ratings on cross-contamination and cleaning  
13 components and *Listeria* spp. levels was statistically significant. This study  
14 demonstrated that overall visual assessment scores would not directly reflect the safety  
15 of salad vegetables and that the significance of microbiological assessment should be  
16 considered in relation to individual inspection components. It is necessary to place

17 effective control measures on cleaning standards and risk of cross-contamination to  
18 improve the microbiological safety of fresh salad vegetables in SMEs.

19

## 20 **1. INTRODUCTION**

21 Fresh vegetables are rich sources of water-soluble vitamins and other nutrients  
22 essentials to improve the nutritional status and decrease the risk of cardiovascular  
23 disease (Su & Arab, 2006). However, when they are not carefully prepared, they can be  
24 subjected to pathogenic contamination and become hazardous to health particularly  
25 when eaten raw (WHO, 2008).

26 Outbreak investigations often indicate that food service establishments (FSE) greatly  
27 contribute to foodborne illnesses involving fresh produce (Jones & Angulo, 2006;  
28 Sodha et al., 2011). Multiple studies revealed that food workers were frequently  
29 engaged in unsafe food handling (Clayton & Griffith, 2004; Manning, 1994; Rajagopal  
30 & Strohbahn, 2013; Sneed, Strohbahn, & Gilmore, 2004) and that microbial  
31 contamination of ready-to-eat (RTE) foods typically occurred in FSEs with food  
32 handlers as asymptomatic carriers of pathogenic microorganisms or with poor personal  
33 hygiene being involved (McEvoy et al., 2004; Todd et al., 2008). Equipment or surfaces  
34 that have not been effectively cleaned or remained wet between cleaning and use also  
35 serve as direct routes for contamination of ready to eat foods (Evans et al., 2004; Gill et  
36 al., 2001), besides inappropriate storage temperatures, and insufficient cooking (Jones  
37 et al., 2008; WHO, 2007).

38 Less information is available on the relative health risks attributed to handling practices  
39 and preparation procedures of raw salad vegetables in SMEs, while other RTE foods  
40 and meats have attracted more attention.

41 Inspection tools are essential for capturing information on the general hygiene standards  
42 and food handlers' practices Although private or local authorities' inspections are an

43 effective mechanism to assure compliance to food safety standards, there is no a clear  
44 indication of a correlation between risk of foodborne illnesses and inspection scores.  
45 There have been many cases when restaurants scored high on inspections and were still  
46 having critical violation in food safety(Jones et al., 2004). The significance of  
47 association of microbiological quality of RTE vegetables to hygiene inspection scores  
48 has not been fully investigated and not sufficiently addressed by researchers. Earlier  
49 attempts to establish direct relationship between the results on microbiological analysis  
50 of food and visual inspections have not been successful and were mostly based on foods  
51 of animal origins(Powell & Attwell, 1995; Tebbutt & Southwell, 1989; Wyatt & Guy,  
52 1980).

53 This study aimed at conducting observational assessment of the fresh produce handling  
54 processes from the receiving stage until display and service to identify risk factors that  
55 may be associated with the microbial safety of fresh produce in SMEs which will  
56 provide further insights to devise effective preventive measures.

57

## 58 **2. MATERIAL AND METHODS**

### 59 **2.1 Observational survey**

60 A convenience sample of fifty SMEs located in Beirut were observationally assessed for  
61 hygiene standards and handling practices of food handlers during the salad vegetable  
62 preparation. The survey checklist comprised 6 constructs of 2-7 components for analysis  
63 in which the good hygienic practices (GHP) and other prerequisites proposed by the  
64 Codex Alimentarius (CAC/RCP 1, 1969) were considered for the visual assessment  
65 (Table 1). Additional components in relation to salad preparation practices were also  
66 included. The criteria for each component were defined to specify limits for  
67 classification. (Supplementary materials).

68 A reliability analysis test was performed to measure the internal consistency in the  
69 survey questionnaire. Cronbach's Alpha was 0.928 which indicates a high level of  
70 internal consistency for our scale.

## 71 **2.2 Additional information**

72 Additional 8 questions on handling practices of fresh vegetables during receiving,  
73 washing and storage were posed to food handlers (n=80) via face-to-face interviews that  
74 were conducted in our earlier study on food safety knowledge, attitudes and practices  
75 (Faour-Klingbeil et al., 2015). The questions were ranked on a five points rating scale  
76 (never = 1, rarely = 2, sometime = 3, often = 4 and always =5).

77 To ensure consistency and unbiased data records, the data collection and visual  
78 assessment were carried out by one of the authors (Dima Faour-Klingbeil).

79

## 80 **2.3 Collection of RTE fresh-cut salads vegetables samples**

### 81 **2.3.1 Management of samples**

82 A total of 118 samples of various fresh cut RTE salad vegetables (lettuce, parsley,  
83 arugula, coriander, cucumber, tomato and radish) prepared in 50 restaurants were  
84 collected after washing and cutting/chopping. On average, 3 types of vegetables were  
85 sampled from each restaurant, being subjected to availability and preparation plans at  
86 times of visits. They were placed in a sterile bag by food handlers at the end of the  
87 preparation process by means of utensils or tools typically used when bringing them  
88 into display or storage containers, taking care that they would not touch the inside of the  
89 bags.

### 90 **2.3.2 Swabs of cutting boards and knives**

91 Before cutting/chopping vegetables, surfaces of cleaned cutting boards and knives  
92 (normally cleaned by assigned cleaners in well-established restaurants, or food workers

93 in less developed restaurants) were swabbed by moistened cotton-tip in buffered  
94 peptone water (BPW) (Bio-rad laboratories Ltd, Hemel Hempstead, UK) in three  
95 different directions: left to right, top to bottom, and diagonal over a 50 cm<sup>2</sup> area for  
96 cutting boards and a length of ca. 10cm on knives. The swabs were placed in tubes of 5  
97 ml buffered peptone water for subsequent analysis.

98

### 99 **2.3.3 Microbiological analysis of samples**

100 Samples of salad vegetables were analysed for the presence of pathogens and hygiene  
101 indicators organisms commonly isolated from RTE fresh vegetables, i.e., *S. aureus*,  
102 *Salmonella* spp., *Listeria* spp., *L. monocytogenes*, in addition to total viable counts  
103 (APC), *E. coli* and TCs (Nguz et al., 2005; Sagoo et al., 2001). For microbiological  
104 analysis, all the media used were obtained from Bio-Rad Laboratories Ltd., Hemel  
105 Hempstead, UK unless otherwise mentioned and samples were analysed according to  
106 ISO 16140. Briefly, 10 g of the samples was weighed into sterile stomacher bags and  
107 homogenized with 90 ml sterile peptone buffered water (BPW) for 2 min at medium  
108 speed. Aliquots of 0.1 ml of appropriate dilutions were spread in duplicates on suitable  
109 media. APC were enumerated on plate-count agar, as for *E. coli* and TC, 1 ml was  
110 dispensed into petri dishes for enumeration by pouring technique using RAPID'*E. coli* 2  
111 agar. The plates were incubated at 37°C for 48 h. *Coagulase-positive Staphylococci*  
112 were enumerated on RAPID'*Staph* Agar supplemented with egg yolk. For the detection  
113 of *S. aureus*, typical presumptive colonies with clear halo resulting from proteolysis of  
114 egg yolk were further tested using a latex agglutination test (Pastorex *Staph* Plus). For  
115 the isolation of *Salmonella* spp., selective enrichment was performed in Rappaport-  
116 Vassiliadis-soya broth to be incubated at 41.5°C. After 24 h of incubation, a 0.1 ml  
117 sample was plated on RAPID'*Salmonella* agar and plates were incubated at 37°C for

118 24h ( $\pm$  2h). While for *L. monocytogenes*, Fraser ½ broth was used in the selective  
119 enrichment and after incubation for 1 h at 20°C, 0.1 ml of the homogenate was  
120 transferred onto RAPID'*L. monocytogenes* agar plates to be incubated at 37°C for 24–  
121 48h. *Listeria* spp. were enumerated and typical *L. monocytogenes* colonies were  
122 afterwards selectively identified and by *Listeria* strips (bioMérieux, Marcy l'Etoile,  
123 France). *Salmonella* spp. colonies were identified biochemically by the lysine iron agar  
124 and tryptic sugar iron agar slants biotyping technique. Additional confirmation for  
125 positive *Salmonella* spp. colonies and for *E. coli* was done by the API 20E bacterial  
126 identification test strip.

127 The counts were reported as means of colony-forming units (CFU) per g and were  
128 converted into Log CFU/g.

129 Additionally, for statistical purposes, *Listeria* spp were ranked into 3 levels (Above 100  
130 CFU/g, Below 100 CFU/g, and Not detected).

### 131 **2.3.4 Swab tests**

132 The swabs in 5 ml tube of BPW were vortexed vigorously for 1 min. Tenfold serial  
133 dilutions were spread-plated onto duplicate plates of PCA, RAPID'*Staph* agar  
134 supplemented with egg yolk and RAPID'*E. coli* 2 agar.(Sneed, Strohbahn, Gilmore, et  
135 al., 2004). Counts were expressed as log CFU/swabbed area.

136

## 137 **3. DATA HANDLING AND STATISTICAL ANALYSIS**

138 All data were analysed using the IBM SPSS Statistics (SPSS) version 22.

139 Observational assessment of each of the 26 components was rated on three units scale  
140 (adequate=3, incomplete=2, inadequate=1). The sum of the total awarded units on  
141 adequacy level (visual assessment scores) was converted to 100 points.

142 Frequency of levels in compliance (adequacy level) for each visually inspected  
143 component was obtained. Bacterial levels differences among different compliance levels  
144 were compared using One-way ANOVA, and independent t-test was performed to  
145 compare results between two groups.

146 The association between bacterial counts and overall visual assessment scores was  
147 assessed by Pearson correlation and multiple linear regression analysis; binomial  
148 regression was performed for *S. aureus*.

149 The percentage variances in bacterial counts (Log CFU/g) explained by individual  
150 inspection components were determined by correlation ratio  $\text{ETA}^2$  ( $\eta^2$  ratio). In the case  
151 *Listeria* and *S. aureus*, Spearman's rho and cross-tabulations Somer'd tests were also  
152 applied.

153

#### 154 **4. RESULTS**

##### 155 **4.1 Overall results on food handlers 'practices and hygiene conditions on premises**

156 Results of the visual inspections of FSEs and food handlers' practices during the  
157 preparation of fresh salads vegetables indicated structural inadequacies and insufficient  
158 fulfilment of hygiene prerequisites with a mean score on overall adequacy level of 55.5  
159  $\pm$  19.0 over 100 possible points (Figure 1), with the majority of locations being below  
160 scores of 50-70. Over half (54%) of the food premises failed to fulfil the basic hygienic  
161 requirements for clean floors, equipment and food contact surfaces, while a third had  
162 limitations in the structural conditions (Figure 2). Recorded incompliances included  
163 open drains, gaps and holes on windows and walls and evidence of pests (cockroaches)  
164 at the time of the survey. Furthermore, 22% had not a completely well maintained  
165 premise. More than a half (52%) of the FSEs had space limitations compromising the  
166 preparation of food safely, whereas only 22% of premises had taken measures to



167 separate areas for the preparation of raw meats and RTE foods. It was notable that the  
168 inappropriate sanitation measures were not applied in 60% of the premises (Figure 2).  
169 Only 8% of FSEs had cleaning schedules, and showed evidence of temperature  
170 monitoring records of salads display and cold storage.

171 In addition, a large percentage of food businesses (64%) lacked hand washing sinks; or  
172 designated sinks for washing fresh fruits and vegetables were either absent (32%) or if  
173 fitted, it was not clean and used for others purposes such as washing hands or  
174 implements used with raw meat and cooked foods (40%). More concerning, gloves were  
175 used correctly and appropriately during the salad preparation in just a fifth (20%) of the  
176 premises.

177 Risks of cross-contamination were detected in 48% of the premises, for example by the  
178 presence of heavily chipped or unclean cutting boards, unfamiliarity of food handlers  
179 with the concept of color-coding or separate use of utensils and cutting boards for raw  
180 meat and fresh vegetables. There was misuse of colour-coded cutting boards in 18% of  
181 FSE's where colour-coded cutting boards were used for several types of food. The  
182 component "frozen foods are thawed properly" was not observed in 74% of the  
183 premises visited, yet it was inadequately performed in 14% of the locations where  
184 frozen fish or chicken soaked in water were noted at the time of the visit.

#### 185 **4.2 Handling practices and the process of salads vegetables preparation**

186 Fresh vegetables were received during the mornings (7-9 a.m.) in plastic crates transported  
187 on open trucks or in vans. The great majority (95%) reported that they received fresh  
188 produce in uncooled vehicles (Table 2). In some cases, the person in charge or business  
189 owner purchased the daily needs from the central market or nearby groceries. More than  
190 two thirds of the respondents reported sourcing the fresh produce from the same

191 supplier (68.4%), and washing the vegetables before cutting (77%). In general,  
192 preparation started early, particularly with bundles of parsley which were finely  
193 chopped for serving later in the day in traditional salads and appetizers. Parsley leaves  
194 were chopped before washing in 34% of FSEs, which is consistent with the typical  
195 preparation sequence at homes (Figure 3), aiming to keep the texture of the leaves  
196 longer, as they would becoming soggy if they are washed ahead of time. About a third  
197 of the food businesses did not sanitize fresh vegetables, and used only water to wash  
198 them. However, a large proportion (84%) reported that the wash water was neither  
199 treated nor filtered. With long-standing shortages of potable water in Lebanon,  
200 restaurants, and homes, purchase water, often of uncertain quality and source, which is  
201 then stored in tanks. Out of the 56% using sanitizers, 21% used sodium  
202 dichloroisocyanurate (NaDCC) and more than a third (45%) applied a post-sanitization  
203 water rinse to remove the remaining taste or odour, respectively. It was noted during  
204 inspection discussions and observations that automated systems regulating the  
205 concentrations of chemical sanitizers in addition to water filters were in place, in some  
206 corporate-managed restaurants. On other places (24 %), incorrect dilutions of sanitiser  
207 was observed, typically as haphazard mixing of vinegar or NaDCC tablets in water. The  
208 majority reported that fresh produce was kept in cold storage, whereas this was actually  
209 only observed in 38% of the premises, with inadequate alternatives including stairways,  
210 kitchen floors of spaces in crowded production areas.

#### 211 **4.3 The microbiological quality of fresh salads vegetables**

212 Results on microbiological analysis of fresh-cut salad vegetables are presented in (Table  
213 3 and 4).

214 The mean APC levels ranged from 2.90 to 7.38 Log CFU/g, with counts above  
215  $10^7$ CFU/g recorded for 17% of the samples. The prevalence rate was substantially high

216 in TCs (79.6%, 94/118). TCs were found between 1.72 - 6.40 Log CFU/g, of which  
217 38% were  $>4$  Log CFU/g. Whereas, *E.coli* was isolated from 31.3% (37/118), with  
218 bacterial loads ranging from less than 1.00 to 7.15 Log CFU/g, and the incidence rate  
219 was 64.8% of the positive samples (24/37) for counts higher than 100 CFU/g.

220 More than two thirds (41.5%) of the samples were found to contain *S. aureus*. In  
221 addition, *Listeria* spp. were isolated from 70.6% of the samples. The overall incidence  
222 level was 53% for counts above 100 CFU/g, with an average of 3.24 Log CFU/g. *L.*  
223 *monocytogenes* had a prevalence rate of 3.7 % mainly in arugula, parsley and lettuce,  
224 whereas *Salmonella* was detected in 0.9%, (lettuce).

225 Results on recovered microorganisms from contact surfaces (cutting boards and knives)  
226 are presented in Table 5. The microbial levels varied from below detection limits (10  
227 CFU/swabbed area) to generally high levels. *E.coli* was isolated from 30.6% (15/49) of  
228 contact surfaces (knives and cutting boards); of those, the mean values were found  
229 between 2.70 - 7.02 Log CFU/swabbed area, whereas the incidence rate in TCs was  
230 higher (53.0%, 26/49) with levels between 4.88 - 8.40 Log CFU/swabbed area. There  
231 was no statistically significant correlation between the microbial counts recovered from  
232 contact surfaces and the ratings on the adequacy level of sanitation of work surfaces  
233 ( $p>0.05$ ).

234 Overall, the analysis of data shows no statistical significant differences and inconsistent  
235 trends in bacterial counts of different visual assessment rankings for each individual  
236 inspection component ( $p>0.05$ ). For instance, higher counts of TCs were observed on  
237 lettuce and parsley obtained from premises with inadequate sanitary conditions and  
238 unsafe handling practices, however this was not the case with cucumbers (Table 6).  
239 Also, the frequency in the distribution of bacterial levels on lettuce and parsley in  
240 relation to hygiene scores shows that high concentration levels were grouped at lower

241 scores (Figure 4). Likewise, the mean levels of coagulase-positive *Staphylococcus* spp.  
242 were higher on all vegetables prepared on premises lacking handwashing sinks (Figure  
243 5).

244 There was no correlation between total visual assessment scores and bacterial levels  
245 ( $p>0.05$ ). However, independent t-test still reveals a significant difference ( $t=-2.198$ , 81,  
246  $p=0.03$ ), between inspection scores for premises with *Listeria* counts above 100 CFU/g  
247 ( $53.44\pm 18.39$ ) and those where the organism was not detected ( $64.48 \pm 26.12$ ). When  
248 Eta correlation and non-parametric tests were further performed for this organism, no  
249 significant correlations of microbial results with all individual inspection component  
250 ( $p>0.05$ ) were shown, while correlation tests and cross tabulations somer'd test  
251 revealed a significantly low and moderate association of *Listeria* levels with the  
252 inspection components related to cross contamination, handling practices, zoning and  
253 availability of handwashing sinks ( $p<0.05$ ) (Figure 6). This association level was  
254 consistent with linear regression establishing that *Listeria* spp levels may be predicted  
255 by the visual assessment scores ( $F_{1,103}=11,614$ ,  $p=0.001$ , but the score accounted for  
256 only 10.5% ( $R^2$ ) of the explained variability in *Listeria* levels in vegetables. Given the  
257 small value of  $R^2$ , the prediction model using the visual assessment scores is not  
258 accurate. However and more interestingly, as we considered each inspected component  
259 individually, Eta<sup>2</sup> coefficients showed higher percentage in variations in *Listeria* spp.  
260 counts (30-34%) which were explained and attributed to cross contamination and  
261 cleaning operations components ( $p<0.05$ ).

## 262 5. DISCUSSION

### 263 5.1 Food safety practices and microbial quality of fresh salads vegetables

264 A number of food safety practices concerns were identified in this study. The general  
265 lack of cleaning and sanitization procedures combined with a clear evidence of cross-  
266 contamination opportunities were generally reflected in the overall unsatisfactory  
267 quality of RTE vegetables. The majority of SMEs seemed to be unaware of the  
268 significance of applying control measures when handling vegetables and of the  
269 fundamental requirements for separate handwashing and vegetables washing sinks. APC  
270 were above the specified limits for RTEs, 7 Log CFU/g, in 17% of the analysed  
271 samples. when APC count is  $>10^6$  CFU/g, it may not necessarily relate to food safety  
272 hazards; in many of these cases, there is a predominant microorganism from an  
273 environmental source (PHLS, 2000) such as the processing stages involving handling,  
274 cutting, slicing and improper storage as well as display conditions (Abadias et al.,  
275 2012); Nguz et al. (2005) showed that chlorine treated fresh-cut organic mixed  
276 vegetables were still found to harbour high levels of TCs (5.9 Log CFU/g) and it was  
277 proposed that high loads of coliforms in RTE vegetables at retails levels is directly  
278 influenced by intense use of untreated manure during pre-harvest, and extensive  
279 handling during postharvest (Aycicek et al., 2006). In our earlier study, TCs  $\geq 5$  Log  
280 CFU/g were isolated from more than two third of the fresh vegetables (69%) coming  
281 from locations with alarming deficits at harvest and post-harvest washing, storage and  
282 distribution stages (Faour-Klingbeil et al., 2016).

283 According to the EC legal food safety criteria and the UK Public Health Laboratory  
284 Service (PHLS) microbiological guidelines for RTE foods sampled at the point of sale,  
285 for category 5 fresh vegetables (HPA, 2009; PHLS, 2000), our study results on  
286 microbial contamination levels of more than half of the RTE salad vegetables were  
287 unsatisfactory due to *E. coli* and *Listeria* spp. counts that exceeded the criteria limits

288  $>10^2$  CFU/g indicating poor hygienic practices and sanitary conditions (Gilbert et al.,  
289 2000).

290 *Listeria* spp. are rarely implicated in illnesses involving produce, however, they may  
291 indicate a significant failure of hygiene standards in the preparation and /or storage of  
292 fresh vegetables(Gilbert et al., 2000) which in turn are considered hazardous for  
293 *L.monocytogenes* contamination (Ponniah et al., 2010). Presence of *L.monocytogenes*  
294 and *Salmonella* spp. were traced back to samples obtained from restaurant that had no  
295 handwashing sinks, fresh vegetable washing sinks, or adequate preparation and storage  
296 areas or surfaces and the corresponding visual assessment score recorded 32 over 100  
297 possible points.

298 The lacking of handwashing sinks explained the fact that proper handwashing before  
299 and after use of gloves were not commonly observed, although many other factors could  
300 interfere as well. High frequency of *S. aureus* indicates poor hygiene practices of food  
301 handlers, the latter being known to be carriers of this pathogen (Todd et al., 2008) and  
302 may contribute in direct contamination of RTE fresh vegetables and contact surfaces via  
303 the hands (Todd et al., 2008).

## 304 **5.2 Food contact surfaces**

305 The PHLS recommended guidelines for cleaned contact surfaces specified levels of  
306 total viable microorganisms less than 80 CFU/cm<sup>2</sup> as satisfactory, 80-10<sup>3</sup>CFU/cm<sup>2</sup> is  
307 borderline, and over 10<sup>3</sup>CFU/cm<sup>2</sup> is unsatisfactory been associated with poor hygiene  
308 practices (Herbert et al., 1990). PCA counts  $\geq 10^3$ CFU/cm<sup>2</sup> was recorded for 33/49  
309 swabbed surface. The overall incidence rate of *E.coli* was 15/49 with counts  $\geq 1$   
310 CFU/cm<sup>2</sup>, whereas *E. coli* counts  $\geq 10^3$ CFU/cm<sup>2</sup> were recorded for 10/49 of swabs. TCs  
311 and *Staphylococcus* spp. were found in 26/49 and 39/49 of swabs with counts  
312  $\geq 10^3$ CFU/cm<sup>2</sup>. In this regard, the high microbial population size on contact surfaces

313 offered an additional assumption for the actual contamination observed on the washed  
314 salad items, particularly that sanitization and cleaning operations were lacking in a great  
315 majority of locations. Sneed, Strohbahn, Gilmore, et al. (2004) indicated that inadequate  
316 sanitation and recontamination problems were actually related to high aerobic plate  
317 counts recovered from cutting boards. Non-sanitized and scratched cutting surfaces,  
318 combined in some cases with misuse of sanitizers dilution, are an appropriate  
319 environment for harbouring pathogens that have the propensity to form biofilm on  
320 surfaces (Pui et al., 2011) and resist washing processes (Ravishankar et al., 2010).

321 As RTE fresh vegetables were obtained after washing, the existing microbiological  
322 characteristics do raise further doubts as to the implication of water quality. It is well  
323 recognized that natural resources and water supply in Lebanon endures a high risk of  
324 chemical and microbial pollution (Houry & El Jeblawi, 2007; Jurdi, 1992), at the same  
325 time, it is substantiated that washing with water of unsatisfactory microbial quality can  
326 serve as a vehicle for dispersion of microorganisms (Holvoet et al., 2013) and was the  
327 primary cause for the homogenous spread of *Salmonella* Enteritidis to fresh-cut  
328 vegetables during processing (Perez-Rodriguez et al., 2014). The quality of water used  
329 for washing or in post-sanitization rinsing process in SMEs should be addressed in  
330 future studies as a critical element to maintain fresh vegetables safety specially when  
331 more restaurants nowadays rely on purchasing water of unknown sources, usually  
332 coming in tankers collected from spring water but may or may not be chlorinated, to  
333 compensate for the shortage in water supply..

### 334 **5.3 Association of microbial counts to visual assessment scores and inspection** 335 **components**

336 Our data revealed an inconsistent association between the bacterial counts and visual  
337 assessment scores of handling practices and hygiene conditions. As we also studied the

338 possibility of association to each single inspection component, the microbiological  
339 quality of salad vegetables did not show any direct correlation with each individual  
340 inspected component. It was found that the cell counts were either corresponding or  
341 conflicting in trend across ranking on adequacy level and types of produce. The  
342 complexity of the interfering factors during sampling of RTE fresh vegetables from  
343 different operational conditions (e.g., environment and storage temperature, receiving  
344 and pre-receiving conditions of fresh vegetables, preparation stages of fresh cut  
345 vegetables, sampling methods) challenges the possibility to detect a clear cut trend and  
346 association. Add to this, large number of samples might be needed to investigate such a  
347 trend. Our findings are in accordance with a study by Powell and Attwell (1995) where  
348 a link between the total viable counts and *S.aureus* on turkey and ham and the  
349 compliance rate to different inspection components was not established. Findings of  
350 earlier studies did not as well confirm such an association with the microbiological  
351 quality of foods of meat origin (Tebbutt & Southwell, 1989; Wyatt & Guy, 1980). Kuri  
352 et al. (1996) found that microbial indicators in meats, including pathogen prevalence,  
353 were not correlated to total hygiene scores of meat retailers, nor to temperature of  
354 samples, but they were related to type of retailer or origin of product.

355 We actually noted higher population size of hygiene indicators on some samples  
356 prepared under inadequate hygiene conditions, although a statistically significant  
357 correlation with the inspection scores failed. According to our results, it may be  
358 reasonable to consider that low visual assessment scores on the hygiene standards and  
359 handling practices probably indicate unsatisfactory microbial quality and likelihood for  
360 risks of salad vegetables contamination with *L.monocytogenes*, however, this  
361 association was only significant in relation to individual components related to cross-  
362 contamination and effective cleaning. The total visual assessment score can be affected



363 by a number of possible combinations of ranking levels of the 26 variables; a low  
364 inspection score might not necessarily indicate low ratings of all the critical components  
365 that have direct impact on the microbiological quality of vegetables. Hence, inspections  
366 should focus upon factors most likely to be responsible for foodborne infection or high  
367 microbial levels associated with RTE vegetables.

368

## 369 **6. CONCLUSION**

370 Links between the visual assessment scores on the overall food safety performance and  
371 the microbiological quality of RTE fresh vegetables are not simple to establish and were  
372 not clearly correlated. The total visual assessment scores per se would not directly  
373 indicate the microbiological safety of RTE vegetables in restaurants. However,  
374 variations in microbial counts and a significant correlation of high *Listeria* levels with  
375 the inadequate cleaning performances and cross-contamination preventive measures  
376 were recorded, which imply that shortfalls in those particular practices may possibly  
377 indicate pathogenic contamination of fresh vegetables.

378 Also, this study found high microbial loads in RTE vegetables that could serve as an  
379 indicator for the need to promote awareness on the critical areas commonly identified in  
380 SMEs and as guidance for local authorities to target those that may mostly affect the  
381 safety of fresh vegetables. It underscored the considerable requisite for improvement in  
382 sanitary and good hygienic practices and for vigilant cleaning and sanitation procedures  
383 to reduce or eliminate contamination and cross-contamination risks that may occur at  
384 pre-farm gate and throughout the supply chain stages. Therefore, applications of critical  
385 control points for the preparation of fresh salad vegetables and personnel training on the  
386 hazards associated with their preparation are fundamentals to improve the food safety of  
387 fresh produce particularly when prepared in small working facilities in SMEs.

**388 7. ACKNOWLEDGEMENTS**

389 The authors gratefully acknowledge the support provided by the Department of  
390 Nutrition and Food Science, American University of Beirut for partially funding this  
391 research work through a grant from the Lebanese National Council for Scientific  
392 Research (CNRS) #102598.

393

## 8. REFERENCES

- 394  
395  
396 Abadias, M., Alegre, I., Oliveira, M., Altisent, R., & Viñas, I. (2012). Growth potential  
397 of *Escherichia coli* O157:H7 on fresh-cut fruits (melon and pineapple) and  
398 vegetables (carrot and escarole) stored under different conditions. *Food Control*,  
399 27(1), 37-44. doi: <http://dx.doi.org/10.1016/j.foodcont.2012.02.032>
- 400 Aycicek, H., Oguz, U., & Karci, K. (2006). Determination of total aerobic and indicator  
401 bacteria on some raw eaten vegetables from wholesalers in Ankara, Turkey.  
402 *International Journal of Hygiene and Environmental Health*, 209(2), 197-201.  
403 doi: 10.1016/j.ijheh.2005.07.006
- 404 CAC/RCP 1(1969). Recommended international code of practice general principles of  
405 food hygiene (Revision 4 - 2003). Retrieved from:  
406 <http://www.mhlw.go.jp/english/topics/importedfoods/guideline/dl/04.pdf>
- 407 Clayton, D., & Griffith, C. (2004). Observation of food safety practices in catering  
408 using notational analysis. *British Food Journal*, 106, 211– 227.
- 409 Evans, J. A., Russell, S. L., James C., & Corry, J. E. L. (2004). Microbial contamination  
410 of food refrigeration equipment *Journal of Food Engineering*, 62, 225-232.
- 411 Faour-Klingbeil, D., Kuri, V., & Todd, E. (2015). Investigating a link of two different  
412 types of food business management to the food safety knowledge, attitudes and  
413 practices of food handlers in Beirut, Lebanon. *Food Control*, 55, 166-175.
- 414 Faour-Klingbeil, D., Murtada, M., Kuri, V., & Todd, E. (2016). Understanding the  
415 routes of contamination of ready-to-eat vegetables in the Middle East. *Food*  
416 *Control*, 62, 125-133. doi: <http://dx.doi.org/10.1016/j.foodcont.2015.10.024>
- 417 Gilbert, R., de Louvois, J., Donovan, T., Little, C., Nye, K., Ribeiro, C., Bolton, F.  
418 (2000). Guidelines for the microbiological quality of some ready-to-eat foods  
419 sampled at the point of sale. PHLS Advisory Committee for Food and Dairy  
420 Products. *Communicable Disease and Public Health*, 3(3), 163-167.
- 421 Gill, C. O., Bryant, J., & Badoni, M. (2001). Effects of hot water pasteurizing  
422 treatments on the microbiological condition of manufacturing beef used for  
423 hamburger patty manufacture. *International Journal of Food Microbiology*,  
424 63(3), 243-256.
- 425 Herbert, M., Donovan, T., & Manger, P. (1990). A study of the microbial contamination  
426 of working surfaces in a variety of food premises using the traditional swabbing  
427 technique and commercial contact slides . In Ashford (Ed.).
- 428 Holvoet, K., Sampers, I., Callens, B., Dewulf, J., & Uyttendaele, M. (2013). Moderate  
429 Prevalence of Antimicrobial Resistance in *Escherichia coli* Isolates from  
430 Lettuce, Irrigation Water, and Soil. *Applied and Environmental Microbiology*  
431 79(21), 6677–6683.
- 432 Hourii, A., & El Jeblawi, S. W. (2007). Water quality assessment of Lebanese coastal  
433 rivers during dry season and pollution load into the Mediterranean Sea. *Journal*  
434 *of Water and Health*, 5(4).
- 435 HPA. (2009). Guidelines for Assessing the microbiological safety of ready-to-eat foods.  
436 *Health Protection Agency*. .
- 437 Jones, S. L., Parry, S. M., O'Brien, S. J., & Palmer, S. R. (2008). Are staff management  
438 practices and inspection risk ratings associated with foodborne disease outbreaks  
439 in the catering industry in England and Wales? *Journal of Food Protection*,  
440 71(3), 550-557.
- 441 Jones, T. F., & Angulo, F. J. (2006). Eating in restaurants: a risk factor for foodborne  
442 disease? *Clin Infect Dis*, 43(10), 1324-1328. doi: 10.1086/508540

- 443 Jones, T. F., Pavlin, B., LaFleur, B., Ingram, L., & Schaffner, W. (2004). Restaurant  
444 inspection Scores and foodborne disease. *Emerging Infectious Diseases* 10(4),  
445 688-692.
- 446 Jurdi, M. (1992). A national study on the quality of potable water in Lebanon.  
447 *Proceedings of the national workshop of the status of water in Lebanon, United*  
448 *Nations International Children's Emergency Fund (UNICEF). Beirut, Lebanon*  
449 pp. 145-173 (in Arabic).
- 450 Kuri, V., Madden, R. H., & Collins, M. A. (1996). Hygienic quality of raw pork and  
451 chorizo (raw pork sausage) on retail sale in Mexico City. *Journal of Food*  
452 *Protection* 59 (2), 141-145.
- 453 Manning, C. K. (1994). Food safety knowledge and attitudes of workers from  
454 institutional and temporary foodservice operations. *Journal of the American*  
455 *Dietetic Association*, 94(8), 895-897. doi: [http://dx.doi.org/10.1016/0002-](http://dx.doi.org/10.1016/0002-8223(94)92372-8)  
456 [8223\(94\)92372-8](http://dx.doi.org/10.1016/0002-8223(94)92372-8)
- 457 McEvoy, J. M., Sheridan, J. J., Blair, I. S., & McDowell, D. A. (2004). Microbiological  
458 contamination of beef carcasses in relation to hygiene assessment based on  
459 criteria used in EU Decision 2001/47/EC. *International Journal of Food*  
460 *Microbiology*, 92, 217-225.
- 461 Nguz, K., Shindano, J., Samapundo, S., & Huyghebaert, A. (2005). Microbiological  
462 evaluation of fresh-cut organic vegetables produced in Zambia. *Food Control*,  
463 16(7), 623-628. doi: <http://dx.doi.org/10.1016/j.foodcont.2004.07.001>
- 464 Perez-Rodriguez, F., Saiz-Abajo, M. J., Garcia-Gimeno, R. M., Moreno, A., Gonzalez,  
465 D., & Vitas, A. I. (2014). Quantitative assessment of the Salmonella distribution  
466 on fresh-cut leafy vegetables due to cross-contamination occurred in an  
467 industrial process simulated at laboratory scale. *International Journal of Food*  
468 *Microbiology*, 184, 86-91. doi: 10.1016/j.ijfoodmicro.2014.05.013
- 469 PHLS. (2000). Microbiological guidelines for some ready-to-eat foods sampled at the  
470 point of sale. . *Communicable Disease and Public Health*, 3, 163-167.
- 471 Ponniah, J., Robin, T., Paie, M., Radu, S., Ghazali, F., Kqueen, C., Malakar, P. (2010).  
472 *Listeria monocytogenes* in raw salad vegetables sold at retail level in Malaysia.  
473 *Food Control*, 21, 774-778.
- 474 Powell, C., & Attwell, R. W. (1995). A comparative study of food retail premises by  
475 means of visual inspection and microbiological quality of food. *Epidemiology*  
476 *and Infection*, 114, 143-151.
- 477 Pui, C. F., Wong, W. C., Chai, L. C., Lee, H. Y., Tang, J. Y. H., Noorlis, A., Son, R.  
478 (2011). Biofilm formation by *Salmonella* Typhi and *Salmonella* Typhimurium  
479 on plastic cutting board and its transfer to dragon fruit. *International Food*  
480 *Research Journal*, 18, 31-38.
- 481 Rajagopal, L., & Strohbehn, C. H. (2013). Observational assessment of glove use  
482 behaviors among foodservice workers in a university dining setting: Testing a  
483 visual intervention tool. *Food Protection Trends*, 33(5), 315-324.
- 484 Ravishankar, S., Zhu, L., & Jaroni, D. (2010). Assessing the cross contamination and  
485 transfer rates of *Salmonella enterica* from chicken to lettuce under different  
486 food-handling scenarios. *Food Microbiology*, 27(6), 791-794. doi:  
487 <http://dx.doi.org/10.1016/j.fm.2010.04.011>
- 488 Sagoo, S. K., Little, C. L., & Mitchell, R. T. (2001). The microbiological examination  
489 of ready-to-eat organic vegetables from retail establishments in the United  
490 Kingdom. *Letters in Applied Microbiology*, 33(6), 434-439.

- 491 Sneed, J., Strohbehn, C., & Gilmore, S. (2004). Food safety practices and readiness to  
492 implement HACCP programs in assisted-living facilities in Iowa. *Journal of*  
493 *American Dietetic Association* 104(11), 1678-1683.
- 494 Sneed, J., Strohbehn, C., Gilmore, S., & Mendonca, A. (2004). Microbiological  
495 evaluation of foodservice contact surfaces in Iowa assisted-living facilities.  
496 *Journal of American Dietetic Associatio.*, 104(11), 1722-1724.
- 497 Sodha, S. V., Lynch, M., Wannemuehler, K., Leeper, M., Malavet, M., Schaffzin, J.,  
498 Braden, C. (2011). Multistate outbreak of Escherichia coli O157:H7 infections  
499 associated with a national fast-food chain, 2006: a study incorporating  
500 epidemiological and food source traceback results. *Epidemiology and Infection*,  
501 139(2), 309-316. doi: 10.1017/S0950268810000920
- 502 Su, L. J., & Arab, L. (2006). Salad and raw vegetable consumption and nutritional status  
503 in the adult US population: results from the Third National Health and Nutrition  
504 Examination Survey. *Journal of American Dietetic Association*, 106(9), 1394-  
505 1404. doi: 10.1016/j.jada.2006.06.004
- 506 Tebbutt, G., & Southwell, J. (1989). Comparative study of visual inspections and  
507 microbiological sampling in premises manufacturing and selling high-risk foods.  
508 *Epidemiology and Infection* 103, 475-486.
- 509 Todd, E. C., Greig, J. D., Bartleson, C. A., & Michaels, B. S. (2008). Outbreaks where  
510 food workers have been implicated in the spread of foodborne disease. Part 5.  
511 Sources of contamination and pathogen excretion from infected persons. *Journal*  
512 *of Food Protection*, 71, 2582-2595.
- 513 WHO. (2007). Food safety and foodborne illness. Retrieved from  
514 WHO. (2008). Microbiological hazards in fresh fruits and vegetables. *WHO report*  
515 *:Microbiological Risk assessment series.*
- 516 Wyatt, C., & Guy, V. (1980). Relationship of microbial quality of retail meat samples  
517 and sanitary conditions. *Journal of Food Protection* 43, 385-389.

518

Table 1. The six different constructs comprised in the visual assessment survey in SMEs

Inspection constructs	Individual Inspection Components
Construct 1: Structural compliance	<ul style="list-style-type: none"> <li>• General maintenance conditions and evidence of pest in the production environment</li> <li>• Zoning (separation of fresh produce from raw meat and poultry)</li> <li>• All major pieces of equipment such fridges, freezers ovens, hot holding equipment, cold holding equipment are fitted with working temperature monitoring gauges</li> <li>• Availability of proper handwashing sink</li> </ul>
Construct 2: Personal Hygiene	<ul style="list-style-type: none"> <li>• Wearing hair cap</li> <li>• Appropriately clean personnel protective clothing</li> </ul>
Construct 3: Sanitation	<ul style="list-style-type: none"> <li>• Clean floors, walls, overall facilities and implements</li> <li>• Waste containers are covered, kept clean</li> <li>• Sanitisers for work surfaces readily available for use during food preparation</li> <li>• Containers used to drain vegetables are kept clean</li> </ul>
Construct 4: Evidence of procedures and management system control	<ul style="list-style-type: none"> <li>• Records keeping for verification of temperature monitoring and system audits (during cooking, cooling, storing)</li> <li>• Cleaning system and schedule</li> <li>• Where a chemical sanitiser is used , there are records to show levels are maintained</li> </ul>
Construct 5: Contamination and Cross contamination control measures	<ul style="list-style-type: none"> <li>• Staff cleaning tools are stored in appropriate manner and not at risk of contaminating food or equipment during preparation</li> <li>• Staff personal belongings are stored in appropriate manner and not at risk of contaminating food or equipment during preparation?</li> <li>• Received fresh vegetable are stored in protected areas</li> <li>• Washing sink designated for fresh produce only</li> <li>• Unprocessed raw vegetables are prepared so that contamination and cross- contamination does not occur (separate cutting boards and utensils)</li> <li>• Visitors or unauthorized staff are granted protective clothing upon entry</li> <li>• Entry for authorized personnel only</li> </ul>
Construct 6: Safe and hygienic handling practices	<ul style="list-style-type: none"> <li>• Appropriate use of gloves and handwashing</li> <li>• Frozen food is properly thawed</li> <li>• Vegetable sanitizers are made up correctly</li> <li>• Food on hold is covered</li> </ul>

Table 2 Frequency of self-reported handling practices of fresh vegetables in foodservice establishments

Process	Frequency of handling practices N (%)				
	Always	Often	Sometimes	Rarely	Never
Are fresh vegetables delivered from one supplier/source?	52(68).	17 (22)	5 (7)	1 (1)	1 (1)
Are fresh leafy vegetables or/and pre-cut vegetables delivered cooled?	2 (3)	0 (0)	2 (3)	0 (0)	72 (94)
Is the washing water used for fresh vegetables and fruits chlorinated?	13(17)	0 (0)	0 (0)	0 (0)	64(83)
Do you wash the vegetables before cutting?	51 (77)	1 (1)	1 (1)	0 (0)	13 (20)
If applicable: how often you record the temperature of the display salad bar?	12 (35)	0 (0)	0 (0)	0 (0)	22 (65)
The received fresh vegetables are kept in the cold storage room/fridge	67 (93)	0 (0)	1 (1)	0 (0)	4 (6)
The washed and cut vegetables for salads and garnishes are held at room temperature before preparation/service	17 (26)	0 (0)	2 (3)	0 (0)	47 (71)

Table 3. Microbial loads of different fresh salads vegetables

<b>Produce</b>	<b>N</b>	<b>PCA<sup>†</sup></b>	<b>Coliforms<sup>†</sup></b>
Lettuce	30	5.50 ± 1.55	3.89 ± 2.19
Parsley	34	5.42 ± 1.32	4.48 ± 2.16
Cucumber	18	4.60 ± 2.01	3.52 ± 2.10
Radish	9	5.09 ± 2.20	1.72 ± 2.68
Mint	11	3.92 ± 2.74	3.93 ± 2.75
Coriander	1	7.38 ± 0.00	6.40 ± 0.00
Aragula	5	3.99 ± 2.44	3.30 ± 3.06
Tomato	3	2.90 ± 2.57	2.13 ± 2.20
Lettuce	4	5.35 ± 1.59	3.20 ± 1.49
Iceberg	3	4.54 ± 0.77	1.46 ± 2.53

<sup>†</sup>Values are mean Log CFU/g ± standard deviation.  
The minimum detection limit was 10 CFU/g.



Table 4. Mean levels of *E. coli* and coagulase-positive *Staphylococcus* spp. on salads vegetables

Produce	N	<i>E. Coli</i>	<i>Staphylococcus</i> spp.
		Log CFU/g $\pm$ SD (min-max)	Log CFU/g $\pm$ SD (min-max)
Lettuce	30	0.92 $\pm$ 1.80 (<1.00 -7.15)	2.89 $\pm$ 2.28 (<1.00 – 7.76)
Parsley	34	0.70 $\pm$ 1.50 (<1.00 - 5.40)	2.93 $\pm$ 1.87 (<1.00 – 6.16)
Cucumber	18	1.30 $\pm$ 1.43 (<1.00 - 3.40)	2.01 $\pm$ 1.99 (<1.00 – 5.45)
Radish	9	0.35 $\pm$ 0.88 (<1.00 -2.65)	2.84 $\pm$ 2.37 (<1.00 – 6.48)
Mint	11	1.36 $\pm$ 1.78 (<1.00 - 4.91)	2.69 $\pm$ 2.08 (<1.00 – 5.62)
Coriander	1	1.30 $\pm$ 0.91 (<1.00 - 1.30)	4.04
Aragula	5	0.92 $\pm$ 1.45 (<1.00 - 3.30)	2.76 $\pm$ 1.67 (<1.00 – 4.15)
Tomato	3	<1.00	2.00 $\pm$ 2.00 (<1.00 – 4.00)
lettuce	4	<1.00	4.47 $\pm$ 1.73 (2.30 – 6.00)
Iceberg	3	0.33 $\pm$ 0.58 (<1.00 – 1.00)	1.83 $\pm$ 1.58 (<1.00 – 2.78)

The minimum detection limit was 10 CFU/g.

Table 5. Bacterial counts recovered from two contact surfaces

Contact surface	N	Mean log CFU/swabbed area (min-max)			
		PCA	<i>Staphylococcus</i> spp	<i>E.coli</i>	Total coliforms
Chopping board†	29	4.99 (<1.00-8.40)	4.42 (<1.00-8.40)	1.19 (<1.00-6.02)	2.62 (<1.00-8.40)
Knife*	20	5.62 (<1.00-8.40)	4.62 (<1.00-7.98)	1.13 (<1.00-5.95)	4.31 (<1.00-8.40)

†Cutting board swabbed area of 50 cm<sup>2</sup>

\*Knife (no defined area – ca.10-20 cm<sup>2</sup>)

Table 6. Distribution of the mean Log CFU/g of bacterial loads on fresh produce according to adequacy level of control measures

Microorganism	Rating†	N	Prevention of cross-contamination	Sanitation	Protected, clean storage of fresh produce	
			Mean ± SD	Mean ± SD	N	Mean ± SD
<b><u>Coliforms</u></b>						
Lettuce	Adequate	9	3.84 ± 3.09	3.67 ± 2.93	11	3.81 ± 2.59
	Inadequate	17	3.86 ± 1.68	4.20 ± 1.98	13	4.42 ± 1.68
Parsley	Adequate	10	3.80 ± 2.20	3.97 ± 2.23	14	3.95 ± 1.94
	Inadequate	20	4.68 ± 2.19	5.35 ± 2.39	13	4.46 ± 2.69
Cucumber	Adequate	6	4.15 ± 2.42	3.92 ± 2.48	7	3.84 ± 2.35
	Inadequate	9	3.79 ± 1.82	3.47 ± 1.99	7	3.61 ± 2.06
<b><u>E.Coli</u></b>						
Lettuce	Adequate	9	1.46 ± 2.50	1.18 ± 2.17	11	1.19 ± 2.31
	Inadequate	17	0.85 ± 1.54	1.23 ± 1.77	13	0.85 ± 1.56
Parsley	Adequate	10	0.54 ± 0.97	0.79 ± 1.55	14	1.15 ± 2.05
	Inadequate	20	0.65 ± 1.48	0.81 ± 1.83	13	0.63 ± 1.15
Cucumber	Adequate	6	1.96 ± 1.47	1.79 ± 1.47	7	1.68 ± 1.53
	Inadequate	9	1.29 ± 1.43	0.91 ± 1.47	7	1.36 ± 1.53
<b><u>PCA</u></b>						
Lettuce	Adequate	9	6.14 ± 1.71	6.10 ± 1.54	11	5.41 ± 1.63
	Inadequate	17	5.21 ± 1.40	5.07 ± 1.32	13	5.41 ± 1.63
Parsley	Adequate	10	5.51 ± 1.51	5.48 ± 1.29	14	5.31 ± 1.28
	Inadequate	20	5.49 ± 1.21	5.30 ± 1.29	13	5.42 ± 1.55
Cucumber	Adequate	6	5.87 ± 1.22	4.36 ± 2.72	7	5.84 ± 1.11
	Inadequate	9	4.09 ± 1.82	4.84 ± 1.11	7	3.87 ± 1.96
<b><u>Staphylococcus</u></b>						
Lettuce	Adequate	9	2.83 ± 1.73	3.36 ± 2.13	11	3.20 ± 1.91
	Inadequate	17	2.67 ± 2.43	2.53 ± 2.55	13	2.84 ± 2.90
Parsley	Adequate	10	2.85 ± 2.17	3.16 ± 1.87	14	3.18 ± 1.89
	Inadequate	20	2.95 ± 1.78	2.26 ± 1.97	13	2.13 ± 2.08
Cucumber	Adequate	6	1.80 ± 2.02	1.56 ± 1.82	7	1.91 ± 1.87
	Inadequate	9	2.53 ± 2.12	3.24 ± 1.97	7	2.86 ± 2.12

†"Incomplete" ranking was omitted for easier presentation of data




Figure 1. The distribution of total score obtained from the overall visual assessment of hygiene conditions and handling practices.




Figure 2. Distribution of food businesses' compliance with basic hygiene requirements and control measures




Figure 3. Distribution of food businesses' adequacy level in relation to washing and storing practices of fresh salads vegetables




Figure 4. The distribution of microorganism levels on fresh vegetables in relation to the different values of visual assessment scores obtained on all inspected components




Figure 5 Distribution of mean levels of *Staphylococcus* spp. in relation to component "Availability of handwashing facilities"






Figure 6. Distribution of *Listeria* spp. in relation to the visual assessment scores on all inspected components during salad vegetables preparation.

**The association of microbiological quality and handling practices of ready-to-eat fresh salad vegetables with food safety environment in restaurants:  
Case study in Lebanon**

**Highlights**

1. Microbial loads on salads vegetables, hygienic conditions/practices were assessed.
2. Association of microbiological quality with visual assessment scores was tested.
3. *Listeria monocytogenes* and *Salmonella* spp. were detected.
4. There was no significant relationship with the total visual assessment scores
5. Correlation of cross-contamination components to *Listeria* levels was significant
6. Poor cleaning can possibly be linked to *Listeria* levels in salads vegetables.