

## Manuscript Details

<b>Manuscript number</b>	AEOLIA_2016_55
<b>Title</b>	Our evolving understanding of aeolian bedforms, based on observation of dunes on different worlds
<b>Article type</b>	Review Article

### Abstract

Dunes, dune fields, and ripples are unique and useful records of the interaction between wind and granular materials – finding such features on a planetary surface immediately suggests certain information about climate and surface conditions (at least during the dunes' formation and evolution). Additionally, studies of dune characteristics under non-Earth conditions allow for “tests” of aeolian process models based primarily on observations of terrestrial features and dynamics, and refinement of the models to include consideration of a wider range of environmental and planetary conditions. To-date, the planetary aeolian community has found and studied dune fields on Mars, Venus, and the Saturnian moon Titan. Additionally, we have observed candidate “aeolian bedforms” on Comet 67P/Churyumov-Gerasimenko, the Jovian moon Io, and – most recently -- Pluto. In this paper, we hypothesize that the progression of investigations of aeolian bedforms and processes on a particular planetary body follows a consistent sequence – primarily set by the acquisition of data of particular types and resolutions, and by the maturation of knowledge about that planetary body. We define that sequence of generated knowledge and new questions (within seven investigation phases) and discuss examples from all of the studied bodies. The aim of such a sequence is to better define our past and current state of understanding about the aeolian bedforms of a particular body, to highlight the related assumptions that require re-analysis with data acquired during later investigations, and to use lessons learned from planetary and terrestrial aeolian studies to predict what types of investigations could be most fruitful in the future.

<b>Keywords</b>	Planetary; terrestrial; aeolian bedforms; aeolian science; dunes; ripples
<b>Corresponding Author</b>	Serina Diniega
<b>Corresponding Author's Institution</b>	Jet Propulsion Laboratory
<b>Order of Authors</b>	Serina Diniega, Mikhail Kreslavsky, Jani Radebaugh, simone silvestro, Matt Telfer, Daniela Tirsch

## Submission Files Included in this PDF

### File Name [File Type]

Replytoreviewers.docx [Response to Reviewers]

Supplement\_yearcounts\_references.docx [Response to Reviewers]

AeolSciEvol\_manuscript\_revise\_v3\_formatted\_noinfo\_acc.docx [Manuscript (without Author Details)]

AeolSciEvol\_manuscript\_revise\_v3\_formatted\_noinfo.pdf [Editor-Annotated Manuscript]

Titlepage\_AeolianSciencef.pdf [Title Page (with Author Details)]

To view all the submission files, including those not included in the PDF, click on the manuscript title on your EVISE Homepage, then click 'Download zip file'.

Manuscript number: AEOLIA\_2016\_55

Title: Our evolving understanding of aeolian bedforms, based on observation of dunes on different worlds

(Previous Title: Our evolving understanding of aeolian bedforms, based on studies of different Worlds)

Article type: Review Article

We thank the two reviewers and the editor for their insightful and helpful comments. In particular, the many reference suggestions and questions about the specific types of investigations we focused on in constructing our “Phase” framework helped point out areas of confusion or incompleteness. We believe this revision much more clearly lays out our inputs, resultant framework, and overall aims.

In the following pages, we have replied in detail to reviewer comments (our replies are in *italics*). Changes can also be found within the tracked-changes manuscript Word document. Additionally, below, we address a couple of large-scale revisions:

- As stated above, thank you all very much for the reference suggestions. We added most of them in (and explain below if we did not). These strengthen the paper and help fill gaps.
- We edited the title and added a paragraph to the Introduction to narrow our focus to dunes on planetary bodies, as studied through observational data. Dunes are usually the first aeolian bedform seen on a body (there’s a slight exception for Venus, where surface observations by the Venera missions pre-dated Magellan radar images – we touch on this in the text) and are a highly useful record of atmosphere and surface conditions (and can provide quantitative constraints on both, via models of their formation processes). We have highlighted more strongly that there are parallel investigations that focus on model development/refinement, on laboratory work, and on field studies (sometimes the feature is studied within the context of model development or sometimes as terrestrial analogs) -- that are touched on but which are not completely described within this paper.
- Although both reviewers asked about TARs (with good reason), as well as other aeolian bedforms -- we have not included discussion of those features. In the particular case of TARs, their formation mechanism is still debated and thus discussion of them is much less straight-forward than for aeolian dunes. However, as noted in our new Introduction paragraph, we would expect that a similar framework would work for them.
- One reviewer also expressed concern that the summaries at the end of each phase section may not work very well in their present format. The goal of these end-portions were to provide easy-to-find-and-read summaries of each Phase. We have added an alternative version which perhaps may be easier to read at the end of Phase 1 – what do you think of

the Table version? If this is an improvement, we can easily turn all summaries into a Table.

Thank you again for your assistance in improving this manuscript!

Thank you,

Serina Diniega and co-authors

### **Editor's (Matt Chojnacki) comments**

- Ln 351-356 and related text: Consider the following reference which used multiple criteria to constrain dune sediment sources: Chojnacki, M., D. M. Burr, J. E. Moersch, and J. J. Wray (2014b), Valles Marineris dune sediment provenance and pathways, *Icarus*, 232(0), 187–219, doi:10.1016/j.icarus.2014.01.011.  
*Reference has been added.*
- Fig. 9's caption needs a citation (Silvestro et al. 2013).  
*Added.*
- Ln 632 and others: "Chojnacki"  
*Apologies! This has been fixed.*
- Ln 640-643 (for sediment fluxes) and Ln 640-643 (for dune ages termed "turnover times") and related text: Consider the following reference: Chojnacki, M., J. R. Johnson, J. E. Moersch, L. K. Fenton, T. I. Michaels, and J. F. Bell III (2015), Persistent aeolian activity at Endeavour crater, Meridiani Planum, Mars; new observations from orbit and the surface, *Icarus*, 251(0), 275–290, doi:10.1016/j.icarus.2014.04.044.  
*Added.*
- Please cite all instruments (HiRISE, THEMIS etc.) appropriately.  
*I assumed this comment was regarding the images used within the figures ... I have checked the captions and now have identified in each what instrument acquired the discussed data.*

### **Reviewer 1's (Lori Fenton) comments:**

The author(s) propose that scientific investigations of planetary aeolian bedforms follow a predictable progression of inquiry, with the aim of understanding how specific data set types either contribute to knowledge gain or leave knowledge gaps. They identify a striking sequence of methods and perspectives used as understanding of planetary bedforms increases, which necessarily differs significantly from the progression of terrestrial aeolian science. The proposed sequence (Phases 1-7) is simpler than the whole available literature would contain (see general and specific comments below). Many studies, aeolian features, and ideas have not been included here, likely leading to the oversimplified nature of the sequence. However, the proposed sequence is essentially consistent with the published literature, and does provide insight for both planetary and terrestrial aeolian scientists. I suggest that, following the comments below, this manuscript should be published with major revisions.

I generally disapprove of not revealing myself to authors, in the hopes that open communication can help clarify concerns and more quickly answer the authors' questions (hopefully leading to the best science and quicker revisions). I encourage the authors to contact me to address any such questions regarding this review.

- Lori Fenton lfenton@seti.org

### **General comments**

1. The work seems to be limited by not including the full available literature for Mars. For example, the first global maps (which define Phase 4) of martian dunes were produced by Thomas (1982) and Ward et al. (1985), which were not included here, and are not shown in Fig. 13 (see specific comments below for these and other references – there are many striking gaps, and I did not include all relevant references). The authors need to include a more thorough search of the rather extensive martian literature, which could be illuminating. Note also that many papers address more than one phase.

*Based on this (thank you, Lori!) and other reviewers' comments, we have added in many more references, especially regarding Mars but also some additional studies for Titan, Venus, and the comet -- and these were valuable additions. However, we note that we cannot aim to include all papers while keeping this manuscript to a reasonable length and focus. We do not aim to present a fully comprehensive review of e.g., Mars aeolian research – although we do try to accurately represent the full scope of work when placing this work within the proposed Phase framework. We have aimed for a reasonable sampling, and hope this is enough to justify our proposal, and to provide a starting ground for any reader interested in learning more on that (or any) particular topic.*

*In general, we have edited the Title and Introduction text to clarify the focus of this paper – especially the new 3<sup>rd</sup> paragraph of the Introduction.*

2. I suspect that the advent of new missions (on Mars that would include three “pulses” of activity: Mariner/Viking in the 1970s, MGS/MPF in the 1990s, and the more recent MO/MRO/MEX/MER/MSL missions in the past 10 years) or new methods (e.g., mesoscale models, use of gross bedform-normal transport, the recent introduction of “fingering” vs. “instability” modes, etc.) can cause renewed activity in lower phases that produces a more thorough understanding.

*Absolutely – this is a point of the paper; by defining a framework, it should be easier to see when renewed activity occurs and thus to identify where updated understanding should flow, with improved constraints on assumptions, a need to re-do a study with increased or improved data, etc. A sentence emphasizing this has been added to the Introduction.*

3. Figure 13 is very informative and clearly demonstrates progression through the phases. However, the horizontal bars do not adequately represent progression within a given phase. I suggest that, using the more thorough literature search from point 1 above, the horizontal bars can be broken into individual years, showing the number (or fraction) of publications that year that addressed each phase. This would nicely show that new methods and data sets produce new pulses of research activity, and reveal the complexity of progression through the phases, as it is not likely to be linear or even monotonic in time.

*This was an intriguing idea. We considered several ways of collecting such information and then displaying it. Our chosen method: we focused first on the papers cited within this manuscript (so that readers have some idea of where the information comes from) and showed what years are included in this reference list with darker colors within Figure 13 (also are listed in the supplemental table). A more general “period” for studies within each Phase was then shown with the light & dark colors. This was based on a looser categorization of the papers (e.g., these included smaller contributions to an additional Phase, versus just the Phase(s) under which the paper was cited) and supplemental checks via google scholar (e.g., we searched for references under keywords “Venus dune” and saw the few years that came up). This latter method was an attempt to mitigate the limitations of our reference list -- as many papers (cited and otherwise) contribute towards more than one Phase, there are many papers that we did not cite, and there were also many abstracts and papers presenting work that was eventually compiled into a more complete/seminal peer-reviewed publication.*

*From all of this, we updated Figure 13 (and its caption).*

4. There is ambiguity over whether the manuscript seeks to address only dunes, or any bedforms. If considering only dunes, then the features on Io and 67P/C-G may not (yet) qualify for discussion, and it may not make sense to discuss the “large martian ripples” in

great detail. The title suggests the authors are discussing any aeolian bedforms, but the main discussion is on dunes (e.g., there is no mention of TARs, the study of which may follow a similar progression of inquiry). Much knowledge on Mars has been gained from ripples (migration rates), TARs, wind tails, and sand drifts (El Dorado) that are not properly classified as “dunes”. While including all aeolian features (e.g., wind streaks, loess deposits, yardangs) would be overly exhaustive (although worth study), I recommend that the manuscript expand to include all (aeolian) bedform types, and thus include, e.g., lee dunes, TARs, coarse-grained ripples, and wind shadows.

*As discussed under point (1) above, to keep this manuscript to a reasonable length, we have focused it onto dunes, so as to present one evolution sequence. Other aeolian features, such as wind shadows, could also be discussed within a similar framework. For TARs specifically: we had previously decided to not mention them as there is not yet a well-accepted description of their formation mechanism (they’ve been identified as transverse dunes forming within a reversing wind regime, granule ripples, and sub-aerial anti-dunes), and so the information these features provide about the environment isn’t as clear or easy to trace as interpretation of dunes. However, as pointed out by both reviewers – this omission without explanation is confusing. So, we have added a note to the Introduction, explaining the focus of this paper and explicitly noting that other “aeolian features” (including TARs) are not included. With mention of the TARs, we have also included the point that there may be no good Earth analogue for these strange features, as this is an interesting idea.*

*In general, we have edited the Title and Introduction text to clarify the focus of this paper.*

5. I suggest including a Phase 0: “Predictions”, made prior to direct observation of bedforms. Some discoveries have been quite surprising (e.g., Io, 67P/C-G), whereas others were considered long before spacecraft data were obtained (e.g., Mars, Titan, Pluto). How correct have the predictions been, and what lessons can be learned to aid predictions for future missions (e.g., Venus)? Given the more surprising locations where putative bedforms have been observed, what other exotic locations can be considered (e.g., on the surfaces of Neptune and Uranus, if any such solid interface exists?).  
*This is a very good idea. We have not added an actual Phase 0 to the discussion (for one thing, describing this in sequence did not seem feasible without having gone through the discussion at least within Phase 1 – and thus seemed to be giving away the punchline before telling the joke), but we have added this intriguing point to the conclusion. And within Phase 1, we discuss predictions that have been made (in either direction, and with different degrees of success).*

6. Finally, I wonder if any similar study has been done for other geologic/atmospheric processes/features found on several planetary worlds. Has anybody considered the sequence of study methods for volcanism or impact cratering throughout the Solar System? If not, then does this investigation of aeolian processes provide a platform for any such studies to work from?

*The authors have not come across any such studies – those would be intriguing to read, and to see if the advancement of science understanding follows a similar progression. (I would imagine so for features that rely upon similar data sets, which would include impact cratering and volcanism as suggested.)*

### **Specific suggestions:**

- Line 72: typo: "... each Phase, s, we use..."  
*Fixed.*
- Table 1: suggest changing "dune" to "bedform"
  - Phase 2, "Unit": "Dune morphology" is not a unit or feature of interest. Suggest: "Dune, aeolian surface materials"  
*We disagree that "morphology" cannot be described as the "feature of interest," although perhaps "characteristic of interest" may be more correct. "Unit" has been removed.*
  - Phase 2: "composition" doesn't seem to fit here as this row is worded. Could also include thermal properties, relief, surface roughness.  
*"Characteristics of surface materials" has been added to include those options.*
  - Phase 3: Suggest "Morphological analysis of the dunes within a field"  
*Phase 3 we have used "Pattern analysis" as is suggested in a later comment (Line 394).*
  - Phase 4: Suggest also including regional studies here. Many informative studies focus on particular regions in greater detail. This could also be added as a subsequent phase. Note that both global and regional analyses of these features often include atmospheric modeling.  
*"Regional" has been added.*
  - Phase 5: could also include erosional features (e.g., gullies).  
*Phase 5 focuses on evidence of wind interaction with the dunes, and so gullies are left in Phase 6 (more general "activity" and dune evolution).*
- Line 104, 143, etc.: Suggest using "saltatable grains" rather than "sand grains". It's possible that in some unusual planetary environments, the saltated grains could be larger or (less likely) smaller than sand (e.g., there are reports of saltating pebbles in katabatic flows in Antarctica, or the >mm-sized grains on comet 67P/C-G).  
*As discussed within Phase 1, "sand" is commonly used to refer to a specific absolute grain size, but in fact can refer to a grain size defined by how the grains are moved by the*



*wind (i.e., are saltatable). We retain use of “sand grains” or “grains” throughout the document as this was much cleaner/smoother in usage than “saltatable grains,” but we have added text to make the meaning clearer.*

- Line 117: See also Edgett, K. S., and N. Lancaster (1993), Volcaniclastic aeolian dunes: terrestrial examples and application to martian sands, *J. Arid Environ.*, 25(3), 271–297, doi:10.1006/jare.1993.1061.

*Reference has been added.*

- Lines 129-130: “direction and velocity”: Suggest “direction and speed”, as velocity inherently incorporates both direction and speed.
- Line 150: “every deeply-studied body with an atmosphere...” Suggest including “and an observable surface”. Jupiter and Saturn are also deeply-studied bodies with atmospheres...
- Lines 154-158: Excellent points, but a very long sentence. Suggest breaking it up.
- Line 195: Use if “either” implies two possibilities, but three are listed.

*Text changed as suggested.*

- Line 202: On comparing venusian bedforms to underwater terrestrial bedforms, see also Neakrase (2015) <http://www.hou.usra.edu/meetings/dunes2015/pdf/8023.pdf>.
- Line 208: See also the description and measurements of ripples and “dune-like structures” in Thomas et al. (2015). They also argue for the airfall mechanism of initiating transport. Thomas, N. et al. (2015), Redistribution of particles across the nucleus of comet 67P/Churyumov-Gerasimenko, *Astron. Astrophys.*, 583(A17), doi:10.1051/0004-6361/201526049.

*References were added.*

- Line 249: typo “imagery”

*Fixed.*

- Line 283: Another question: “If conditions are unusual (e.g., tenuous atmosphere), how does sustained saltation and/or reptation occur on this world?” (This comes from considering the unusually low air pressures on Io and 67P/C-G.)

*Question added.*

- Line 302: Suggest “350 m/px”
- Line 309: Remove “via”

*Text changed as suggested.*

- Line 309-312: See Rubin and Hunter (1987) and Rubin and Ikeda (1990). Longitudinal dunes form when two winds of roughly equal transport strength (i.e., they move the same amount of sand) are  $>\sim 90^\circ$  apart. Essentially the crestline is trying to be as normal to the incident wind as possible. When two winds are  $<\sim 90^\circ$  apart they produce transverse dunes (i.e., they are “parallel enough” to work constructively to produce the same crestline alignment). Inbetween these extremes are oblique dunes, which have characteristics of both types – most dunes likely have some oblique component to them. Rubin, D. M., and R. E. Hunter (1987), Bedform alignment in directionally varying flows, *Science*, 237(4812), 276–8, doi:10.1126/science.237.4812.276. Rubin, D. M., and H. Ikeda (1990), Flume experiments on the alignment of transverse, oblique, and longitudinal dunes in directionally varying flows, *Sedimentology*, 37, 673–684. *References added. (Thank you, I was looking for these!)*
- See also Courrech du Pont et al. (2014) and other related publications, who found that dunes align either in a “bed instability mode”, which can approximate the gross bedform-normal transport alignment found by Rubin and Hunter (1987) and Rubin and Ikeda (1990), or (in conditions of low sediment supply/availability), in a “fingering mode”, which can approximate a longitudinal direction parallel to the resultant drift direction. Courrech du Pont, S., C. Narteau, and X. Gao (2014), Two modes for dune orientation, *Geology*, 42(9), doi:10.1130/G35657.1. The two modes identified in China by Rubin and Hesp (2009) and applied to Titan dunes (as described in Sec. 3.2) likely occur for reasons of changing sediment availability (i.e., the “sticky” sediment leads to lower sediment availability, so these dunes formed in “fingering mode”).
- This is an example of my general comment #2 above: that introduction of a new model for dune alignment has redefined aeolian science, and prompted reanalysis of previous data sets. In this case, the study of Titan dunes occurred as this recent understanding was being developed. *A very good point. This reference and description has been added within Section 3.2.*
- Lines 327-328: The text (outside the Fig. 6 caption) does not describe what processing has been done to this image. Suggest rewording and/or a brief description in the text (or removing this frame from the figure). *This image (6b) has not really been "processed", aside from the fact that it's a SAR image. Text has been added to the caption to better explain this – “The right image (b), a Synthetic Aperture RADAR (SAR) image, shows bedrock as bright, because it is rough, and unorganized dune sands as dark, because dunes are smooth at the SAR wavelength of 6 cm (black areas are regions devoid of data returned to the SAR antenna).”*
- Line 336: Surely images from Viking 1 have been used for mapping as well? Also see Ward et al. (1985) and Thomas (1982). Ward, A. W., K. B. Doyle, P. J. Helm, M. K.

Weisman, and N. E. Witbeck (1985), Global Map of Eolian Features on Mars, *J. Geophys. Res.*, 90(B2), 2038–2056, doi:10.1029/JB090iB02p02038. Thomas, P. (1982), Present wind activity on Mars - Relation to large latitudinally zoned sediment deposits, *J. Geophys. Res.*, 87(B12), 9999–10,008, doi:10.1029/JB087iB12p09999.

*Very good point – references have been added and this start on Phases 3-4 has been added to Figure 13.*

- Lines 344-347: The work by Fenton et al. (2005) was done using MOC NA images, worth mentioning so readers don't assume it was done with Mariner 9 and Viking images (as stated in the first sentence of the paragraph). Also, this work used a mesoscale model, not a GCM.

*Text has been fixed.*

- Note also that the first comparisons with atmospheric models were done before MGS: see Greeley, R., A. Skyepeck, and J. B. Pollack (1993), Martian aeolian features and deposits: comparisons with general circulation model results, *J. Geophys. Res.*, 98(E2), 3183–3196.

*Thank you for the pointer. However, this reference has not been added as the aeolian features considered for this comparison did not include dunes. The study focused instead on bright wind streaks, dark wind streaks, and yardangs.*

- Line 353: Suggest “proposed” instead of “identified”, as these sources are not confirmed. See also Geissler, P. E., N. W. Stantz, N. T. Bridges, M. C. Bourke, S. Silvestro, and L. K. Fenton (2013), Shifting sands on Mars: insights from tropical intra-crater dunes, *Earth Surf. Process. Landforms*, 38(4), 407–412, doi:10.1002/esp.3331.

*Text has been fixed. And the reference has been added to that section (in addition to where it was already cited).*

- Line 361: Suggest “proposed” instead of “identified”; the presence of microdunes are by no means verified.
- Line 369: typo “withing”
- Lines 375-381: Suggest that Phase 2 can continue *concurrently* with initiation of Phases 3-5, with the introduction of new observations and the development of models, and analysis methods.

*Text changed as suggested.*

- Table 2:

- Suggest “low albedo” rather than “dark albedo”.

*Fixed.*

- If not including Io and Pluto here, then why include 67P/C-G?

*Analysis of the comet has passed peer review, while analysis of the Pluto and Io features have not yet. (This distinction is made within the title of Table 2: “as presented in the literature.”)*

- Line 394: Suggest calling this phase “Pattern analysis”, as simply “analysis” alone could mean anything, and is used to describe other phases.  
*Done (related: see Table 1 comment above).*
- Line 403: Suggest “upwind margin” rather than “start” (which could imply time rather than space) and “lateral margins” rather than “boundaries”. Note that not all dune fields have a single upwind and downwind margin (e.g., many intracrater dune fields on Mars), complicating this issue.
- Line 438: Suggest “uniformity” rather than “consistency”.
- Line 446: The knowledge gain is considerably more than just the dune field pattern and shape. The pattern reveals the maturity state, and perhaps relative age, of the bedforms (defined by defect density, which is not directly discussed but hinted at in lines 436-440) and possible temporal changes in the sediment state (e.g., sediment supply, wind patterns).  
*Text changed/added to as suggested.*
- Line 456: In this section, consider including discussion of the martian global (or nearly global) inventories of Thomas (1982), Ward et al. (1985), Hayward et al. (2007; 2014) and the included discussions of world-scale processes and factors (e.g., sediment sources, characteristic dune field sizes, location relative to topography and geology, relation to global-scale wind circulation, etc.). Hayward, R. K., L. K. Fenton, and T. N. Titus (2014), Mars Global Digital Dune Database (MGD3): Global dune distribution and wind pattern observations, *Icarus*, 230, 38–46, doi:10.1016/j.icarus.2013.04.011.  
*References have been added, along with a bit of discussion about global studies of martian aeolian bedforms.*
- Lines 464-465: Suggest “...terrestrial dunes can rely on, for example, detailed petrographic...” Other studies have done the same with different methods.  
*Changed “can” to “may” to address this point.*
- Line 470: Suggest “proposed” rather than “identified”, as the source of the NPSS is still debated. Some mention should also be made of the extensive aeolian deposit underlying the current polar cap (which is the most recent major sand source), making the extent of the sand reservoir even more vast. See Byrne and Murray (2002). Byrne, S., and B. C. Murray (2002), North polar stratigraphy and the paleo-erg of Mars, *J. Geophys. Res.*, 107(E6, 5044).  
*Text changed and references has been added.*
- Line 493: Suggest “incorporated” rather than “absorbed”.
- Line 505: Suggest “crater retention age of <10 000 years”, as the dunes may be much older.  
*Text has been fixed.*
- Lines 531-532: Note that some early global-scale studies (limited simply by the low resolution of earlier data sets) may produce the larger-scale sediment transport pathways

or atmospheric circulation interpretations (Phase 4) sooner than the “field-specific results of Phase 2 and 3”. (e.g., Ward et al., 1985). The progression through the phases is not necessarily directly in order.

*This point has been added to the summary.*

- Line 539: Crater-retention age, rather than actual age.  
*Text has been fixed.*
- Line 543: “Details” as described here refers to “superposed bedforms”, suggest being more specific.  
*Phase name is now “Analysis of superposed bedforms on the dune formed due to wind interaction with the dune.”*
- Line 548: Suggest “as fine as” in place of “up to” (0.25 m/pixel...)  
*Text changed as suggested.*
- Lines 631-636: Actually there were a few publications indicating observations of bedform movement/change prior to the HiRISE overlap studies (Fenton, 2006; Bourke et al., 2008; Sullivan et al., 2008). Fenton, L. K. (2006), Dune migration and slip face advancement in the Rabe Crater dune field, Mars, *Geophys. Res. Lett.*, 33(20), 1–5, doi:10.1029/2006GL027133. Bourke, M. C., K. S. Edgett, and B. A. Cantor (2008), Recent aeolian dune change on Mars, *Geomorphology*, 94(1-2), 247–255, doi:10.1016/j.geomorph.2007.05.012. Sullivan, R. et al. (2008), Wind-driven particle mobility on Mars: Insights from Mars Exploration Rover observations at “El Dorado” and surroundings at Gusev Crater, *J. Geophys. Res.*, 113, E06S07, doi:10.1029/2008JE003101.  
*Thank you – references have been added.*
- Line 630: Suggest also including features indicative of inactivity, which is also revealing about bedform migration rates (e.g., fissures on north polar dunes from Portyankina et al., 2012, pits and other features on southern midlatitude dunes from Fenton and Hayward, 2010). Portyankina, G., A. Pommerol, K.-M. Aye, C. J. Hansen, and N. Thomas (2012), Polygonal cracks in the seasonal semi-translucent CO<sub>2</sub> ice layer in Martian polar areas, *J. Geophys. Res.*, 117(E2006), doi:10.1029/2011JE003917.  
*This is a very good point -- it has been added to the end of Phase 6 (with the suggested reference).*
- Lines 642-643: Note that the sand fluxes are comparable to those in the Antarctic Dry Valleys, which are low compared to those elsewhere on Earth.  
*Text has been fixed.*
- Line 658: Suggest also Kereszturi, A., D. Möhlmann, S. Berczi, T. Ganti, A. Horvath, A. Kuti, A. Sik, and E. Szathmary (2010), Indications of brine related local seepage phenomena on the northern hemisphere of Mars, *Icarus*, 207(1), 149–164, doi:10.1016/j.icarus.2009.10.012.

*Within the paper, we discuss features that move significant movement of sand over the dune lee slopes. This paper discusses a different type of activity on the dunes, but over a scale that is much smaller and thus is not as relevant for analysis of present-day dune evolution. Thus, this reference has not been added.*

- Line 685: typo “dunes slopes”, “levels of aeolian activity”
- Line 692: “were” in place of “was”
- Line 719: Suggest adding “How can observations of sediment grain size and bedform morphology provide insight regarding transport processes and the nature/frequency of mobilization events?”  
*Text has been changed as suggested.*
- Line 776-778: Sullivan et al. (2008) would be a good reference for this. They proposed that the meter-scale bedforms on the El Dorado deposit in Gusev crater were sand ripples.  
*Reference has been added.*
- Line 905: typo “transverse”  
*Text has been fixed.*
- Line 932-950: The case of the Amazonian paleo-dune fields seems to be more of a lesson that terrestrial dune researchers can learn from planetary dune researchers than the reverse. Suggest relocation to the following section.  
*The text has been moved, as suggested, to the end of the next section.*
- Line 954-955: “no global catalogue of dunes for Earth” See Lancaster et al. (2015). However, it is telling that this database was assembled many years after the first planetary examples. Lancaster, N. et al. (2016), The INQUA Dunes Atlas chronologic database, in press at *Quat. Int.*, doi:10.1016/j.quaint.2015.10.044.  
*This reference has been added, with some explanation of what it is -- despite the name, this paper is a database of dated dune samples, rather than a spatial atlas of dune extent, morphology, etc. (For instance, currently very-active regions such as the Sahara scarcely feature in this database.) It is telling that a comprehensive global database of the type to be expected within planetary science has not yet been created for Earth, and even partial catalogs have been put together only in the current decade!*
- Line 955: typo “editions”  
*Text has been fixed.*

## **Reviewer 2's comments:**

This is an interesting review of progress in planetary dune studies, with a valuable conceptual framework to organize the recent history of investigations and to refer to the often scattered literature on planetary dunes.

- The conceptual organization into 7 phases provides a good way to organize understanding of studies. It seems to me that phases 6 and 7 could be combined, as both emphasize the "details" - which I would prefer to call ""dynamics". There is a natural progression from inferring dynamics from morphology, to actual observations of processes and dynamics.

*While we agree that there is a natural progression between and partial overlap of studies fitting into Phases 6 and 7, we want to highlight the different temporal/spatial scale, frequency, and mode of observation that is used in the two Phases, as these lead into different types of investigations. Phase 6 can rely solely on orbital images (even if that leads down the wrong path), although of course in situ observations can assist; and so the dynamics observed within Phase 6 can span a much longer time period than in situ observations may allow for, with coarser temporal resolution. Phase 7 may involve no observation of dynamics, or only evidence of sand transport and not bedform evolution; but instead can yield groundtruth information about grain characteristics and bedform scale. Thus we maintain them as separate phases.*

- I am not sure that the summaries at the end of each phase section work very well in their present format. I suggest that they be re-written as continuous prose, so that each provides a lead into the next section.

*The goal of these end-portion were to provide easy-to-find-and-read summaries of each Phase. Turning them into smooth prose would lengthen them significantly, and thus somewhat negate this intent. We have added an alternative version which perhaps may be easier to read, at the end of Phase 1 – what do you think of the Table version? If this is an improvement, we can easily turn all summaries into a Table.*

- Inevitably, such a review is selective, and one can quibble about the selection of material and examples. One example is the use of spectral information to examine dune sediment composition. There is a lot more that could be said here (especially for Mars), because compositional information is key to understanding many aspects of planetary geologic and climatic evolution. Another example is a bed form that has proved difficult to understand is the transverse aeolian bedforms (TARs) of Mars - this is a good example of where Earth analogues have failed to help.

*Based on other reviewers' comments, we have added in a few more references relevant to compositional studies on Mars. We agree we cannot include them all, while keeping this manuscript to a reasonable length! We have aimed for a reasonable sampling, and hope*

*this is enough to provide a starting ground for any reader interested in learning more on that (or any) particular topic.*

*As for TARs, we had previously decided to not mention them as there is not yet a well-accepted description of their formation mechanism (they've been identified as transverse dunes forming within a reversing wind regime, granule ripples, and sub-aerial anti-dunes), and so the information these features provide about the environment isn't as clear or easy to trace as interpretation of dunes. However, as pointed out by both reviewers – this omission without explanation is confusing. So, we have added a note to the Introduction, explaining the focus of this paper and explicitly noting that other “aeolian features” (including TARs) are not included. With mention of the TARs, we have also included the point that there may be no good Earth analogue for these strange features, as this is an interesting idea.*

*In general, we have edited the Title and Introduction text to clarify the focus of this paper – especially the new 3<sup>rd</sup> paragraph of the Introduction.*

- One aspect that could be emphasized more is the linkages between planetary and terrestrial aeolian studies. It is clear that in many cases questions raised by planetary studies have resulted in a re-examination of the basic physics of particle movement (e.g Greeley and Iverson work in the 1970's on thresholds for particle movement), in addition to prompting further study of terrestrial aeolian features.  
*More discussion and references around this point have been added to various parts of the manuscript (in particular, within the caption for Figure 1), to emphasize this point.*

### **Some specific comments**

- Some additional sub headings within each section would be very helpful - to emphasize the examples discussed - and to break up the lengthy text  
*This was tried (as we agree the text is lengthy). However, while a few sections had natural subsection break points, others did not; and so we decided to leave subsection headers out rather than create an inconsistent organizational framework.*

*We also note that part of the difficulty that we had with this is that we did not want to create subsections that divided discussion about different planets -- Part of the goal of this paper was to show that all planetary studies fall onto a common framework.*

- In the introduction, the conceptual framework for dune field dynamics is perhaps best exemplified by Kocurek, G., Lancaster, N., 1999. Aeolian system sediment state: theory and Mojave Desert Kelso dune field example. *Sedimentology* 46, 505 - 515.  
*This is an intriguing summary of dune field dynamics, and a pointer towards it is now included within the Phase 1 discussions (as a more technical description of the fairly*




*basic points discussed within this manuscript). We did not add it to the Introduction as there are no other citations within that portion of the manuscript.*

- A good summary of the status of knowledge of Martian aeolian and dune studies at the end of the Viking era is provided by Greeley R., Lancaster, N., Lee, S., Thomas, P., 1992. Martian aeolian processes, sediments and features, in: Kieffer, H., Jakosky, B.M., Snyder, C.W., Matthews, M.S. (Eds.), Mars. University of Arizona Press, Tucson, pp. 730-767.

*Reference has been added.*

Phase of aeolian bedform study on a planetary body		Mars	Venus	Titan
1	<b>Recognition of dune(s)</b>	71 - 1, 72 - 2, 73 - 1, 76 - 1	92 - 1, 94 - 1	06 - 1
2	<b>Analysis of gross individual dune characteristics: e.g., morphology and composition</b>	79 - 2, 81 - 1, 82 - 1 92 - 1 05 - 1 10 - 1, 11 - 1, 13 - 1, 16 - 2	92 - 1, 95 - 1, 97 - 1, 99 - 1 15 - 1	08 - 2, 09 - 1, 10 - 1, 13 - 1, 14 - 1, 15 - 1
3	<b>Pattern analysis of the dunes within a field, including variations due to e.g., sediment supply and wind variations</b>	79 - 1, 81 - 1 05 - 2, 06 - 1 10 - 3, 11 - 1, 12 - 2, 14 - 1, 15 - 1, 16 - 3	94 - 1 06 - 1	11 - 1, 14 - 1
4	<b>Regional and global surveys and aggregate-analysis of dune characteristics, with a re-aggregation of data for e.g., estimates of age or sand volumes, identification of large-scale sediment transport pathways, or identification/estimation of the effect of location-related non-aeolian processes</b>	82 - 1, 83 - 1, 85 - 1 92 - 1 00 - 1 03 - 1 07 - 2, 09 - 1, 10 - 1, 11 - 2, 12 - 1, 14 - 1	94 - 1	09 - 1, 11 - 1, 12 - 1, 13 - 1, 14 - 1, 15 - 3, 16 - 2
5	<b>Analysis of dune superposed bedforms on the (such as ripples) formed due to wind interaction with the dune</b>	06 - 1 10 - 1, 11 - 1, 12 - 3, 13 - 1, 14 - 1, 15 - 1, 16 - 6		
6	<b>Observation of activity on the dune, including non-aeolian activity</b>	(non-activity) 00 - 1 06 - 1, (dome dunes disappearing) 08 - 1 10 - 1, 11 - 2, 12 - 4, 14 - 1, 15 - 3, 16 - 3		
7	<b>Groundtruth data</b>	16 - 5		



1 **Title: Our evolving understanding of aeolian bedforms, based on observation of dunes on different**  
2 **worlds**

3 © 2016. All rights reserved

4

5 **Abstract (253 words)**

6 Dunes, dune fields, and ripples are unique and useful records of the interaction between wind and  
7 granular materials – finding such features on a planetary surface immediately suggests certain  
8 information about climate and surface conditions (at least during the dunes’ formation and evolution).  
9 Additionally, studies of dune characteristics under non-Earth conditions allow for “tests” of aeolian  
10 process models based primarily on observations of terrestrial features and dynamics, and refinement of  
11 the models to include consideration of a wider range of environmental and planetary conditions. To-  
12 date, the planetary aeolian community has found and studied dune fields on Mars, Venus, and the  
13 Saturnian moon Titan. Additionally, we have observed candidate “aeolian bedforms” on Comet  
14 67P/Churyumov-Gerasimenko, the Jovian moon Io, and – most recently -- Pluto. In this paper, we  
15 hypothesize that the progression of investigations of aeolian bedforms and processes on a particular  
16 planetary body follows a consistent sequence – primarily set by the acquisition of data of particular  
17 types and resolutions, and by the maturation of knowledge about that planetary body. We define that  
18 sequence of generated knowledge and new questions (within seven investigation phases) and discuss  
19 examples from all of the studied bodies. The aim of such a sequence is to better define our past and  
20 current state of understanding about the aeolian bedforms of a particular body, to highlight the related  
21 assumptions that require re-analysis with data acquired during later investigations, and to use lessons  
22 learned from planetary and terrestrial aeolian studies to predict what types of investigations could be  
23 most fruitful in the future.

24

25 **Keywords (at least 6):** Planetary; terrestrial; aeolian bedforms; aeolian science; dunes; ripples

26

27 **Highlights** (max 85 characters, including spaces)

- 28 • Planetary dunes yield useful/unique information about climate & surface conditions.
- 29 • Aeolian bedform studies progress logically in questions/assumptions/new knowledge.
- 30 • Considering this progression exposes gaps/assumptions to be reviewed with new data.
- 31 • Comparing planetary progression with Earth aeolian studies yields lessons for each.

32

### 33 **1. Introduction**

34 Over the past couple of centuries, explorers and scientists of terrestrial dune fields have shown  
35 that the interaction between wind and granular material results in regular geometries and rates of  
36 evolution. Field observations and studies have inspired modeling and experimental works that have  
37 aided in the interpretation of measurable ripples and dunes as proxy indicators of wind speed and  
38 direction(s), grain sizes and sources, and underlying topography. The study of such landforms has been  
39 greatly extended and advanced by observation of analogous features on other planetary bodies. The  
40 comparison of these extraterrestrial features with aeolian process models has increased our  
41 understanding of aeolian bedform evolution in both directions – observations of (potential) aeolian  
42 bedforms generate investigations into the wind regime and granulometrics of surface materials on a  
43 planetary body, and also enable refinement of bedform evolution models as hypotheses about  
44 dominant effects are “tested” outside of Earth-conditions.

45 In this paper, we will review how our understanding (or assumed understanding) of sand dunes  
46 and/or ripples on a planetary body, and the information those aeolian bedforms yield about planetary  
47 conditions and processes, has progressed on different bodies. We hypothesize that the progression of  
48 investigations of these types of aeolian bedforms on a particular planetary body follows a consistent  
49 sequence – primarily set by the acquisition of data of particular types and resolutions, and by the  
50 maturation of knowledge about that planetary body. Our aim is to define that progression so we can

51 better constrain our level of knowledge about the aeolian bedforms of a particular body, highlight the  
52 gaps in our knowledge (i.e., our assumptions), and predict what type of future investigations could be  
53 most useful in addressing new questions and/or enabling improvement over an assumption.

54 In the interests of space and focus, most discussion (and cited literature) will focus on dunes and  
55 dune fields - i.e., the larger aeolian bedforms and thus usually the first seen on a planetary body. We do  
56 delve into ripples (and martian mega-ripples) in portions of the discussion, but primarily as features seen  
57 on dunes and that complement dune analysis; and we acknowledge that far more could be said about  
58 these smaller-scale bedforms and that studies of these bedforms on Mars has contributed much more  
59 towards our understanding of the aeolian environment and processes than is presented here. We also  
60 do not generally discuss other types of aeolian bedforms within this paper. In particular, we do not  
61 include discussion of Transverse Aeolian Ridges (TARs) on Mars as there is still much debate about their  
62 formation mechanism (perhaps as they are an example of a feature that does not have a good terrestrial  
63 analog?). It is likely, however, that one could trace advancements in our understanding of TARs or other  
64 aeolian features along a similar progression of ideas as is presented here for dunes, as all of these  
65 features are studied via similar observation types and their dynamics and morphologies tie into similar  
66 questions about atmospheric and surface conditions. We also consider only the observation and analysis  
67 of bedforms on the surface of a planet, not e.g., evidence of past bedforms recorded within sandstone  
68 stratigraphy (and thus, while dunes and ripples can also form due to the flow of other fluids, such as  
69 water, in this paper we focus on aeolian dunes). Thirdly, in discussing our evolution in thinking about  
70 aeolian bedforms and processes on other planets, we focus on observation-driven science  
71 advancements; we touch on but do not delve as deeply into the parallel lines of investigation focused on  
72 model development and validation, empirical studies, and (analog) terrestrial field work - investigations  
73 that feed into advancements within (and between) the Phases that we outline here. Finally, we  
74 recognize that we present only a sampling of relevant studies - we aimed for enough to map out

75 advances in understanding, to justify our proposed framework, and to provide a starting ground for any  
76 reader interested in learning more on a more specific topic.

77 In defining the “progression” of understanding (Section 2), we focus on Mars, Venus, and the  
78 Saturnian moon Titan – all planetary bodies where aeolian bedforms have primarily been explored with  
79 remotely acquired data. We also will comment on recently discovered candidate “aeolian bedforms” on  
80 Comet 67P/Churyumov-Gerasimenko and possible dune-like landforms on Io and Pluto. Within each  
81 phase of investigation (Subsections 2.1-7), we aim to identify the type of observations generally needed  
82 and connect these to the primary knowledge, assumptions, and questions that result, and then lead into  
83 future investigations (summarized at the end of each section). Furthermore, we identify the typical  
84 investigations (outside of direct studies of the aeolian bedforms) that follow each gain in knowledge, to  
85 show how aeolian bedform studies contribute to the larger study of that planetary body.

86 Our proposed framework of phases, regarding investigation of aeolian bedforms on a planetary  
87 body, is summarized in Table 1. We again note that this framework is not meant to be fully  
88 comprehensive for aeolian dune studies. We also note that progression in investigations and  
89 understanding is not necessarily linear/sequential – for example, planetary missions are generally  
90 focused on objectives other than aeolian bedform investigations, so observation types can be acquired  
91 in a “mixed” order. Additionally, the advent of new missions, methods, or models can lead to renewed  
92 activity within “lower” phases along with advances into higher phases. Science questions also often end  
93 up circling back as an assumption becomes superseded by new information and derived products and  
94 assumptions must be re-thought. Thus, in addition to identifying typical assumptions associated with  
95 each Phase, we use that framework to identify some example areas of knowledge gaps or the types of  
96 typical assumptions and results that need re-evaluation when new data becomes available (Section 3.1).

97 We also discuss how modeling (Section 3.2) and terrestrial studies (Section 3.3) relate to  
98 planetary aeolian studies. In particular, we highlight lessons learned regarding our understanding of

99 aeolian processes and their drivers, as well as in methodologies employed. These lessons translate (or  
100 could translate) to improved results within other areas of aeolian science.



**Table 1. Summary of the investigation phases**

Phase of aeolian bedform study on a planetary body	Area of interest	Characteristic(s)/Feature(s) of interest	Data needed to move to this phase (from an earlier phase)	Complementary science investigations
<b>1 Recognition of dune(s)</b>	Dune (possibly a dune field)	Dune morphology (i.e., recognizable, distinctive gross dune shape or crestline patterns within a field)	Images (visible, radar, spectral, etc.) with sufficient resolution to distinguish dune from non-dune surroundings, e.g., linear/arcuate and isolated/repeated morphology	Knowledge of and about analog features (terrestrial or planetary)
<b>2 Analysis of gross individual dune characteristics: e.g., morphology and composition</b>	Dune	Dune morphology, characteristics of surface materials	Images (visible, radar, spectral, etc.) with sufficient resolution to identify/correlate with dune margin and/or crestline patterns (possibly same data as Phase 1)	Global/regional-scale climate models (specifically: wind velocity, direction, and variation); Dune formation models
<b>3 Pattern analysis of the dunes within a field, including variations due to e.g., sediment supply and wind variations</b>	Dune field	Dune shapes throughout the field	Coverage (of images, see above) throughout dune field	Regional/local-scale climate models (specifically: wind); Regional/local-scale topography; Studies of non-aeolian dune-modifying processes (e.g., polar or surface crust forming processes); Maps of other aeolian features around the dune field
<b>4 Regional and global surveys and aggregate-analysis of dune characteristics; e.g., estimates of age or sand volumes, identification of large-scale sediment transport pathways, or identification/ estimation of the effect of location-related non-aeolian processes</b>	Regional or Global (i.e., multiple dune fields)	Dune field characteristics (including morphology of field and dunes within each field) and spatial distribution	At least regional coverage of images or (often coarser and/or less definitive) proxy data (e.g., thermal inertia)	Expansive composition maps for identification of potential sand sources; Maps of other aeolian features; Global or regional-scale climate models
<b>5 Analysis of superposed</b>	Dune	Within/on-dune	(Very) high-resolution images	Ripple formation models;

	<b>bedforms on the dune (such as ripples) formed due to wind interaction with the dune</b>		features (e.g., ripples)		High-resolution topography (dune); Very high resolution climate model (CFD) (specifically: wind)
<b>6</b>	<b>Observation of activity</b> on the dune, including non-aeolian activity	Dune	Smaller-scale evidence of change (e.g., ripple crestlines, dune margins)	Repeat images for seeing planform changes (e.g., movement of material); these images need sufficient resolution and temporal baseline for changes to be observable	Ripple and dune migration models
<b>7</b>	<b>Groundtruth data</b>	Dune	All features and components of the dune, especially sand size/composition	In situ observations of the dune, sampling and analysis	In situ observations of potential sediment sources for fluvio-sedimentary landforms

## 101 2. The Phases of Investigation

### 102 2.1. Phase 1: Recognition of dune(s)

103 In this first phase of aeolian studies, we focus on the occurrence of the *first observation of a*  
104 *dune* (or, more likely, a dune field). This has geologic significance as aeolian bedforms – dunes and  
105 ripples – directly record an interaction between the atmosphere and surface: specifically, the movement  
106 of granular material due to wind. Furthermore, a dune or ripple is more than a pile of sand – it is a  
107 distinctive landform that requires certain conditions to organize, and that appears to evolve and move  
108 “as a unit” through the aggregation of the actual movement of individual grains of sand, onto and off of  
109 the dune. Specifically, the existence of an aeolian bedform implies:

- 110 • A sufficient supply of saltatable (sand) grains,
- 111 • A wind of sufficient velocity and consistency to move those grains, and
- 112 • A period of consistent wind blowing, long enough for the bedform to form and evolve.

113 (We now examine what each of those underlined terms imply about the planetary body’s environment,  
114 focusing on the larger-scale dunes that are typically observed first. A more technical summary of the  
115 conceptual framework for dune field dynamics and how this is affected by the sediment state of a dune  
116 field -- related to sediment supply, sediment availability, and transport capacity of the wind – is  
117 described within Kocurek and Lancaster (1999).)


118 A sufficient supply means much more than the volume of the dune – for most dunes to form  
119 and evolve, sand must be able to move onto and off of the dune (possible exceptions would be climbing  
120 dunes or other such features where the sand accumulates due to blockage). Barchans in particular are  
121 an inefficient dune shape due to sand leakage from the horns (Hersen, 2004). Thus, an important  
122 implication with the first recognition of a dune feature on a planetary body is that a process must exist  
123 that will yield a significant amount of sand (discussed in within an example in Phase 4). Depending on  
124 the body, that process may involve erosion of larger rocks (e.g., as is common on the Earth) or a process

125 that directly forms grains of that size. For instance, martian volcanic activity has been proposed to  
126 create sand-sized particles (Edgett and Lancaster, 1993; Wilson and Head, 1994) and photochemical  
127 processes in the Titan atmosphere may eventually lead to saltatable grains, perhaps via an intermediate  
128 evaporite or sedimentary location (Soderblom et al. 2007; Radebaugh, 2013; Barnes et al. 2015).

129 On the Earth, nearly all dunes and ripples are comprised of sand grains – and this refers to a  
130 specific size. (e.g., the Canada Dept. Agriculture (1976) lists sand as grains 0.05-2 mm in diameter).  
131 However, “sand grains” can also be defined based on dynamics. “Sand” is the size of grains most easily  
132 moved by a fluid (Bagnold, 1941) – smaller grains are held together by interparticle, cohesive forces and  
133 larger grains have more mass and so are held down more by gravity. Under the Earth’s atmosphere and  
134 gravity, grains ~1 mm in diameter are able to saltate, and thus are the most easily moved by the wind.  
135 However, under the influence of a different atmospheric (or fluid) density and gravity, the grain size  
136 most easily moved by the wind could be a different size (Figure 1; Edgett and Christensen, 1991; Greeley  
137 et al., 1974; 1980; 1992a; Moore et al., 2015). Throughout this discussion, when discussing “sand  
138 grains,” we mean “the grain most easily moved by the wind (or fluid)” and not a fixed size range. Thus,  
139 the existence of a dune (i.e., a landform composed of sand grains) on a planetary surface yields a  
140 coupled constraint on the grains and the wind velocity.

141 Even if the wind reaches sufficient strength to transport sand, if it is not consistent (in direction  
142 and speed) over a sufficient period of time, the wind would just move small amounts of sand back-and-  
143 forth until that sand became trapped into depressions, sheltered areas, and other sand-traps; that sand  
144 would not be able to organize into a bedform. Models have shown that sand dunes have a minimum size  
145 (e.g., Claudin and Andreotti, 2006; Parteli et al., 2007); below this size the slipface is unable to develop.  
146 A slipface is necessary to stabilize the dune (as sand then can be captured on the sheltered, lee slope) to  
147 allow it to continue growing and migrating.

148



149

150 **Figure 1: Wind shear velocity needed to move grains of different sizes, on different planets.** Plot

151 showing the estimated threshold shear velocity for wind-driven transport of a grain of a specific diameter

152 for (from top) Pluto, Mars, Earth, and Titan; curves are taken from Moore et al. (2015; Fig. 17). The

153 general shape of the curve is reflective of smaller particles experiencing stronger interparticle forces

154 (such as electrostatic forces), while larger particles have more mass – either effect thus requiring more

155 shear velocity to initiate and sustain grain movement. The curve’s minimum indicates the expected size

156 of “sand grains” (i.e., the grains most easily lifted and moved by a shearing fluid – by saltation) on that

157 planetary body, that would be involved in the formation of aeolian bedforms. On Earth, sand grains are

158 commonly ~0.1mm in diameter. On Mars observations of saltatable grains (“sand grains”) in aeolian

159 deposits such as dunes (e.g., [Figure 11](#)) yield comparable diameters, which is consistent with the curves

160 shown. The differences in shear velocity needed to initiate motion are due primarily to differences in the

161 estimated air (fluid) and grain densities on each planetary body. The first investigations into these curves

162 and how they shift under different planetary conditions (Greeley et al., 1974) were initiated based on

163 observation of dunes on Mars (as described in the text: Belcher et al., 1971; McCauley et al., 1972; Cutts

164 and Smith, 1973).

165

166           To date, we have seen potential dunes on every deeply-studied body with an atmosphere and  
167 observable surface (including Titan, where dunes were considered unlikely: Lorenz et al., 1995), as well  
168 as a few bodies with no known atmosphere (Table 2). Based on the connections outlined above, this first  
169 “sighting” suggests the *accumulation of a lot of sand* (leading to questions about where the sand is  
170 coming from and why it is accumulating) and implications about *wind strength, direction, and*  
171 *consistency*. This yields a “groundtruth” observation for comparison with atmospheric models in both  
172 wind strength and direction (although it may be unclear when the bedform was created and thus what  
173 input conditions should be used for the model, or how the bedforms may have since been modified by  
174 non-aeolian processes).


175           Two classic examples of this are Mars and Titan. On Mars, signs of aeolian processes had been  
176 seen in cyclic, large-scale albedo changes and Mariner 6 imaged crescent-shaped features that were  
177 hypothesized to be very large barchan or parabolic dunes (Belcher et al., 1971). The first clear example  
178 of martian dunes was observed by Mariner 9 (McCauley et al., 1972; Cutts and Smith, 1973). Those  
179 observations suggested a wind regime that would allow for transport and collection of, as well as the  
180 presence of, a large amount of granular materials<sup>1</sup>, leading into laboratory studies of aeolian granular  
181 transport (Greeley et al., 1974; 1980). When Viking 2 imaged the north polar erg (Figure 2), this led to  
182 investigations of martian erosional processes (acting on polar layered deposits or soils of lower  
183 latitudes?) and climate models (Cutts et al., 1976). (A summary of results from Viking and Mariner-based  
184 aeolian studies can be found in Greeley et al., 1992a.) On Titan, “cat-scratch” features had been  
185 observed circumnavigating its equator, but were not immediately recognized as dunes until Vic Baker

---

<sup>1</sup> As Cutts and Smith [1973, p4151] put it: “The principal implication of dunes is a supply of noncohesive particles in the Martian surface environment and wind velocities sufficient for saltation transport. ... Dunes are not amenable to an alternative explanation of this sort. Thus we feel that we can now confidently assert the existence of a saltation regime on Mars”, which leads to “many implications of a saltation regime such as wind abrasion, wind scour, and dust production.”

186 brought the large draa of Saharan/Arabian/Namib deserts to the attention of the Cassini RADAR Team.  
187 The presence of dunes was a surprise as it had been hypothesized that while Titan's atmosphere may be  
188 capable of moving sand grains, it seemed unlikely that grains of the right size would exist (Lorenz et al.,  
189 1995). Observation of the dunes (Lorenz et al., 2006) led immediately to detailed investigations of what  
190 grains could be made of and how they would form (furthering studies of the chemistry on this Saturnian  
191 moon: Lorenz et al. 2006; Soderblom et al. 2007; Barnes et al. 2015) as well as leading to attempts to  
192 reconcile the observed dune morphologies with the model-predicted wind regime around the equator  
193 (Lorenz and Radebaugh 2009).

194




195

196 **Figure 2. An early image of the martian north polar erg.** (a) These linear features, imaged near the  
197 martian north polar cap by Viking 2 (frame 59B32: 62 km x 104 km), were hypothesized to be dune fields  
198 based on their consistent orientation and wavelength, and low sinuosity, branching and merging. Image  
199 and description are taken from Cutts et al. (1976; Fig. 7). (b) Higher-resolution images have proven that  
200 these are dune fields, with a wavelength (between primary crestlines) of approximately 0.4 km. A few

201 more orders of aeolian bedforms (e.g., the smaller crestlines, transverse to the primary crestlines) are  
202 also visible. Image is a portion of HiRISE PSP\_007115\_2600 (MRO/NASA/UA).

203



204

205 **Figure 3. A ~5 m boulder and potential aeolian features at Philae lander's touch-down-1 site. This**  
206 **ROLIS descent camera image shows a depression partly surrounding the boulder and a triangular-shaped**  
207 **apron on the opposite side have been interpreted as a moat and a windtail, indicating transport of**  
208 **granular material across the comet's surface (Mottola et al., 2015). Initial studies evaluated possible**  
209 **aeolian mechanisms for this transport. © ESA/Rosetta/Philae/ROLIS/DLR**

210

211 Contrarily, while Venus has a dense atmosphere, *only* two potential dune fields and a few  
212 possible microdune fields have been identified within Magellan radar data ( Greeley et al., 1992b; 1995;  
213 Weitz et al., 1994),, which covers 98% of the surface with 100 - 200 m resolution and shows wide  
214 coverage in other aeolian features such as windstreaks and potential yardangs (Greeley et al., 1995).  
215 This confirmed the hypothesis that aeolian bedform development on Venus must be limited, based on  
216 Venera 13 and 14 observations of the venusian surface that showed a dearth of aeolian ripples within  
217 loose material (Basilevsky et al., 1985; Florensky et al., 1983). (Note that for Venus, surface observations  
218 were first, before the mapping of surface topography from orbit.) The implication was that venusian




219 conditions and processes commonly obliterate features, the conditions for dune formation are not  
220 common on Venus (e.g., that there are few sands available on Venus or wind is not sustained at the  
221 surface), or that dunes are not generally visible via the radar images (Greeley and Arvidson, 1990; Weitz  
222 et al., 1994). Wind tunnel (Greeley et al., 1984a; 1984b; Marshall and Greeley, 1992; Williams and  
223 Greeley, 1994) and sand flux modeling (Kok et al., 2012) has shown that saltation under venusian  
224 conditions may occur in a very thin near-surface layer with very low velocity, which does not favor  
225 formation of large dunes. Thus, venusian sand transport appears more comparable to terrestrial  
226 bedform formation under water (Marshall and Greeley, 1992; Kok et al. 2012; Neakrase, 2015).  
227 Unfortunately, no new data about Venus has been acquired since Magellan, so Venus dune  
228 investigations remain stuck just past Phase 1 (with a small start within Phases 2-4, see discussion of  
229 Phase 2; Figure 13).

230           Potential aeolian bedforms have also been seen on planetary bodies lacking an atmosphere. For  
231 example, planetary scientists were recently very surprised to see features that looked like aeolian  
232 bedforms (i.e. moats, wind tails, and dune-like ridges) on comet 67P/Churyumov-Gerasimenko (Figure 3;  
233 Mottola et al. 2015; Thomas et al. 2015a; 2015b). A comet seemed clearly to be a planetary body that  
234 would lack an atmosphere, and thus any wind -- yet the features were observed. This immediately led to  
235 studies trying to determine how a "wind" could exist on this comet, if even transiently. One mechanism  
236 proposed to explain particle mobilization on comets was gas outflow from reservoirs of subsurface  
237 sublimating ice that emerges and erodes particles from channel walls (Cheng et al., 2013). However,  
238 since this process would only affect localized regions and the dune-like features on 67P have been  
239 observed in a much wider area, "splashing" initiated by airfall, i.e. ejection of particles by incoming  
240 projectiles, has been suggested as the most significant mechanism to explain particle mobility (Mottola  
241 et al, 2015; Thomas et al., 2015b). A three-dimensional cellular automaton model has proven that moats  
242 can result from abrasion of the surface by impinging particles, whereas wind tails develop where

243 granular surface materials were shielded by obstacles from particle transport (Mottola et al., 2015). The  
244 results of this study put forward the explanation that the aeolian bedform-like features on comet 67P  
245 are of erosional nature, rather than depositional – but the questions and investigations that arose in  
246 response to the recognition of features that resembled aeolian bedforms were consistent with typical  
247 Phase 1 discussions.

248           The surface of Io is covered in a ubiquitous frost of SO<sub>2</sub> as seen by the Galileo Near Infrared  
249 Mapping Spectrometer (NIMS), likely mixed with dust and fine-grained materials, all ejected from the  
250 continuously erupting volcanic plumes and explosive volcanic eruptions (Kieffer et al. 2000, Milazzo et  
251 al. 2001). The surface as seen by the Imaging Science Subsystem (ISS) instrument on Galileo is mostly  
252 uniformly light-colored from this frost and generally smooth, with some fractures, slumps and pits  
253 (McEwen et al. 2000). In a few regions, there are landforms with dune-like characteristics: regular  
254 spacing, a slightly meandering form, “crestline” defects, and apparent topography visible through the  
255 uneven collection of frosts (not possible to confirm with Galileo’s instruments). In one location, the  
256 dune-like landforms are found near a particularly active volcanic plume source, the Prometheus plume,  
257 which is sourced by advancing lava flows over vaporizing frosts (Figure 4; Kieffer et al. 2000; Milazzo et  
258 al. 2001). It is possible this plume forms a localized atmosphere dense enough to loft particles from the  
259 surface and deposit them nearby in dunes, much like one of the processes hypothesized for forming the  
260 features on comet 67P.

261




262

263 **Figure 4. Potential aeolian bedforms on Io.** This ridged terrain has been postulated as potential aeolian  
264 bedforms, formed from volcanic plume deposits. Lava is erupting from a fissure about 40 km east (right)  
265 of the edge of this mosaic, and the 100 km tall Prometheus plume is erupting from somewhere near the  
266 western (left) end of this mosaic. The bright streaks radiating from the area where the lava flows (the  
267 dark features) overrun the field are where the hot lava recently vaporized the sulfur dioxide, which then  
268 coated the lava-facing sides of the ridges. These images were taken by Galileo during a flyby of Io on  
269 February 22, 2000, with a resolution of 12 m/pixel. Image and description are taken from NASA  
270 Photojournal PIA02568.

271

272 The flyby of Pluto by New Horizons in July 2015 produced one of the most striking increases in  
273 image quality of a planetary surface in the history of planetary exploration (Moore et al., 2016; New  
274 Horizons, 2015; Stern et al., 2016). The landscape imaged during the flyby revealed a surprising diversity  
275 of landforms, which suggest varied geological and geomorphological processes active within recent  
276 geological history surface (Moore et al., 2016; Stern et al. 2015; Trilling et al., 2016). Mountains, glaciers,

277 plains, possible cryovolcanism and a surprisingly low density of craters covered much of the surface.  
278 Initial science results from the flyby noted the possible existence of 'windstreaks' (Stern et al. 2015) on  
279 Sputnik Planum, apparently extending in the lee of dark hills protruding through the nitrogen ice of  
280 which the plains are composed. Aeolian bedforms were speculated on before New Horizons arrived  
281 (Moore et al., 2015), and dunes have since been posited in the Baré Montes and enigmatic features in  
282 the Tartarus Dorsa have been interpreted variously as dunes or erosional aeolian features (Fenton,  
283 2016; New Horizons, 2016). However, recent imagery from Sputnik Planum (Figure 5) provides perhaps  
284 some of the most convincing examples of potential aeolian bedforms, with continuous linear features  
285 with a spacing of ~400-600 m. These features extend across the polygonal dark features, which have  
286 been interpreted as convectional cells within the ice, suggesting that they are the result of surface  
287 processes not related to the convective movement within the ice. Pluto is thus tentatively within Phase  
288 1 of the progression, and if an aeolian origin can be shown to be feasible, then available data may be  
289 sufficient for progression into Phases 2-3. The dune-like landforms on comet 67P, Io and Pluto are well in  
290 Phase 1 discussions and primary work remains to be done for these features to determine their ultimate  
291 origin.  
292



293

294 **Figure 5. Possible Pluto aeolian bedforms at the margins of the nitrogen ice of Sputnik Planum.** These  
295 features are oriented approximately parallel to the 'shore' where the ice abuts mountains. Image width  
296 is approximately 75 km. Image credit: NASA/JHUAPL/SwRI.

297

#### 298 **Summary of Phase 1**

299 **Data needed:** Images of the surface topography, of sufficient resolution to identify the distinctive  
300 shapes of dunes -- images could be visible or spectral imagery or radar scans of the planetary body's  
301 surface; identified analog (usually terrestrial) aeolian bedforms.

302 **Knowledge gain (from that data):** Existence of a potential aeolian bedform.

303 **Assumptions generated:** Conditions (wind conditions and grain size/supply) conducive to dune  
304 formation and evolution exist or have existed - note that this is a coupled constraint, and further  
305 information is needed to estimate each individual measurement.

306 **Questions:** What is the composition of the grains? How were sand-sized grains formed? Why do the  
 307 grains accumulation in that particular location (related to winds, topography, sand source)? Are the  
 308 dunes active? If conditions are unusual (e.g., tenuous atmosphere), how does sustained saltation and/or  
 309 reptation occur on this world?

310 **Lead to investigations of:** Models of surface processes and rates (chemical, erosional, etc.) that could  
 311 create “sand” grains; comparison to (global) atmospheric models; independent measurements of  
 312 surface conditions/composition for comparison to grain-formation models; independent studies of wind  
 313 speeds and/or grain sizes (e.g., thermal inertia) to decouple saltation conditions.

314 (Alternatively: ) Table N. Summary of Phase 1

Data needed within Phase 1	Images of the surface topography, of sufficient resolution to identify the distinctive shapes of dunes -- images could be visible or spectral imagery or radar scans of the planetary body's surface; Identified analog (usually terrestrial) aeolian bedforms.
Knowledge gain (from that data)	Existence of a potential aeolian bedform.
Assumptions generated	Conditions (wind conditions and grain size/supply) conducive to dune formation and evolution exist or have existed – note that this is a coupled constraint, and further information is needed to estimate each individual measurement.
Questions generated	What is the composition of the grains? How were sand-sized grains formed? Why do the grains accumulation in that particular location (related to winds, topography, sand source)? Are the dunes active? If conditions are unusual (e.g., tenuous atmosphere), how does sustained saltation and/or reptation occur on this world?
Leads to investigations of	Models of surface processes and rates (chemical, erosional, etc.) that could create “sand” grains; Comparison to (global) atmospheric models; Independent measurements of surface conditions/composition for comparison to grain-formation models; Independent studies of wind speeds and/or grain sizes (e.g., thermal inertia) to decouple saltation conditions.

315

## 316 2.2. Phase 2: Analysis of gross individual dune characteristics

317 Analysis of the dune morphology (Phase 2) typically closely follows Phase 1, often via the same  
 318 medium of an image of the surface. As models (e.g., Pelletier, 2009; Sauermaun et al., 2001; Werner,

319 1995) and laboratory studies (e.g., Andreotti et al., 2006; Parteli et al., 2009) have established, a dune's  
320 overall shape and orientation yields additional information about wind conditions when the dunes were  
321 forming and evolving<sup>2</sup>. Thus, this investigation stage generally involves detailed mapping of dune shape  
322 (outline and crestline) and comparison to analog terrestrial dunes. Numerical models of sand transport  
323 and dune evolution are also used for comparison between dune-generated predicted wind directions  
324 and atmospheric models (possibly down to a mesoscale or regional scale and including the effects of  
325 large-scale topography).


326         For example, once it was recognized that the long, generally linear cat-scratch features on Titan  
327 were dunes (Lorenz et al. 2006), researchers began investigating what type of dunes were observed and  
328 what wind conditions were required for their formation and persistence (in addition to questions about  
329 what the sand may be made of). From the morphology of the dunes, visible in 350 m/px Synthetic  
330 Aperture RADAR (SAR) data from the Cassini spacecraft (slightly hyper-resolution in nature in some  
331 locations because of the high contrast between SAR-absorbing sands and fractured, signal-scattering  
332 bedrock), it was determined they are longitudinal in type (also called linear; Lorenz et al. 2006;  
333 Radebaugh et al. 2008). Cassini Visual and Infrared Mapping Spectrometer (VIMS) data in select, high-  
334 resolution regions confirmed the general morphology of Titan's longitudinal dunes as well as their  
335 spectral contrast between sand and substrate (Barnes et al. 2008). On the Earth, longitudinal dunes are  
336 typically formed when several alternating winds of roughly equal transport strength (i.e., they move the  
337 same amount of sand) are >90° apart, yielding a single sand transport direction that is along the dune  
338 crestline (Fryberger and Dean 1979; Parteli et al., 2009; Rubin and Hunter, 1987; Rubin and Ikeda, 1990;  
339 Tsoar, 1983). (An alternate hypothesis that had been put forth connecting longitudinal dune morphology  
340 and wind directionality for Titan and some Earth dunes is discussed in Subsection 3.2.)

---

<sup>2</sup> However, establishing a connection between wind directions and dune slipface orientations sometimes is not a straightforward process – such studies often rely on many assumptions about timing of the winds and their consistency, and results are usually non-unique. Thus additional information is usually needed to evaluate a proposed interpretation. See Phase 3 for more discussion of the complexity that can be found in many dune fields.

341 Further information about these dune fields was then used to also predict the dominant wind  
342 directions, for comparison with climate models (discussed here, but merging into Phase 3  
343 investigations). Titan's dunes are found strictly between 30°N and S, ringing the equator, and are  
344 oriented roughly parallel to the equator (Lorenz et al. 2006; Radebaugh et al. 2008). Their morphology,  
345 and especially behavior around topographic obstacles (wherein grains are piled up at the upwind margin  
346 and are more sparse at the downwind margin), indicated a general sand transport direction from west  
347 to east (Figure 6; Courrech du Pont et al., 2014; Lorenz and Radebaugh 2009; Lucas et al., 2014;  
348 Radebaugh et al. 2010;). However, this was found to be at odds with the global atmospheric transport  
349 direction of east-to-west predicted in the equatorial zone by global climate models. Subsequent  
350 modeling studies revealed that seasonal or storm-driven winds could produce fast westerlies near the  
351 equator (Tokano 2010; Charnay et al., 2015), possibly resolving this inconsistency.  
352





353

354 **Figure 6: Earth and Titan dunes, diverging around topographic obstacles.** (a-b) Landsat 7 ETM+ and  
 355 SRTM C-band images of dunes in Namibia (centered on 25° 23' S 15° 16' E). The right image (b), a  
 356 Synthetic Aperture RADAR (SAR) image of the same dunes, shows bedrock as bright, because it is rough,  
 357 and unorganized dune sands as dark, because dunes are smooth at the SAR wavelength of 6 cm (black  
 358 areas are regions devoid of data returned to the SAR antenna). The winds blow SE to NW (bottom L to  
 359 upper R), as evidenced by the dune-free regions in the lee of the bedrock topography and the diversion of  
 360 dunes around the upwind sides of the topography. In the lower image (c), dunes similarly divert around  
 361 topographic obstacles and resume on the downwind side, within this region of the Belet sand sea, Titan.

362 *This is a Cassini Radar image (300 m resolution) centered on 6.5° S, 251° W, with winds interpreted to*  
363 *blow SW-NE (L-R). Images and description are from Radebaugh et al. (2009).*


364

365           On Mars, mapping of the duneforms observed by Mariner 9 and Viking orbiters, combined with  
366 information about the surrounding surface topography and composition (including other potential  
367 aeolian features), was used to determine the wind direction(s) that would yield the dune crestline  
368 orientation(s), the wind and topography that would yield sand accumulation within that area and its  
369 likely stability (i.e., was this a stable sink for sand, or a temporary repository), and possible sand sources  
370 (Cutts and Smith, 1973; Greeley et al., 1992a; Thomas, 1981; 1982; Tsoar, 1979). For example, the large  
371 number of transverse dunes observed within southern mid-latitudes and north polar region on Mars (vs.  
372 longitudinal dunes), in Viking and Mariner images, implied that dune fields were forming within regions  
373 of near unidirectional winds, and the asymmetry in dune field complexity between the north and south  
374 hemisphere was seen as evidence of a more complex southern wind regime (Greeley et al., 1992a;  
375 Thomas, 1981). After higher resolution images became available and atmospheric modeling became  
376 more refined in technique and topography/boundary inputs, a more detailed comparison was done to  
377 see if atmospheric models could reproduce the observed dune shapes and orientations. For example,  
378 mesoscale modeling of dunes within Proctor Crater on Mars based on MOC NA images matched two  
379 sets of dune slipface orientations (the primary and tertiary) to seasonal winds that were impacted by  
380 daily and seasonal insolation patterns and the crater topography, as predicted by a mesoscale model  
381 (Fenton et al., 2005; however, the secondary dune slipface orientation remained unexplained). This  
382 study validated the mesoscale atmospheric model by providing a reasonable explanation for the range  
383 of slipface orientations seen within that dune field, and thus advanced the use of these models within  
384 model-observation comparison studies for understanding aeolian processes on Mars.

385 Coupled with analysis of wind direction, observations of the dune's local surroundings can also  
386 be studied to identify sediment sources and sand transport pathways. In some areas, distinct sand  
387 transport pathways leading from the sediment layer to the dune bodies have been revealed by the  
388 detection of congruent material composition. For example, on Mars, local sediment sources for intra-  
389 crater dunes have been proposed at impact crater walls by comparative analyses of high resolution  
390 image and spectral data of dune bodies and the sediment layers exposed (e.g., Fenton, 2005; Geissler et  
391 al., 2013; Silvestro et al., 2010a; Tirsch et al., 2011). Within the Valles Marineris rift system, spectral  
392 analysis, morphological evidence of erosion and sand transport, and topographic information was used  
393 to show that diverse and distinct sediment sources serve as local and regional sources (Chojnacki et al.,  
394 2014).

395 On Venus, analysis of dune morphology is almost impossible due to lack of data. Dunes in both  
396 recognized dune fields are at the resolution limit of radar images obtained by Magellan mission (Figure  
397 7), the only adequate data source. Therefore, results of attempted of detailed analysis are not reliable  
398 and are controversial (Greeley et al., 1997; Lorenz, 2015). Modeling of anisotropic radar scattering  
399 (Kreslavsky and Vdovichenko, 1999) indicated that the microdune fields proposed by Weitz et al. (1994)  
400 possess abundant unresolved steep slipfaces.

401



402

403 **Figure 7: Venusian dune field.** This Magellan SAR image of the Aglaonice dune field (25°S, 340°E) shows  
404 a field of white dots that are interpreted as specular reflections from the slopes of transverse dunes  
405 (between the arrows). This type of reflection occurs if the slope is mostly smooth, and is oriented near-  
406 normal to the incidence angle of the radar (which is 35° or near the angle of repose); similar features are  
407 observed within Seasat and space shuttle radar images of terrestrial sand dunes. The implied wind  
408 direction of these features, based on their orientation, is also consistent with nearby bright and dark  
409 wind streaks extending from behind cones (not included in this cropped image). Radar illumination is  
410 from the left, and north is at the top. Image is from NASA Photojournal PIA00483, and description is from  
411 Photojournal and Weitz et al. (1994).

412

413 In general, Phase 2-type investigations can continue for a long period with refinement of the  
414 investigations as higher-resolution images of the dunes are acquired, more information becomes  
415 available about the local topography and other evidence of aeolian processes and conditions, and/or  
416 atmospheric or bedform formation models are improved. This investigation phase may eventually grade  
417 into (and occur concurrently with) Phases 3 and 5 which involve, respectively, higher-resolution analysis

418 of features within the dune field and features on the dunes formed by the dune-wind interaction (such  
419 as ripple patterns on the dune slopes). Additionally, as new observations, models, and analysis methods  
420 are developed, Phase 2 investigations can be renewed and revisited (as with all Phases beyond Phase 1).

421

## 422 **Summary of Phase 2**

423 **Data needed:** *same as Phase 1.*

424 **Knowledge gain:** Morphology of the potential crestlines and general dune shape; composition of the  
425 dunes and surroundings; possibly identification of local sand sources.

426 **Assumptions generated:** Wind direction and consistency *hypothesized* to generate the observed shapes;  
427 variations in wind speed implied by changes in sinuosity and shape through the dune field.

428 **Questions:** What sets the wind direction and causes its variations (e.g., daily or seasonal cycles)? Are  
429 these representative of present-day wind conditions, conditions during a past period, or a convolution  
430 of conditions during different past periods?

431 **Lead to investigations of:** Comparison over one to several dune fields with (global/mesoscale)  
432 atmosphere models; reliably identify wind direction(s) consistent with the dunes' forms.

**Table 2: Recognition and first analysis of dunes and dune fields on planetary bodies (Phases 1-2), as presented in the literature**

Planet. body	“Aeolian” bedforms first sighted	Data used	Immediate Implications	Immediate Questions	Reference
<b>Mars</b>	<b>Mariner 9</b> , Hellespontus region of Mars, dense and large transverse dune field	Visual image of surface, <1 km/pixel	Dune material is dark (so some low albedo areas are regions of deposition, not erosion; and some dark material will saltate); Due to the lower atmospheric density on Mars, wind velocities may need to be much higher to move sand.	Which “dark splotch[s] or streak[s]” are due to deposition of material (vs. deflation of overlying bright material)? What is the source of the dark material? Does the wind reach transonic velocities? Is high-velocity sand-blasting resulting in highly-efficient wind erosion?	Sagan et al., 1972
		above; comparisons to other albedo markings indicative of wind direction	Presence of lots of sand and saltation processes; Dune material accretion directions and influence of topography (craters) on field location and dune morphologies; comparable scale and shapes as terrestrial dunes	What is the composition of the sand? Why is it so dark?	Cutts and Smith, 1973
	<b>Viking 2</b> , north polar erg, transverse and barchan dune fields	Visual image of surface, 30-60 m/pixel	Lots of sand → some erosional process; Variability in wind regime; two wind directions in portions	What is the composition and source of sand? Why is it accumulated around north polar cap?	Cutts et al., 1976
		Strong winds; Two wind directions, thought to be seasonal; grains may be eroded from the northern plains	Are the dunes active, and how mature is the dune field? Are the dunes modified during the winter/early spring, when the entire region is covered by CO <sub>2</sub> ice?	Tsoar et al., 1979	
<b>Venus</b>	<b>Magellan</b> , transverse dunes, two fields	Radar images, 75 m/pixel; compared with orientation of other aeolian features in same dataset	Lots of sand in specific areas → some erosional process (perhaps impacts?)	What is the composition and source of the sand? What are the saltation dynamics under a much denser atmosphere?	Greeley et al., 1992b
<b>Titan</b>	<b>Cassini</b> , longitudinal dunes, large	Synthetic Aperture Radar images, 175 m/pixel	Lots of sand → some “grain” formation process; one dominant or at least two converging wind	What is the composition and source of the sand? What is the underlying topography, causing accumulation in equatorial region	Lorenz et al., 2006

	field around equator		directions throughout equatorial region; pristine appearance and superposition over geologic features → young and possibly currently active	and diversion of dunes? Is there any connection with the potential fluvial channels? Has the sand circumnavigated the globe several times (implying a lack of sand-sinks in the area)?	
		Visual and Infrared Mapping Spectrometer (VIMS) observations, 500 m/pixel	Observations of interdunes → recent activity and overall dune field maturity; spectral information → constraints on the composition of the dunes and interdune regions; photoclinometry yielded height and wavelength estimates	What information does the variability in dune coverage and height, and in the terrain that they cover, yield for the evolution history and conditions for these mature and recently active dunes? What are the dune grains made of, given their lower relative waterice content than Titan's average?	Barnes et al., 2008
<b>Comet 67P/Churyumov-Gerasimenko</b>	<b>Rosetta</b> , moats, wind tails, and aeolian-like ridges/ripples	optical imagery: OSIRIS orbiter camera images ( $\geq 0.29$ m/px), ROLIS decent camera images ( $\geq 1$ cm/px)	"Sustained" granular transport along the surface (so as to form aeolian bedform-like features) exists on a comet	How does granular transport work on a body without atmosphere? What is the moving agents? Are these bedforms accumulative or erosive? What is the grain size of the bedform materials? What are the material sources? How long does it take to form bedforms?	Mottola et al., 2015; Thomas et al., 2015a; 2015b

### 433 2.3. Phase 3: Pattern analysis of the dunes within a field

434 Dune field evolution is related to the evolution of its constituent dunes, but occurs on a larger  
435 spatial and temporal scale and involves areas of investigation that are different from (and can be larger-  
436 than) the sum of its parts. For example, as dunes evolve within a field they exchange sand between each  
437 other through both sand flux and collisions, and environmental boundary conditions such as the sand  
438 influx geometry can affect dune field pattern development (Diniega et al., 2010a; Ewing and Kocurek,  
439 2010). As such, it is necessary to model dune field evolution as more than just a collection of individually  
440 evolving dunes, and to recognize that the large-scale dune field pattern can reflect conditions (and  
441 changes in those conditions) around and throughout the field. For example, sand and dune influx  
442 conditions will be different near the upwind margin of the dune field than near the terminus or lateral  
443 margins (Ewing and Kocurek, 2010) due to proximity to sand sources or other dunes, or changes in  
444 topography or “cementing” influences (e.g., chemical duricrust or, on the Earth, vegetation) within the  
445 field (Kocurek and Ewing, 2005) (Figure 6: examples of dune interactions with topography on Titan and  
446 Earth). Such changes can result in different dune sizes, spacing, or defect frequency (Diniega et al.,  
447 2010a; Ewing and Kocurek, 2010).


448 The effect of underlying topography is also a key parameter affecting dune characteristics at the  
449 dune field scale (Ewing and Kocurek, 2010). On Earth, bedrock topography has been linked to the effect  
450 of roughness variations induced by the dune field itself producing an internal boundary layer decreasing  
451 the shear stress downwind (Jerolmack et al., 2012) and/or to the feedback mechanism between long-  
452 wavelength topography and the dunes (Pelletier, 2015). The role of topography in enhancing and  
453 deflecting regional winds has also been invoked to explain complex dune field pattern on Mars in  
454 Olympia Undae (Ewing et al., 2010) and complex dune arrangements in Moreux (Cardinale et al., 2012)  
455 and Matara crater (Diniega et al., 2010b; Silvestro et al., 2012). However, it was only thanks to the  
456 availability of high resolution DTMs from the HiRISE instrument that the effect of underlying topography



457 could be more precisely linked to different dune characteristics such as migration rates, dune heights  
458 and density (Cardinale et al., 2016, Vaz et al., submitted). In particular, in Herschel crater dune density,  
459 slip face advancements and migration rates are all controlled by two major topographic highs on the  
460 crater floor (Vaz et al., submitted).

461         The dune field may also record changes in conditions over a longer-timescale than that recorded  
462 within any individual dune. Multiple patterns (e.g., different types of dunes) can be superimposed  
463 (creating a complex, versus a simple, dune field) as smaller dunes migrate and change in response to the  
464 new environment faster than larger dunes (Ewing and Kocurek, 2010; Hugenholtz and Barchyn, 2010;  
465 Kocurek and Ewing, 2005). We note that this possible complexity within dune fields can complicate  
466 analysis of the dune morphology (Phase 2). For example, even identification of the dominant (or most  
467 recent?) slipface orientations can be non-trivial. This is especially true within planetary dune fields  
468 where datasets may be limited to remote images, so dune slope angles and potential activity have to be  
469 interpreted from images of the dunes' planform appearance, possibly under suboptimal illumination  
470 conditions for this type of image analysis. For example, within the north polar erg on Mars, many dunes  
471 contain slipfaces pointing in opposite directions (sometimes on the same dune). One interpretation is  
472 that some of these fields may contain both active and fossil dunes (Gardin et al., 2012). Within the Mars  
473 southern mid-latitudes, at least two periods of dune-building (or dune-building occurring over 2  
474 different timescales) are apparent as within the same field one can often find a dense collection of  
475 transverse dunes (with slipface towards the east) and then barchans clearly climbing up and over the  
476 transverse dunes on the east side (with slipfaces towards the west) (Figure 8; Diniega et al., 2010b).

477



478

479 **Figure 8: The complex dune patterns found in along the eastern edge of Matara dune field, Mars**  
480 (49.5°S, 34.8°E). This dune field, like many others in the Mars southern mid-latitudes, is a dense  
481 transverse dune field, captured within a crater. The transverse dune crestlines are oriented north-south  
482 with the clearest slipfaces towards the east (white arrows extend up the possible stoss slope, towards the  
483 dune brink). However, along the eastern side of this field, many smaller dunes (mostly barchans – yellow  
484 arrows, some possible transverse crestlines near the bottom of the image) are oriented with towards the  
485 northwest. This potentially reflects two periods (or just two timescales?) of dune evolution, with a

486 *change in the dominant wind direction. North is up and illumination is from the left. Image is a portion of*  
487 *HiRISE PSP\_006648\_1300 (MRO/NASA/UA).*

488

489 A lack of variations can also yield information about the field's and planetary body's history. On  
490 Titan, dune width and spacing measurements over more than 7000 linear dunes showed a high level of  
491 uniformity around the moon, with no signs of compound or complex dunes (Savage et al., 2014). This,  
492 coupled with the dunes' large sizes, indicates that Titan's dunes are mature features that have evolved  
493 within consistent and stable environmental conditions for a long period of time.

494

#### 495 **Summary of (3)**

496 **Data needed:** Observations of dunes fields, of sufficient spatial coverage and resolution to note changes  
497 in dune patterns throughout the field, especially in tracing crestlines; possibly need knowledge of  
498 topography.

499 **Knowledge gain:** The dune field pattern and shape; maturity state (and possibly relative age) of the  
500 bedforms; possible temporal changes in e.g., sediment supply and wind patterns

501 **Assumptions generated:** Changes in the environmental conditions, in space or time.

502 **Questions:** For a given dune field, is sand sourced from one or several locations? Is the dune evolving  
503 through one dominant wind pattern, or several? Have the dominant effects (sand source or wind  
504 pattern) changed over the lifetime of the dune field?

505 **Lead to investigations of:** Explore influences on dune shapes beyond current dominant winds (*Phase 2*)  
506 – such as the location of sand source(s) or of sand-starved regions of the field, a transition between  
507 wind regimes, interactions between dunes (such as dune collisions), or other environmental  
508 influences/processes.

509

#### 510 **2.4. Phase 4: Regional and global surveys and aggregate-analysis of dune characteristics**

511           As we gather information about dunes in more and more different dune fields around a  
512 planetary body, it becomes possible to aggregate data to deal with high-level, large timescale questions  
513 about aeolian processes and sediment supply, such as “How much sand is available in total?” and “Are  
514 there primary types/locations of sediment sources that can yield information about how that sand has  
515 been created, how it is transported, and whether it has been recycled?” Addressing such big-picture  
516 questions can provide important information for investigations of grain-producing processes (e.g.,  
517 surface erosion) and planetary surface history over the lifetime of the involved sand grains. Note that  
518 while studies of grain history and sediment transport pathways involving terrestrial dunes may rely on  
519 detailed petrographic and heavy-mineral techniques, with geochronology (e.g., Garzanti et al., 2013),  
520 studies of planetary bodies often are based only on surface topography and, possibly, coarse  
521 compositional information.

522           For example, on Mars, a near-global map of sediment deposits (including dunes) and wind  
523 streaks to generate early estimates of sediment transport pathways/source regions (Thomas, 1982). An  
524 early global map of aeolian features showed variations in time and space in the large-scale wind  
525 directions recorded by the orientation of dunes, wind streaks, yardangs, wind grooves, and deflation pits  
526 (Ward et al., 1985). Such studies have since been updated with increased coverage and image resolution  
527 (e.g., Hayward et al., 2007; 2014), and still provide important information about direction and variability  
528 in the wind patterns (down to intra-field scales), the influence of topography and local geology on wind-  
529 flow and bedform development, and likely sediment sources for the observed deposits. At a regional  
530 scale, the martian north polar erg volume has been estimated as ~1130-3250 km<sup>3</sup> of dark sand (Greeley  
531 et al., 1992a; Hayward, 2011), which is significant as the icy layers of the north polar cap has been  
532 proposed as the source of the circumpolar dune fields (Byrne and Murray, 2002; Tanaka et al, 2008).  
533 These deposits appear composed of recycled aeolian sediments, which were likely transported poleward

534 and deposited (Breed et al., 1979; Byrne and Murray, 2002). This suggests that a huge volume of sand  
535 may have formed on Mars during an earlier epoch and that these sand grains have survived at least a  
536 couple of sustained dune-forming periods.

537         The Titan dune fields provide an example of how analysis of the distribution of dune field  
538 locations (on the planetary body, or relative position within regional topography) and morphologies (i.e.,  
539 field outline or crestline patterns) can yield additional information about larger-scale atmospheric and  
540 topographic/surface conditions. Mapping of 16,000 Titan dune segments (covering 8% of Titan's surface  
541 which suggested that dunes cover a total of 20% of the global surface: Lorenz and Radebaugh, 2009)  
542 showed general dune field orientation and spacing patterns and confirmed that these features are  
543 within a global field with few longitudinal trends, but with latitudinal trends in orientation and limited to  
544 within 30° of Titan's equator. Although dunes on Titan are organized into several separate sand seas  
545 across the equator, all have some level of broad interconnectedness (Le Gall et al. 2012; Radebaugh  
546 2013; Savage et al. 2014). As such, studies of the Titan sand sources, sediment transport pathways, and  
547 deposition patterns are best analyzed from a "global" perspective.

548         Titan sands may be derived directly from the atmosphere, perhaps through clumping on the  
549 surface, though it is perhaps more likely the sand has been processed through erosion of organic  
550 sedimentary layers (Radebaugh 2013), possibly close to the equator where fluvial channels have been  
551 imaged (Burr et al., 2013; Lorenz et al., 2008; Radebaugh et al., 2016). Other possible sources include  
552 erosion of the SAR-uniform mid-latitudes, a possible sedimentary deposit (Malaska et al., 2016), and the  
553 northern dry lakebed evaporite deposits, which have similar spectral characteristics to the VIMS  
554 instrument (Barnes et al., 2015). Once the materials are transported into the Titan sand seas, they are  
555 incorporated into the giant linear dunes, and either stay confined to one sand sea or contribute to a  
556 global system of west-to-east sediment transport that persists over time (Savage et al., 2014).  
557 Topography appears to play an important role, as it does for sand seas on Earth, in that it can help

558 confine the sands to certain regions or preclude them from others, like from the rugged Xanadu region  
559 (Lorenz et al., 2006; Radebaugh et al., 2011). Decreases in dune density within radar-bright and elevated  
560 regions may provide regional-scale constraints on Titan's winds for atmospheric models (Lucas et al.,  
561 2014). Furthermore, topographic obstacles can cause diversion of dunes and dune/topography  
562 relationships and perhaps reveal longer-term climatic changes (Ewing et al. 2015).

563         Consideration of the dune fields in aggregate can also allow for analyses that require a larger  
564 area or more numerous measurements to reflect larger-scale temporal or spatial trends. For example,  
565 dune fields on Mars appear very young as they lack craters, but constraints on their age had large  
566 uncertainties due to their low individual areal coverage. Adding the dune fields together allowed for a  
567 more robust estimated crater-retention age of <10 000 years (Fenton and Hayward, 2010). These dune  
568 fields also exhibit latitude-dependent morphological trends in crestline sharpness/pattern, dune slopes,  
569 and field shapes, so considering the dunes over the hemisphere enables studies of influence from polar  
570 as well as aeolian processes (Fenton and Hayward, 2010). Another study of southern intracrater dune  
571 fields on Mars compared dune field centroid locations, relative to the crater center, with mesoscale  
572 atmospheric modeling to look at broad-scale atmospheric trends (over a much longer time period than  
573 that recorded in dune slipface orientations within an individual dune field) (Hayward et al., 2009).

574         Although only two dune fields and a few microdune fields were identified with some certainty in  
575 the whole set of Magellan radar images of Venus, a few lines of indirect evidence suggest that  
576 unresolved small-scale anisotropic topographic features are ubiquitous; such features have been  
577 interpreted as unresolved gently sloping aeolian bedforms (Kreslavsky and Vdovichenko, 1999;  
578 Bondarenko et al., 2006). A comprehensive global inventory of aeolian bedforms on Venus will require  
579 global imaging data set(s) of a higher resolution than presently exists.

580         Beyond global imaging of the data-type first used to identify the dunes, proxy measurements  
581 can sometimes be used to supplement limited image coverage. For example, thermal inertia can be used

582 to identify large deposits of unconsolidated, granular material. On Mars, further evidence that these  
583 dark patches with high-thermal inertia were aeolian deposits were that these were found downwind of  
584 topographic depressions (Christensen, 1983; Mellon et al., 2000). Thus global maps of thermal inertia  
585 with resolution ~100 m/pixel have been used to map dune fields around Mars and estimate the number  
586 of dune fields and their surface areal extent (Christensen et al., 2003; Hayward et al., 2007; Hayward,  
587 2011).

588

#### 589 **Summary of (4)**

590 **Data needed:** Identification of dunes around globe (from the data used in Phases 1-3, and possibly from  
591 proxy data such as thermal inertia).

592 **Knowledge gain:** Dune field location and (possibly) morphology/type distributions; variations in location  
593 and morphology related to sediment supply, climate history, and/or and other active processes (e.g.,  
594 related to latitude, regional topography); identification of large-scale sediment transport pathways  
595 (larger-scale than field-specific results of *Phase 2 and 3*; and possibly first produced earlier based on  
596 low-resolution, but high-coverage datasets) based on (global/mesoscale) atmospheric models and  
597 observation of sediment sources.

598 **Assumptions generated:** Correlations between dunes and proxy data; feasibility of extrapolation from  
599 studies of individual dune fields/sand sources to a global model.

600 **Questions:** How much sand is there, where is it from/stored, and how did it get there? Over what spatial  
601 and temporal scale is the sand being transported (i.e., what is the lifetime of a sand grain and is  
602 sand/bedforms being recycled)?

603 **Lead to investigations of:** Age estimation of dune fields (as can aggregate together land-areas to  
604 statistical significance; likely to be a relative or crater-retention age); identification/investigation of

605 large-scale sediment sources (locations and/or processes; perhaps updated from *Phase 2*); global  
606 surface areal coverage of dunes/volume of sand.

607

## 608 **2.5. Phase 5: Analysis of superposed bedforms on the dune formed due to wind interaction with the** 609 **dune**

610         Ripples, like dunes, form spontaneously within sand beds due to wind (or fluid) flow and record  
611 wind and sediment conditions through their period of formation and evolution. However, as these are  
612 much smaller features, they record conditions over smaller temporal and spatial scales and thus can be  
613 reflective of a different set of environmental conditions than dunes. To-date, ripple-like features have  
614 been only observed on Mars, where HiRISE images of the martian surface have resolution as fine as 0.25  
615 m/pixel (McEwen et al., 2007). These features have wavelengths of 1-to-a-few meters, have been found  
616 within sandy regions including on the slopes of dunes, and have been individually mapped and  
617 monitored for movement (Phase 6) (Figure 9; Bridges et al., 2012; Silvestro et al., 2011). The study of  
618 ripple morphologies and dynamics on Mars yields information about the wind flow over the dunes,  
619 under the influence of the local wind patterns as well as the dune topography. This yields information  
620 about the recent, local wind regime within several areas on Mars (Bridges et al., 2012b). Such  
621 information about the temporally and spatially small-scale surface wind dynamics can be compared with  
622 meso and microscale climatic models and in-situ wind measurements (e.g., Jackson et al., 2015; Silvestro  
623 et al., 2013). In addition, because ripple morphology and migration rates are controlled by the  
624 topographic and wind flow boundary conditions imposed by the dune morphology (Kocurek and Ewing,  
625 2012), studies of the ripples' form and variation provide insights to the underlying dune's evolution  
626 (Ewing et al., 2010; Vaz et al., submitted).

627         Ripple mapping and monitoring have been an important tool within recent martian studies,  
628 where the crestline orientations and migration rates and directions of the large martian ripples are



629 commonly used to reconstruct the wind regime over the dunes and to estimate sand fluxes (Ayoub et  
630 al., 2014; Bridges et al., 2012a; 2012b; Cardinale et al., 2016; Silvestro et al., 2010b, 2011, 2013).  
631 Automatic approaches have been developed to derive ripple trend and migration rates, enabling high-  
632 resolution wind regime estimations and sand flux measurements to be computed over large areas  
633 (Ayoub et al., 2014; Bridges et al., 2012a; 2012b; Silvestro et al., 2011; Vaz and Silvestro, 2014).

634 However, all of these studies have assumed that that the observed “smaller” bedforms on the  
635 dunes are analogous to terrestrial sand ripples, and that ripple trends and migrations are normal to the  
636 last wind of sufficient strength to move sand, as is typically the case for aeolian ripples on Earth. Recent  
637 work has drawn those assumptions into question:

- 638 • Most ripple patterns on Mars are dominated by sinuous crestlines (Vaz et al., submitted), while on  
639 Earth ripple crestlines are typically straight (Rubin, 2012) (Figure 9). In some areas, ripple patterns  
640 observed on Mars show complex arrangements with two crestlines intersecting at right angles  
641 (Figure 10; Silvestro et al., 2011, 2013). This suggests that some of the ripples on Mars might not be  
642 in equilibrium with the last sand-moving winds or that the two sets of crestlines are  
643 contemporaneous, but oblique to the formative winds (Silvestro et al., 2016).
- 644 • Additionally, unusual longitudinal displacement of crest-line defect terminations and oblique crest  
645 migrations have been observed within orbital data in Gale and Herschel crater, suggesting that the  
646 large ripples of Mars are different from terrestrial impact ripples (Silvestro et al., in press; Vaz et al.,  
647 submitted). This hypothesis is in agreement with recent in situ observations from the NASA MSL  
648 Curiosity rover, which shows that large ripples have sinuous and sharp crests and slip faces with  
649 evident grainfall and grainflow structures (Bridges et al., 2016; Lapotre et al., 2016a) (Figure 10) that  
650 are not common within terrestrial impact ripples. Superposing these large bedforms are smaller  
651 “terrestrial-like” impact ripples of ~10 cm in wavelength (Bridges et al., 2016; Lapotre et al., 2016a).

652           These observations suggest that terrestrial aeolian impact ripples might not be good analogs for  
653 the Martian large ripples (Lapotre et al., 2016a; Silvestro et al., 2016; Vaz et al., submitted). As this gets  
654 worked out, previous studies will need to be carefully reviewed, such as where the interpretation has  
655 been that a multidirectional wind regime exists, perhaps triggered by the local dune topography or by  
656 larger topographic features (e.g., Jackson et al., 2015; Silvestro et al., 2011). Also, the presence of such  
657 large ripples on the dune's stoss side and their migration across the slipface (Figures 9-10) may alter the  
658 wind profile above the dune and the slipface dynamics, beyond the way that these processes are  
659 typically captured in dune evolution models applied to terrestrial dunes and their ripples (e.g., Ewing et  
660 al., 2016). Increased coverage of high-resolution images coupled with in-situ observations by rovers are  
661 necessary to progress understanding of the nature and dynamic of the martian large ripples. This is  
662 fundamental for understanding how these ripples can be used to constrain local wind directions and to  
663 tune sand flux estimations over the dunes.

664

#### 665 **Summary of (5)**


666 **Data needed:** Higher-resolution images of dune field, reflecting variation over the dune, including in  
667 composition or granulometrics; mapping and analysis of second-order and higher-order bedforms (e.g.,  
668 ripples) and how these reflect the wind pattern around the dune.

669 **Knowledge gain:** Measurements of ripple movement and characteristics over the dune.

670 **Assumptions generated:** Use of the right analog features/models for interpretation of the smaller-scale  
671 features.


672 **Questions:** What is the local sand flux and wind patterns over the dunes (as reflected in ripple  
673 movement)? Are grains sorted within the ripples, and if so, why? Is ripple movement coupled  
674 with/connected with current dune evolution, or e.g. does ripple movement reflect a surficial mobile  
675 layer of sand over a relict dune core?

676 **Lead to investigations of:** Wind diversion around dune topography; observation/better understanding  
677 of local source regions.  
678



679  
680 **Figure 9. Observed ripple movement on Mars.** Images show (a) a rippled dome dune in the Bagnold  
681 dune field, Gale Crater, with (b) ripple migration over the dune stoss side between Mars years 28 (2006)  
682 and 29 (2008) (Silvestro et al., 2013). (c: T1-T2) The zoom-in shows one ripple (white arrow) moving over  
683 the dune brink, reflecting grain transport onto the slipface and suggesting that dune migration may also  
684 be occurring. HiRISE images shown: (a,c/T1) PSP\_001488\_1750 (taken 20 November 2006), (c/T2)  
685 PSP\_009650\_1755 (17 August 2008) (MRO/NASA/UA).

686



687

688 **Figure 10. Ripple morphology on Mars.** (a-b) HiRISE images of High Dune, an active dune within the  
 689 Bagnold dune field, Gale Crater, that has also been investigated by MSL Curiosity (Bridges et al., 2016;  
 690 Lapotre et al., 2016a). Visible in these orbital images is a complex ripple pattern on High dune's stoss  
 691 slope, with what appears to be two sets of large ripples intersecting at right angles (Silvestro et al.,  
 692 2016). This ripple configuration is typical of the Bagnold dunes and seems to be common on Mars. HiRISE  
 693 images are: (a) ESP\_042682\_1755, (b) PSP\_009294\_1750. (MRO/NASA/UA) (c) This image of the High  
 694 Dune stoss slope was taken by the Mast Camera on Curiosity, showing the complex morphology of these  
 695 large ripples, which in this closer-inspection perhaps do not appear analogous to terrestrial sand ripples.  
 696 The scalebar (one meter length) is for the lower portion of this cropped image. Image is from NASA  
 697 Photojournal PIA20168.

698

699 **2.6. Phase 6: Observation of dune activity (aeolian or otherwise)**

700 Only recently has it been observed that martian dunes and ripples are very actively migrating  
 701 and evolving within the present-day climate (Figure 9; Bourke et al., 2008; Bridges et al., 2012a; 2012b;  
 702 Chojnacki et al., 2011; 2015; Fenton, 2006; Geissler et al., 2013; Silvestro et al., 2010b; 2011; 2013).  
 703 Previously, an incongruence appeared in our understanding of present-day martian sand transport, as  
 704 the morphology of many aeolian bedforms (apparently sharp crestlines of dunes and ripples) and a  
 705 surface observation of saltation (Greeley et al., 2006) and ripple movement (Sullivan et al., 2008)

706 suggested that some aeolian bedforms should be active. However climate models did not produce the  
707 wind velocities predicted for saltation processes to occur under present conditions and no bedform  
708 motion was observed within higher-resolution images (although some dome dunes were seen to  
709 disappear (Bourke et al., 2008)). This was taken to imply that martian dunes may be stabilized (e.g.,  
710 Zimbelman, 2000) and possibly relict features of a past climate with a denser atmosphere (e.g., Breed et  
711 al., 1979), and that surface degradation processes must be slow. However, acquisition of a sufficient  
712 temporal baseline and careful comparison of overlapping high-resolution images now yield measurable  
713 and consistent changes in dune margin and ripple crestline locations through several fields (e.g.,  
714 Endeavor Crater: Chojnacki et al., 2015), and show that sand fluxes on Mars are comparable to sand  
715 fluxes in the Antarctic Dry Valleys (Bridges et al., 2012b). Within Endeavor Crater, these martian sand  
716 fluxes are sufficient for dune turnover times to be much less than the time since known large climatic  
717 shifts (e.g., an obliquity shift or increased atmosphere density), implying that these dunes are not  
718 records of paleo-climate conditions (Chojnacki et al., 2015).

719         These new observations, proving that sand is currently moving on Mars in large volumes and  
720 that at least some aeolian bedforms are presently active, were helpful in the advance of sediment flux  
721 models and understanding how sediment flux dynamics may vary on different planetary bodies. For  
722 example, an update to the model of steady state saltation (Kok and Renno, 2009) and application to  
723 Earth and Mars conditions (Kok, 2010) showed that saltation can be maintained on Mars by wind speeds  
724 an order of magnitude less than that required to initiate it, while nearly the same wind speed is needed  
725 to both initiate and maintain saltation on Earth. This provides a viable explanation for why aeolian  
726 bedforms appear to evolve at lower-than-predicted wind velocities (as well as an explanation for the  
727 smaller-than-expected minimum dune size on Mars: Kok, 2010). Estimates of aeolian sand flux (in the  
728 present or past) are important as they feed into models of surface erosion rates (e.g., Golombek et al.,  
729 2006; 2014).

730 Sand dunes on Mars are also subject to other processes in the present-day. For example, alcove-  
731 apron and alcove-channel-apron (i.e., gully) formation has been observed in southern mid-latitude dune  
732 fields (Figure 8; Diniega et al., 2010b; Dundas et al., 2012; 2015) and similar activity has been observed  
733 in the north polar dune fields (Hansen et al., 2011; 2015; Horgan and Bell, 2012), moving large volumes  
734 of sand downslope and possibly contributing to the overall migration of the dunes. Some have proposed  
735 that this activity may have aeolian drivers (Horgan and Bell, 2012; Treiman, 2003), but most studies have  
736 shown a seasonal control on the timing of feature formation and evolution, possibly related to CO<sub>2</sub> frost  
737 processes (Diniega et al., 2010b; Dundas et al., 2012; 2015; Hansen et al., 2011). It is also possible that  
738 both aeolian and seasonal frost processes have an influence on these types of dune modification  
739 activities (Hansen et al., 2015). Regardless of underlying process, these changes are actively modifying  
740 the dune slopes (Allen et al., 2016; Diniega et al., 2016; Hansen et al., 2011) and thus need to be  
741 investigated and explained to form a complete story for the martian dune evolution and accurate  
742 interpretation of observed dune morphology.

743 It is also important to note that some dunes have features indicative of a lack of activity, such as  
744 fissures on north polar dunes (Portyankina et al., 2012) and pits and softened topography on southern  
745 mid-latitude dunes (Fenton and Hayward, 2010). Such evidence for stability can provide constraints on  
746 the current availability of mobile material and the near-surface wind environment, as well as a contrast  
747 with the conditions when the (now inactive) bedform had evolved.

748

749 **Summary of (6)**

750 **Data needed:** Repeat images of sufficient spatial and temporal resolution to detect (and measure)  
751 changes in surface morphology (or lack thereof).

752 **Knowledge gain:** Observation and constraints on the estimated (average/net) amount of of sediment  
753 transport.

754 **Assumptions generated:** Activity rates observed in the present-day can be extrapolated to past times  
755 and conditions.

756 **Questions:** What other processes are contributing to dune evolution? How much sediment is moving  
757 within the present climate? Has that amount of aeolian sediment transport changed since a previous  
758 climate?

759 **Lead to investigations of:** How the estimated sediment transport may affect surface erosion rates  
760 (including formation of sand) and formation of other aeolian features such as yardangs; How the climate  
761 has shifted, if changes in sediment transport are apparent.

762

## 763 **2.7. Phase 7: Groundtruth measurements**

764 To-date, we have only visited – at ground-level and up-close – dunes on one planet other than  
765 the Earth. While various Mars rovers have in situ imaged sand deposits and ripples (e.g., Greeley et al.,  
766 2006; JPL, 2012; 2014; Sullivan et al., 2005), Curiosity’s visit to Bagnold Dune Field is the first in situ  
767 observation of dunes and dune sand (JPL, 2015; Bridges et al., 2016). This rover has examined dune sand  
768 on several different slopes on and around dune thought to be undergoing different levels of aeolian  
769 activity (based on orbital observations of ripple migration and the strength of spectral signatures of dust  
770 cover (Lapotre et al., 2016b)). Within even the first observations of dune sand (scooped from the stoss  
771 slope and imaged on the lee slope; Figure 11), grain size differences have been noted that are perhaps  
772 correlated with differences in grain composition (as grains of different sizes appear to correspond to  
773 different materials) (Achilles et al., 2016; Cousin et al., 2016; Ehlmann et al., 2016). Images of the lee  
774 slope of the more active “High dune” have yielded many grainflow features and potentially evidence of  
775 some level of induration (Figure 12; Ewing et al., 2016) – none of which were visible in the orbital  
776 images. The first results of this work are currently being put together, and we look forward to learning  
777 more about the first in situ investigated extraterrestrial dunes.

778



779

780 **Figure 11. Image of the undisturbed surface within the base a Martian sand dune called "High Dune"**  
781 *visited by NASA's Curiosity rover. The image covers an area 3.6 by 2.7 centimeters. Grain sizes show*  
782 *some range, but a fairly consistent size – comparable to dune sand on Earth. It was taken by the Mars*  
783 *Hand Lens Imager (MAHLI) camera on the Curiosity rover's arm on Dec. 5, 2015. Image and description*  
784 *are from JPL (2015).*

785





786

787 **Figure 12: An active dune slipface on Mars, imaged by NASA's Curiosity Mars rover. Multiple grain**  
788 **flows, slumps, and ripples are visible on the slipface of "Namib Dune," a dune within the "Bagnold**  
789 **Dunes" field along the northwestern flank of Mount Sharp, Gale Crater. None of these fine details are**  
790 **visible from orbital (HiRISE) images. The overall slope is 26-28°, and 4-5 m in height. This view combines**  
791 **multiple images from the telephoto-lens camera of the Mast Camera (Mastcam), taken on Dec. 21, 2015.**  
792 **The scene is presented with a color adjustment that approximates white balancing, to resemble how the**  
793 **sand would appear under daytime lighting conditions on Earth. Image and description are from NASA**  
794 **Photojournal PIA20283.**

795

796 **Summary of (7)**

797 **Data needed:** In situ observations of the dune and dune sand (possibly from different portions of the  
798 dune); possibly observations of saltation on the dune or grainflow on the slipface.

799 **Knowledge gain:** Size, composition, and other characteristics of grains involved in saltation.

800 **Assumptions generated:** That the observed characteristics and activity are not anomalous, in time and  
801 space (i.e., the observation didn't catch a rare circumstance/event).

802 **Questions:** Why do the grains look as they do, and what causes the variation/distribution in grain size?  
803 How representative is this observed dune's characteristics and activity? What causes the dune's  
804 features? How can observations of sediment grain size and bedform morphology provide insight  
805 regarding transport processes and the nature/frequency of mobilization events?

806 **Lead to investigations of:** Models of dune activity and evolution, and generation of the observed sand  
807 grains (extending or perhaps redirecting previous work); Based on in situ observation of features that  
808 may not be visible from orbit, what is implied about dune activity and characteristics and how does  
809 information that feed back into models of dune evolution (and what assumptions and related results  
810 should be re-evaluated).

811


### 812 3. Discussion

813       There are some differences between the aeolian science investigations undertaken on each  
814 planetary body - in particular as study methods of more recently studied bodies can build from lessons  
815 learned in aeolian bedform studies of a previously observed body, and as overall our understanding of  
816 aeolian processes becomes more refined as models are forced to reconcile with a wider range of  
817 environmental and planetary conditions. But, as has been described, studies of aeolian bedforms on  
818 planets (other than Earth) broadly tend to follow a similar pattern of gained knowledge, generated  
819 assumptions, and follow-up investigations (that rely on the new knowledge and assumptions). The  
820 similarities in the history of aeolian science over different planetary bodies (Figure 13) are due partially  
821 to the knowledge-advancement at each body being based on the same types of data. Such data is  
822 usually acquired in the same order, which is based on the way in which higher-resolution and increased

823 coverage are acquired during extended or subsequent missions, and as concepts and investigations  
 824 mature and become more specific within all areas of planetary exploration.

825 Within that progression, we focus here on the gaps that seem likely to occur for any planetary  
 826 body. We then move beyond planetary aeolian studies, to look at the interplay of planetary aeolian  
 827 bedform studies with investigative fields that follow their own sequences of discovery and refinement:  
 828 aeolian process modeling and terrestrial aeolian studies.

829



830  
 831 **Figure 13. Timelines showing movement into and through different investigation phases, for Mars,**  
 832 **Titan, and Venus.** Darker colors indicate publication dates for relevant studies (as referenced within this  
 833 paper), and lighter colors indicate the general period of activity (again, based references within this  
 834 paper and checked against google.scholar search results with keywords e.g., “Venus dunes”). Under  
 835 Mars, Phase 6 is divided into observations of “no dune activity due to aeolian bedform processes” (6/n)  
 836 and of such activity (6/a). Within Phase 1 for Titan and Venus, publications predicting a limit on aeolian  
 837 bedform formation on those bodies are highlighted with a P – and the first observation of a bedform  
 838 occurs later. Arrival dates of relevant spacecraft, to the planetary body, are included – for obvious  
 839 reasons, these often initiate or re-invigorate investigations begun during a previous mission.

840

841 **3.1. Gaps that can form within the planetary aeolian science sequence of investigations**

842 Over time and as more data is acquired, our understanding of aeolian processes and  
843 interpretation of the aeolian landforms builds. However, as that understanding builds, it is important to  
844 keep track of which building blocks are assumptions and not actual observations. It is necessary for  
845 assumptions to be made to keep the science investigations moving forward and to guide development  
846 of the next set of investigations, but an assumption that is treated like an “observation” can lead to  
847 models with unrecognized limitations, which in turn can lead to incorrect interpretations of new  
848 observations or even a lack of attention paid to “contradictory” observations. Thus, assumptions should  
849 be recognized as such (and not treated as data) and be re-evaluated for consistency with new and  
850 different data, until direct measurement of the assumed variable or process is possible – doing this can  
851 make it easier to identify and investigate intriguing new understandings about processes and conditions.

852 Several examples of areas where new information has supplanted previous assumptions have  
853 been mentioned within the discussion of the investigation phases (Subsections 2.1-7). Some additional  
854 examples:

- 855 • As higher-resolution and more detailed studies are completed about specific dune fields, results of  
856 these studies (Phases 5-7) should be inserted into field (Phases 2 & 3) and global studies (Phase 4)  
857 that previously relied on lower-resolution or less complete data and assumptions about form and  
858 process uniformity (in time and space) through the field. As was discussed under Phase 7, the  
859 martian large ripples are a new example of this -- where in situ observations are drawing into  
860 question previous work done regarding the scaling of aeolian bedform size between Earth and Mars  
861 and interpretations of ripple crestline complexity, that had been based on interpretation of orbital  
862 images. In general, as more detailed studies are conducted over specific martian dune fields, it is  
863 important to regularly consider how those results fit within the results of larger-scale studies.

864 • These observations have also led into a model of a potential new mode of subaerial bedform  
865 migration and evolution (Lapotre et al., 2016a). As discussed in Phase 5, on the stoss slope of Namib  
866 dune, two types of ripples have been imaged: The large ripples (few m-wavelength) were previously  
867 observed in orbital images, and were thought to be analogous to the wind ripples that we see on  
868 the slopes of terrestrial dunes (e.g., Sullivan et al. 2008). However these large ripples have very  
869 different morphology and dynamics (Silvestro et al., 2016) and in fact are superimposed by small  
870 ripples (~10s cm wavelength) that have morphology more similar to terrestrial impact ripples (but  
871 were not visible within orbital images; Figure 10). The large ripples are now hypothesized to be fluid-  
872 drag ripples (Bagnold, 1951; Wilson, 1973), which on Earth form under water, but on Mars are able  
873 to develop sub-aerially because of the higher kinematic viscosity of the low density atmosphere  
874 (Lapotre et al., 2016a). This example shows the limitations of analysis from only orbital imagery,  
875 even when we think that we understand what we are looking at. Additionally, the limits of  
876 comparative planetology can mean we misinterpret observations where we lack a terrestrial analog.

877 • Dune sand grain sizes on Mars have been estimated since dunes were first seen, based on  
878 assumptions about conditions for dune formation (Phases 1 & 2). Efforts to “measure” grain sizes  
879 from proxy thermal inertia estimates have also been undertaken (mentioned within Phase 4), and  
880 compared to and debated against the assumptions about the ability of the wind to move grains of  
881 different sizes. Studies based on these estimations, and their results, now should be re-evaluated as  
882 Curiosity has recently completed the first in situ investigation of a dune located on a planet other  
883 than Earth (Phase 7), yielding the first direct measurements of martian dune grain sizes (Figure 11).  
884 While on Titan sand grain sizes have not yet been measured in situ (and won't be in the near-  
885 future), studies have explored what grain sizes can be reached via feasible physical processes which  
886 puts constraints on models of dune formation conditions, and visa-versa.

887            Additionally, it is important to recognize the gaps and limitations that can occur in aeolian  
888 studies if only the “standard” aeolian science inputs are considered (e.g., the “complementary sciences”  
889 listed in Table 1 also need to be considered). As in all areas of planetary science and geology, it is  
890 important to consider many pieces of information (and observations, as possible), and all need to be  
891 consistent with the model for the model to be validated. For example, while potential sediment sources  
892 can possibly be tracked from visible imagery, climate models, and/or topography models, compositional  
893 information about the dune grains and the potential sand sources is needed to check that the model is  
894 consistent with the full environment. This may extend beyond compositional information in the local  
895 environment (which was included in Phase 2), as grains may have been transported over large distances  
896 or have been recycled a few times – and this history may not be apparent without a broad-swath of  
897 environmental information. Additionally, processes outside of standard, dune-forming aeolian processes  
898 may be playing a role in dune evolution and observed morphology. For example, the dunes in the  
899 martian polar regions have morphologies and features different from those in the equatorial region,  
900 making it clear that polar processes are altering the aeolian bedforms and thus must be considered in  
901 their interpretation (e.g., in the north: Hansen et al., 2011; 2015; in the south: Fenton et al., 2010).

902

### 903 **3.2. Connections to modeling of the physical processes**

904            As discussed above, looking at aeolian bedforms on other planets allows models to be tested  
905 against a range of environmental and planetary conditions. From that, we refine our understanding of  
906 aeolian processes without assumption of Earth-conditions. This can especially have a large impact on  
907 models of the small-scale and complicated dynamics of sand-wind and sand-sand interactions. For  
908 example, as discussed under Phases 2 and 6, our understanding of the way in which sand is picked up by  
909 the wind, causing or continuing saltation, has now been “tested” under terrestrial, martian, and

910 venusian conditions (Kok, 2010; 2012), resulting in an updated model of how saltation and reptation are  
911 initiated and interact.

912           On Titan, questions about how “sticky” organic sand particles would interact with the wind  
913 were part of an investigation to explain how the dunes had formed, and from that to connect the  
914 crestline orientation to the forming-wind direction(s). The Titan dune sand color appears consistent with  
915 a composition of organics, and such long-chain molecules (of as-yet undetermined exact composition)  
916 could be derived from the atmospheric photodissociation of methane, which creates small particles  
917 (Carl Sagan’s “tholins”) that snow down from the atmosphere (and then perhaps get incorporated into  
918 surface sedimentary layers or clump together into larger granules, that are eroded and transported to  
919 the dune-forming regions) (Radebaugh, 2013; Barnes et al., 2015). Studies of clay-rich dunes in China  
920 had revealed that “sticky” particles could form dunes, but would anchor themselves to the downwind  
921 edge of a longitudinal dune and thus grow and migrate the dune along the dune crestline; this was  
922 proposed as a potential analog to the Titan dunes (Rubin and Hesp, 2009). Although the Titan dune  
923 morphologies were overall found to be more consistent with freely-moving particles (i.e., the saltation  
924 more usually observed on Earth) and thus this longitudinal dune formation model is less favored than  
925 the model discussed in Subsection 2.2. This type of questioning highlighted a different type of terrestrial  
926 dune-formation mechanism and “tested” behavior of the traditional dune formation model if one does  
927 not assume a non-cohesive sand grain. This led to further development of a dune-wind alignment model  
928 that brought these two hypotheses together as well as explained how bedforms with different  
929 alignments can exist within the same multidirectional wind regime (Courrech du Pont et al., 2014).  
930 Within this single model, dune alignment reflects growth via either a “bed instability mode” (which  
931 approximates the longitudinal dune growth process proposed by Rubin and Hunter (1987) and Rubin  
932 and Ikeda (1990)) or a “fingering mode” (the growth process proposed by Rubin and Hesp (2009)),  
933 depending on sediment availability.

934 Models that examine larger-scale dynamics can also be tested through application to different  
935 planetary surfaces. For example, it was in studying martian dunes that a discrepancy was noticed  
936 between the minimum dune size expected on that planet (~100x the minimum Earth dune size) and that  
937 observed (~10x), thus driving new models of dune formation to explain the scaling factor. Model studies  
938 aiming to replicate the observed minimum barchan dune size on Earth and Mars addressed this  
939 question, and tested assumptions about how saltation, reptation, and wind drag interact in setting  
940 characteristic sand trajectory distances, and from this the generation of instability within a sand bed  
941 under a moving fluid (Claudin and Andreotti, 2006).

942

### 943 **3.3. Connections to terrestrial studies and knowledge gain**

944 The trajectory of terrestrial dune studies has differed markedly from the framework proposed  
945 here for planetary dune studies. In essence, the difference is one of top-down vs. bottom-up approaches  
946 as in situ observations of terrestrial landforms, conditions, and activity are significantly easier to carry  
947 out. However, this has not resulted in the history of terrestrial dune fields being an opposite to the  
948 sequence suggested as being characteristic of planetary dune research. The earliest published studies of  
949 terrestrial dune fields were linked with exploration by non-indigenous people, and many of the founding  
950 points of contemporary dune science can be traced to these expeditions. The exploration of the  
951 southern African and Australian interior (mid-19th century), the Sahara (around the beginning of the  
952 20th century, mostly by the French in the west and the English in the east) and the Arabian Rub al'Khali  
953 (most notably by Wilfred Thesiger in the late 1940s) all had exploration as their primary goals. As with  
954 contemporary rover exploration of the martian dune fields, many dune fields were approached with  
955 trepidation due to the hazards they posed. Despite science being incidental rather than implicit to most  
956 of the explorations, there was, nonetheless, early recognition of the great spatial extent of many dune



957 fields, the remarkably organized nature of dunes and the fact that dunes could exist at differing activity  
958 levels.

959           Although Bagnold's work in the 1930s and 1940s is most commonly cited as being the  
960 foundation of modern understanding of aeolian processes and landforms, there were significant  
961 precursors. George Perkins Marsh (1864) considered geoengineering problems associated with drifting  
962 sand, and the role of vegetation in stabilizing dunes, and Russian geologist Nikolay Sokolów had  
963 discussed dune sedimentology and theories of dune formation in a 300 page book devoted to the  
964 subject (1894). Georges Rolland, a French mining engineer, set out a series of propositions in 1890  
965 based on fieldwork in the Algerian Sahara which addressed such issues as sediment source, the  
966 distribution of dune fields, varying levels of dune activity and the relationship between wind regime and  
967 different dune shapes (Burt et al., 2008). At this point, the role of the wind in dune formation was still  
968 contested by many, and it was widely held that dunes would prove to have rocky cores (Goudie, 2002).  
969 Many other aspects of contemporary aeolian science date from surprisingly early studies. Aerial imagery  
970 was used to examine dune planform morphology in the 1930s (Aufrère 1932, Madigan 1936), and the  
971 recognition of dunes as a particulate waveform in a fluid medium can be traced to the work of Cornish  
972 (1914). Bagnold's work, utilizing field and wind tunnel experimentation, is an early example of the  
973 'quantitative revolution' widely recognized in geosciences in the middle of the 20th century. This directly  
974 influenced the next half-century of research, via fieldwork and laboratory experimentation, in a phase  
975 perhaps best summarized by Lancaster's (1995) state-of-the-art textbook. Coincidentally, the same year  
976 saw the publication of Werner's (1995) application of cellular automata models to aeolian bedforms,  
977 which accepted that dunes formed as an emergent property of a complex system, one of the first  
978 indications of the failure of reductionist approaches to fully explain aeolian landscapes (Livingstone et  
979 al., 2007). The same period saw the rise of the use of luminescence dating to provide ages for dune

980 emplacement, since described as having has a transformative effect on studies of dryland science  
981 (Singhvi and Porat, 2008).

982           Planetary studies of aeolian dunes therefore have the advantage of decades of terrestrial work  
983 to draw upon, and this is reflected in the very rapid progress made on newly-discovered dunes (e.g.  
984 Titan, Comet 67P). Terrestrial science, conversely, has had the advantage of a relatively steady  
985 progression in the quality of the available data -- although the related understanding of aeolian systems  
986 has not progressed as steadily. The progress made in understanding terrestrial dunes has not been  
987 without challenges, and it is instructive to reflect on whether there are lessons for the planetary  
988 community can be drawn from progress on terrestrial dune fields, and conversely whether the evolution  
989 of extraterrestrial dune research can inform the research strategies of Earth's dune studies.

990

991 ***What can planetary science learn from the history of terrestrial dune studies?***

992           Much of planetary dune science is already directly influenced by the methods, theory and  
993 process understanding derived from terrestrial studies, manifest in the numerous analog studies.  
994 However, there are some less well-discussed points that are worthy of consideration.

995           As was noted in section 3.1, close attention must be paid to the difference between assumed  
996 and observed knowledge. Cautionary tales can be drawn from terrestrial dune studies, and this is  
997 perhaps best illustrated by the roll vortex hypothesis for longitudinal (linear) dune formation. First  
998 proposed by Bagnold (1953), and promoted subsequently (e.g. Hanna, 1969) this suggested that  
999 thermally induced vortices in regional wind-flow would lead to the development of helical horizontal  
1000 flow cells that might lead to sand accumulation in linear bedforms extending downwind. The theory is  
1001 strikingly devoid of empirical supporting evidence, and yet still persists in the literature. Quite simply,  
1002 vortices of sizes that might explain dune spacing have never been observed despite numerous  
1003 experimental attempts, and the transverse component of roll vortices does not appear to have sufficient

1004 velocity to move sand (Lancaster 1995). Planetary studies should be careful to question existing  
1005 paradigms and theories, and be willing to point out when data do not support these hypotheses.

1006         Bagnold's great advances in aeolian science can be largely attributed to willingness and  
1007 fearlessness towards innovation, in terms of methods and physical exploration. The novel application of  
1008 wind tunnels to aeolian transport and sedimentation and the methods developed to enable remote  
1009 desert travel directly enabled the advances in understanding that Bagnold brought. Planetary  
1010 perspectives support this, with the radical advances in data brought from missions such as MSL,  
1011 Cassini/Huygens, Rosetta/Philae and New Horizons. Such evidence supports the potential knowledge  
1012 gains from similarly ambitious mission concepts of other planetary surface exploration missions, such as  
1013 AVIATR (Barnes et al. 2012) and VISE (NRC, 2013) which have been considered for Titan and Venus  
1014 exploration, respectively. The evidence from both terrestrial and planetary dune studies suggests that  
1015 high-risk, innovative research has led to some of the greatest advances.

1016         The discrepancy between the timescales of aeolian process and the timescales evident in  
1017 aeolian landscapes is also very evident – possibly even more so – on some planetary bodies. Despite  
1018 processes operating within dune landscape on timescales of seconds to hours, the resultant landscape  
1019 development frequently operates on timescales of  $>10^3$  years. Dating of aeolian sediment, primarily via  
1020 the suite of luminescence dating methods, has been adopted very widely on terrestrial dune studies,  
1021 and has played a crucial role in linking the short-term process understanding with the long-term  
1022 geomorphological record. It has enabled calculation of rates of landform evolution beyond that possible  
1023 using observational records (e.g. Kocurek et al., 2007; Telfer, 2011), revealed complex spatial variability  
1024 in aeolian accumulation (Telfer and Thomas, 2007) and frequently been used to infer external drivers of  
1025 dune activity (e.g. climatic changes). Experimentation with luminescence readers suitable for Mars  
1026 missions has been explored (e.g. McKeever et al., 2003; Jain et al. 2006), and if the substantial technical  
1027 challenges can be overcome (Doran et al. 2004), martian luminescence dating offers the potential to

1028 extend understanding of accumulation beyond the period of direct observation. Recent progress  
1029 suggests that solutions may exist to these challenges (e.g. Sohbaty et al., 2012).

1030

1031 ***What can terrestrial dune studies learn from the history of planetary science?***

1032           Although at least parts of Phases 1-3 and 5-7 have been investigated on Earth for 70 years or  
1033 more, a striking difference between planetary and terrestrial dune studies is that currently there is no  
1034 global catalogue of dunes for Earth (Phase 4). The first edition of the Mars Global Digital Dune Database  
1035 (Hayward et al., 2007) was published within six years of the start of THEMIS data collection (which are  
1036 used to identify the thermal inertia proxy identifiers for dune fields), and global mapping of Titan's  
1037 dunes within the constraints of the available data (Lorenz and Radebaugh, 2009) was published within a  
1038 similar timeframe since the arrival of Cassini at Titan. Although some terrestrial regions have been  
1039 mapped and duneforms catalogued (e.g., Namib; Livingstone et al., 2010), and a global database of dunes  
1040 with dating constraints has recently been compiled (Lancaster et al., in press), global-scale consideration  
1041 of terrestrial dune fields is currently lagging behind planetary science. Efforts in this direction are  
1042 currently in progress (Hesse et al., 2015) -- but it has been over 40 years since the advent of global  
1043 terrestrial satellite coverage. The focus of planetary global catalogues of dunes has been on  
1044 understanding global circulation patterns (e.g. Charnay et al. 2015, Ewing et al. 2015), sediment sources  
1045 (e.g. Tirsch et al., 2011), identification of large-scale variations in dune form due to different evolution  
1046 processes or rates (e.g., Hayward and Fenton, 2010; Savage et al., 2014), and targeting areas for detailed  
1047 study (e.g., Hayward, 2011). Whilst it is not necessary to use dune morphology to understand modern  
1048 circulation patterns on Earth, applications of such a database would include quantification of aeolian  
1049 sediment volumes and flux, improved understanding of regions where dunes are currently stabilized,  
1050 and potential for monitoring change in environmentally-sensitive, dynamic landscapes.

1051 Livingstone et al. (2007), reviewing the state of understanding of terrestrial dune  
1052 geomorphology, concluded that decades of largely inductive, and increasingly reductionist, study had  
1053 not brought the completeness of understanding that had been hoped, and that integration of  
1054 methodologies (field, modeling and remote sensing) offered the best prospects for knowledge. Perhaps  
1055 due to the difficulties in conducting 'field study' of extraterrestrial dunes (i.e., Phase 7), which are only  
1056 very recently being overcome on Mars, such combined strategies are often exemplified by planetary  
1057 aeolian studies, where studies employing a wide range of methodologies including numerical modeling,  
1058 laboratory experimentation, field study (presently via analog environments), and remote sensing are  
1059 commonplace (e.g., Lucas et al. 2014). Although some terrestrial studies do synthesize such diverse  
1060 methodologies, the example set by many planetary studies is a good one for terrestrial dune studies.

1061 Much of the focus of this paper has been on the increasing availability, resolution, and coverage  
1062 of new remotely sensed data. The same has been true for Earth, and here lessons learned in planetary  
1063 studies help guide interpretation of terrestrial images at the margins of the spectral and/or spatial  
1064 resolution of the imagery. This is perhaps best illustrated with the example of the highly contested issue  
1065 of extensive palaeo-dune fields in the Amazon Basin, presently covered with extensive tropical forest.  
1066 Tricart (1974), working with recently-released first-generation Landsat imagery, identified widespread  
1067 stabilized aeolian landforms in the tropical Amazon Basin. This interpretation was repeated numerous  
1068 times (e.g. Klammer, 1982) and led to a widespread belief in arid phases accompanied by vegetation  
1069 loss during the late Quaternary evolution of the region, with huge implications for understanding of  
1070 regional biogeography in one of the world's most biodiverse regions. However, whilst subsequent  
1071 reanalysis of high-resolution data has revealed aeolian dune fields around the margins of the Amazon  
1072 basin (e.g. Teeuw and Rhodes, 2004; May, 2013) and/or immediately adjacent to large rivers where the  
1073 sediment supply has at times been the dominant control (e.g. Carneiro, 2002), the existence of wide  
1074 swathes of paleodunes across the Amazon basin has not withstood closer scrutiny (Colinvaux et al.,

1075 2001). Tripaldi and Zarate (in press) reviewed the evidence, and demonstrated the importance of  
1076 groundtruthing when image interpretation is challenging. Planetary geomorphologists are usually  
1077 admirably conservative in terms of implying process from (apparent) landform, especially when imagery  
1078 is at the limit of its spatial or spectral resolution, and terrestrial incidents such as the question of an arid  
1079 Amazon suggest that such conservatism is wise.

1080

#### 1081 **4. Conclusion/Summary**

1082         Studies of aeolian bedforms over a wide range of planetary bodies have resulted in significant  
1083 progress in our understanding of past and present climate and surface conditions, physical processes,  
1084 and the interconnectivity of dynamics over a range of spatial and temporal scales. These studies  
1085 contribute, in meaningful and often unique ways, towards a range of planetary science investigations.  
1086 For example, as discussed, interpretation of dune morphology often provides unique, if proxy,  
1087 groundtruth data about past or present wind conditions, and the proven presence of a large amount of  
1088 sand grains can drive investigations about processes responsible for creating such grains. Beyond studies  
1089 that involve this type of direct interpretation of the aeolian bedforms, aeolian science studies also yield  
1090 information about many tangentially-related areas of investigation. In particular, aeolian-driven sand  
1091 flux appears to be an important force in erosional modification of a planetary surface. Quantitative  
1092 estimations of wind speeds and sand flux and identification of sediment transport pathways yield  
1093 quantitative estimates of erosional process rates. This can, for example, lead to improved interpretation  
1094 of observed landforms – such as yardangs (e.g., Ward 1979), or the rate of crater degradation by aeolian  
1095 processes which is important for accurately estimating the age of a planetary surface (e.g., Golombek et  
1096 al., 2014; Grant et al., 2006; 2008; 2016). This can also provide bounds on surface-ages of exposed rock  
1097 surfaces, which is can be of importance to rover missions – such as Mars missions searching for

1098 reachable-environments near the surface that may have been habitable and that may include preserved  
1099 biosignatures (e.g., Arvidson et al., 2015).

1100         As discussed, planetary aeolian studies have also made key contributions towards improving the  
1101 methodologies employed in aeolian science, and in challenging assumptions built (perhaps  
1102 inadvertently) into aeolian process models based on terrestrial observations. To-date, this has resulted  
1103 in the refinement of several models of dune-field forming processes, from interactions between sand  
1104 grains and the wind or with each other, up through interactions between dunes and topography and  
1105 climate shifts.

1106         Given all of the ways in which our aeolian study results impact our understanding of planetary  
1107 surface conditions and histories (as well as the Earth's), it is thus very important to critically look at how  
1108 we progress in planetary aeolian science, and in particular to consider carefully which results (and  
1109 resultant models) are based on assumptions versus observations – and then to revisit those results  
1110 when new information becomes available. Here, we have proposed one framework for identifying  
1111 progress within planetary aeolian studies, and have used that framework to chart the progression of  
1112 data, assumptions, and generated knowledge. We hope this framework, and our identification of gaps,  
1113 will help future planetary aeolian researchers strategically fill knowledge gaps or at least carefully  
1114 recognize where assumptions are being used to progress a study.

1115         Additionally, this framework may help identify the types of data that would be most useful for  
1116 future planetary missions. Pluto, Io, and Comet 67P were all discussed as having reached Phase 1, where  
1117 at least a potential aeolian bedform has been observed. On Titan, global datasets exist and have  
1118 contributed to large shifts in our understanding of the Titan climate and organic cycles. Venus also has a  
1119 global topography dataset, but the low resolution and apparent lack of dune fields stalled progress in its  
1120 aeolian science investigations (and thus related advancements in planetary surface studies).

1121 Unfortunately for Venus and Titan, further progression within Phases 2-4 (and movement into Phases 5  
1122 and beyond) will likely need to wait for new and higher-resolution surface datasets.

1123 Mars' aeolian bedforms are the best studied within planetary aeolian science (outside of  
1124 Earth's), with both widespread coverage in certain data-types and many regions with high-resolution  
1125 data regarding the dunes' and dune field environment's morphology and composition, collected over  
1126 the past 43 years. However, with the progress that has been made, we cannot lose sight of the fact that  
1127 much of it has been built on interpretations of remote data. (As discussed under Phase 7, in situ dune  
1128 field observations have not been possible until just recently.) Furthermore, much of the work involves a  
1129 meshing of coarse global data with a few more-deeply monitored and studied dune fields, and thus  
1130 much extrapolation is done that assumes certain types of consistency between fields. This is an odd  
1131 contrast with Earth dune field studies, where the global dataset (Phase 4) is what is missing.

1132 For all planetary bodies (including Earth), we look forward to further advancements in the  
1133 interpretation of aeolian bedforms and what interpretations about those bedforms will imply about the  
1134 environmental conditions and processes. If history is to be any guide, with each advance into a new  
1135 phase (due to acquisition of a new type of data and/or enablement of a new type of analysis), we find  
1136 exciting new understandings about that planetary body and the general understanding of aeolian  
1137 processes. One area of intriguing advancement is the prediction of where dunes and/or ripples could be  
1138 found (which could be thought of as a "Phase 0" within our framework). As we explore more bodies and  
1139 learn more about the conditions under which bedforms resembling aeolian dunes are found, we can  
1140 wonder about the next place where we may expect to find potential dunes, as well as identify lessons to  
1141 aid in such predictions (e.g., when we return to Venus). In addition, perhaps in the near future, we will  
1142 move into a yet-undefined Phase 8 (e.g., through comparison between in situ measurements of some  
1143 very different types of aeolian bedforms? Hints of that are starting with sand grain comparisons (e.g.,  
1144 O'Connell-Cooper, 2016)), yielding a new type of data that can supersede assumptions made in Phases



1145 1-7, further expanding our broad understanding of aeolian processes and bedforms, and increasing the  
1146 overall information gained from planetary aeolian studies.

1147

#### 1148 **Acknowledgements**

1149 [Mostly removed for review.] We thank the two reviewers for their very helpful comments, and in  
1150 particular for the suggestions of additional references.

1151

#### 1152 **References**

- 1153 Achilles, C.N., D.T. Vaniman, D.F. Blake, T.F. Bristow, E.B. Rampe, D.W. Ming, S.J. Chipera, R.V. Morris,  
1154 S.M. Morrison, R.T. Downs, K.V. Fendrich, B.L. Ehlmann, A.S. Yen, P.C. Sarrazin, A.H. Treiman, P.I.  
1155 Craig, M.G.A. Lapotre, K.S. Edgett, R. Gellert, J.A. Crisp, J.M. Morookian, J.P. Grotzinger, D.J. Des  
1156 Marais, J.D. Farmer, 2016. Mineralogy of eolian sands at Gale crater. 47<sup>th</sup> LPSC, Ab. 2532.
- 1157 Allen, A., S. Diniega, C. Hansen, 2016. Gully and aeolian activity within the “Tleilax” dune field in the  
1158 Olympia Undae, Mars. 47<sup>th</sup> LPSC, Ab. 1759.
- 1159 Andreotti, B., P. Claudin, O. Pouliquen, 2006. Aeolian Sand Ripples: Experimental Study of Fully  
1160 Developed States. *Physical Review Letters*, v. 96, 028001, doi:10.1103/PhysRevLett.96.028001.
- 1161 Arvidson, R. E., J.F. Bell III, J.G. Catalano, B.C. Clark, V.K. Fox, R. Gellert, J.P. Grotzinger, E. A. Guinness,  
1162 K.E. Herkenhoff, A.H. Knoll, M. G. A. Lapotre, S.M. McLennan, D.W. Ming, R.V. Morris, S.L. Murchie,  
1163 K.E. Powell, M.D. Smith, S.W. Squyres, M.J. Wolff, J.J. Wray, 2015. Mars Reconnaissance Orbiter and  
1164 Opportunity Observations of Burns Formation and Underlying Strata: Crater-Hopping at Meridiani  
1165 Planum. *J. Geophys. Res. - Planets*, v. 120, issue 3, 429-451, doi:10.1002/2014JE004686.
- 1166 Aufrère, L., 1932. Morphologie dunaire et météorologie saharienne. *Association de Géographes Français*  
1167 *Bulletin* 56, 34-47.
- 1168 Ayoub, F., J-P. Avouac, C.E. Newman, M.I. Richardson, A. Lucas, S. Leprince, N.T Bridges, 2014. Threshold  
1169 for sand mobility on Mars calibrated from seasonal variations of sand flux. *Nature Communications*  
1170 5, doi:10.1038/ncomms6096.
- 1171 Bagnold, R.A., 1941. *The Physics of Blown Sand and Desert Dunes* (New York: Methuen).
- 1172 Bagnold, R.A, 1953. The surface movement of blown sand in relation to meteorology. *Research Council*  
1173 *of Israel, Special Publication*, 2, 89-93.
- 1174 Barnes, J.W., R.H. Brown, L. Soderblom, C. Sotin, S. Le Mouelic, S. Rodriguez, R. Jaumann, R.A. Beyer, R.  
1175 Clark, P. Nicholson, 2008. Spectroscopy, morphometry, and photoclinometry of Titan’s dune fields  
1176 from Cassini/VIMS. *Icarus* 195, 400-414.
- 1177 Barnes, J.W., L. Lemke, R. Foch, C.P. McKay, R.A. Beyer, J.Radebaugh, D.H. Atkinson, R.D. Lorenz, S. Le  
1178 Mouélic, S. Rodriguez, J. Gundlach, 2012. AVIATR—Aerial Vehicle for In-situ and Airborne Titan  
1179 Reconnaissance. *Experimental Astronomy*, 33(1), pp.55-127.
- 1180 Barnes, J.W., R.D. Lorenz, J. Radebaugh, A.G. Hayes, K. Arnold, C. Chandler, 2015. Production and global  
1181 transport of Titan's sand particles. *Planetary Science* 4, 1-19.

1182 Basilevsky, A.T., R.O. Kuzmin, O.V. Nikolaeva, A.A. Pronin, L.B. Ronca, V.S. Avduevsky, G.R. Uspensky, Z.P.  
1183 Cheremukhina, V.V. Semenchenko, V.M. Ladygin, 1985. The surface of Venus as revealed by the  
1184 Venera landings: Part II. Geological Society of America Bulletin, 96, 137-144.

1185 Belcher, D., J. Veverka, C. Sagan, 1971. Mariner photography of Mars and aerial photography of Earth:  
1186 Some analogies, Icarus, vol. 15, 241-252.

1187 Bondarenko, N. V., M. A. Kreslavsky, J. W. Head, 2006. North-south roughness anisotropy on Venus from  
1188 the Magellan Radar Altimeter: Correlation with geology, Journal of Geophysical Research (Planets)  
1189 111, E06S12.

1190 Bourke, M.C., K.S. Edgett, B.A. Cantor, 2008. Recent aeolian dune change on Mars. *Geomorphology*,  
1191 94(1-2), 247-255, doi:10.1016/j.geomorph.2007.05.012.

1192 Breed, C., M. J. Grolier, J.F. McCauley, 1979. Morphology and Distribution of Common 'Sand' Dunes on  
1193 Mars: Comparison With the Earth, J. Geophys. Res., vol. 84, no. B14, 8183-8204.

1194 Bridges, N.T., M.C. Bourke, P.E. Geissler, M.E. Banks, C. Colon, S. Diniega, M.P. Golombek, C.J. Hansen, S.  
1195 Mattson, A.S. McEwen, M.T. Mellon, N. Stantzos, B.J. Thomson (2012a), Planet-wide sand motion on  
1196 Mars. *Geology* 40(1), 31-34, doi:10.1130/G32373.1.

1197 Bridges, N.T., F. Ayoub, J-P. Avouac, S. Leprince, A. Lucas, S. Mattson (2012b), Earth-like sand fluxes on  
1198 Mars. *Nature* 485, 339-342, doi:10.1038/nature11022.

1199 Bridges, N.T., B.L. Ehlmann, R.C. Ewing, C.E. Newman, R. Sullivan, P.G. Conrad, A. Cousin, K.S. Edgett, M.R.  
1200 Fisk, A.A. Fraeman, J.R. Johnson, M. Lamb, M. Lapotre, S. Le Mouélic, G.M. Martinez, P-Y. Meslin, P.  
1201 Pinet, L.M. Thompson, J. van Beek, A.R. Vasavada, R.C. Wiens, 2016. Investigation of the Bagnold  
1202 Dunes by the Curiosity rover: Overview of initial results from the first study of an active dune field  
1203 on another planet, 47th LPSC, Ab. 2298.

1204 Burt et al. 2008 (eds). The history of the study of landforms, Volume 4: A history of geomorphology  
1205 between the 1890s and the 1960s. London, The Geological Society.

1206 Burr, D. M., S.A. Drummond, R. Cartwright, B.A. Black, J.T. Perron (2013). Morphology of fluvial networks  
1207 on Titan: Evidence for structural control, Icarus, vol. 226(1), 742-759,  
1208 doi:10.1016/j.icarus.2013.06.016.

1209 Canada Dept. Agriculture (1976), Glossary of terms in soil science, © Minister of Supply and Services,  
1210 Canada, ISBN 0-662-01533-9. [http://sis.agr.gc.ca/cansis/publications/manuals/1976-](http://sis.agr.gc.ca/cansis/publications/manuals/1976-glossary/pub1459_report.pdf)  
1211 [glossary/pub1459\\_report.pdf](http://sis.agr.gc.ca/cansis/publications/manuals/1976-glossary/pub1459_report.pdf)

1212 Cardinale, M., G. Komatsu, S. silvestro, D. Tirsch, 2012. The influence of local topography for wind  
1213 direction on Mars: two examples of dune fields in crater basins. *Earth Surface Processes and*  
1214 *Landforms*, 37, 13, 1437-1443, DOI: 10.1002/esp.3289.

1215 Cardinale, M., S. Silvestro, D.A. Vaz, T. Michaels, M.C. Bourke, G. Komatsu, L. Marinangeli, 2016. Present-  
1216 day aeolian activity in Herschel Crater, Mars. *Icarus* 265, 139-148, doi:10.1038/ncomms6096.

1217 Carneiro Filho, A., D. Schwartz, S.H. Tatum, T. Rosique, 2002. Amazonian paleodunes provide evidence  
1218 for drier climate phases during the Late Pleistocene-Holocene. *Quaternary Research*, 58(2), pp.205-  
1219 209.

1220 Charnay, B., E. Barth, S. Rafkin, C. Narteau, S. Lebonnois, S. Rodriguez, S.C. Du Pont, A. Lucas, 2015.  
1221 Methane storms as a driver of Titan's dune orientation. *Nature Geoscience*, 8(5), pp.362-366.

1222 Cheng, A.F., C.M. Lisse, M. A'Hearn, 2013. Surface geomorphology of Jupiter Family Comets: A geologic  
1223 process perspective. *Icarus* 222, 808-817, doi:10.1016/j.icarus.2012.10.004.

1224 Christensen, P.R., 1983. Eolian intracrater deposits on Mars: Physical properties and global distribution.  
1225 *Icarus* 56(3), 496-518.

1226 Christensen, P.R., J.L. Bandfield, J.F. Bell III, N. Gorelick, V.E. Hamilton, A. Ivanov, B.M. Jakosky, H.H.  
1227 Kieffer, M.D. Lane, M.C. Malin, T. McConnochie, 2003. Morphology and composition of the surface  
1228 of Mars: Mars Odyssey THEMIS results. *Science* 300(5628), pp.2056-2061.

1229 Chojnacki, M., D.M. Burr, J.E. Moersch, T.I. Michaels, 2011. Orbital observations of contemporary dune  
1230 activity in Endeavour Crater, Meridiani Planum, Mars. *J. Geophys. Res.* 116, E00F19. Chojnacki, M.,  
1231 D. M. Burr, J. E. Moersch, J. J. Wray, 2014. Valles Marineris dune sediment provenance and  
1232 pathways. *Icarus*, 232(0), 187–219, doi:10.1016/j.icarus.2014.01.011.

1233 Chojnacki, M., J.R. Johnson, J.E. Moersch, L.K. Fenton, T.I. Michaels, J.F. Bell III, 2015. Persistent aeolian  
1234 activity at Endeavour crater, Meridiani Planum, Mars; new observations from orbit and the surface.  
1235 *Icarus*, 251(0), 275–290, doi:10.1016/j.icarus.2014.04.044.

1236 Claudin, P., B. Andreotti, 2006. A scaling law for aeolian dunes on Mars, Venus, Earth, and for  
1237 subaqueous ripples. *Earth and Planetary Science Letters*, vol. 252, 30–44,  
1238 doi:10.1016/j.epsl.2006.09.004.

1239 Colinvaux, P.A., G. Irion, M.E. Räsänen, M.B. Bush, J.N. De Mello, 2001. A paradigm to be discarded:  
1240 geological and paleoecological data falsify the Haffer & Prance refuge hypothesis of Amazonian  
1241 speciation. *Amazoniana*, 16(3), pp.609-646.

1242 Courrech du Pont, S., C. Narteau, X. Gao, 2014. Two modes for dune orientation. *Geology*, 42(9),  
1243 doi:10.1130/G35657.1.

1244 Cousin, A., O. Forni, P.-Y. Meslin, S. Schröder, O. Gasnault, N. Bridges, B. Elhmann, S. Maurice, R. Wiens,  
1245 2016. Chemical diversity among fine-grained soils at Gale (Mars): a chemical transition as the rover  
1246 is approaching the Bagnold Dunes? 47<sup>th</sup> LPSC, Ab. 2044.

1247 Cutts, J.A., R.S.U. Smith, 1973. Eolian deposit and Dunes on Mars. *J. Geophys. Res.* 78(20), 4139-4154.

1248 Cutts, J.A., K.R. Blasius, G.A. Briggs, M.H. Carr, R. Greeley, H. Masursky, 1976. North Polar Region of  
1249 Mars: Imaging Results from Viking 2. *Science*, vol. 194(4271), 1329-37.

1250 Diniega, S., K. Glasner, S. Byrne, 2010a. Long scale evolution of aeolian sand dune fields: influences of  
1251 initial conditions and dune collisions. *Geomorphology (special edition: Planetary Dunes)* 121, 55-68,  
1252 doi:10.1016/j.geomorph.2009.02.010.

1253 Diniega, S., S. Byrne, N.T. Bridges, C.M. Dundas, A.S. McEwen, 2010b. Seasonality of present-day Martian  
1254 dune-gully activity. *Geology* 38, no. 11, 1047-1050, doi:10.1130/G31287.1.

1255 Diniega, S., A. Allen, T. Perez, C.J. Hansen, 2016. Tracking gully activity within the north polar erg, Mars.  
1256 47<sup>th</sup> LPSC, Ab. 1740.

1257 Doran, P.T., S.M. Clifford, S.L. Forman, L. Nyquist, D.A. Papanastassiou, B.W. Stewart, N.C. Sturchio, T.D.  
1258 Swindle, T. Cerling, J. Kargel, G. McDonald, 2004. Mars chronology: assessing techniques for  
1259 quantifying surficial processes. *Earth-Science Reviews*, 67(3), pp.313-337.

1260 Dundas C.M., S. Diniega, C.J. Hansen, S. Byrne, A.S. McEwen, 2012. HiRISE observations of seasonal  
1261 activity and morphological changes in Martian gullies. *Icarus* 220(1), 124-143,  
1262 doi:10.1016/j.icarus.2012.04.005.

1263 Dundas, C.M., S. Diniega, A.S. McEwen, 2015. Long-Term Monitoring of Martian Gully Formation on  
1264 Mars and Evolution with HiRISE. *Icarus (special issue: Dynamic Mars)* 251, 244-263,  
1265 doi:10.1016/j.icarus.2014.05.013.

1266 Edgett, K.S., P.R. Christensen, 1991. The particle size of Martian aeolian dunes. *JGR* 96(E5), 22765-22776.

1267 Edgett, K. S., N. Lancaster (1993), Volcaniclastic aeolian dunes: terrestrial examples and application to

1268 martian sands, *J. Arid Environ.*, 25(3), 271-297, doi:10.1006/jare.1993.1061.

1269 Ehlmann, B.L., N. Bridges, A.A. Fraeman, M.G.A. Lapotre, K. Edgett, J.R. Johnson, A. Cousin, A. Yen, P.

1270 Conrad, L. Thompson, J. Van Beek, D. Vaniman, S. Schröder, A. Vasavada, 2016. Chemistry and

1271 mineralogy in situ at the Bagnold sand dunes: Evidence for aeolian sorting and size-dependence in

1272 sand composition. 47<sup>th</sup> LPSC, Ab. 1536.

1273 Ewing, R.C., G. Kocurek, 2010. Aeolian dune-field pattern boundary conditions. *Geomorphology* 114,

1274 175-187, doi:10.1016/j.geomorph.2009.06.015.

1275 Ewing, R.C. A. Peyret, G. Kocurek and M.C. Bourke, 2010. Dune-field pattern formation and recent

1276 transporting winds in the Olympia Undae dune field, north polar region of Mars. *Journal of*

1277 *Geophysical Research*, 115, E08005.

1278 Ewing, R.C., A.G. Hayes, A. Lucas, 2015. Sand dune patterns on Titan controlled by long-term climate

1279 cycles. *Nature* 10.1038/ngeo2323.

1280 Ewing, R.C., N.T. Bridges, R. Sullivan, M.G.A. Lapotre, W.W. Fisher, M.P. Lamb, D. Rubin, K. Lewis, S.

1281 Gupta, 2016. Aeolian Sedimentary Processes at the Bagnold Dunes, Mars: Implications for Modern

1282 Dune Dynamics and Sedimentary Structures in the Aeolian Stratigraphic Record of Mars. 47<sup>th</sup> LPSC,

1283 Ab. 2783.

1284 Fenton, L.K., 2005. Potential sand sources for the dune fields in Noachis Terra, Mars. *J. Geophys. Res.*,

1285 vol. 110, E11, doi:10.1029/2005JE002436.

1286 Fenton, L. K., 2006. Dune migration and slip face advancement in the Rabe Crater dune field, Mars.

1287 *Geophys. Res. Lett.*, 33(20), 1-5, doi:10.1029/2006GL027133.

1288 Fenton, L., 2016. Dunes on Pluto? Blog on SETI website: [http://www.seti.org/seti-institute/dunes-on-](http://www.seti.org/seti-institute/dunes-on-pluto)

1289 [pluto](http://www.seti.org/seti-institute/dunes-on-pluto).

1290 Fenton, L.K., R.K. Hayward, 2010. Southern high latitude dune fields on Mars: Morphology, aeolian

1291 inactivity, and climate change, *Geomorphology* 121, Issues 1-2, 98-121,

1292 doi:10.1016/j.geomorph.2009.11.006.

1293 Fenton, L.K., A.D. Toigo, M.I. Richardson, 2005. Aeolian processes in Proctor Crater on Mars: Mesoscale

1294 modeling of dune-forming winds. *J. Geophys. Res.* 110, E06005, doi:10.1029/2004JE002309.

1295 Florensky, C.P., A.T. Basilevsky, V.P. Krychkov, R.O. Kusmin, O.V. Nikolaeva, A.A. Pronin, I.M. Chernaya,

1296 Yu.S. Tyufin, A.S. Selivanov, M.K. Naraeva, L.B. Ronca, 1983. Venera 13 and Venera 14: Sedimentary

1297 rocks on Venus? *Science* 221, 57-589.

1298 Fryberger, S.G., G. Dean, 1979. Dune forms and wind regime, in, E. D. McKee, ed., *A study of global sand*

1299 *seas*. U.S. Geol. Surv. Prof. Pap. 1052, 137-169.

1300 Gardin, E., P. Allemand, C. Quantin, S. Silvestro, C. Delacourt, 2012. Dune fields on Mars: Recorders of a

1301 climate change? *Planet. Space Sci.* 60, 314-321, doi:10.1016/j.pss.2011.10.004.

1302 Garzanti, E., P. Vermeesch, S. Andò, G. Vezzoli, M. Valagussa, K. Allen, K.A. Kadi, A.I.A. Al-Juboury, 2013.

1303 Provenance and recycling of Arabian desert sand. *Earth-Sci. Rev.* 120, 1-19,

1304 doi:10.1016/j.earscirev.2013.01.005.

1305 Geissler, P.E., N.W. Stantzos, N.T. Bridges, M.C. Bourke, S. Silvestro, L.K. Fenton, 2013. Shifting sands on

1306 Mars: Insights from tropical intra-crater dunes. *ESP&L*, vol. 38, no. 4, 407-412,

1307 doi:10.1002/esp.3331.

1308 Golombek, M. P., J.A. Grant, L.S. Crumpler, R. Greeley, R.E. Arvidson, J.F. Bell III, C.M. Weitz, R. Sullivan,  
1309 P.R. Christensen, L.A. Soderblom, S.W. Squyres, 2006. Erosion rates at the Mars Exploration Rover  
1310 landing sites and long-term climate change on Mars. *J. Geophys. Res.*, 111, E12S10,  
1311 doi:10.1029/2006JE002754.

1312 Golombek, M.P., N.H. Warner, V. Ganti, M.P. Lamb, T.J. Parker, R.L. Fergason, R. Sullivan, 2014. Small  
1313 crater modification on Meridiani Planum and implications for erosion rates and climate change on  
1314 Mars. *J. Geophys. Res. Planets*, 119, doi:10.1002/2014JE004658.

1315 Goudie, A., 2002. Great warm deserts of the world: landscapes and evolution (Vol. 1). Oxford University  
1316 Press on Demand.

1317 Grant, J. A., R.E. Arvidson, L.S. Crumpler, M.P. Golombek, B. Hahn, A.F.C. Haldemann, R. Li, L.A.  
1318 Soderblom, S. W. Squyres, S.P. Wright, W.A. Watters, 2006. Crater gradation in Gusev crater and  
1319 Meridiani Planum, Mars. *J. Geophys. Res.*, 111, E02S08, doi:10.1029/2005JE002465.

1320 Grant, J. A., S. A. Wilson, B. A. Cohen, M. P. Golombek, P. E. Geissler, R. J. Sullivan, R. L. Kirk, and T. J.  
1321 Parker (2008), Degradation of Victoria crater, Mars, *J. Geophys. Res.*, 113, E11010,  
1322 doi:10.1029/2008JE003155.

1323 Grant, J. A., T. J. Parker, L. S. Crumpler, S. A. Wilson, M. P. Golombek, and D. W. Mittlefehldt, (2016), The  
1324 degradational history of Endeavour crater, Mars: *Icarus*, doi:10.1016/j.icarus.2015.08.019, in press.

1325 Greeley, R., R.E. Arvidson, 1990. Aeolian processes on Venus. *Earth, Moon, and Planets*, 50/51, 127-157.

1326 Greeley, R., J.D. Iversen, J.B. Pollack, N. Udovich, B. White, 1974. Wind tunnel studies of martian aeolian  
1327 processes. *Proc. Royal Soc. London*, 341, 331-360, doi: 10.1098/rspa.1974.0191.

1328 Greeley, R., R. Leach, B. White, J. Iversen, J. Pollack, 1980. Threshold windspeeds for sand on Mars: Wind  
1329 tunnel simulations. *Geophysical Research Letters*, vol. 7, no. 2, 121-124.

1330 Greeley, R., J. Iversen, R. Leach, J. Marshall, B. White, S. Williams, 1984a. Windblown sand on Venus:  
1331 Preliminary results of laboratory simulations. *Icarus* 57(1), 112-124.

1332 Greeley, R., J.R. Marshall, R.N. Leach, 1984b. Microdunes and other aeolian bedforms on Venus: Wind  
1333 tunnel experiments. *Icarus* 60, 152-160. Greeley R., N. Lancaster, S. Lee, P. Thomas, 1992a. Martian  
1334 aeolian processes, sediments and features, in: Kieffer, H., Jakosky, B.M., Snyder, C.W., Matthews,  
1335 M.S. (Eds.), Mars. University of Arizona Press, Tucson, pp. 730-767.

1336 Greeley, R., R.E. Arvidson, C. Elachi, M.A. Geringer, J.J. Plaut, R.S. Saunders, G. Schubert, E.R. Stofan,  
1337 E.J.P. Thouvenot, S.D. Wall, C.M. Weitz, 1992b. Aeolian features on Venus: Preliminary Magellan  
1338 results. *J. Geophys. Res.* 97(E8), 13319-13345.

1339 Greeley, R. K. Bender, P.E. Thomas, G. Schubert, D. Limonadi, C.M. Weitz, 1995. Wind-related features  
1340 and processes on Venus: Summary of Magellan results. *Icarus*, 115, 399-420.

1341 Greeley, R., K.C. Bender, R.S. Saunders, G. Schubert, C. M. Weitz, 1997. Aeolian processes and features  
1342 on Venus, in *Venus II: Geology, Geophysics, Atmosphere, and Solar Wind Environment*, edited by S.  
1343 W. Bougher, D. M. Hunten, and R. J. Philips, pp. 547- 590, Univ. of Ariz. Press, Tucson.

1344 Greeley, R., R.E. Arvidson, P.W. Barlett, D. Blaney, N.A. Cabrol, P.R. Christensen, R.L. Fergason, M.P.  
1345 Golombek, G.A. Landis, M.T. Lemmon, S.M. McLennan, J.N. Maki, T. Michaels, J.E. Moersch, L.D.V.  
1346 Neakrase, S.C.R. Rafkin, L. Richter, S.W. Squyres, P.A. de Souza Jr., R.J. Sullivan, S.D. Thompson, P.L.  
1347 Whelley, 2006. Gusev crater: Wind-related features and processes observed by the Mars Exploration  
1348 Rover Spirit. *J. Geophys. Res.* 111, E02S09, doi:10.1029/2005JE002491.

1349 Hanna, S.R., 1969. The formation of longitudinal sand dunes by large helical eddies in the atmosphere.  
1350 Journal of Applied Meteorology, 8(6), pp.874-883.

1351 Hansen, C. J., M. Bourke, N.T. Bridges, S. Byrne, C. Colon, S. Diniega, C. Dundas, K. Herkenhoff, A.  
1352 McEwen, M. Mellon, G. Portyankina, N. Thomas, 2011. Seasonal erosion and restoration of Mars'  
1353 northern polar dunes. Science 331(6017), 575-578, doi:10.1126/science.1197636.

1354 Hansen, C.J., S. Diniega, N. Bridges, S. Byrne, C. Dundas, A. McEwen, G. Portyankina, 2015. Agents of  
1355 change on Mars' northern dunes: CO2 ice and wind, Icarus (special issue: Dynamic Mars) 251, 264-  
1356 274. doi:10.1016/j.icarus.2014.11.015.

1357 Hayward, R.K., 2011. Mars Global Digital Dune Database (MGD3): north polar region (MC-1) distribution,  
1358 applications, and volume estimates. Earth Surface Processes and Landforms, doi:10.1002/esp.2219.

1359 Hayward, R.K., K.F. Mullins, L.K. Fenton, T.M. Hare, T.N. Titus, M.C. Bourke, A. Colaprete, P.R.  
1360 Christensen, 2007. Mars global digital dune database and initial science results. J. Geophys. Res.  
1361 Planets 112, E11007, doi:10.1029/2007JE002943.

1362 Hayward, R.K., T.N. Titus, T.I. Michaels, L.K. Fenton, A. Colaprete, P.R. Christensen, 2009. Aeolian dunes  
1363 as ground truth for atmospheric modeling on Mars. J. Geophys. Res., 114, E11012,  
1364 doi:10.1029/2009JE003428.

1365 Hayward, R.K., L.K. Fenton, T.N. Titus, 2014. Mars Global Digital Dune Database (MGD3): Global dune  
1366 distribution and wind pattern observations. Icarus, 230, 38-46, doi:10.1016/j.icarus.2013.04.011.

1367 Hersen, P., 2004. On the crescentic shape of barchan dunes. Eur. Phys. J. B., vol. 37, 507-514,  
1368 doi:10.1140/epjb/e2004-00087-y.

1369 Hesse, P., N. Lancaster, M.W. Telfer, 2015. Digital mapping of the extent of global dune systems.  
1370 Geophysical Research Abstracts, vol. 17, EGU2015-3638.

1371 Horgan, B.H.N., J.F. Bell, 2012. Seasonally active slipface avalanches in the north polar sand sea of Mars:  
1372 Evidence for a wind-related origin. Geophys. Res. Lett. 39, L09201, doi:10.1029/2012GL051329.

1373 Hugenholtz, C.H., T.E. Barchyn, 2010. Spatial analysis of sand dunes with a new global topographic  
1374 dataset: new approaches and opportunities. Earth Surf. Process. Landforms 35, 986-992,  
1375 doi:10.1002/esp.2013.

1376 Jackson, D.W.T., M.C. Bourke, T.A.G. Smyth, 2015. The dune effect on sand-transporting winds on Mars.  
1377 Nature Communications, DOI: 10.1038/ncomms9796.

1378 Jain, M., C.E. Andersen, L. Bøtter-Jensen, A.S. Murray, H. Haack, J.C. Bridges, 2006. Luminescence dating  
1379 on Mars: OSL characteristics of Martian analogue materials and GCR dosimetry. Radiation  
1380 Measurements, 41(7), pp.755-761.

1381 Jerolmack, D.J., R.C. Ewing, F. Falcini, R.L. Martin, C. Masteller, C. Phillips, M.D. Reitz, I. Buynevich, 2012.  
1382 Internal boundary layer model for the evolution of desert dune fields. Nature Geoscience, 5, 3, 1-4,  
1383 DOI: 10.1038/ngeo1381.

1384 JPL, Oct. 4, 2012. NASA Mars Curiosity Rover Prepares to Study Martian Soil, Press Release  
1385 <http://www.jpl.nasa.gov/news/news.php?feature=3540>.

1386 JPL, Feb. 3, 2014. Through the Gap: Curiosity Mars Rover Crosses Dune, Press Release  
1387 <http://www.jpl.nasa.gov/news/news.php?release=2014-035>.

1388 JPL, Dec. 10, 2015. NASA Mars Rover Curiosity Reaches Sand Dunes, Press Release  
1389 <http://www.jpl.nasa.gov/news/news.php?feature=4787>.

1390 Klammer, G., 1982. Die Paleoste des Pantanal von Mato Grosso und die pleistozene Klimageschichte der  
1391 brasilianischen Randtropen. Zeitschrift. der Geomorphologie N.F., vol. 26, 393-416

1392 Kieffer, S.W., R. Lopes-Gautier, A. McEwen, W. Smythe, L. Keszthelyi, R. Carlson, 2000. Prometheus: Io's  
1393 wandering plume. *Science* 288, 1204-1208.

1394 Kocurek, G., R.C. Ewing, 2005. Aeolian dune field self-organization – implications for the formation of  
1395 simple versus complex dune-field patterns. *Geomorphology* 72, 94-105,  
1396 doi:10.1016/j.geomorph.2005.05.005.

1397 Kocurek, G., N. Lancaster, 1999. Aeolian system sediment state: theory and Mojave Desert Kelso dune  
1398 field example. *Sedimentology* 46, 505 - 515.

1399 Kocurek, G., R.C. Ewing, 2012. Source-to-sink: An Earth/Mars comparison of boundary conditions for  
1400 eolian dune systems. *Sedimentary Geology of Mars*, SEPM Special Publication No. 102, Print ISBN  
1401 978-1-56576-312-8, CD/DVD ISBN 978-1-56576-313-5, p. 151–168.

1402 Kocurek, G., M. Carr, R. Ewing, K.G. Havholm, Y.C. Nagar, A.K. Singhvi, 2007. White Sands Dune Field,  
1403 New Mexico: age, dune dynamics and recent accumulations. *Sedimentary Geology*, vol. 197, no. 3,  
1404 313-331.

1405 Kok, J. F., N.O. Renno, 2009. A comprehensive numerical model of steady state saltation (COMSALT). *J.*  
1406 *Geophys. Res.*, 114, D17204, doi:10.1029/2009JD011702.

1407 Kok, J.F., 2010. Difference in the wind speeds required for initiation versus continuation of sand  
1408 transport on Mars: Implications for dunes and dust Storms. *Phys. Rev. Lett.* 104, 074502,  
1409 doi:10.1103/PhysRevLett.104.074502.

1410 Kok J.F., E.J.R. Parteli, T.I. Michaels, D. Bou Karam, 2012. The physics of windblown sand and dust. *Rep.*  
1411 *Prog. Phys.* 75, 106901, doi:10.1088/0034-4885/75/10/106901.

1412 Kreslavsky, M.A., R.V. Vdovichenko, 1999. Anisotropy of Venus Surface Properties Inferred from  
1413 Magellan Side-Looking-Radar Data. *Solar System Research* 33, 110.

1414 Lancaster, N., 1995. *Geomorphology of Desert Dunes* (Routledge Physical Environment Series).  
1415 Routledge. ISBN-13: 978-0415060943.

1416 Lancaster, N., S. Wolfe, D. Thomas, C. Bristow, O. Bubenzer, S. Burrough, G. Duller, A. Halfen, P. Hesse, J.  
1417 Roskin, A. Singhvi, H. Tsoar, A. Tripaldi, X. Yang, M. Zárate, M., in press. The INQUA Dunes Atlas  
1418 chronologic database. *Quaternary International*, doi:10.1016/j.quaint.2015.10.044.

1419 Lapotre, M.G.A., R.C. Ewing, M.P. Lamb, W.W. Fischer, J.P. Grotzinger, D.M. Rubin, K.W. Lewis, M.  
1420 Ballard, M. Day, S. Gupta, S.G. Banham, N.T. Bridges, D.J. Des Marais, A.A. Fraeman, J.A. Grant, K.E.  
1421 Herkenhoff, D.W. Ming, M.A. Mischna, M.S. Rice, D.A. Summer, A.R. Vasavada, R.A. Yingst, 2016a.  
1422 Large wind ripples on Mars: A record of atmospheric evolution. *Science*, 353, 6294, 55-58, DOI:  
1423 10.1126/science.aaf3206.

1424 Lapotre, M.G.A., B.L. Ehlmann, A.A. Fraeman, S.E. Minson, F. Ayoub, R.C. Ewing, R.E. Arvidson, N.T.  
1425 Bridges, MSL Science Team, 2016b. A quantitative assessment of aeolian fractionation at the  
1426 Bagnold dunes of Gale crater, Mars, from orbit to the ground. 47<sup>th</sup> LPSC, Ab. 1513.

1427 Le Gall, A., A.G. Hayes, R. Ewing, M.A. Janssen, J. Radebaugh, C. Savage, P. Encrenaz, 2012. Latitudinal  
1428 and altitudinal controls of Titan's dune field morphometry. *Icarus*, vol. 217, issue 1, 231–242.

1429 Livingstone, I., G.F. Wiggs, C.M. Weaver, 2007. Geomorphology of desert sand dunes: a review of recent  
1430 progress. *Earth-Science Reviews*, 80(3), 239-257.

1431 Livingstone, I., C. Bristow, R.G. Bryant, J. Bullard, K. White, G.F. Wiggs, A.C. Baas, M.D. Bateman, D.S.  
1432 Thomas, 2010. The Namib Sand Sea digital database of aeolian dunes and key forcing variables.  
1433 *Aeolian Research*, 2(2), 93-104.

1434 Lorenz, R.D., 2015. Heights of Fortuna-Meshknet Dunes (Al-Uzza Undae), Venus, from Magellan  
1435 Radarclinometry. Fourth Annual International Planetary Dunes Workshop, abstract # 8004.

1436 Lorenz, R.D., J. Radebaugh, 2009. Global pattern of Titan's dunes: Radar survey from the Cassini prime  
1437 mission. *Geophys. Res. Lett.* 36, L03202, doi:10.1029/2008/GL036850.

1438 Lorenz, R.D., J.I. Lunine, J.A. Grier, M.A. Fisher, 1995. Prediction of aeolian features on planets:  
1439 Application to Titan paleoclimatology. *JGR* 100(E12), 26377-26386.

1440 Lorenz, R.D., S. Wall, J. Radebaugh, G. Boubin, E. Reffet, M. Janssen, E. Stofan, R. Lopes, R. Kirk, C. Elachi,  
1441 J. Lunine, K. Mitchell, F. Paganelli, L. Soderblom, C. Wood, L. Wye, H. Zebker, Y. Anderson, S. Ostro,  
1442 M. Allison, R. Boehmer, P. Callahan, P. Encrenaz, G.G. Ori, G. Francescetti, Y. Gim, G. Hamilton, S.  
1443 Hensley, W. Johnson, K. Kelleher, D. Muhleman, G. Picardi, F. Posa, L. Roth, R. Seu, S. Shaffer, B.  
1444 Stiles, S. Vetrella, E. Flamini, R. West, 2006. The sand seas of Titan: Cassini RADAR observations of  
1445 longitudinal dunes. *Science* 312, 724-727, doi: 10.1126/science.1123257.

1446 Lorenz, R. D., R.M. Lopes, F. Paganelli, J.I. Lunine, R.L. Kirk, K.L. Mitchell, L.A. Soderblom, E.R. Stofan, G.  
1447 Ori, M. Myers, H. Miyamoto, J. Radebaugh, B. Stiles, S.D. Wall, C.A. Wood and the Cassini RADAR  
1448 Team, 2008. Fluvial channels on Titan: Initial Cassini RADAR observations. *Planet. Space Sci.* 56,  
1449 1132-1144.

1450 Lucas, A., S. Rodriguez, C. Nartean, B. Charnay, S.C. Pont, T. Tokano, A. Garcia, M. Thiriet, A.G. Hayes,  
1451 R.D. Lorenz, O. Aharonson, 2014. Growth mechanisms and dune orientation on Titan.  
1452 *Geophysical Research Letters*, 41(17), 6093-6100.

1453 Madigan, C.T., 1936. The Australian sand-ridge deserts. *Geographical Review*, 26(2), 205-227.

1454 Malaska, M.J., R.M. Lopes, A.G. Hayes, J. Radebaugh, R.D. Lorenz and E.P. Turtle, 2016. Material  
1455 transport map of Titan: The fate of dunes. *Icarus*, 270, 183-196,  
1456 doi:10.1016/j.icarus.2015.09.029.

1457 Marsh, G.P., 1864. *Man and Nature*. (Reprint, 1965) Cambridge, MA: Harvard University Press.

1458 Marshall, J.R., R. Greeley, 1992. An experimental study of aeolian structures on Venus. *J. Geophys. Res.*  
1459 97(E1), 1007-1016.

1460 May, J.H., 2013. Dunes and dune fields in the Bolivian Chaco as potential records of environmental  
1461 change. *Aeolian Research*, vol. 10, 89-102.

1462 McCauley, J.F., M.H. Carr, J.A. Cutts, W.K. Hartmann, H. Masursky, D.J. Milton, R.P. Sharp, D.E.  
1463 Wilhelms, 1972. Preliminary Mariner 9 Report on the Geology of Mars, *Icarus* 17(2), 289-327,  
1464 doi:10.1016/0019-1035(72)90003-6.

1465 McEwen, A.S., M.J.S. Belton, H.H. Breneman, S.A. Fagents, P. Geissler, R. Greeley, J.W. Head, G. Hoppa,  
1466 W.L. Jaeger, T.V. Johnson, L. Keszthelyi, 2000. Galileo at Io: Results from high-resolution imaging.  
1467 *Science*, vol. 288, no. 5469, 1193-1198.

1468 McEwen, A.S., E.M. Eliason, J.W. Bergstrom, N.T. Bridges, C.J. Hansen, W.A. Delamere, J.A. Grant, V.C.  
1469 Gulick, K.E. Herkenhoff, L. Keszthelyi, R.L. Kirk, 2007. Mars Reconnaissance Orbiter's High Resolution  
1470 Imaging Science Experiment (HiRISE). *J. Geophys. Res.*, 112, issue E5, doi:10.1029/2005JE002605.



1471 McKeever, S.W.S., D. Banerjee, M. Blair, S.M. Clifford, M.S. Cloudsley, S.S. Kim, M. Lamothe, K. Lepper,  
1472 M. Leuschen, K.J. McKeever, M. Prather, 2003. Concepts and approaches to in situ luminescence  
1473 dating of martian sediments. *Radiation Measurements*, 37(4), pp.527-534.

1474 Mellon, M.T., B.M. Jakovsky, H.H. Kieffer, P.R. Christensen, 2000. High-resolution thermal inertia  
1475 mapping from the Mars Global Surveyor Thermal Emission Spectrometer. *Icarus*, 148, 437-455,  
1476 doi:10.1006/icar.2000.6503.

1477 Milazzo, M.P., L.P. Keszthelyi, A.S. McEwen, 2001. Observations and initial modeling of lava-SO<sub>2</sub>  
1478 interactions at Prometheus, Io. *J. Geophys. Res.* 106, 33,121-33,127.

1479 Moore, J.M., A.D. Howard, P.M. Schenk, W.B. McKinnon, R.T. Pappalardo, R.C. Ewing, E.B. Bierhaus, V.J.  
1480 Bray, J.R. Spencer, R.P. Binzel, B. Buratti, 2015. Geology before Pluto: Pre-encounter considerations.  
1481 *Icarus*, 246, 65-81, doi:10.1016/j.icarus.2014.04.028.

1482 Moore, J.M., W.B. McKinnon, J.R. Spencer, A.D. Howard, P.M. Schenk, R.A. Beyer, F. Nimmo, K.N. Singer,  
1483 O.M. Umurhan, O.L. White, S.A. Stern, K. Ennico, C.B. Olkin, H.A. Weaver, L.A. Young, R.P. Binzel,  
1484 M.W. Buie, B.J. Buratti, A.F. Cheng, D.P. Cruikshank, W.M. Grundy, I.R. Linscott, H.J. Reitsema, D.C.  
1485 Reuter, M.R. Showalter, V.J. Bray, C.L. Chavez, C.J.A. Howett, T.R. Lauer, C.M. Lisse, A.H. Parker, S.B.  
1486 Porter, S.J. Robbins, K. Runyon, T. Stryk, H.B. Throop, C.C.C. Tsang, A.J. Verbiscer, A.M. Zangari, A.L.  
1487 Chaikin, D.E. Wilhelms, New Horizons Science Team, 2016. The geology of Pluto and Charon through  
1488 the eyes of New Horizons. *Science* v. 351, Is. 6279, 1284-93. doi:10.1126/science.aad7055.

1489 Mottola, S., Arnold, G., Grothues, H.-G., Jaumann, R., Michaelis, H., Neukum, G., Bibring, J.-P., Schröder,  
1490 S.E., Hamm, M., Otto, K.A., Pelivan, I., Proffe, G., Scholten, F., Tirsch, D., Kreslavsky, M., Remetean,  
1491 E., Souvannavong, F., Dolives, B., 2015. The structure of the regolith on 67P/Churyumov-  
1492 Gerasimenko from ROLIS descent imaging. *Science* 349, 10.1126/science.aab0232.

1493 Neakrase, L.D.V. (2015), Terrestrial Analogues for Venusian dunes: Sub-aqueous, seafloor dune fields.  
1494 Fourth International Planetary Dunes Workshop. Ab. 8023.

1495 New Horizons, 2015. [http://www.nasa.gov/mission\\_pages/newhorizons/images/index.html](http://www.nasa.gov/mission_pages/newhorizons/images/index.html).

1496 New Horizons, 2016. The Tartarus Dorsa Mountains Rise Up Along Pluto, [http://www.nasa.gov/image-](http://www.nasa.gov/image-feature/the-tartarus-dorsa-mountains-rise-up-along-pluto)  
1497 [feature/the-tartarus-dorsa-mountains-rise-up-along-pluto](http://www.nasa.gov/image-feature/the-tartarus-dorsa-mountains-rise-up-along-pluto).

1498 NRC, 2013. Vision and Voyages for Planetary Science in the Decade 2013-2022. Steering Group Chair:  
1499 Squyres, National Academies Press, Washington, D.C., ISBN: 0-309-20955-2.

1500 O'Connell-Cooper, C.D., L.M. Thompson, J.G. Spray, J.A. Berger, E.D. Desouza, M. Schmidt, A. Yen, N.I.  
1501 Boyd, R. Gellert, G. Perrett, S.J. VanBommel, the MSL-APXS Team, 2016. Preliminary comparison of  
1502 soils within Gale crater to those from Gusev Crater and Meridiani Planum. 47<sup>th</sup> LPSC, Ab. 2477.

1503 Parteli, E.J.R., O. Durán, H.J. Herrmann, 2007. Minimal size of a barchan dune. *Phys. Rev. E* 75, 011301,  
1504 doi:10.1103/PhysRevE.75.011301.

1505 Parteli, E.J.R., O. Durán, H. Tsoar, V. Schwämmle, H.J. Herrmann, 2009. Dune formation under bimodal  
1506 winds. *PNAS* 106(52), 22085-22089, doi:10.1073/pnas.0808646106.

1507 Pelletier, J.D., 2009. Controls on the height and spacing of eolian ripples and transverse dunes: A  
1508 numerical modeling investigation. *Geomorphology*, v. 105, 322-333,  
1509 doi:10.1016/j.geomorph.2008.10.010.

1510 Pelletier, J.D., 2015. Controls on the large-scale spatial variations of dune field properties in the  
1511 barchanoid portion of White Sands dune field, New Mexico. *J. Geophys. Res. Earth Surf.*, 120, 453-  
1512 473, doi:10.1002/2014JF003314.

1513 Portyankina, G., A. Pommerol, K.-M. Aye, C. J. Hansen, N. Thomas, 2012. Polygonal cracks in the  
1514 seasonal semi-translucent CO<sub>2</sub> ice layer in Martian polar areas. *J. Geophys. Res.*, 117(E2006),  
1515 doi:10.1029/2011JE003917.

1516 Radebaugh, J., 2013. Dunes on Saturn's moon Titan at the end of the Cassini Equinox Mission. *Aeolian*  
1517 *Research* 11, 23-41.

1518 Radebaugh, J., R. Lorenz, J. Lunine, S. Wall, G. Boubin, E. Reffet, R. Kirk, R. Lopes, E. Stofan, L. Soderblom,  
1519 M. Allison, M. Janssen, P. Paillou, P. Callahan and the Cassini Radar Team, 2008. Dunes on Titan  
1520 observed by Cassini Radar. *Icarus* 194, 690-703, doi:10.1016/j.icarus.2007.10.015.

1521 Radebaugh, J., R. Lorenz, T. Farr, P. Paillou, C. Savage, C. Spencer, 2010. Linear dunes on Titan and earth:  
1522 Initial remote sensing comparisons. *Geomorphology* 121, 122-132,  
1523 doi:10.1016/j.geomorph.2009.02.022.

1524 Radebaugh, J., R.D. Lorenz, S.D. Wall, R.L. Kirk, C.A. Wood, J.I. Lunine, E.R. Stofan, R.M.C. Lopes, P.  
1525 Valora, T.G. Farr, A.G. Hayes, B. Stiles, G. Mitri, H. Zebker, M. Janssen, L. Wye, A. Le Gall, K.L.  
1526 Mitchell, F. Paganelli, R.D. West, E.L. Schaller and the Cassini RADAR Team, 2011. Regional  
1527 geomorphology and history of Titan's Xanadu province. *Icarus* 211, 672-685.

1528 Radebaugh, J., D. Ventra, R.D. Lorenz, T. Farr, R. Kirk, A. Hayes, M.J. Malaska, S. Birch, Z. Y-C. Liu, J.  
1529 Lunine, J. Barnes, A. Le Gall, R. Lopes, E. Stofan, S. Wall, P. Paillou, 2016. Alluvial and fluvial fans on  
1530 Saturn's moon Titan reveal processes, materials and regional geology, in, Ventra, D. & Clarke, L. E.  
1531 (eds) *Geology and Geomorphology of Alluvial and Fluvial Fans: Terrestrial and Planetary*  
1532 *Perspectives*. Geological Society, London, Special Publications, 440, doi: 10.1144/SP440.6.

1533 Rubin, D.M., 2012. A unifying model for planform straightness of ripples and dunes in air and water.  
1534 *Earth-Science Reviews*, 113(3), 176-185, doi:10.1016/j.earscirev.2012.03.010.

1535 Rubin, D.M., P.A. Hesp, 2009. Multiple origins of linear dunes on Earth and Titan. *Nature Geoscience*,  
1536 vol. 2, 653-658, doi:10.1038/ngeo610.

1537 Rubin, D.M., R.E. Hunter, 1987. Bedform alignment in directionally varying flows. *Science*, 237(4812),  
1538 276-8, doi:10.1126/science.237.4812.276.

1539 Rubin, D. M., H. Ikeda, 1990. Flume experiments on the alignment of transverse, oblique, and  
1540 longitudinal dunes in directionally varying flows. *Sedimentology*, 37, 673-684.

1541 Sagan, C., J. Veverka, P. Fox, R. Dubisch, J. Lederberg, E. Levinthal, L. Quam, R. Tucker, J.B. Pollack, B.A.  
1542 Smith, 1972. Variable features on Mars: Preliminary Mariner 9 television results. *Icarus* 17(2), 346-  
1543 372, doi:10.1016/0019-1035(72)90005-X.

1544 Sauermann, G., K. Kroy, H.J. Herrmann, 2001. Continuum saltation model for sand dunes, *Physical*  
1545 *Review E*, v. 64, 031305, doi:10.1103/PhysRevE.64.031305.

1546 Savage, C.J., J. Radebaugh, E.H. Christiansen, R.D. Lorenz, 2014. Implications of dune pattern analysis for  
1547 Titan's surface history. *Icarus* 230, 180-190.

1548 Silvestro, S., G. Di Achille, G.G. Ori, 2010a. Dune morphology, sand transport pathways and possible  
1549 source areas in east Thaumasia Region (Mars), *Geomorphology*, vol. 121, 84-97,  
1550 doi:10.1016/j.geomorph.2009.07.019.

1551 Silvestro, S., L.K. Fenton, D.A. Vaz, N.T. Bridges, G.G. Ori, 2010b. Ripple migration and dune activity on  
1552 Mars: Evidence for dynamic processes. *Geophys. Res. Lett.* 37, L20203 (2010).

1553 Silvestro, S., D.A. Vaz, L.K. Fenton and P.E. Geissler, 2011. Active Aeolian processes on Mars: a regional  
1554 study in Arabia and Meridiani Terrae, *Geophysical Research Letters*, 38, L20201,  
1555 doi:10.1029/2011GL048955.

1556 Silvestro S., L.K. Fenton, T.I. Michaels, A. Valdez, G.G. Ori, 2012. Interpretation of the complex dune  
1557 morphology on Mars: dune activity, modeling and a terrestrial analogue. *Earth Surface Processes*  
1558 *and Landforms*, 37, 13, 1424-1436, DOI: 10.1002/esp.3286.

1559 Silvestro, S., D.A. Vaz, R.C. Ewing, A.P. Rossi, L.K. Fenton, T.I. Michaels, J. Flahaut and P.E.  
1560 Geissler, 2013. Pervasive aeolian activity along Curiosity's traverse in Gale Crater, Mars.  
1561 *Geology*, 41 (4), 483-486, doi:10.1130/G34162.1.

1562 Silvestro S., D.A. Vaz, H. Yizhaq, F. Esposito, 2016. Dune-like dynamic of martian aeolian large ripples.  
1563 *Geophysical Research Letters*, doi:10.1002/2016GL070014.

1564 Singhvi, A.K., N. Porat, 2008. Impact of luminescence dating on geomorphological and palaeoclimate  
1565 research in drylands. *Boreas*, 37(4), 536-558.

1566 Soderblom, L., J. Anderson, K. Baines, J. Barnes, J. Barrett, R. Brown, B. Buratti, R. Clark, D. Cruikshank, C.  
1567 Elachi, M. Janssen, R. Jaumann, R. Kirk, E. Karkoschka, S. Lemouelic, R. Lopes, R. Lorenz, J. Lunine, T.  
1568 McCord, P. Nicholson, J. Radebaugh, B. Rizk, C. Sotin, E. Stofan, T. Sucharski, M. Tomasko, S. Wall,  
1569 2007. Correlations between Cassini VIMS spectra and RADAR SAR images: Implications for Titan's  
1570 surface composition and the character of the Huygens Probe landing site. *Planetary and Space*  
1571 *Science*, 55, 2025-2036.

1572 Sohbaty, R., A.S. Murray, J.P. Buylaert, M. Ortuño, P.P. Cunha, E. Masana, 2012. Luminescence dating of  
1573 Pleistocene alluvial sediments affected by the Alhama de Murcia fault (eastern Betics, Spain)-a  
1574 comparison between OSL, IRSL and post-IR IRSL ages. *Boreas*, 41(2), 250-262, doi:10.1111/j.1502-  
1575 3885.2011.00230.x

1576 Sokolów, N. A., 1894. *Die Dunen; Bildung, Entwicklung, and innerer Bau* (German transl. from Russian  
1577 4th ed., 1884, assisted by A. Arzruni). Berlin, Springer.

1578 Sullivan, R., R. Arvidson, J.F. Bell III, R. Gellert, M. Golombek, R. Greeley, K. Herkenhoff, J. Johnson, S.  
1579 Thompson, P. Whelley, J. Wray, 2008. Wind-driven particle mobility on Mars: Insights from Mars  
1580 Exploration Rover observations at "El Dorado" and surroundings at Gusev Crater. *J. Geophys. Res.*,  
1581 113, E06S07, doi:10.1029/2008JE003101.

1582 Stern and the New Horizons team, 2016. *New Horizons: The Exploration of the Pluto System and the*  
1583 *Kuiper Belt beyond*. 47<sup>th</sup> LPSC, Ab. 1317.

1584 Sullivan, R., D. Banfield, J. F. Bell III, W. Calvin, D. Fike, M. Golombek, R. Greeley, J. Grotzinger, K.  
1585 Herkenhoff, D. Jerolmack, M. Malin, D. Ming, L. A. Soderblom, S. W. Squyres, S. Thompson, W.A.  
1586 Watters, C.M. Weitz, A. Yen, 2005. Aeolian processes at the Mars Exploration Rover Meridiani  
1587 Planum landing site. *Nature* 436, 58-61, doi:10.1038/nature03641.

1588 Tanaka, K.L., Rodriguezb, J.A.P., Skinner Jr., J.A., Bourke, M.C., Fortezzo, C.M., Herkenhoff, K.E., Kolb, E.J.,  
1589 Okubo, C.H., 2008. North polar region of Mars: Advances in stratigraphy, structure, and erosional  
1590 modification. *Icarus* 192, 318-358, 10.1016/j.icarus.2008.01.021.

1591 Teeuw, R.M., E.J. Rhodes, 2004. Aeolian activity in northern Amazonia: optical dating of Late Pleistocene  
1592 and Holocene palaeodunes. *Journal of Quaternary Science*, 19(1), pp.49-54.

1593 Telfer, M.W., 2011. Growth by extension, and reworking, of a south-western Kalahari linear dune. *Earth*  
1594 *Surface Processes and Landforms*, 36(8), pp.1125-1135.

1595 Telfer, M.W. and Thomas, D.S.G., 2007. Late Quaternary linear dune accumulation and  
1596 chronostratigraphy of the southwestern Kalahari: implications for aeolian palaeoclimatic  
1597 reconstructions and predictions of future dynamics. *Quaternary Science Reviews*, 26(19), pp.2617-  
1598 2630.

1599 Tirsch, D., R. Jaumann, A. Pacifici, F. Poulet, 2011. Dark aeolian sediments in Martian craters:  
1600 Composition and sources. *J Geophys Res-Planet* 116, doi: 10.1029/2009je003562.

1601 Thomas, P., 1981. North-south asymmetry of eolian features in martian polar regions: Analysis based on  
1602 crater-related wind markers. *Icarus*, 48, 76-90.

1603 Thomas, P., 1982. Present wind activity on Mars - Relation to large latitudinally zoned sediment  
1604 deposits. *J. Geophys. Res.*, 87(B12), 9999-10,008, doi:10.1029/JB087iB12p09999.

1605 Thomas, N., H. Sierks, C. Barbieri, P.L. Lamy, R. Rodrigo, H. Rickman, D. Koschny, H.U. Keller, J. Agarwal,  
1606 M.F. A'Hearn, F. Angrilli, A.-T. Auger, M.A. Barucci, J.-L. Bertaux, I. Bertini, S. Besse, D. Bodewits, G.  
1607 Cremonese, V. Da Deppo, B. Davidsson, M. De Cecco, S. Debei, S., M.R. El-Maarry, F. Ferri, S.  
1608 Fornasier, M. Fulle, L. Giacomini, O. Groussin, P.J. Gutierrez, C. Güttler, S.F. Hviid, W.-H. Ip, L. Jorda,  
1609 J. Knollenberg, J.-R. Kramm, E. Kührt, M. Küppers, F. La Forgia, L.M. Lara, M. Lazzarin, J.J.L. Moreno,  
1610 S. Magrin, S. Marchi, F. Marzari, M. Massironi, H. Michalik, R. Moissl, S. Mottola, G. Naletto, N.  
1611 Ookay, M. Pajola, A. Pommerol, F. Preusker, L. Sabau, F. Scholten, C. Snodgrass, C. Tubiana, J.-B.  
1612 Vincent, K.-P. Wenzel, 2015a. The morphological diversity of comet 67P/Churyumov-Gerasimenko.  
1613 *Science* 347, doi:10.1126/science.aaa0440.

1614 Thomas, N., B. Davidsson, M.R. El-Maarry, S. Fornasier, L. Giacomini, A.G. Gracia-Berná, S.F. Hviid, W.-H.  
1615 Ip, L. Jorda, H.U. Keller, J. Knollenberg, E. Kührt, F. La Forgia, I.L. Lai, Y. Liao, R. Marschall, M.  
1616 Massironi, S. Mottola, M. Pajola, O. Poch, A. Pommerol, F. Preusker, F. Scholten, C.C. Su, J.S. Wu, J.-  
1617 B. Vincent, H. Sierks, C. Barbieri, P.L. Lamy, R. Rodrigo, D. Koschny, H. Rickman, M.F. A'Hearn, M.A.  
1618 Barucci, J.-L. Bertaux, I. Bertini, G. Cremonese, V. Da Deppo, S. Debei, M. de Cecco, M. Fulle, O.  
1619 Groussin, P.J. Gutierrez, J.-R. Kramm, M. Küppers, L.M. Lara, M. Lazzarin, J.J. Lopez Moreno, F.  
1620 Marzari, H. Michalik, G. Naletto, J. Agarwal, C. Güttler, N. Ookay, C. Tubiana, 2015b. Redistribution of  
1621 particles across the nucleus of comet 67P/Churyumov-Gerasimenko. *Astronomy and Astrophysics*,  
1622 583, A17, doi:10.1051/0004-6361/201526049.

1623 Treiman, A.H., 2003. Geologic settings of Martian gullies: Implications for their origins. *J. Geophys. Res.*  
1624 108 (E4), 8031, doi:10.1029/2002JE001900.

1625 Tricart, J., 1974. Existence de periodes seches au Quaternaire en Amazonie et dans les regions voisines.  
1626 *Révue de Géomorphologie Dynamique* 23, 145-158

1627 Trilling, D.E., 2016. The surface age of Sputnik Planum, Pluto, must be less than 10 million years. *PLOS*  
1628 *ONE*, doi:10.1371/journal.pone.0147386.

1629 Tripaldi, A., M.A. Zárata, in press. A review of Late Quaternary inland dune systems of South America  
1630 east of the Andes. *Quaternary International*, available online 24 July 2014,  
1631 doi:10.1016/j.quaint.2014.06.069.

1632 Tokano, T., 2008. Dune-forming winds on Titan and the influence of topography. *Icarus* 194, 243-262,  
1633 doi:10.1016/j.icarus.2007.10.007.

1634 Tsoar, H., R. Greeley, A.R. Peterfreund, 1979. Mars: The north polar sand seas and related wind patterns.  
1635 *J. Geophys. Res.*, vol. 84, no. B14, 8167-8180.

1636 Tsoar, H., 1983. Dynamic processes acting on a longitudinal (seif) dune. *Sedimentology* 30, 567-578.

- 1637 Vaz, D.A., S. Silvestro, 2014. Mapping and characterization of small-scale aeolian structures on Mars: An  
1638 example from the MSL landing site in Gale Crater. *Icarus*, vol. 230, 151-161.
- 1639 Vaz, D.A., S. Silvestro, P.T.K. Sarmiento, M. Cardinale, submitted. Migrating meter-scale bedforms on  
1640 Martian dark dunes: are terrestrial aeolian ripples good analogs? To *Aeolian Research* (this issue).
- 1641 Ward, A.W., 1979. Yardangs on Mars: Evidence of recent wind erosion. *J. Geophys. Res* 84, 8147-8165.
- 1642 Ward, A.W., K.B. Doyle, P.J. Helm, M.K. Weisman, N.E. Witbeck, 1985. Global map of eolian features on  
1643 Mars. *J. Geophys. Res.*, 90(B2), 2038-2056.
- 1644 Weitz, C.M., J.J. Plaut, R. Greeley, R.S. Saunders, 1994. Dunes and microdunes on Venus: Why were so  
1645 few found in the Magellan data? *Icarus* 112, 282-295.
- 1646 Werner, B.T., 1995. Eolian dunes: Computer simulations and attractor interpretation. *Geology* 23(12),  
1647 1107-1110.
- 1648 Williams, S.H., R. Greeley, 1994. Windblown sand on Venus: The effect of high atmospheric density.  
1649 *Geophys. Res. Lett.* 21(25), 2825-2828.
- 1650 Wilson, L., J.W. Head, 1994. Mars: Review and analysis of volcanic eruption theory and relationships to  
1651 observed landforms. *Reviews of Geophysics*, vol. 32, no. 3, 221-263.
- 1652 Zimbelman, J.R., 2000. Non-active dunes in the Acheron Fossae Region of Mars between the Viking and  
1653 Mars Global Surveyor eras. *Geophys. Res. Lett.* 27, 1069-1072, doi:10.1029/1999GL008399.

1 | **Title: Our evolving understanding of aeolian bedforms, based on observation of dunes studies on different worlds**

2 | © 2016. All rights reserved

3 | **Abstract (253 words)**

4 |  
5 |  
6 | Dunes, dune fields, and smaller aeolian bedforms ripples are unique and useful records of the interaction  
7 | between wind and granular materials – finding a dune field such features on a planetary surface  
8 | immediately suggests certain information about climate and surface conditions (at least during the  
9 | dunes’ formation and evolution). Additionally, studies of dune characteristics under non-Earth  
10 | conditions allow for “tests” of aeolian process models based primarily on observations of terrestrial  
11 | features and dynamics, and refinement of the models to include consideration of a wider range of  
12 | environmental and planetary conditions. To-date, the planetary aeolian community has found and  
13 | studied dune fields on Mars, Venus, and the Saturnian moon Titan. Additionally, we have observed  
14 | candidate “aeolian bedforms” on Comet 67P/Churyumov-Gerasimenko, the Jovian moon Io, and – most  
15 | recently -- Pluto. In this paper, we hypothesize that the progression of investigations of aeolian  
16 | bedforms and processes on a particular planetary body follows a consistent sequence – primarily set by  
17 | the acquisition of data of particular types and resolutions, and by the maturation of knowledge about  
18 | that planetary body. We define that sequence of generated knowledge and new questions (within seven  
19 | investigation phases) and discuss examples from all of the studied bodies. The —with an aim of such a  
20 | sequence is to better define our past and current state of understanding about the aeolian bedforms of  
21 | a particular body, to highlight the related assumptions that require re-analysis with data acquired during  
22 | later investigations, and to use lessons learned from planetary and terrestrial aeolian studies to predict  
23 | what types of investigations could be most fruitful in the future.

24 |  
25 | **Keywords (at least 6):** Planetary; terrestrial; aeolian bedforms; aeolian science; dunes; ripples

27 **Highlights (max 85 characters, including spaces)**

- 28 • Planetary dunes yield useful/unique information about climate & surface conditions.
- 29 • Aeolian bedform studies progress logically in questions/assumptions/new knowledge.
- 30 • Considering this progression exposes gaps/assumptions to be reviewed with new data.
- 31 • Comparinge planetary progression with Earth aeolian studies ~~to~~ yields lessons for each.

32

33 **1. Introduction**

34 Over the past couple of centuries, explorers and scientists of terrestrial dune fields have shown  
35 that the interaction between wind and granular material results in regular geometries and rates of  
36 evolution. Field observations and studies have inspired modeling and experimental works that have  
37 aided in the interpretation of measurable ripples and dunes as proxy indicators of wind speed and  
38 direction(s), grain sizes and sources, and underlying topography. The study of such landforms has been  
39 greatly extended and advanced by observation of analogous features on other planetary bodies. The  
40 comparison of these extraterrestrial features with aeolian process models has increased our  
41 understanding of aeolian bedform evolution in both directions – observations of (potential) aeolian  
42 bedforms generate investigations into the wind regime and granulometrics of surface materials on a  
43 planetary body, and also enable refinement of bedform evolution models as hypotheses about  
44 dominant effects are “tested” outside of Earth-conditions.

45 In this paper, we will review how our understanding (or assumed understanding) of ~~the aeolian~~  
46 ~~bedforms~~sand dunes and/or ripples on a planetary body, and the information those aeolian bedforms  
47 yield about planetary conditions and processes, has progressed on different bodies. We hypothesize  
48 that the progression of investigations of these types of aeolian bedforms on a particular planetary body  
49 follows a consistent sequence – primarily set by the acquisition of data of particular types and  
50 resolutions, and by the maturation of knowledge about that planetary body. Our aim is to define that

51 | progression so we can better constrain our ~~understanding level of knowledge about~~of the aeolian  
52 | bedforms of a particular body, highlight the gaps in our knowledge (i.e., our assumptions), and predict  
53 | what type of future investigations could be most useful in addressing new questions and/or enabling  
54 | improvement over an assumption.

55 | In the interests of space and focus, most discussion (and cited literature) will focus on dunes and  
56 | dune fields – i.e., the larger aeolian bedforms and thus usually the first seen on a planetary body. We do  
57 | delve into ripples (and martian mega-ripples) in portions of the discussion, but primarily as features seen  
58 | on dunes and that complement dune analysis; and we acknowledge that far more could be said about  
59 | these smaller-scale bedforms and that studies of these bedforms on Mars has contributed much more  
60 | towards our understanding of the aeolian environment and processes than is presented here. We also  
61 | do not generally discuss other types of aeolian bedforms within this paper. In particular, we do not  
62 | include discussion of Transverse Aeolian Ridges (TARs) on Mars as there is still much debate about their  
63 | formation mechanism (perhaps as they are an example of a feature that does not have a good terrestrial  
64 | analog?). It is likely, however, that one could trace advancements in our understanding of TARs or other  
65 | aeolian features along a similar progression of ideas as is presented here for dunes, as all of these  
66 | features are studied via similar observation types and their dynamics and morphologies tie into similar  
67 | questions about atmospheric and surface conditions. We also consider only the observation and analysis  
68 | of ~~aeolian~~ bedforms on the surface of a planet, not e.g., evidence of past bedforms recorded within  
69 | sandstone stratigraphy (and thus, while dunes and ripples can also form due to the flow of other fluids,  
70 | such as water, in this paper we focus on aeolian dunes). Thirdly, in discussing our evolution in thinking  
71 | about aeolian bedforms and processes on other planets, we focus on observation-driven science  
72 | advancements; we touch on but do not delve as deeply into the parallel lines of investigation focused on  
73 | model development and validation, empirical studies, and (analog) terrestrial field work – investigations  
74 | that feed into advancements within (and between) the Phases that we outline here. Finally, we



75 | recognize that we present only a sampling of relevant studies – we aimed for enough to map out  
76 | advances in understanding, to justify our proposed framework, and to provide a starting ground for any  
77 | reader interested in learning more on a more specific topic.

78 |         In defining the “progression” of understanding (Section 2), we focus on Mars, Venus, and the  
79 | Saturnian moon Titan – all planetary bodies where aeolian bedforms have primarily been explored with  
80 | remotely acquired data. We also will comment on recently discovered candidate “aeolian bedforms” on  
81 | Comet 67P/Churyumov-Gerasimenko and possible dune-like landforms on Io and Pluto. Within each  
82 | phase of investigation (Subsections 2.1-7), we aim to identify the type of observations generally needed  
83 | and connect these to the primary knowledge, assumptions, and questions that result, and then lead into  
84 | future investigations (summarized at the end of each section). Furthermore, we identify the typical  
85 | investigations (outside of direct studies of the aeolian bedforms) that follow each gain in knowledge, to  
86 | show how aeolian bedform studies contribute to the larger study of that planetary body.

87 |         Our proposed framework of phases, regarding investigation of aeolian bedforms on a planetary  
88 | body, is summarized in Table 1. We again note that this framework is not meant to be fully  
89 | comprehensive for aeolian bedform-dune studies ~~—we focus here on investigations relying on remote~~  
90 | ~~observations (exception: Phase 7) and on planetary studies, not the (parallel) development and~~  
91 | ~~validation of the models and terrestrial in-situ studies that often feed information into those studies.~~ We  
92 | also note that progression in investigations and understanding is not necessarily linear/sequential – for  
93 | example, planetary missions are generally focused on objectives other than aeolian bedform  
94 | investigations, so observation types can be acquired in a “mixed” order. Additionally, the advent of new  
95 | missions, methods, or models can lead to renewed activity within “lower” phases along with advances  
96 | into higher phases. Science questions also often end up circling back as an assumption becomes  
97 | superseded by new information and derived products and assumptions must be re-thought. Thus, in  
98 | addition to identifying typical assumptions associated with each Phase~~s~~, we use that framework to

99 identify some example areas of knowledge gaps or the types of typical assumptions and results that  
100 need re-evaluation when new data becomes available (Section 3.1).

101 | We also ~~then~~ discuss how modeling (Section 3.2) and terrestrial studies (Section 3.3) relate to  
102 planetary aeolian studies. In particular, we highlight lessons learned regarding our understanding of  
103 aeolian processes and their drivers, as well as in methodologies employed. These lessons translate (or  
104 could translate) to improved results within other areas of aeolian science.

**Table 1. Summary of the investigation phases**

Phase of aeolian bedform study on a planetary body	Area of interest	<u>“Unit”/Characteristic (s)/Feature(s) of interest</u>	Data needed to move to this phase (from an earlier phase)	Complementary science investigations
1 <b>Recognition of dune(s)</b>	Dune (possibly a dune field)	Dune <u>morphology</u> ( <del>and i.e.</del> , recognizable, distinctive gross dune shape or crestline patterns within a field)	Images (visible, radar, spectral, etc.) with sufficient resolution to distinguish dune from non-dune surroundings, e.g., linear/arcuate and isolated/repeated morphology	Knowledge of and about analog features (terrestrial or planetary)
2 <b>Analysis of gross individual dune characteristics: e.g., morphology and composition</b>	Dune	Dune morphology, <u>characteristics of surface materials</u>	Images (visible, radar, spectral, etc.) with sufficient resolution to identify/ <u>correlate with</u> dune margin and/or crestline patterns (possibly same data as Phase 1)	Global/regional-scale climate models (specifically: wind velocity, direction, and variation); Dune formation models
3 <b><u>Pattern Analysis of the dunes within a field</u></b> , including variations due to e.g., sediment supply and wind variations	Dune field	Dune <u>shapes</u> throughout the field	Coverage (of images, see above) throughout dune field	Regional/local-scale climate models (specifically: wind); Regional/local-scale topography; Studies of non-aeolian dune-modifying processes (e.g., polar or surface crust forming processes); <u>Maps of other aeolian features around the dune field</u>
4 <b><u>Regional and global surveys and aggregate-analysis of dune characteristics;</u></b> <del>with a re-aggregation of data for</del> e.g., estimates of age or sand volumes, identification of large-scale sediment transport pathways, or identification/	<u>Regional or Global (i.e., multiple dune fields)</u>	Dune field characteristics (including morphology <u>of field and dunes within each field</u> ) and spatial distribution	<u>At least regional Global</u> coverage of images or (often coarser and/or less definitive) proxy data (e.g., thermal inertia)	Expansive composition maps for identification of potential sand sources; <u>Maps of other aeolian features; Global or regional-scale climate models</u>

	estimation of the effect of location-related non-aeolian processes				
5	<b>Analysis of <u>superposed bedforms on the dune</u> “<del>details</del>” (such as ripples) formed due to wind interaction with the dune</b>	Dune	Within/on-dune features (e.g., ripples)	(Very) high-resolution images	Ripple formation models; High-resolution topography (dune); Very high resolution climate model (CFD) (specifically: wind)
6	<b>Observation of activity</b> on the dune, including non-aeolian activity	Dune	Smaller-scale evidence of change (e.g., ripple crestlines, dune margins)	Repeat images for seeing planform changes (e.g., movement of material); these images need sufficient resolution and temporal baseline for changes to be observable	Ripple and dune migration models
7	<b>Groundtruth data</b>	Dune	All features and components of the dune, especially sand size/composition	In situ observations of the dune, sampling and analysis	In situ observations of potential sediment sources for fluvio-sedimentary landforms

95 **2. The Phases of Investigation**

96 **2.1. Phase 1: Recognition of dune(s)**

97 In this first phase of aeolian studies, we focus on the occurrence of the *first observation of a*  
98 *dune* (or, more likely, a dune field). This has geologic significance as aeolian bedforms – dunes and  
99 ripples – directly record an interaction between the atmosphere and surface: specifically, the movement  
100 of granular material due to wind<sup>4</sup>. Furthermore, a dune or ripple is more than a pile of sand – it is a  
101 distinctive landform that requires certain conditions to organize, and that appears to evolve and move  
102 “as a unit” through the aggregation of the actual movement of individual grains of sand, onto and off of  
103 the dune. Specifically, the existence of an aeolian bedform implies:

- 104 • A sufficient supply of ~~sand~~ saltatable (sand) grains,
- 105 • A wind of sufficient velocity and consistency to move those grains, and
- 106 • A period of consistent wind blowing, long enough for the bedform to form and evolve.

107 (We now examine what each of those underlined terms imply about the planetary body’s environment,  
108 focusing on the larger-scale dunes that are typically observed first. A more technical summary of the  
109 conceptual framework for dune field dynamics and how this is affected by the sediment state of a dune  
110 field -- related to sediment supply, sediment availability, and transport capacity of the wind – is  
111 described within Kocurek and Lancaster (1999).)

112 A sufficient supply means much more than the volume of the dune – for most dunes to form  
113 and evolve, sand must be able to move onto and off of the dune (possible exceptions would be climbing  
114 dunes or other such features where the sand accumulates due to blockage). Barchans in particular are  
115 an inefficient dune shape due to sand leakage from the horns (Hersen, 2004). Thus, an important  
116 implication with the first recognition of a dune feature on a planetary body is that a process must exist  
117 that will yield a significant amount of sand (discussed in within an example more detail in Phase 43).

---

<sup>4</sup> ~~Dunes and ripples can also form due to the flow of other fluids, such as water. In this paper, we focus on aeolian dunes.~~


118 Depending on the body, that process may involve erosion of larger rocks (e.g., as is common on the  
119 Earth) or a process that directly forms grains of that size. For instance, martian volcanic activity has been  
120 proposed to create sand-sized particles ([Edgett and Lancaster, 1993](#); Wilson and Head, 1994) and  
121 photochemical processes in the Titan atmosphere may eventually lead to ~~sand-saltatable~~ grains, perhaps  
122 via an intermediate evaporite or sedimentary location (Soderblom et al. 2007; Radebaugh, 2013; Barnes  
123 et al. 2015).

124 On the Earth, nearly all dunes and ripples are comprised of sand grains – and this refers to a  
125 specific size. Although sand is commonly defined based on size (e.g., the Canada Dept. Agriculture (1976)  
126 lists sand as grains 0.05-2 mm in diameter). However, “sand grains”, it can also be defined as a size class  
127 defined based on dynamics. “Sand” is the size of grains most easily moved by a fluid (Bagnold, 1941) –  
128 smaller grains are held together by interparticle, cohesive forces and larger grains have more mass and  
129 so are held down more by gravity. Under the Earth’s atmosphere and gravity, grains ~1 mm in diameter  
130 are able to saltate, and thus are the most easily moved by the wind. On Earth, this correlates with the  
131 size range generally considered as sand; but on other bodies (with However, under the influence of a  
132 different atmospheric (or fluid) density and gravity), the grain size most easily moved by the wind could  
133 be a different size (**Figure 1**; Edgett and Christensen, 1991; Greeley et al., 1974; 1980; [1992a](#); Moore et  
134 al., 2015). Thus, throughout this discussion, when discussing “sand grains,” we mean “the grain most  
135 easily moved by the wind (or fluid)” and not a fixed size range. Thus, the existence of a dune (i.e., a  
136 landform composed of sand grains) on a planetary surface gives us yields a coupled constraint on the  
137 grains and the wind velocity.

138 Even if the wind reaches sufficient strength to transport sand, if it is not consistent (in direction  
139 and velocityspeed) over a sufficient period of time, the wind would just move small amounts of sand  
140 back-and-forth until that sand became trapped into depressions, sheltered areas, and other sand-traps;  
141 that sand would not be able to organize into a bedform. Models have shown that sand dunes have a

142 minimum size (e.g., Claudin and Andreotti, 2006; Parteli et al., 2007); below this size the slipface is  
 143 unable to develop. A slipface is necessary to stabilize the dune (as sand then can be captured on the  
 144 sheltered, lee slope) to allow it to continue growing and migrating.

145



146

147 **Figure 1: Wind shear velocity needed to move grains of different sizes, on different planets. Plot**

148 showing the estimated threshold shear velocity for moving wind-driven transport of a grain of a specific  
 149 diameter for (from top) Pluto, Mars, Earth, and Titan; curves are taken from Moore et al. (2015; Fig. 17).

150 The general shape of the curve is reflective of smaller particles experiencing stronger interparticle forces  
 151 (such as electrostatic forces), while larger particles have more mass – either effect thus requiring more  
 152 shear velocity to initiate and sustain grain movement. The curve’s minimum indicates the expected size  
 153 of “sand grains” (i.e., the grains most easily lifted and moved by a shearing fluid – by saltation) on that  
 154 planetary body, that would be involved in the formation of aeolian bedforms. On Earth, sand grains are

155 commonly ~0.1mm in diameter. On Mars observations of saltatable grains (“sand grains”) in sand  
 156 aeolian deposits and such as dunes (e.g., Figure 11) yield comparable diameters, which is consistent with

157 the curves shown. The differences in shear velocity needed to initiate motion are due primarily to

158 | differences in the estimated air (fluid) and grain densities on each planetary body. The first investigations  
159 | into these curves and how they shift under different planetary conditions (Greeley et al., 1974) were  
160 | initiated based on observation of dunes on Mars (as described in the text: Belcher et al., 1971; McCauley  
161 | et al., 1972; Cutts and Smith, 1973).

162 |  
163 | To date, we have seen potential dunes on every deeply-studied body with an atmosphere and  
164 | observable surface (including ~~one~~ Titan, where dunes were considered unlikely: Lorenz et al., 1995), as  
165 | well as a few bodies with no known atmosphere (Table 2). Based on the connections outlined above,  
166 | this first “sighting” suggests the *accumulation of a lot of sand* (leading to questions about where the  
167 | sand is coming from and why it is accumulating) and implications about *wind strength, direction, and*  
168 | *consistency*. This yields ~~(yielding~~ a “groundtruth” observation for comparison with atmospheric models  
169 | in both wind strength and direction (—although it may be unclear when the bedform was created and  
170 | thus what input conditions should be used for the model, or how the bedforms may have since been  
171 | modified by non-aeolian processes).

172 | Two classic examples of this are Mars and Titan. On Mars, signs of aeolian processes had been  
173 | seen in cyclic, large-scale albedo changes and Mariner 6 imaged crescent-shaped features that were  
174 | hypothesized to be very large barchan or parabolic dunes (Belcher et al., 1971). The first clear example  
175 | of martian dunes was observed by Mariner 9 (McCauley et al., 1972; Cutts and Smith, 1973). Those  
176 | observations suggested a wind regime that would allow for transport and collection of, as well as the  
177 | presence of, a large amount of granular materials<sup>2</sup>, leading into laboratory studies of aeolian granular  
178 | transport (Greeley et al., 1974; 1980). When Viking 2 imaged the north polar erg (Figure 2), this led to


---

<sup>2</sup> As Cutts and Smith [1973, p4151] put it: “The principal implication of dunes is a supply of noncohesive particles in the Martian surface environment and wind velocities sufficient for saltation transport. ... Dunes are not amenable to an alternative explanation of this sort. Thus we feel that we can now confidently assert the existence of a saltation regime on Mars”, which leads to “many implications of a saltation regime such as wind abrasion, wind scour, and dust production.”



179 investigations of martian erosional processes (acting on polar layered deposits or soils of lower  
180 latitudes?) and climate models (Cutts et al., 1976). (A summary of results from Viking and Mariner-based  
181 aeolian studies can be found in Greeley et al., 1992a.) On Titan, “cat-scratch” features had been  
182 observed circumnavigating its equator, but were not immediately recognized as dunes until Vic Baker  
183 brought the large draa of Saharan/Arabian/Namib deserts to the attention of the Cassini RADAR Team.  
184 The presence of dunes was a surprise as it had been hypothesized that while Titan’s atmosphere may be  
185 capable of moving sand grains, it seemed unlikely that grains of the right size would exist (Lorenz et al.,  
186 1995). Observation of the dunes (Lorenz et al., 2006) led immediately to detailed investigations of what  
187 grains could be made of and how they would form (furthering studies of the chemistry on this Saturnian  
188 moon: Lorenz et al. 2006; Soderblom et al. 2007; Barnes et al. 2015) as well as leading to attempts to  
189 reconcile the observed dune morphologies with the model-predicted wind regime around the equator  
190 (Lorenz and Radebaugh 2009).

191




192

193 **Figure 2. An early image of the martian north polar erg.** (a) These linear features, imaged near the  
194 martian north polar cap by Viking 2 (frame 59B32: 62 km x 104 km), were hypothesized to be dune fields

195 based on their consistent orientation and wavelength, and low sinuosity, branching and merging. Image  
196 and description are taken from Cutts et al. (1976; Fig. 7). (b) Higher-resolution images have proven that  
197 these are dune fields, with a wavelength (between primary crestlines) of approximately 0.4 km. A few  
198 more orders of aeolian bedforms (e.g., the smaller crestlines, transverse to the primary crestlines) are  
199 also visible. Image is a portion of HiRISE PSP\_007115\_2600 (MRO/NASA/UA).

200



201

202 **Figure 3. A ~5 m boulder and potential aeolian features at Philae lander's touch-down-1 site.** This  
203 ROLIS descent camera image shows a depression partly surrounding the boulder and a triangular-shaped  
204 apron on the opposite side have been interpreted as a moat and a windtail, indicating transport of  
205 granular material across the comet's surface (Mottola et al., 2015). Initial studies evaluated possible  
206 aeolian mechanisms for this transport. © ESA/Rosetta/Philae/ROLIS/DLR

207

208 Contrarily, while Venus has a dense atmosphere, only two potential dune fields and a few  
209 possible microdune fields have been identified within Magellan radar data ([Greeley et al., 1992b; 1995;](#)  
210 [Weitz et al., 1994](#)), which covers 98% of the surface with 100 - 200 m resolution ([Greeley et al., 1992;](#)  
211 [Weitz et al., 1994](#)) and shows wide coverage in other aeolian features such as windstreaks and potential  
212 yardangs ([Greeley et al., 1995](#)). This confirmed the hypothesis that aeolian bedform development on


213 Venus must be limited, based on Venera 13 and 14 observations of the venusian surface that showed a  
214 dearth of aeolian ripples within loose material (Basilevsky et al., 1985; Florensky et al., 1983). (Note that  
215 for Venus, surface observations were first, before the mapping of surface topography from orbit.) The  
216 implication was implies that ~~either~~ venusian conditions and processes commonly obliterate features,  
217 the conditions for dune formation are not common on Venus (e.g., that there are few sands available on  
218 Venus or wind is not sustained at the surface), or that dunes are not generally visible via the radar  
219 images (Greeley and Arvidson, 1990; Weitz et al., 1994). Wind tunnel (Greeley et al., 1984a; 1984b;  
220 Marshall and Greeley, 1992; Williams and Greeley, 1994) and sand flux modeling (Kok et al., 2012) has  
221 shown that saltation under venusian conditions may occur in a very thin near-surface layer with very low  
222 velocity, which does not favor formation of large dunes. Thus, venusian sand transport appears more  
223 comparable to terrestrial bedform formation under water (Marshall and Greeley, 1992; Kok et al. 2012;  
224 Neakrase, 2015). Unfortunately, no new data about Venus has been acquired since Magellan, so Venus  
225 dune investigations remain stuck just past Phase 1 (with a small start ~~withinat~~ Phases 2-4, see discussion  
226 of Phase 2; Figure 13).

227 Potential aeolian bedforms have also been seen on planetary bodies lacking an atmosphere. For  
228 example, planetary scientists were recently very surprised to see features that looked like aeolian  
229 bedforms (i.e. moats, wind tails, and aeoliandune-like ridges) on comet 67P/Churyumov-Gerasimenko  
230 (Figure 3; Mottola et al. 2015; Thomas et al. 2015a; 2015b). A comet seemed clearly to be a planetary  
231 body that would lack an atmosphere, and thus any wind -- yet the features were observed. This  
232 immediately led to studies trying to determine how a “wind” could exist on this comet, if even  
233 transiently. One mechanism proposed to explain particle mobilization on comets was gas outflow from  
234 reservoirs of subsurface sublimating ice that emerges and erodes particles from channel walls (Cheng et  
235 al., 2013). However, since this process would only affect localized regions and the aeoliandune-like  
236 features on 67P have been observed in a much wider area, ~~Mottola et al. (2015) suggested~~ “splashing”

237 initiated by airfall, i.e. ejection of particles by incoming projectiles, ~~has been suggested as to be~~ the most  
238 significant mechanism to explain particle mobility (Mottola et al, 2015; Thomas et al., 2015b). A three-  
239 dimensional cellular automaton model has proven that moats can result from abrasion of the surface by  
240 impinging particles, whereas wind tails develop where granular surface materials were shielded by  
241 obstacles from particle transport (Mottola et al., 2015). The results of this study put forward the  
242 explanation that the aeolian bedform-like features on comet 67P are of erosional nature, rather than  
243 depositional – but the questions and investigations that arose in response to the recognition of features  
244 that resembled aeolian bedforms were consistent with typical Phase 1 discussions.

245 The surface of Io is covered in a ubiquitous frost of SO<sub>2</sub> as seen by the Galileo Near Infrared  
246 Mapping Spectrometer (NIMS), likely mixed with dust and fine-grained materials, all ejected from the  
247 continuously erupting volcanic plumes and explosive volcanic eruptions (Kieffer et al. 2000, Milazzo et  
248 al. 2001). The surface as seen by the Imaging Science Subsystem (ISS) instrument on Galileo is mostly  
249 uniformly light-colored from this frost and generally smooth, with some fractures, slumps and pits  
250 (McEwen et al. 2000). In a few regions, there are landforms with dune-like characteristics: regular  
251 spacing, a slightly meandering form, “crestline” defects, and apparent topography visible through the  
252 uneven collection of frosts (not possible to confirm with Galileo’s instruments). In one location, the  
253 dune-like landforms are found near a particularly active volcanic plume source, the Prometheus plume,  
254 which is sourced by advancing lava flows over vaporizing frosts (Figure 4; Kieffer et al. 2000; Milazzo et  
255 al. 2001). It is possible this plume forms a localized atmosphere dense enough to loft particles from the  
256 surface and deposit them nearby in dunes, much like one of the processes hypothesized for forming the  
257 features on comet 67P.

258



259


260 **Figure 4. Potential aeolian bedforms on Io.** This ridged terrain has been postulated as potential aeolian  
261 bedforms, formed from volcanic plume deposits. Lava is erupting from a fissure about 40 km east (right)  
262 of the edge of this mosaic, and the 100 km tall Prometheus plume is erupting from somewhere near the  
263 western (left) end of this mosaic. The bright streaks radiating from the area where the lava flows (the  
264 dark features) overrun the field are where the hot lava recently vaporized the sulfur dioxide, which then  
265 coated the lava-facing sides of the ridges. These images were taken by Galileo during a flyby of Io on  
266 February 22, 2000, with a resolution of 12 m/pixel. Image and description are taken from NASA  
267 Photojournal PIA02568.

268

269 The flyby of Pluto by New Horizons in July 2015 produced one of the most striking increases in  
270 image quality of a planetary surface in the history of planetary exploration (Moore et al., 2016; New  
271 Horizons, 2015; Stern et al., 2016). The landscape revealed imaged by during the flyby revealed a  
272 surprising diversity of landforms, which suggest varied geological and geomorphological processes active  
273 within recent geological history surface (Moore et al., 2016; Stern et al. 2015; Trilling et al., 2016).

274 Mountains, glaciers, plains, possible cryovolcanism and a surprisingly low density of craters covered  
275 much of the surface. Initial science results from the flyby noted the possible existence of 'windstreaks'  
276 (Stern et al. 2015) on Sputnik Planum, apparently extending in the lee of dark hills protruding through  
277 the nitrogen ice of which the plains are composed. Aeolian bedforms were speculated on before New  
278 Horizons arrived (Moore et al., 2015), and dunes have since been posited in the Baré Montes and  
279 enigmatic features in the Tartarus Dorsa have been interpreted variously as dunes or erosional aeolian  
280 features (Fenton, 2016; New Horizons, 2016). However, recent imagery from Sputnik Planum (Figure 5)  
281 provides perhaps some of the most convincing examples of potential aeolian bedforms, with continuous  
282 linear features with a spacing of ~400-600 m. These features extend across the polygonal dark features,  
283 which have been interpreted as convectonal cells within the ice, suggesting that they are the result of  
284 surface processes not related to the convective movement within the ice. Pluto is thus tentatively within  
285 Phase 1 of the progression, and if an aeolian origin can be shown to be feasible, then available data may  
286 be sufficient for progression into Phases 2-3. The dune-like landforms on comet 67P, Io and Pluto are  
287 well in Phase 1 discussions and primary work remains to be done for these features to determine their  
288 ultimate origin.

289



290

291 **Figure 5. Possible Pluto aeolian bedforms at the margins of the nitrogen ice of Sputnik Planum.** These  
292 features are oriented approximately parallel to the 'shore' where the ice abuts mountains. Image width  
293 is approximately 75 km. Image credit: NASA/JHUAPL/SwRI.

294

#### 295 **Summary of Phase 1**

296 **Data needed:** Images of the surface topography, of sufficient resolution to identify the distinctive  
297 shapes of dunes -- images could be visible or spectral imagery or radar scans of the planetary body's  
298 surface; identified analog (usually terrestrial) aeolian bedforms.

299 **Knowledge gain (from that data):** Existence of a potential aeolian bedform.

300 **Assumptions generated:** Conditions (wind conditions and grain size/supply) conducive to dune  
301 formation and evolution exist or have existed – note that this is a coupled constraint, and further  
302 information is needed to estimate each individual measurement.

303 **Questions:** What is the composition of the grains? How were sand-sized grains formed? Why do the  
 304 grains accumulation in that particular location (related to winds, topography, sand source)? Are the  
 305 dunes active? If conditions are unusual (e.g., tenuous atmosphere), how does sustained saltation and/or  
 306 reptation occur on this world?

307 **Lead to investigations of:** Models of surface processes and rates (chemical, erosional, etc.) that could  
 308 create “sand” grains; comparison to (global) atmospheric models; independent measurements of  
 309 surface conditions/composition for comparison to grain-formation models; independent studies of wind  
 310 speeds and/or grain sizes (e.g., thermal inertia) to decouple saltation conditions.

311 (Alternatively: ) Table N. Summary of Phase 1

<u>Data needed within Phase 1</u>	<u>Images of the surface topography, of sufficient resolution to identify the distinctive shapes of dunes -- images could be visible or spectral imagery or radar scans of the planetary body's surface;</u> <u>Identified analog (usually terrestrial) aeolian bedforms.</u>
<u>Knowledge gain (from that data)</u>	<u>Existence of a potential aeolian bedform.</u>
<u>Assumptions generated</u>	<u>Conditions (wind conditions and grain size/supply) conducive to dune formation and evolution exist or have existed – note that this is a coupled constraint, and further information is needed to estimate each individual measurement.</u>
<u>Questions generated</u>	<u>What is the composition of the grains?</u> <u>How were sand-sized grains formed?</u> <u>Why do the grains accumulation in that particular location (related to winds, topography, sand source)?</u> <u>Are the dunes active?</u> <u>If conditions are unusual (e.g., tenuous atmosphere), how does sustained saltation and/or reptation occur on this world?</u>
<u>Leads to investigations of</u>	<u>Models of surface processes and rates (chemical, erosional, etc.) that could create “sand” grains;</u> <u>Comparison to (global) atmospheric models;</u> <u>Independent measurements of surface conditions/composition for comparison to grain-formation models;</u> <u>Independent studies of wind speeds and/or grain sizes (e.g., thermal inertia) to decouple saltation conditions.</u>

312

313 **2.2. Phase 2: Analysis of gross individual dune characteristics**

314 Analysis of the dune morphology (Phase 2) typically closely follows Phase 1, often via the same  
 315 medium of an image of the surface. As models (e.g., Pelletier, 2009; Sauermaun et al., 2001; Werner,



1995) and laboratory studies (e.g., Andreotti et al., 2006; Parteli et al., 2009) have established, a dune's overall shape and orientation yields additional information about wind conditions when the dunes were forming and evolving<sup>3</sup>. Thus, this investigation stage generally involves detailed mapping of dune shape (outline and crestline) and comparison to analog terrestrial dunes. Numerical models of sand transport and dune evolution are also used for comparison between dune-generated predicted wind directions and atmospheric models (possibly down to a mesoscale or regional scale and including the effects of large-scale topography).

For example, once it was recognized that the long, generally linear cat-scratch features on Titan were dunes (Lorenz et al. 2006), researchers began investigating what type of dunes were observed and what wind conditions were required for their formation and persistence (in addition to questions about what the sand may be made of). From the morphology of the dunes, visible in 350 m/pixel Synthetic Aperture RADAR (SAR) data from the Cassini spacecraft (slightly hyper-resolution in nature in some locations because of the high contrast between SAR-absorbing sands and fractured, signal-scattering bedrock), it was determined they are longitudinal in type (also called linear; Lorenz et al. 2006; Radebaugh et al. 2008). Cassini Visual and Infrared Mapping Spectrometer (VIMS) data in select, high-resolution regions confirmed the general morphology of Titan's longitudinal dunes as well as their spectral contrast between sand and substrate (Barnes et al. 2008). On the Earth, longitudinal dunes are typically formed ~~via~~ when several alternating winds of roughly equal transport strength (i.e., they move the same amount of sand) are >90° apart ~~blow slightly off-axis (within ~15°) to the dune crestlines,~~ yielding a single sand transport direction that is along the dune crestline (Fryberger and Dean 1979; Parteli et al., 2009; Rubin and Hunter, 1987; Rubin and Ikeda, 1990; Tsoar, 1983). (An alternate


---

<sup>3</sup> However, establishing a connection between wind directions and dune slipface orientations sometimes is not a straightforward process – such studies often rely on many assumptions about timing of the winds and their consistency, and results are usually non-unique. Thus additional information is usually needed to evaluate a proposed interpretation. See Phase 3 for more discussion of the complexity that can be found in many dune fields.

337 hypothesis that had been put forth connecting longitudinal dune morphology and wind directionality for  
338 Titan and some Earth dunes is discussed in Subsection 3.2.)

339 Further information about these dune fields was then used to also predict the dominant wind  
340 | directions, for comparison with climate models (discussed here, but merging into Phase 3-~~type~~  
341 | investigations). Titan's dunes are found strictly between 30°N and S, ringing the equator, and are  
342 | oriented roughly parallel to the equator (Lorenz et al. 2006; Radebaugh et al. 2008). Their morphology,  
343 | and especially behavior around topographic obstacles (wherein sands-grains are piled up at the upwind  
344 | margin and are more sparse at the downwind margin), indicated a general sand transport direction from  
345 | west to east (Figure 6; Courrech du Pont et al., 2014; Lorenz and Radebaugh 2009; Lucas et al., 2014;  
346 | Radebaugh et al. 2010;). However, this was found to be at odds with the global atmospheric transport  
347 | direction of east-to-west predicted in the equatorial zone by global climate models. Subsequent  
348 | modeling studies revealed that seasonal or storm-driven winds could produce fast westerlies near the  
349 | equator (Tokano 2010; Charnay et al., 2015), possibly resolving this inconsistency.

350



351  
 352 **Figure 6: Earth and Titan dunes, diverging around topographic obstacles.** (a-b) Landsat 7 ETM+ and  
 353 SRTM C-band images of dunes in Namibia (centered on 25° 23' S 15° 16' E). The right image (b), a  
 354 Synthetic Aperture RADAR (SAR) image of the same dunes, shows bedrock as bright, because it is rough,  
 355 and unorganized dune sands as dark, because dunes are smooth at the SAR wavelength of 6 cm (black  
 356 areas are regions devoid of data returned to the SAR antenna).The right image (b) shows bedrock as  
 357 bright and unorganized dune sands as dark (black areas are unprocessed voids). The winds blow SE to  
 358 NW (bottom L to upper R), as evidenced by the dune-free regions in the lee of the bedrock topography  
 359 and the diversion of dunes around the upwind sides of the topography. In the lower image (c), dunes  
 360 similarly divert around topographic obstacles and resume on the downwind side, within this region of the

361 *Belet sand sea, Titan. This is a Cassini Radar image (300 m resolution) centered on 6.5° S, 251° W, with*  
362 *winds interpreted to blow SW–NE (L–R). Images and description are from Radebaugh et al. (2009).*


363

364 |           On Mars, mapping of the duneforms observed by Mariner 9 and Viking orbiters2, combined with  
365 | information about the surrounding surface topography and composition (including other potential  
366 | aeolian features), was used to determine the wind direction(s) that would yield the dune crestline  
367 | orientation(s), the wind and topography that would yield sand accumulation within that area and its  
368 | likely stability (i.e., was this a stable sink for sand, or a temporary repository), and possible sand sources  
369 | (Cutts and Smith, 1973; Greeley et al., 1992a; Thomas, 1981; 1982; Tsoar, 1979). For example, the large  
370 | number of transverse dunes observed within southern mid-latitudes and north polar region on Mars (vs.  
371 | longitudinal dunes), in Viking and Mariner images, implied that dune fields were forming within regions  
372 | of near unidirectional winds, and the asymmetry in dune field complexity between the north and south  
373 | hemisphere was seen as evidence of a more complex southern wind regime (Greeley et al., 1992a;  
374 | Thomas, 1981). Afters higher resolution images became available and atmospheric modeling became  
375 | more refined in technique and topography/boundary inputs, a more detailed comparison was ~~then~~ done  
376 | to see if atmospheric models could reproduce the observed dune shapes and orientations. For example,  
377 | mesoscale modeling of dunes within Proctor Crater on Mars based on MOC NA images matched two  
378 | sets of dune slipface orientations (the primary and tertiary) to seasonal winds that were impacted by  
379 | daily and seasonal insolation patterns and the crater topography, as predicted by a Global Circulation  
380 | ~~Model~~ mesoscale model (Fenton et al., 2005; however, the secondary dune slipface orientation  
381 | remained unexplained). This study validated the mesoscale atmospheric model by providing a  
382 | reasonable explanation for the range of slipface orientations seen within that dune field, and thus  
383 | advanced the use of these models within model-observation comparison studies for understanding  
384 | aeolian processes on Mars.

385 Coupled with analysis of wind direction, observations of the dune's local surroundings can also  
386 be studied to identify sediment sources and sand transport pathways. In some areas, distinct sand  
387 transport pathways leading from the sediment layer to the dune bodies have been revealed by the  
388 detection of congruent material composition. For example, on Mars, local sediment sources for intra-  
389 crater dunes have been ~~identified~~ proposed at impact crater walls by comparative analyses of high  
390 resolution image and spectral data of dune bodies and the sediment layers exposed (e.g., Fenton, 2005;  
391 Geissler et al., 2013; Silvestro et al., 2010a; Tirsch et al., 2011). Within the Valles Marineris rift system,  
392 spectral analysis, morphological evidence of erosion and sand transport, and topographic information  
393 was used to show that diverse and distinct sediment sources serve as local and regional sources  
394 (Chojnacki et al., 2014). ~~Distinct sand transport pathways leading from the sediment layer to the dune~~  
395 ~~bodies have been revealed by congruent material composition.~~

396 On Venus, analysis of dune morphology is almost impossible due to lack of data. Dunes in both  
397 recognized dune fields are at the resolution limit of radar images obtained by Magellan mission (Figure  
398 7), the only adequate data source. Therefore, results of attempted of detailed analysis are not reliable  
399 and are controversial (Greeley et al., 1997; Lorenz, 2015). Modeling of anisotropic radar scattering  
400 (Kreslavsky and Vdovichenko, 1999) indicated that the microdune fields ~~identified~~ proposed by Weitz et  
401 al. (1994) possess abundant unresolved steep slipfaces.

402



403

404 **Figure 7: Venusian dune field.** This Magellan SAR image of the Aglaonice dune field (25°S, 340°E) shows  
 405 a field of white dots that are interpreted as specular reflections from the slopes of transverse dunes  
 406 (between the arrows). This type of reflection occurs if the slope is mostly smooth, and is oriented near-  
 407 normal to the incidence angle of the radar (which is 35° or near the angle of repose); similar features are  
 408 observed with ~~g~~ Seasat and space shuttle radar images of terrestrial sand dunes. The implied wind  
 409 direction of these features, based on their orientation, is also consistent with nearby bright and dark  
 410 wind streaks extending from behind cones (not included in this cropped image). Radar illumination is  
 411 from the left, and north is at the top. Image is from NASA Photojournal PIA00483, and description is from  
 412 Photojournal and Weitz et al. (1994).

413

414 In general, Phase 2-type investigations can continue for a long period with refinement of the  
 415 investigations as higher-resolution images of the dunes are acquired, more information becomes  
 416 available about the local topography and other evidence of aeolian processes and conditions, and/or  
 417 atmospheric or bedform formation models are improved. This investigation phase may eventually grade  
 418 into (and occur concurrently with) Phases 3 and 5 which involve, respectively, higher-resolution analysis

419 of features within the dune field and features on the dunes formed by the dune-wind interaction (such  
420 as ripple patterns on the ~~stoss or lee dune~~ slopes ~~of dunes~~). Additionally, as new observations, models,  
421 and analysis methods are developed, Phase 2 investigations can be renewed and revisited (as with all  
422 Phases beyond Phase 1).

423

#### 424 **Summary of Phase 2**

425 **Data needed:** *same as Phase 1.*

426 **Knowledge gain:** Morphology of the potential crestlines and general dune shape; composition of the  
427 dunes and surroundings; possibly identification of local sand sources.

428 **Assumptions generated:** Wind direction and consistency *hypothesized* to generate the observed shapes;  
429 variations in wind speed implied by changes in sinuosity and shape through the dune field.

430 **Questions:** What sets the wind direction and causes its variations (e.g., daily or seasonal cycles)? Are  
431 these representative of present-day wind conditions, conditions during a past period, or a convolution  
432 of conditions during different past periods?

433 **Lead to investigations of:** Comparison over one to several dune fields with (global/mesoscale)  
434 atmosphere models; reliably identify wind direction(s) consistent with the dunes' forms.

**Table 2: Recognition and first analysis of dunes and dune fields on planetary bodies (Phases 1-2), as presented in the literature**

Planet. body	“Aeolian” bedforms first sighted	Data used	Immediate Implications	Immediate Questions	Reference
Mars	Mariner 9, Hellespontus region of Mars, dense and large transverse dune field	Visual image of surface, <1 km/pixel	Dune material is dark (so some <del>dark</del> -low albedo areas are regions of deposition, not erosion; and some dark material will saltate); Due to the lower atmospheric density on Mars, wind velocities may need to be much higher to move sand.	Which “dark splotch[s] or streak[s]” are due to deposition of material (vs. deflation of overlying bright material)? What is the source of the dark material? Does the wind reach transonic velocities? Is high-velocity sand-blasting resulting in highly-efficient wind erosion?	Sagan et al., 1972
		above; comparisons to other albedo markings indicative of wind direction	Presence of lots of sand and saltation processes; Dune material accretion directions and influence of topography (craters) on field location and dune morphologies; comparable scale and shapes as terrestrial dunes	What is the composition of the sand? Why is it so dark?	Cutts and Smith, 1973
	Viking 2, north polar erg, transverse and barchan dune fields	Visual image of surface, 30-60 m/pixel	Lots of sand → some erosional process; Variability in wind regime; two wind directions in portions Strong winds; Two wind directions, thought to be seasonal; grains may be eroded from the northern plains	What is the composition and source of sand? Why is it accumulated around north polar cap? Are the dunes active, and how mature is the dune field? Are the dunes modified during the winter/early spring, when the entire region is covered by CO <sub>2</sub> ice?	Cutts et al., 1976 Tsoar et al., 1979
Venus	Magellan, transverse dunes, two fields	Radar images, 75 m/pixel; compared with orientation of other aeolian features in same dataset	Lots of sand in specific areas → some erosional process (perhaps impacts?)	What is the composition and source of the sand? What are the saltation dynamics under a much denser atmosphere?	Greeley et al., 1992 <b>b</b>
Titan	Cassini, longitudinal	Synthetic Aperture Radar images, 175	Lots of sand → some “grain” formation process; one dominant	What is the composition and source of the sand? What is the underlying topography,	Lorenz et al., 2006



	dunes, large field around equator	m/pixel	or at least two converging wind directions throughout equatorial region; pristine appearance and superposition over geologic features → young and possibly currently active	causing accumulation in equatorial region and diversion of dunes? Is there any connection with the potential fluvial channels? Has the sand circumnavigated the globe several times (implying a lack of sand-sinks in the area)?	
		Visual and Infrared Mapping Spectrometer (VIMS) observations, 500 m/pixel	Observations of interdunes → recent activity and overall dune field maturity; spectral information → constraints on the composition of the dunes and interdune regions; photogrammetry yielded height and wavelength estimates	What information does the variability in dune coverage and height, and in the terrain that they cover, yield for the evolution history and conditions for these mature and recently active dunes? What are the dune grains made of, given their lower relative waterice content than Titan's average?	Barnes et al., 2008
<b>Comet 67P/Churyumov-Gerasimenko</b>	<b>Rosetta</b> , moats, wind tails, and aeolian-like ridges/ripples	optical imagery: OSIRIS orbiter camera images ( $\geq 0.29$ m/px), ROLIS decent camera images ( $\geq 1$ cm/px)	"Sustained" granular transport along the surface (so as to form aeolian bedform-like features) exists on a comet	How does granular transport work on a body without atmosphere? What is the moving agents? Are these bedforms accumulative or erosive? What is the grainsize of the bedform materials? What are the material sources? How long does it take to form bedforms?	Mottola et al., 2015; Thomas et al., 2015a; <a href="#">2015b</a>


### 394 | 2.3. Phase 3: PatternA analysis of the dunes within a field

395 | Dune field evolution is related to the evolution of its constituent dunes, but occurs on a larger  
396 | spatial and temporal scale and involves areas of investigation that are different from (and can be larger-  
397 | than) the sum of its parts. For example, as dunes evolve within a field they exchange sand between each  
398 | other through both sand flux and collisions, and environmental boundary conditions such as the sand  
399 | influx geometry can affect dune field pattern development (Diniaga et al., 2010a; Ewing and Kocurek,  
400 | 2010). As such, it is necessary to model dune field evolution as more than just a collection of individually  
401 | evolving dunes, and to recognize that the large-scale dune field pattern can reflect conditions (and  
402 | changes in those conditions) around and throughout the field. For example, sand and dune influx  
403 | conditions will be different near the start-upwind margin of the dune field than near the terminus or  
404 | boundaries-lateral margins (Ewing and Kocurek, 2010) due to proximity to sand sources or other dunes,  
405 | or changes in topography or “cementing” influences (e.g., chemical duricrust or, on the Earth,  
406 | vegetation) within the field (Kocurek and Ewing, 2005) (Figure 6: examples of dune interactions with  
407 | topography on Titan and Earth). Such changes can result in different dune sizes, spacing, or defect  
408 | frequency (Diniaga et al., 2010a; Ewing and Kocurek, 2010).

409 | The effect of underlying topography is also a key parameter affecting dune characteristics at the  
410 | dune field scale (Ewing and Kocurek, 2010). On Earth, bedrock topography has been linked to the effect  
411 | of roughness variations induced by the dune field itself producing an internal boundary layer decreasing  
412 | the shear stress downwind (Jerolmack et al., 2012) and/or to the feedback mechanism between long-  
413 | wavelength topography and the dunes (Pelletier, 2015). The role of topography in enhancing and  
414 | deflecting regional winds has also been invoked to explain complex dune field pattern on Mars in  
415 | Olympia Undae (Ewing et al., 2010) and complex dune arrangements in Moreux (Cardinale et al., 2012)  
416 | and Matara crater (Diniaga et al., 2010b; Silvestro et al., 2012). However, it was only thanks to the  
417 | availability of high resolution DTMs from the HiRISE instrument that the effect of underlying topography

418 could be more precisely linked to different dune characteristics such as migration rates, dune heights  
419 and density (Cardinale et al., 2016, Vaz et al., submitted). In particular, in Herschel crater dune density,  
420 slip face advancements and migration rates are all controlled by two major topographic highs on the  
421 crater floor (Vaz et al., submitted).

422         The dune field may also record changes in conditions over a longer-timescale than that recorded  
423 within any individual dune. Multiple patterns (e.g., different types of dunes) can be superimposed  
424 (creating a complex, versus a simple, dune field) as smaller dunes migrate and change in response to the  
425 new environment faster than larger dunes (Ewing and Kocurek, 2010; Hugenholz and Barchyn, 2010;  
426 Kocurek and Ewing, 2005). We note that this possible complexity within dune fields can complicate  
427 analysis of the dune morphology (Phase 2). For example, even identification of the dominant (or most  
428 recent?) slipface orientations can be non-trivial. This is especially true within planetary dune fields  
429 where datasets may be limited to remote images, so dune slope angles and potential activity have to be  
430 interpreted from images of the dunes' planform appearance, possibly under suboptimal illumination  
431 conditions for this type of image analysis. For example, within the north polar erg on Mars, many dunes  
432 contain slipfaces pointing in opposite directions (sometimes on the same dune). One interpretation is  
433 that some of these fields may contain both active and fossil dunes (Gardin et al., 2012). Within the Mars  
434 southern mid-latitudes, at least two periods of dune-building (or dune-building occurring over 2  
435 different timescales) are apparent as within the same field one can often find a dense collection of  
436 transverse dunes (with slipface towards the east) and then barchans clearly climbing up and over the  
437 transverse dunes on the east side (with slipfaces towards the west) (Figure 8; Diniega et al., 2010b).  
438



439

440 **Figure 8: The complex dune patterns found in along the eastern edge of Matara dune field, Mars**  
441 *(49.5°S, 34.8°E). This dune field, like many others in the Mars southern mid-latitudes, is a dense*  
442 *transverse dune field, captured within a crater. The transverse dune crestlines are oriented north-south*  
443 *with the clearest slipfaces towards the east (white arrows extend up the possible stoss slope, towards the*  
444 *dune brink). However, along the eastern side of this field, many smaller dunes (mostly barchans – yellow*  
445 *arrows, some possible transverse crestlines near the bottom of the image) are oriented with towards the*  
446 *northwest. This potentially reflects two periods (or just two timescales?) of dune evolution, with a*

447 *change in the dominant wind direction. North is up and illumination is from the left. Image is a portion of*  
448 *HiRISE PSP\_006648\_1300 (MRO/NASA/UA).*

449

450 A lack of variations can also yield information about the field's and planetary body's history. On  
451 Titan, dune width and spacing measurements over more than 7000 linear dunes showed a high level of  
452 consistency-uniformity around the moon, with no signs of compound or complex dunes (Savage et al.,  
453 2014). This, coupled with the dunes' large sizes, indicates that Titan's dunes are mature features that  
454 have evolved within consistent and stable environmental conditions for a long period of time.

455

### 456 **Summary of (3)**

457 **Data needed:** Observations of dunes fields, of sufficient spatial coverage and resolution to note changes  
458 in dune patterns throughout the field, especially in tracing crestlines; possibly need knowledge of  
459 topography.

460 **Knowledge gain:** The dune field pattern and shape; maturity state (and possibly relative age) of the  
461 bedforms; possible temporal changes in e.g., sediment supply and wind patterns-

462 **Assumptions generated:** Changes in the environmental conditions, in space or time.

463 **Questions:** For a given dune field, is sand sourced from one or several locations? Is the dune evolving  
464 through one dominant wind pattern, or several? Have the dominant effects (sand source or wind  
465 pattern) changed over the lifetime of the dune field?

466 **Lead to investigations of:** Explore influences on dune shapes beyond current dominant winds (*Phase 2*)  
467 – such as the location of sand source(s) or of sand-starved regions of the field, a transition between  
468 wind regimes, interactions between dunes (such as dune collisions), or other environmental  
469 influences/processes.

470

471 | **2.4. Phase 4: ~~Regional and Global-global~~ surveys and ~~aggregate~~-analysis of dune ~~characteristics~~**

472 |         As we gather information about dunes in more and more different dune fields around a  
473 | planetary body, it becomes possible to aggregate data to deal with high-level, large timescale questions  
474 | about aeolian processes and sediment supply, such as “How much sand is available in total?” and “Are  
475 | there primary types/locations of sediment sources that can yield information about how that sand has  
476 | been created, how it is transported, and whether it has been recycled?” Addressing such big-picture  
477 | questions can provide important information for investigations of grain-producing processes (e.g.,  
478 | surface erosion) and planetary surface history over the lifetime of the involved sand grains. Note that  
479 | while studies of grain history and sediment transport pathways involving terrestrial dunes ~~can~~may rely  
480 | on detailed petrographic and heavy-mineral techniques, with geochronology (e.g., Garzanti et al., 2013),  
481 | studies of planetary bodies often are based only on surface topography and, possibly, coarse  
482 | compositional information.

483 |         For example, on Mars, a near-global map of sediment deposits (including dunes) and wind  
484 | streaks to generate early estimates of sediment transport pathways/source regions (Thomas, 1982). ~~the~~  
485 | An early global map of aeolian features showed variations in time and space in the large-scale wind  
486 | directions recorded by the orientation of dunes, wind streaks, yardangs, wind grooves, and deflation pits  
487 | (Ward et al., 1985). Such studies have since been updated with increased coverage and image resolution  
488 | (e.g., Hayward et al., 2007; 2014), and still provide important information about direction and variability  
489 | in the wind patterns (down to intra-field scales), the influence of topography and local geology on wind-  
490 | flow and bedform development, and likely sediment sources for the observed deposits. At a regional  
491 | scale, the martian north polar erg volume has been estimated as ~1130-3250 km<sup>3</sup> of dark sand (Greeley  
492 | et al., 1992a; Hayward, 2011), which is significant as the Planum Boreum Cavy unit, which is part of the  
493 | icy layers of the north polar cap, has been ~~identified~~proposed as the source of the circumpolar dune  
494 | fields (Byrne and Murray, 2002; Tanaka et al, 2008). These deposits ~~are~~appear composed of recycled

495 | aeolian sediments, which were likely transported poleward and deposited ~~there~~ (Breed et al., 1979;  
496 | [Byrne and Murray, 2002](#)). This suggests that a huge volume of sand may have formed on Mars during an  
497 | earlier epoch and that these sand grains have survived at least a couple of sustained dune-forming  
498 | periods.

499 |         The Titan dune fields provide an example of how ~~an~~ analysis of the distribution of dune field  
500 | locations (on the planetary body, or relative position within regional topography) and morphologies (i.e.,  
501 | field outline or crestline patterns) can yield additional information about larger-scale atmospheric and  
502 | topographic/surface conditions. Mapping of 16,000 Titan dune segments (covering 8% of Titan’s surface  
503 | which suggested that dunes cover a total of 20% of the global surface: Lorenz and Radebaugh, 2009)  
504 | showed general dune field orientation and spacing patterns and confirmed that these features are  
505 | within a global field with few longitudinal trends, but with latitudinal trends in orientation and limited to  
506 | within 30° of Titan’s equator. Although dunes on Titan are organized into several separate sand seas  
507 | across the equator, all have some level of broad interconnectedness (Le Gall et al. 2012; Radebaugh  
508 | 2013; Savage et al. 2014). As such, studies of the Titan sand sources, sediment transport pathways, and  
509 | deposition patterns are best analyzed from a “global” perspective.

510 |         Titan sands may be derived directly from the atmosphere, perhaps through clumping on the  
511 | surface, though it is perhaps more likely the sand has been processed through erosion of organic  
512 | sedimentary layers (Radebaugh 2013), possibly close to the equator, ~~7~~ where fluvial channels have been  
513 | imaged (Burr et al., 2013; Lorenz et al., 2008; Radebaugh et al., 2016). Other possible sources include  
514 | erosion of the SAR-uniform mid-latitudes, a possible sedimentary deposit (Malaska et al., 2016), and the  
515 | northern dry lakebed evaporite deposits, which have similar spectral characteristics to the VIMS  
516 | instrument (Barnes et al., 2015). Once the materials are ~~incorporated-transported~~ into the Titan sand  
517 | seas, they are ~~absorbed-incorporated~~ into the giant linear dunes, and either stay confined to one sand  
518 | sea ~~or another~~ or contribute to a global system of west-to-east sediment transport that persists over

519 time (Savage et al., 2014). Topography appears to play an important role, as it does for sand seas on  
520 Earth, in that it can help confine the sands to certain regions or preclude them from others, like from the  
521 rugged Xanadu region (Lorenz et al., 2006; Radebaugh et al., 2011). Decreases in dune density within  
522 radar-bright and elevated regions may provide regional-scale constraints on Titan's winds for  
523 atmospheric models (Lucas et al., 2014). Furthermore, topographic obstacles can cause diversion of  
524 dunes and dune/topography relationships and perhaps reveal longer-term climatic changes (Ewing et al.  
525 2015).

526 |           Consideration of the dune fields in aggregate can also allow for analyses that require a larger  
527 | area or more numerous measurements to reflect larger-scale temporal or spatial trends. For example,  
528 | dune fields on Mars appear very young as they lack craters, but constraints on their age had large  
529 | uncertainties due to their low individual areal coverage. Adding the dune fields together allowed for a  
530 | more robust estimated crater-retention age of <10 000 years (Fenton and Hayward, 2010). These dune  
531 | fields also exhibit latitude-dependent morphological trends in crestline sharpness/pattern, dune slopes,  
532 | and field shapes, so considering the dunes over the hemisphere enables studies of which reflect different  
533 | degrees of influence from polar as well as aeolian processes (Fenton and Hayward, 2010). Another study  
534 | of southern intracrater dune fields on Mars compared dune field centroid locations, relative to the  
535 | crater center, with mesoscale atmospheric modeling to look at broad-scale atmospheric trends (over a  
536 | much longer time period than that recorded in dune slipface orientations within an individual dune  
537 | field) (Hayward et al., 2009).

538           Although only two dune fields and a few microdune fields were identified with some certainty in  
539 the whole set of Magellan radar images of Venus, a few lines of indirect evidence suggest that  
540 unresolved small-scale anisotropic topographic features are ubiquitous; such features have been  
541 interpreted as unresolved gently sloping aeolian bedforms (Kreslavsky and Vdovichenko, 1999;



542 Bondarenko et al., 2006). A comprehensive global inventory of aeolian bedforms on Venus will require  
543 global imaging data set(s) of a higher resolution than presently exists.

544 Beyond global imaging of the data-type first used to identify the dunes, proxy measurements  
545 can sometimes be used to supplement limited image coverage. For example, thermal inertia can be used  
546 to identify large deposits of unconsolidated, granular material. On Mars, further evidence that these  
547 dark patches with high-thermal inertia were aeolian deposits were that these were found downwind of  
548 topographic depressions (Christensen, 1983; [Mellon et al., 2000](#)). Thus global maps of thermal inertia  
549 with resolution ~100 m/pixel have been used to map dune fields around Mars and estimate the number  
550 of dune fields and their surface areal extent (Christensen et al., 2003; Hayward et al., 2007; Hayward,  
551 2011).

552

#### 553 **Summary of (4)**

554 **Data needed:** Identification of dunes around globe (from the data used in Phases 1-3, and possibly from  
555 proxy data such as thermal inertia).

556 **Knowledge gain:** Dune field location and (possibly) morphology/type distributions; variations in location  
557 and morphology related to sediment supply, climate history, and/or and other active processes (e.g.,  
558 related to latitude, regional topography); identification of large-scale sediment transport pathways  
559 (larger-scale than field-specific results of *Phase 2 and 3*; [and possibly first produced earlier based on](#)  
560 [low-resolution, but high-coverage datasets](#)) based on (global/mesoscale) atmospheric models and  
561 observation of sediment sources.

562 **Assumptions generated:** Correlations between dunes and proxy data; feasibility of extrapolation from  
563 studies of individual dune fields/sand sources to a global model.

564 **Questions:** How much sand is there, where is it from/stored, and how did it get there? Over what spatial  
565 and temporal scale is the sand being transported (i.e., what is the lifetime of a sand grain and is  
566 sand/bedforms being recycled)?

567 **Lead to investigations of:** Age estimation of dune fields (as can aggregate together land-areas to  
568 statistical significance; likely to be a relative or crater-retention age); identification/investigation of  
569 large-scale sediment sources (locations and/or processes; perhaps updated from *Phase 2*); global  
570 surface areal coverage of dunes/volume of sand.

571

572 **2.5. Phase 5: Analysis of superposed bedforms on the dune “details” formed due to wind interaction**  
573 **with the dune**

574         Ripples, like dunes, form spontaneously within sand beds due to wind (or fluid) flow and record  
575 wind and sediment conditions through their period of formation and evolution. However, as these are  
576 much smaller features, they record conditions over smaller temporal and spatial scales and thus can be  
577 reflective of a different set of environmental conditions than dunes. To-date, ripple-like features have  
578 been only observed on Mars, where HiRISE images of the martian surface have resolution up to as fine as  
579 0.25 m/pixel (McEwen et al., 2007). These features have wavelengths of 1-to-a-few meters, have been  
580 found within sandy regions including on the slopes of dunes, and have been individually mapped and  
581 monitored for movement (Phase 6) (**Figure 9**; Bridges et al., 2012; Silvestro et al., 2011). The study of  
582 ripple morphologies and dynamics on Mars yields information about the wind flow over the dunes,  
583 under the influence of the local wind patterns as well as the dune topography. This yields information  
584 about the recent, local wind regime within several areas on Mars (Bridges et al., 2012b). This Such  
585 information about the temporally and spatially small-scale surface wind dynamics can be compared with  
586 meso and microscale climatic models and in-situ wind measurements (e.g., Jackson et al., 2015; Silvestro  
587 et al., 2013). In addition, because ripple morphology and migration rates are controlled by the

588 topographic and wind flow boundary conditions imposed by the dune morphology (Kocurek and Ewing,  
589 2012), studies of the ripples' form and variation provide insights to the underlying dune's evolution  
590 (Ewing et al., 2010; Vaz et al., [2016submitted](#)).

591 Ripple mapping and monitoring have been an important tool within recent martian studies,  
592 where the crestline orientations and migration rates and directions of the large martian ripples are  
593 [commonly](#) used to reconstruct the wind regime over the dunes and to estimate sand fluxes (Ayoub et  
594 al., 2014; Bridges et al., 2012a; 2012b; Cardinale et al., 2016; Silvestro et al., 2010b, 2011, 2013).  
595 Automatic approaches have been developed to derive ripple trend and migration rates, enabling high-  
596 resolution wind regime estimations and sand flux measurements to be computed over large areas  
597 (Ayoub et al., 2014; Bridges et al., 2012a; 2012b; Silvestro et al., 2011; Vaz and Silvestro, 2014).

598 However, all of these studies have assumed that that the observed "smaller" bedforms on the  
599 dunes are analogous to terrestrial sand ripples, and that ripple trends and migrations are normal to the  
600 last wind of sufficient strength to move sand, as is typically the case for [aeolian](#) ripples on Earth. Recent  
601 work has drawn those assumptions into question:

- 602 • Most ripple patterns [on Mars](#) are dominated by sinuous crestlines (Vaz et al., [2016submitted](#)), while  
603 on Earth ripple crestlines are typically straight (Rubin, 2012) ([Figure 9](#)). In some areas, ripple  
604 patterns observed on Mars show complex arrangements with two crestlines intersecting at right  
605 angles ([Figure 10](#); Silvestro et al., 2011, 2013). This suggests that some of the ripples on Mars might  
606 not be in equilibrium with the last sand-moving winds [or that the two sets of crestlines are](#)  
607 [contemporaneous, but oblique to the formative winds \(Silvestro et al., 2016\)](#).
- 608 • Additionally, unusual [longitudinal](#) displacement of crest-line defect terminations and oblique crest  
609 migrations have been observed within orbital data [in Gale and Herschel crater](#), suggesting that the  
610 large ripples of Mars [might be](#) different from terrestrial impact ripples (Silvestro et al., [201in](#)  
611 [press5](#); Vaz et al., [2016submitted](#)). This hypothesis [may be](#) in agreement with recent in situ

612 observations from the NASA MSL Curiosity rover, which shows that large ripples have sinuous and  
613 sharp crests and slip faces with evident grainfall and grainflow structures (Bridges et al., 2016;  
614 Lapotre et al., 2016a) (Figure 10) that are not common within terrestrial impact ripples. Superposing  
615 these large bedforms are smaller "terrestrial-like" impact ripples of ~10 cm in wavelength (Bridges  
616 et al., 2016; [Lapotre et al., 2016a](#)).

617 These observations suggest that terrestrial aeolian impact ripples might not be good analogs for  
618 the Martian large ripples ([Lapotre et al., 2016a](#); [Silvestro et al., 2016](#); Vaz et al., [2016submitted](#)). As this  
619 gets worked out, previous studies will need to be carefully reviewed, such as where the interpretation  
620 has been that a multidirectional wind regime exists, perhaps triggered by the local dune topography or  
621 by larger topographic features (e.g., Jackson et al., 2015; Silvestro et al., 2011). Also, the presence of  
622 such large ripples on the dune's stoss side and their migration across the slipface (Figures 9-10) may  
623 alter the wind profile above the dune and the slipface dynamics, beyond the way that these processes  
624 are typically captured in dune evolution models applied to terrestrial dunes and their ripples (e.g., Ewing  
625 et al., 2016). Increased coverage of high-resolution images coupled with in-situ observations by rovers  
626 are necessary to progress understanding of the nature and dynamic of the martian large ripples. This is  
627 fundamental for understanding how these ripples can be used to constrain local wind directions and to  
628 tune sand flux estimations over the dunes.

629

### 630 **Summary of (5)**

631 **Data needed:** Higher-resolution images of dune field, reflecting variation over the dune, including in  
632 composition or granulometrics; mapping and analysis of second-order and higher-order bedforms (e.g.,  
633 ripples) and how these reflect the wind pattern around the dune.


634 **Knowledge gain:** Measurements of ripple movement and characteristics over the dune.

635 **Assumptions generated:** Use of the right analog features/models for interpretation of the smaller-scale  
636 features.

637 **Questions:** What is the local sand flux and wind patterns over the dunes (as reflected in ripple  
638 movement)? Are grains sorted within the ripples, and if so, why? Is ripple movement coupled  
639 with/connected with current dune evolution, or e.g. does ripple movement reflect a surficial mobile  
640 layer of sand over a relict dune core?

641 **Lead to investigations of:** Wind diversion around dune topography; observation/better understanding  
642 of local source regions.

643




644

645 **Figure 9. Observed ripple movement on Mars.** Images show (a) a rippled dome dune in the Bagnold  
646 dune field, Gale Crater, with (b) ripple migration over the dune stoss side between Mars years 28 (2006)

647 | and 29 (2008) [\(Silvestro et al., 2013\)](#). (c: T1-T2) The zoom-in shows one ripple (white arrow) moving over  
 648 | the dune brink, reflecting grain transport onto the slipface and suggesting that dune migration may also  
 649 | be occurring. HiRISE images shown: (a,c/T1) PSP\_001488\_1750 (taken 20 November 2006), (c/T2)  
 650 | PSP\_009650\_1755 (17 August 2008) (MRO/NASA/UA).

651



652

653 | **Figure 10. Ripple morphology on Mars.** (a-b) HiRISE images of High Dune, an active dune within the  
 654 | Bagnold dune field, Gale Crater, that has also been investigated by MSL Curiosity ([Bridges et al., 2016](#);  
 655 | [Lapotre et al., 2016a](#)). Visible in these orbital images is a complex ripple pattern on High dune's stoss  
 656 | slope, with what appears to be two sets of large ripples intersecting at right angles [\(Silvestro et al.,](#)  
 657 | [2016\)](#). This ripple configuration is typical of the Bagnold dunes and seems to be common on Mars. HiRISE  
 658 | images are: (a) ESP\_042682\_1755, (b) PSP\_009294\_1750. (MRO/NASA/UA) (c) This image of the High  
 659 | Dune stoss slope was taken by the Mast Camera on Curiosity, showing the complex morphology of these  
 660 | large ripples, which in this closer-inspection perhaps do not appear analogous to terrestrial sand ripples.  
 661 | The scalebar (one meter length) is for the lower portion of this cropped image. Image is from NASA  
 662 | Photojournal PIA20168.

663

664 | **2.6. Phase 6: Observation of dune activity (aeolian or otherwise)**

665 Only recently has it been observed that martian dunes and ripples are very actively migrating  
666 and evolving within the present-day climate ([Figure 9](#); [Bourke et al., 2008](#); Bridges et al., 2012a; 2012b;  
667 [Chojnacki et al., 2011; 2015](#); [Fenton, 2006](#); Geissler et al., 2013~~2~~; Silvestro et al., 2010b; 2011; 2013).  
668 Previously, an incongruence appeared in our understanding of present-day martian sand transport, as  
669 the morphology of many aeolian bedforms (apparently sharp crestlines of dunes and ripples) and a  
670 surface observation of saltation (Greeley et al., 2006) [and ripple movement \(Sullivan et al., 2008\)](#)  
671 suggested that some aeolian bedforms should be active. However climate models did not produce the  
672 wind velocities predicted for saltation processes to occur under present conditions and no bedform  
673 motion was observed within high~~er~~-resolution images [\(although some dome dunes were seen to](#)  
674 [disappear \(Bourke et al., 2008\)\)](#). This was taken to imply that martian dunes may be [stabilized \(e.g.,](#)  
675 [Zimelman, 2000\) and possibly](#) relict features of a past climate with a denser atmosphere (e.g., Breed et  
676 al., 1979), [and that surface](#) degradation processes must be slow. However, acquisition of a sufficient  
677 temporal baseline and careful comparison of overlapping high-resolution images now yield measurable  
678 and consistent changes in dune margin and ripple crestline locations through several fields [\(e.g.,](#)  
679 [Endeavor Crater: Chojnacki et al., 2015\)](#), and show that sand fluxes on Mars are comparable to ~~and in~~  
680 ~~some cases exceeding, terrestrial~~ sand fluxes [in the Antarctic Dry Valleys](#) (Bridges et al., 2012b). [Within](#)  
681 [Endeavor Crater, these martian sand fluxes are sufficient for dune turnover times to be much less than](#)  
682 [the time since known large climatic shifts \(e.g., an obliquity shift or increased atmosphere density\),](#)  
683 [implying that these dunes are not records of paleo-climate conditions \(Chojnacki et al., 2015\).](#)

684 These new observations, proving that sand is currently moving on Mars in large volumes and  
685 that at least some aeolian bedforms are presently active, were helpful in the advance of sediment flux  
686 models and understanding how sediment flux dynamics may vary on different planetary bodies. For  
687 example, an update to the model of steady state saltation (Kok and Renno, 2009) and application to  
688 Earth and Mars conditions (Kok, 2010) showed that saltation can be maintained on Mars by wind speeds

689 | an order of magnitude less than that required to initiate it, while nearly the same wind speed is needed  
690 | to both initiate and maintain saltation on Earth. This provides a viable explanation for why aeolian  
691 | bedforms appear to evolve at lower-than-predicted wind velocities (as well as an explanation for the  
692 | smaller-than-expected minimum dune size on Mars: Kok, 2010). Estimates of aeolian sand flux (in the  
693 | present or past) are important as they feed into models of surface erosion rates (e.g., Golombek et al.,  
694 | 2006; 2014).

695 | Sand dunes on Mars are also subject to other processes in the present-day. For example, in  
696 | particular, small- and large-scale gully alcove-apron and alcove-channel-apron (i.e., gully) formation  
697 | activity has been observed in southern mid-latitude dune fields (Figure 8; Diniega et al., 2010b; Dundas  
698 | et al., 2012; 2015) and similar alcove-fan formation activity has been observed in the north polar dune  
699 | fields (Hansen et al., 2011; 2015; Horgan and Bell, 2012), moving large volumes of sand downslope and  
700 | possibly contributing to the overall migration of the dunes. Some have proposed that martian gully this  
701 | activity may have aeolian drivers (Horgan and Bell, 2012; Treiman, 2003), but most studies have shown a  
702 | seasonal control on the timing of gully activity feature formation and evolution, possibly related to CO<sub>2</sub>  
703 | frost processes (Diniega et al., 2010b; Dundas et al., 2012; 2015; Hansen et al., 2011). It is also possible  
704 | that both aeolian and seasonal frost processes have an influence on these types of dune modification  
705 | activities (Hansen et al., 2015). Regardless of underlying process, these changes are actively modifying  
706 | the dune slopes (Allen et al., 2016; Diniega et al., 2016; Hansen et al., 2011) and thus need to be  
707 | investigated and explained to form a complete story for the martian dune evolution and accurate  
708 | interpretation of observed dune morphology.

709 | It is also important to note that some dunes have features indicative of a lack of activity, such as  
710 | fissures on north polar dunes (Portyankina et al., 2012) and pits and softened topography on southern  
711 | mid-latitude dunes (Fenton and Hayward, 2010). Such evidence for stability can provide constraints on



712 | the current availability of mobile material and the near-surface wind environment, as well as a contrast  
713 | with the conditions when the (now inactive) bedform had evolved.

714

## 715 | **Summary of (6)**

716 | **Data needed:** Repeat images of sufficient spatial and temporal resolution to detect (and measure)

717 | changes in surface morphology (or lack thereof).

718 | **Knowledge gain:** Observation and constraints on the estimated (average/net) amount of (and possibly,  
719 | measurement) of sediment transport.

720 | **Assumptions generated:** Activity rates observed in the present-day can be extrapolated to past times  
721 | and conditions.

722 | **Questions:** What other processes are contributing to dune evolution? How much sediment is moving  
723 | within the present climate? Has that amount of aeolian sediment transport changed since a previous  
724 | climate?

725 | **Lead to investigations of:** How the estimated sediment transport may affect surface erosion rates  
726 | (including formation of sand) and formation of other aeolian features such as yardangs; How the climate  
727 | has shifted, if changes in sediment transport are apparent.


728

## 729 | **2.7. Phase 7: Groundtruth measurements**

730 | To-date, we have only visited – at ground-level and up-close – dunes on one planet other than  
731 | the Earth. While various Mars rovers have in situ imaged sand deposits and ripples (e.g., Greeley et al.,  
732 | 2006; JPL, 2012; 2014; Sullivan et al., 2005), Curiosity’s visit to Bagnold Dune Field is the first in situ  
733 | observation of dunes and dune sand (JPL, 2015; Bridges et al., 2016). This rover has examined dune sand  
734 | on several different slopes on and around dunes slopes thought to be undergoing different levels of  
735 | aeolian activity (based on orbital observations of ripple migration and the strength of spectral signatures

736 of dust cover (Lapotre et al., 2016b)). Within even the first observations of dune sand (scooped from the  
737 stoss slope and imaged on the lee slope; Figure 11), grain size differences have been noted that are  
738 perhaps correlated with differences in grain composition (as grains of different sizes appear to  
739 correspond to different materials) (Achilles et al., 2016; Cousin et al., 2016; Ehlmann et al., 2016).  
740 Images of the lee slope of the more active “High dune” have yielded many grainflow features and  
741 potentially evidence of some level of induration (Figure 12; Ewing et al., 2016) – none of which ~~was~~were  
742 visible in the orbital images. The first results of this work are currently being put together, and we look  
743 forward to learning more about the first in situ investigated extraterrestrial dunes.

744



745

746 **Figure 11. Image of the undisturbed surface within the base a Martian sand dune called "High Dune"**  
747 *visited by NASA's Curiosity rover. The image covers an area 3.6 by 2.7 centimeters. Grain sizes show*  
748 *some range, but a fairly consistent size – comparable to dune sand on Earth. It was taken by the Mars*  
749 *Hand Lens Imager (MAHLI) camera on the Curiosity rover's arm on Dec. 5, 2015. Image and description*  
750 *are from JPL (2015).*

751



752

753 **Figure 12: An active dune slipface on Mars, imaged by NASA's Curiosity Mars rover. Multiple grain**  
754 **flows, slumps, and ripples are visible on the slipface of "Namib Dune," a dune within the "Bagnold**  
755 **Dunes" field along the northwestern flank of Mount Sharp, Gale Crater. None of these fine details are**  
756 **visible from orbital (HiRISE) images. The overall slope is 26-28°, and 4-5 m in height. This view combines**  
757 **multiple images from the telephoto-lens camera of the Mast Camera (Mastcam), taken on Dec. 21, 2015.**  
758 **The scene is presented with a color adjustment that approximates white balancing, to resemble how the**  
759 **sand would appear under daytime lighting conditions on Earth. Image and description are from NASA**  
760 **Photojournal PIA20283.**

761

## 762 **Summary of (7)**

763 **Data needed:** In situ observations of the dune and dune sand (possibly from different portions of the  
764 dune); possibly observations of saltation on the dune or grainflow on the slipface.

765 **Knowledge gain:** Size, composition, and other characteristics of grains involved in saltation.

766 **Assumptions generated:** That the observed characteristics and activity are not anomalous, in time and  
767 space (i.e., the observation didn't catch a rare circumstance/event).

768 **Questions:** Why do the grains look as they do, and what causes the variation/distribution in grain size?  
769 How representative is this observed dune's characteristics and activity? What causes the dune's  
770 features? How can observations of sediment grain size and bedform morphology provide insight  
771 regarding transport processes and the nature/frequency of mobilization events?

772 **Lead to investigations of:** Models of dune activity and evolution, and generation of the observed sand  
773 grains (extending or perhaps redirecting previous work); Based on in situ observation of features that  
774 may not be visible from orbit, what is implied about dune activity and characteristics and how does  
775 information that feed back into models of dune evolution (and what assumptions and related results  
776 should be re-evaluated).

777


### 778 3. Discussion

779 There are some differences between the aeolian science investigations undertaken on each  
780 planetary body – in particular as study methods of more recently studied bodies can build from lessons  
781 learned in aeolian bedform studies of a previously observed body, and as overall our understanding of  
782 aeolian processes becomes more refined as models are forced to reconcile with a wider range of  
783 environmental and planetary conditions. But, as has been described, studies of aeolian bedforms on  
784 planets (other than Earth) broadly tend to follow a similar pattern of gained knowledge, generated  
785 assumptions, and follow-up investigations (that rely on the new knowledge and assumptions). The  
786 similarities in the history of aeolian science over different planetary bodies (Figure 13) are due partially  
787 to the knowledge-advancement at each body being based on the same types of data. Such data is  
788 usually acquired in the same order, which is based on the way in which higher-resolution and increased

789 coverage are acquired during extended or subsequent missions, and as concepts and investigations  
 790 mature and become more specific within all areas of planetary exploration.

791 Within that progression, we focus here on the gaps that seem likely to occur for any planetary  
 792 body. We then move beyond planetary aeolian studies, to look at the interplay of planetary aeolian  
 793 bedform studies with investigative fields that follow their own sequences of discovery and refinement:  
 794 aeolian process modeling and terrestrial aeolian studies.

795



796

797 **Figure 13. Timelines showing movement into and through different investigation phases, for Mars,**  
 798 **Titan, and Venus.** *Darker colors indicate publication dates for relevant studies (as referenced within this*  
 799 *paper), and lighter colors indicate the general period of activity (again, based The timespans for the*  
 800 *investigations phases are based on publication dates references within this paper and checked against*  
 801 *google.scholar search results with keywords e.g., “Venus dunes”). Under Mars, Phase 6 is divided into*  
 802 *observations of “no dune activity due to aeolian bedform processes” (6/n) and of such activity (6/a).*  
 803 *Within Phase 1 for Titan and Venus, publications predicting a limit on aeolian bedform formation on*  
 804 *those bodies are highlighted with a P – and the first observation of a bedform occurs later. Arrival dates*

805 | *of relevant spacecraft, to the planetary body, are included – for obvious reasons, these often initiate or*  
806 | *re-invigorate investigations begun during a previous mission.*

807

### 808 **3.1. Gaps that can form within the planetary aeolian science sequence of investigations**

809           Over time and as more data is acquired, our understanding of aeolian processes and  
810 interpretation of the aeolian landforms builds. However, as that understanding builds, it is important to  
811 | keep track of which building blocks are assumptions and not actual observations. ~~While~~ it is necessary  
812 | for assumptions to be made to keep the science investigations moving forward and to guide  
813 | development of the next set of investigations, but an assumption that is treated like an “observation”  
814 | can lead to models with unrecognized limitations, which in turn can lead to incorrect interpretations of  
815 new observations or even a lack of attention paid to “contradictory” observations. Thus, assumptions  
816 should be recognized as such (and not treated as data) and be re-evaluated for consistency with new  
817 | and different data, until direct measurement of the assumed variable or process is possible – doing this  
818 | can make it easier to identify and investigate intriguing new understandings about processes and  
819 conditions.

820           Several examples of areas where new information has supplanted previous assumptions have  
821 been mentioned within the discussion of the investigation phases (Subsections 2.1-7). Some additional  
822 examples:

- 823 • As higher-resolution and more detailed studies are completed about specific dune fields, results of  
824 these studies (Phases 5-7) should be inserted into field (Phases 2 & 3) and global studies (Phase 4)  
825 that previously relied on lower-resolution or less complete data and assumptions about form and  
826 process uniformity (in time and space) through the field. As was discussed under Phase 7, the  
827 martian large ripples are a new example of this -- where in situ observations are drawing into  
828 question previous work done regarding the scaling of aeolian bedform size between Earth and Mars

829 and interpretations of ripple crestline complexity, that had been based on interpretation of orbital  
830 images. In general, as more detailed studies are conducted over specific martian dune fields, it is  
831 important to regularly consider how those results fit within the results of larger-scale studies.

- 832 • These observations have also led into a model of a potential new mode of subaerial bedform  
833 migration and evolution (Lapotre et al., 2016a). As discussed in Phase 5, on the stoss slope of Namib  
834 dune, two types of ripples have been imaged: The large ripples (few m-wavelength) were previously  
835 observed in orbital images, and were thought to be analogous to the wind ripples that we see on  
836 the slopes of terrestrial dunes (e.g., Sullivan et al. 2008). However these large ripples have very  
837 different morphology and dynamics (Silvestro et al., 2016) and in fact are superimposed by small  
838 ripples (~10s cm wavelength) that have morphology more similar to terrestrial sand-impact ripples  
839 (but were not visible within orbital images; Figure 10). The large ripples are now hypothesized to be  
840 fluid-drag ripples (Bagnold, 1951; Wilson, 1973), which on Earth form under water, but on Mars are  
841 able to develop sub-aerially because of the higher kinematic viscosity of the low density atmosphere  
842 (Lapotre et al., 2016a) a form of aeolian bedform that does not form on the Earth, due to our much  
843 thicker atmosphere. Although this model is still being worked out and debated, this example has  
844 already shownshows that the limitations of analysis from only orbital imagery resolution need to be  
845 remembered, even when we think that we understand what we are looking at. Additionally, the  
846 limits of comparative planetology can mean we misinterpret observations where we lack a  
847 terrestrial analog.
- 848 • Dune sSand grain sizes on Mars have been estimated since dunes were first seen, based on  
849 assumptions about conditions for dune formation (Phases 1 & 2). Efforts to “measure” grain sizes  
850 from proxy thermal inertia estimates have also been undertaken (mentioned within Phase 4), and  
851 compared to and debated against the assumptions about the ability of the wind to move grains of  
852 different sizes. Studies based on these estimations, and their results, now should be re-evaluated as

853 Curiosity has recently completed the first in situ investigation of a dune located on a planet other  
854 than Earth (Phase 7), yielding the first direct measurements of martian dune grain sizes (Figure 11).  
855 While on Titan sand grain sizes have not yet been measured in situ (and won't be in the near-  
856 future), studies have explored what grain sizes can be reached via feasible physical processes which  
857 puts constraints on models of dune formation conditions, and visa-versa.

858 Additionally, it is important to recognize the gaps and limitations that can occur in aeolian  
859 studies if only the "standard" aeolian science inputs are considered (e.g., the "complementary sciences"  
860 listed in Table 1 also need to be considered). As in all areas of planetary science and geology, it is  
861 important to consider many pieces of information (and observations, as possible), and all need to be  
862 consistent with the model for the model to be validated. For example, while potential sediment sources  
863 can possibly be tracked from visible imagery, climate models, and/or topography models, compositional  
864 information about the dune grains and the potential sand sources is needed to check that the model is  
865 consistent with the full environment. This may extend beyond compositional information in the local  
866 environment (which was included in Phase 2), as grains may have been transported over large distances  
867 or have been recycled a few times – and this history may not be apparent without a broad-swath of  
868 environmental information. Additionally, processes outside of standard, dune-forming aeolian  
869 processes may be playing a role in dune evolution and observed morphology. For example, the dunes in  
870 the martian polar regions have morphologies and features different from those in the equatorial region,  
871 making it clear that polar processes are altering the aeolian bedforms and thus must be considered in  
872 their interpretation (e.g., in the north: Hansen et al., 2011; 2015; in the south: Fenton et al., 2010).

873

### 874 **3.2. Connections to modeling of the physical processes**

875 As discussed above, looking at aeolian bedforms on other planets allows models to be tested  
876 against a range of environmental and planetary conditions. From that, we refine our understanding of



877 aeolian processes without assumption of Earth-conditions. This can especially have a large impact on  
878 models of the small-scale and complicated dynamics of sand-wind and sand-sand interactions. For  
879 example, as discussed under Phases 2 and 6, our understanding of the way in which sand is picked up by  
880 the wind, causing or continuing saltation, has now been “tested” under terrestrial, martian, and  
881 venusian conditions (Kok, 2010; 2012), resulting in an updated model of how saltation and reptation are  
882 initiated and interact.

883         On Titan, questions about how “sticky” organic sand particles would interact with the wind  
884 were part of an investigation to explain how the dunes had formed, and from that to connect the  
885 crestline orientation to the forming-wind direction(s). The Titan dune sand color appears consistent with  
886 a composition of organics, and such long-chain molecules (of as-yet undetermined exact composition)  
887 could be derived from the atmospheric photodissociation of methane, which creates small particles  
888 (Carl Sagan’s “tholins”) that snow down from the atmosphere (and then perhaps get incorporated into  
889 surface sedimentary layers or clump together into larger granules, that are eroded and transported to  
890 the dune-forming regions) (Radebaugh, 2013; Barnes et al., 2015). Studies of clay-rich dunes in China  
891 had revealed that “sticky” particles could form dunes, but would anchor themselves to the downwind  
892 | edge of a longitudinal dune and thus progress-grow and migrate the dune -in their transport direction,  
893 | along the dune crestline; this was proposed as a potential analog to the Titan dunes (Rubin and Hesp,  
894 | 2009). Although the Titan dune morphologies were overall found to be generally more consistent with  
895 | freely-moving particles (i.e., the saltation more usually observed on Earth) and thus this longitudinal  
896 | dune formation model is less favored than the model discussed in Subsection 2.2. This type of  
897 | questioning highlighted a different type of terrestrial dune-formation mechanism and “tested” behavior  
898 | of the traditional dune formation model if one does not assume a non-cohesive sand grain. This led to  
899 | further development of a dune-wind alignment model that brought these two hypotheses together as  
900 | well as explained how bedforms with different alignments can exist within the same multidirectional

901 wind regime (Courrech du Pont et al., 2014). Within this single model, dune alignment reflects growth  
902 via either a “bed instability mode” (which approximates the longitudinal dune growth process proposed  
903 by Rubin and Hunter (1987) and Rubin and Ikeda (1990)) or a “fingering mode” (the growth process  
904 proposed by Rubin and Hesp (2009)), depending on sediment availability.

905         Models that examine larger-scale dynamics can also be tested through application to different  
906 planetary surfaces. For example, it was in studying martian dunes that a discrepancy was noticed  
907 between the minimum dune size expected on that planet (~100x the minimum Earth dune size) and that  
908 observed (~10x), thus driving new models of dune formation to explain the scaling factor. Model studies  
909 aiming to replicate the observed minimum barchan dune size on Earth and Mars addressed this  
910 question, and tested assumptions about how saltation, reptation, and wind drag interact in setting  
911 characteristic sand trajectory distances, and from this the generation of instability within a sand bed  
912 under a moving fluid (Claudin and Andreotti, 2006).

913

### 914 **3.3. Connections to terrestrial studies and knowledge gain**

915         The trajectory of terrestrial dune studies has differed markedly from the framework proposed  
916 here for planetary dune studies. In essence, the difference is one of top-down vs. bottom-up approaches  
917 as in situ observations of terrestrial landforms, conditions, and activity are significantly easier to carry  
918 out. However, this has not resulted in the history of terrestrial dune fields being an opposite to the  
919 sequence suggested as being characteristic of planetary dune research. The earliest published studies of  
920 terrestrial dune fields were linked with exploration by non-indigenous people, and many of the founding  
921 points of contemporary dune science can be traced to these expeditions. The exploration of the  
922 southern African and Australian interior (mid-19th century), the Sahara (around the beginning of the  
923 20th century, mostly by the French in the west and the English in the east) and the Arabian Rub al’Khali  
924 (most notably by Wilfred Thesiger in the late 1940s) all had exploration as their primary goals. As with

925 contemporary rover exploration of the martian dune fields, many dune fields were approached with  
926 trepidation due to the hazards they posed. Despite science being incidental rather than implicit to most  
927 of the explorations, there was, nonetheless, early recognition of the great spatial extent of many dune  
928 fields, the remarkably organized nature of dunes and the fact that dunes could exist at differing activity  
929 levels.

930           Although Bagnold's work in the 1930s and 1940s is most commonly cited as being the  
931 foundation of modern understanding of aeolian processes and landforms, there were significant  
932 precursors. George Perkins Marsh (1864) considered geoengineering problems associated with drifting  
933 sand, and the role of vegetation in stabilizing dunes, and Russian geologist Nikolay Sokolów had  
934 discussed dune sedimentology and theories of dune formation in a 300 page book devoted to the  
935 subject (1894). Georges Rolland, a French mining engineer, set out a series of propositions in 1890  
936 based on fieldwork in the Algerian Sahara which addressed such issues as sediment source, the  
937 distribution of dune fields, varying levels of dune activity and the relationship between wind regime and  
938 different dune shapes (Burt et al., 2008). At this point, the role of the wind in dune formation was still  
939 contested by many, and it was widely held that dunes would prove to have rocky cores (Goudie, 2002).  
940 Many other aspects of contemporary aeolian science date from surprisingly early studies. Aerial imagery  
941 was used to examine dune planform morphology in the 1930s (Aufrère 1932, Madigan 1936), and the  
942 recognition of dunes as a particulate waveform in a fluid medium can be traced to the work of Cornish  
943 (1914). Bagnold's work, utilizing field and wind tunnel experimentation, is an early example of the  
944 'quantitative revolution' widely recognized in geosciences in the middle of the 20th century. This directly  
945 influenced the next half-century of research, via fieldwork and laboratory experimentation, in a phase  
946 perhaps best summarized by Lancaster's (1995) state-of-the-art textbook. Coincidentally, the same year  
947 saw the publication of Werner's (1995) application of cellular automata models to aeolian bedforms,  
948 which accepted that dunes formed as an emergent property of a complex system, one of the first

949 indications of the failure of reductionist approaches to fully explain aeolian landscapes (Livingstone et  
950 al., 2007). The same period saw the rise of the use of luminescence dating to provide ages for dune  
951 emplacement, since described as having had a transformative effect on studies of dryland science  
952 (Singhvi and Porat, 2008).

953           Planetary studies of aeolian dunes therefore have the advantage of decades of terrestrial work  
954 to draw upon, and this is reflected in the very rapid progress made on newly-discovered dunes (e.g.  
955 Titan, Comet 67P). Terrestrial science, conversely, has had the advantage of a relatively steady  
956 progression in the quality of the available data -- although the related understanding of aeolian systems  
957 has not progressed as steadily. The progress made in understanding terrestrial dunes has not been  
958 without challenges, and it is instructive to reflect on whether there are lessons for the planetary  
959 community can be drawn from progress on terrestrial dune fields, and conversely whether the evolution  
960 of extraterrestrial dune research can inform the research strategies of Earth's dune studies.

961

### 962 ***What can planetary science learn from the history of terrestrial dune studies?***

963           Much of planetary dune science is already directly influenced by the methods, theory and  
964 process understanding derived from terrestrial studies, manifest in the numerous analog studies.  
965 However, there are some less well-discussed points that are worthy of consideration.

966           As was noted in section 3.1, close attention must be paid to the difference between assumed  
967 and observed knowledge. Cautionary tales can be drawn from terrestrial dune studies, and this is  
968 perhaps best illustrated by the roll vortex hypothesis for longitudinal (linear) dune formation. First  
969 proposed by Bagnold (1953), and promoted subsequently (e.g. Hanna, 1969) this suggested that  
970 thermally induced vortices in regional wind-flow would lead to the development of helical horizontal  
971 flow cells that might lead to sand accumulation in linear bedforms extending downwind. The theory is  
972 strikingly devoid of empirical supporting evidence, and yet still persists in the literature. Quite simply,

973 vortices of sizes that might explain dune spacing have never been observed despite numerous  
974 experimental attempts, and the transverse component of roll vortices does not appear to have sufficient  
975 velocity to move sand (Lancaster 1995). Planetary studies should be careful to question existing  
976 paradigms and theories, and be willing to point out when data do not support these hypotheses.

977 Bagnold's great advances in aeolian science can be largely attributed to willingness and  
978 fearlessness towards innovation, in terms of methods and physical exploration. The novel application of  
979 wind tunnels to aeolian transport and sedimentation and the methods developed to enable remote  
980 desert travel directly enabled the advances in understanding that Bagnold brought. Planetary  
981 perspectives support this, with the radical advances in data brought from missions such as MSL,  
982 Cassini/Huygens, Rosetta/Philae and New Horizons. Such evidence supports the potential knowledge  
983 gains from similarly ambitious mission concepts of other planetary surface exploration missions, such as  
984 AVIATR (Barnes et al. 2012) and VISE (NRC, 2013) which have been considered for Titan and Venus  
985 exploration, respectively. The evidence from both terrestrial and planetary dune studies suggests that  
986 high-risk, innovative research has led to some of the greatest advances.

987 The discrepancy between the timescales of aeolian process and the timescales evident in  
988 aeolian landscapes is also very evident – possibly even more so – on some planetary bodies. Despite  
989 processes operating within dune landscape on timescales of seconds to hours, the resultant landscape  
990 development frequently operates on timescales of  $>10^3$  years. Dating of aeolian sediment, primarily via  
991 the suite of luminescence dating methods, has been adopted very widely on terrestrial dune studies,  
992 and has played a crucial role in linking the short-term process understanding with the long-term  
993 geomorphological record. It has enabled calculation of rates of landform evolution beyond that possible  
994 using observational records (e.g. Kocurek et al., 2007; Telfer, 2011), revealed complex spatial variability  
995 in aeolian accumulation (Telfer and Thomas, 2007) and frequently been used to infer external drivers of  
996 dune activity (e.g. climatic changes). Experimentation with luminescence readers suitable for Mars

997 missions has been explored (e.g. McKeever et al., 2003; Jain et al. 2006), and if the substantial technical  
998 challenges can be overcome (Doran et al. 2004), martian luminescence dating offers the potential to  
999 extend understanding of accumulation beyond the period of direct observation. Recent progress  
1000 suggests that solutions may exist to these challenges (e.g. Sohbaty et al., 2012).

1001 ~~————— Much of the focus of this paper has been on the increasing availability, resolution, and coverage~~  
1002 ~~of new remotely sensed data. The same has been true for Earth, and here lessons learned in planetary~~  
1003 ~~studies help guide interpretation of terrestrial images at the margins of the spectral and/or spatial~~  
1004 ~~resolution of the imagery. This is perhaps best illustrated with the example of the highly contested issue~~  
1005 ~~of extensive palaeo-dune fields in the Amazon Basin, presently covered with extensive tropical forest.~~  
1006 ~~Tricart (1974), working with recently released first-generation Landsat imagery, identified widespread~~  
1007 ~~stabilized aeolian landforms in the tropical Amazon Basin. This interpretation was repeated numerous~~  
1008 ~~times (e.g. Klammer, 1982) and led to a widespread belief in arid phases accompanied by vegetation~~  
1009 ~~loss during the late Quaternary evolution of the region, with huge implications for understanding of~~  
1010 ~~regional biogeography in one of the world's most biodiverse regions. However, whilst subsequent~~  
1011 ~~reanalysis of high-resolution data has revealed aeolian dune fields around the margins of the Amazon~~  
1012 ~~basin (e.g. Teeuw and Rhodes, 2004; May, 2013) and/or immediately adjacent to large rivers where the~~  
1013 ~~sediment supply has at times been the dominant control (e.g. Carneiro, 2002), the existence of wide~~  
1014 ~~swathes of paleodunes across the Amazon basin has not withstood closer scrutiny (Colinvaux et al.,~~  
1015 ~~2001). Tripaldi and Zarate (2014) reviewed the evidence, and demonstrated the importance of~~  
1016 ~~groundtruthing when image interpretation is challenging. Planetary geomorphologists are usually~~  
1017 ~~admirably conservative in terms of implying process from (apparent) landform, especially when imagery~~  
1018 ~~is at the limit of its spatial or spectral resolution, and terrestrial incidents such as the question of an arid~~  
1019 ~~Amazon suggest that such conservatism is wise.~~

1020

1021 ***What can terrestrial dune studies learn from the history of planetary science?***

1022           Although at least parts of Phases 1-3 and 5-7 have been investigated on Earth for 70 years or  
1023 more, a striking difference between planetary and terrestrial dune studies is that currently there is no  
1024 global catalogue of dunes for Earth (Phase 4). The first editions of the Mars Global Digital Dune Database  
1025 (Hayward et al., 2007) was published within six years of the start of THEMIS data collection (which are  
1026 used to identify the thermal inertia proxy identifiers for dune fields), and global mapping of Titan's  
1027 dunes within the constraints of the available data (Lorenz and Radebaugh, 2009) was published within a  
1028 similar timeframe since the arrival of Cassini at Titan. Although some terrestrial regions have been  
1029 mapped and duneforms catalogued (e.g., Namib; Livingstone et al., 2010), and a global database of  
1030 dunes with dating constraints has recently been compiled (Lancaster et al., in press), global-scale  
1031 consideration of terrestrial dune fields is currently lagging behind planetary science. Efforts in this  
1032 direction are currently in progress (Hesse et al., 2015) -- but it has been over 40 years since the advent  
1033 of global terrestrial satellite coverage. The focus of planetary global catalogues of dunes has been on  
1034 understanding global circulation patterns (e.g. Charnay et al. 2015, Ewing et al. 2015), sediment sources  
1035 (e.g. Tirsch et al., 2011), identification of large-scale variations in dune form due to different evolution  
1036 processes or rates (e.g., Hayward and Fenton, 2010; Savage et al., 2014), and targeting areas for detailed  
1037 study (e.g., Hayward, 2011). Whilst it is not necessary to use dune morphology to understand modern  
1038 circulation patterns on Earth, applications of such a database would include quantification of aeolian  
1039 sediment volumes and flux, improved understanding of regions where dunes are currently stabilized,  
1040 and potential for monitoring change in environmentally-sensitive, dynamic landscapes.

1041           Livingstone et al. (2007), reviewing the state of understanding of terrestrial dune  
1042 geomorphology, concluded that decades of largely inductive, and increasingly reductionist, study had  
1043 not brought the completeness of understanding that had been hoped, and that integration of  
1044 methodologies (field, modeling and remote sensing) offered the best prospects for knowledge. Perhaps

1045 due to the difficulties in conducting 'field study' of extraterrestrial dunes (i.e., Phase 7), which are only  
1046 very recently being overcome on Mars, such combined strategies are often exemplified by planetary  
1047 aeolian studies, where studies employing a wide range of methodologies including numerical modeling,  
1048 laboratory experimentation, field study (presently via analog environments), and remote sensing are  
1049 commonplace (e.g., Lucas et al. 2014). Although some terrestrial studies do synthesize such diverse  
1050 methodologies, the example set by many planetary studies is a good one for terrestrial dune studies.

1051 Much of the focus of this paper has been on the increasing availability, resolution, and coverage  
1052 of new remotely sensed data. The same has been true for Earth, and here lessons learned in planetary  
1053 studies help guide interpretation of terrestrial images at the margins of the spectral and/or spatial  
1054 resolution of the imagery. This is perhaps best illustrated with the example of the highly contested issue  
1055 of extensive palaeo-dune fields in the Amazon Basin, presently covered with extensive tropical forest.  
1056 Tricart (1974), working with recently-released first-generation Landsat imagery, identified widespread  
1057 stabilized aeolian landforms in the tropical Amazon Basin. This interpretation was repeated numerous  
1058 times (e.g. Klammer, 1982) and lead to a widespread belief in arid phases accompanied by vegetation  
1059 loss during the late Quaternary evolution of the region, with huge implications for understanding of  
1060 regional biogeography in one of the world's most biodiverse regions. However, whilst subsequent  
1061 reanalysis of high-resolution data has revealed aeolian dune fields around the margins of the Amazon  
1062 basin (e.g. Teeuw and Rhodes, 2004; May, 2013) and/or immediately adjacent to large rivers where the  
1063 sediment supply has at times been the dominant control (e.g. Carneiro, 2002), the existence of wide  
1064 swathes of paleodunes across the Amazon basin has not withstood closer scrutiny (Colinvaux et al.,  
1065 2001). Tripaldi and Zarate (in press) reviewed the evidence, and demonstrated the importance of  
1066 groundtruthing when image interpretation is challenging. Planetary geomorphologists are usually  
1067 admirably conservative in terms of implying process from (apparent) landform, especially when imagery



1068 [is at the limit of its spatial or spectral resolution, and terrestrial incidents such as the question of an arid](#)  
1069 [Amazon suggest that such conservatism is wise.](#)

1070

#### 1071 **4. Conclusion/Summary**

1072           Studies of aeolian bedforms over a wide range of planetary bodies have resulted in significant  
1073 progress in our understanding of past and present climate and surface conditions, physical processes,  
1074 and the interconnectivity of dynamics over a range of spatial and temporal scales. These studies  
1075 contribute, in meaningful and often unique ways, towards a range of planetary science investigations.  
1076 For example, as discussed, interpretation of dune morphology often provides unique, if proxy,  
1077 groundtruth data about past or present wind conditions, and the proven presence of a large amount of  
1078 sand grains can drive investigations about processes responsible for creating such grains. Beyond studies  
1079 that involve this type of direct interpretation of the aeolian bedforms, aeolian science studies also yield  
1080 information about many tangentially-related areas of investigation. In particular, aeolian-driven sand  
1081 flux appears to be an important force in erosional modification of a planetary surface. Quantitative  
1082 estimations of wind speeds and sand flux and identification of sediment transport pathways yield  
1083 quantitative estimates of erosional process rates. This can, for example, lead to improved interpretation  
1084 of observed landforms – such as yardangs (e.g., Ward 1979), or the rate of crater degradation by aeolian  
1085 processes which is important for accurately estimating the age of a planetary surface (e.g., Golombek et  
1086 al., 2014; Grant et al., 2006; 2008; 2016). This can also provide bounds on surface-ages of exposed rock  
1087 surfaces, which is can be of importance to rover missions – such as Mars missions searching for  
1088 reachable-environments near the surface that may have been habitable and that may include preserved  
1089 biosignatures (e.g., Arvidson et al., 2015).

1090           As discussed, planetary aeolian studies have also made key contributions towards improving the  
1091 methodologies employed in aeolian science, and in challenging assumptions built (perhaps

1092 inadvertently) into aeolian process models based on terrestrial observations. To-date, this has resulted  
1093 in the refinement of several models of dune-field forming processes, from interactions between sand  
1094 grains and the wind or with each other, up through interactions between dunes and topography and  
1095 climate shifts.

1096           Given all of the ways in which our aeolian study results impact our understanding of planetary  
1097 surface conditions and histories (as well as the Earth's), it is thus very important to critically look at how  
1098 we progress in planetary aeolian science, and in particular to consider carefully which results (and  
1099 resultant models) are based on assumptions versus observations – and then to revisit those results  
1100 when new information becomes available. Here, we have proposed one framework for identifying  
1101 progress within planetary aeolian studies, and have used that framework to chart the progression of  
1102 data, assumptions, and generated knowledge. We hope this framework, and our identification of gaps,  
1103 will help future planetary aeolian researchers strategically fill knowledge gaps or at least carefully  
1104 recognize where assumptions are being used to progress a study.

1105           Additionally, this framework may help identify the types of data that would be most useful for  
1106 future planetary missions. Pluto, Io, and Comet 67P were all discussed as having reached Phase 1, where  
1107 at least a potential aeolian bedform has been observed. On Titan, global datasets exist and have  
1108 contributed to large shifts in our understanding of the Titan climate and organic cycles. Venus also has a  
1109 global topography dataset, but the low resolution and apparent lack of dune fields stalled progress in its  
1110 aeolian science investigations (and thus related advancements in planetary surface studies).  
1111 Unfortunately for Venus and Titan, further progression within Phases 2-4 (and movement into Phases 5  
1112 and beyond) will likely need to wait for new and higher-resolution surface datasets.

1113           Mars' aeolian bedforms are the best studied within planetary aeolian science (outside of  
1114 Earth's), with both widespread coverage in certain data-types and many regions with high-resolution  
1115 data regarding the dunes' and dune field environment's morphology and composition, collected over

1116 the past 43 years. However, with the progress that has been made, we cannot lose sight of the fact that  
1117 much of it has been built on interpretations of remote data. (As discussed under Phase 7, in situ duene  
1118 field observations have not been possible until just recently.) Furthermore, much of the work involves a  
1119 meshing of coarse global data with a few more-deeply monitored and studied dune fields, and thus  
1120 much extrapolation is done that assumes certain types of consistency between fields. This is an odd  
1121 contrast with Earth dune field studies, where the global dataset (Phase 4) is what is missing.

1122 For all planetary bodies (including Earth), we look forward to further advancements in the  
1123 interpretation of aeolian bedforms and what interpretations about those bedforms will imply about the  
1124 environmental conditions and processes. If history is to be any guide, with each advance into a new  
1125 phase (due to acquisition of a new type of data and/or enablement of a new type of analysis), we find  
1126 exciting new understandings about that planetary body and the general understanding of aeolian  
1127 processes. One area of intriguing advancement is the prediction of where dunes and/or ripples could be  
1128 found (which could be thought of as a “Phase 0” within our framework). As we explore more bodies and  
1129 learn more about the conditions under which bedforms resembling aeolian dunes are found, we can  
1130 wonder about the next place where we may expect to find potential dunes, as well as identify lessons to  
1131 aid in such predictions (e.g., when we return to Venus). In addition, Perhaps in the near future, we will  
1132 move into a yet-undefined Phase 8 (e.g., through comparison between in situ measurements of some  
1133 very different types of aeolian bedforms? Hints of that are starting with sand grain comparisons (e.g.,  
1134 O’Connell-Cooper, 2016)), yielding a new type of data that can supersede assumptions made in Phases  
1135 1-7, further expanding our broad understanding of aeolian processes and bedforms, and increasing the  
1136 overall information gained from planetary aeolian studies.

1137

#### 1138 **Acknowledgements**

1139 [Mostly removed for review.] We thank the two reviewers for their very helpful comments, and in  
1140 particular for the suggestions of additional references.

1141

1142 **References**

- 1143 Achilles, C.N., D.T. Vaniman, D.F. Blake, T.F. Bristow, E.B. Rampe, D.W. Ming, S.J. Chipera, R.V. Morris,  
1144 S.M. Morrison, R.T. Downs, K.V. Fendrich, B.L. Ehlmann, A.S. Yen, P.C. Sarrazin, A.H. Treiman, P.I.  
1145 Craig, M.G.A. Lapotre, K.S. Edgett, R. Gellert, J.A. Crisp, J.M. Morookian, J.P. Grotzinger, D.J. Des  
1146 Marais, J.D. Farmer, 2016. Mineralogy of eolian sands at Gale crater. 47<sup>th</sup> LPSC, Ab. 2532.
- 1147 Allen, A., S. Diniega, C. Hansen, 2016. Gully and aeolian activity within the “Tleilax” dune field in the  
1148 Olympia Undae, Mars. 47<sup>th</sup> LPSC, Ab. 1759.
- 1149 Andreotti, B., P. Claudin, O. Pouliquen, 2006. Aeolian Sand Ripples: Experimental Study of Fully  
1150 Developed States. Physical Review Letters, v. 96, 028001, doi:10.1103/PhysRevLett.96.028001.
- 1151 Arvidson, R. E., J.F. Bell III, J.G. Catalano, B.C. Clark, V.K. Fox, R. Gellert, J.P. Grotzinger, E. A. Guinness,  
1152 K.E. Herkenhoff, A.H. Knoll, M. G. A. Lapotre, S.M. McLennan, D.W. Ming, R.V. Morris, S.L. Murchie,  
1153 K.E. Powell, M.D. Smith, S.W. Squyres, M.J. Wolff, J.J. Wray, 2015. Mars Reconnaissance Orbiter and  
1154 Opportunity Observations of Burns Formation and Underlying Strata: Crater-Hopping at Meridiani  
1155 Planum. J. Geophys. Res. - Planets, v. 120, issue 3, 429-451, doi:10.1002/2014JE004686.
- 1156 Aufrère, L., 1932. Morphologie dunaire et météorologie saharienne. Association de Géographes Français  
1157 Bulletin 56, 34-47.
- 1158 Ayoub, F., J-P. Avouac, C.E. Newman, M.I. Richardson, A. Lucas, S. Leprince, N.T Bridges, 2014. Threshold  
1159 for sand mobility on Mars calibrated from seasonal variations of sand flux. Nature Communications  
1160 5, doi:10.1038/ncomms6096.
- 1161 Bagnold, R.A., 1941. The Physics of Blown Sand and Desert Dunes (New York: Methuen).
- 1162 Bagnold, R.A, 1953. The surface movement of blown sand in relation to meteorology. Research Council  
1163 of Israel, Special Publication, 2, 89-93.
- 1164 Barnes, J.W., R.H. Brown, L. Soderblom, C. Sotin, S. Le Mouelic, S. Rodriguez, R. Jaumann, R.A. Beyer, R.  
1165 Clark, P. Nicholson, 2008. Spectroscopy, morphometry, and photoclinometry of Titan’s dune fields  
1166 from Cassini/VIMS. Icarus 195, 400-414.
- 1167 Barnes, J.W., L. Lemke, R. Foch, C.P. McKay, R.A. Beyer, J.Radebaugh, D.H. Atkinson, R.D. Lorenz, S. Le  
1168 Mouélic, S. Rodriguez, J. Gundlach, 2012. AVIATR—Aerial Vehicle for In-situ and Airborne Titan  
1169 Reconnaissance. Experimental Astronomy, 33(1), pp.55-127.
- 1170 Barnes, J.W., R.D. Lorenz, J. Radebaugh, A.G. Hayes, K. Arnold, C. Chandler, 2015. Production and global  
1171 transport of Titan's sand particles. Planetary Science 4, 1-19.
- 1172 [Basilevsky, A.T., R.O. Kuzmin, O.V. Nikolaeva, A.A. Pronin, L.B. Ronca, V.S. Avduevsky, G.R. Uspensky, Z.P.](#)  
1173 [Cheremukhina, V.V. Semenchenko, V.M. Ladygin, 1985. The surface of Venus as revealed by the](#)  
1174 [Venera landings: Part II. Geological Society of America Bulletin, 96, 137-144.](#)
- 1175 Belcher, D., J. Veverka, C. Sagan, 1971. Mariner photography of Mars and aerial photography of Earth:  
1176 Some analogies, Icarus, vol. 15, 241 252.
- 1177 Bondarenko, N. V., M. A. Kreslavsky, J. W. Head, 2006. North-south roughness anisotropy on Venus from  
1178 the Magellan Radar Altimeter: Correlation with geology, Journal of Geophysical Research (Planets)  
1179 111, E06S12.

1180 | [Bourke, M.C., K.S. Edgett, B.A. Cantor, 2008. Recent aeolian dune change on Mars. \*Geomorphology\*,](#)  
1181 | [94\(1-2\), 247–255, doi:10.1016/j.geomorph.2007.05.012.](#)

1182 | Breed, C., M. J. Grolier, J.F. McCauley, 1979. Morphology and Distribution of Common 'Sand' Dunes on  
1183 | Mars: Comparison With the Earth, *J. Geophys. Res.*, vol. 84, no. B14, 8183-8204.

1184 | Bridges, N.T., M.C. Bourke, P.E. Geissler, M.E. Banks, C. Colon, S. Diniega, M.P. Golombek, C.J. Hansen, S.  
1185 | Mattson, A.S. McEwen, M.T. Mellon, N. Stantz, B.J. Thomson (2012a), Planet-wide sand motion on  
1186 | Mars. *Geology* 40(1), 31–34, doi:10.1130/G32373.1.

1187 | Bridges, N.T., F. Ayoub, J-P. Avouac, S. Leprince, A. Lucas, S. Mattson (2012b), Earth-like sand fluxes on  
1188 | Mars. *Nature* 485, 339-342, doi:10.1038/nature11022.

1189 | Bridges, N.T., B.L. Ehlmann, R.C. Ewing, C.E. Newman, R. Sullivan, P.G. Conrad, A. Cousin, K.S. Edgett, M.R.  
1190 | Fisk, A.A. Fraeman, J.R. Johnson, M. Lamb, M. Lapotre, S. Le Mouélic, G.M. Martinez, P-Y. Meslin, P.  
1191 | Pinet, L.M. Thompson, J. van Beek, A.R. Vasavada, R.C. Wiens, 2016. Investigation of the Bagnold  
1192 | Dunes by the Curiosity rover: Overview of initial results from the first study of an active dune field  
1193 | on another planet, 47th LPSC, Ab. 2298.

1194 | Burt et al. 2008 (eds). The history of the study of landforms, Volume 4: A history of geomorphology  
1195 | between the 1890s and the 1960s. London, The Geological Society.

1196 | Burr, D. M., S.A. Drummond, R. Cartwright, B.A. Black, J.T. Perron (2013). Morphology of fluvial networks  
1197 | on Titan: Evidence for structural control, *Icarus*, vol. 226(1), 742-759,  
1198 | doi:10.1016/j.icarus.2013.06.016.

1199 | Canada Dept. Agriculture (1976), Glossary of terms in soil science, © Minister of Supply and Services,  
1200 | Canada, ISBN 0-662-01533-9. [http://sis.agr.gc.ca/cansis/publications/manuals/1976-](http://sis.agr.gc.ca/cansis/publications/manuals/1976-glossary/pub1459_report.pdf)  
1201 | [glossary/pub1459\\_report.pdf](http://sis.agr.gc.ca/cansis/publications/manuals/1976-glossary/pub1459_report.pdf)

1202 | [Cardinale, M., G. Komatsu, S. silvestro, D. Tirsch, 2012. The influence of local topography for wind](#)  
1203 | [direction on Mars: two examples of dune fields in crater basins. \*Earth Surface Processes and\*](#)  
1204 | [Landforms, 37, 13, 1437-1443, DOI: 10.1002/esp.3289.](#)

1205 | Cardinale, M., S. Silvestro, D.A. Vaz, T. Michaels, M.C. Bourke, G. Komatsu, L. Marinangeli, 2016. Present-  
1206 | day aeolian activity in Herschel Crater, Mars. *Icarus* 265, 139-148, doi:10.1038/ncomms6096.

1207 | Carneiro Filho, A., D. Schwartz, S.H. Tatum, T. Rosique, 2002. Amazonian paleodunes provide evidence  
1208 | for drier climate phases during the Late Pleistocene–Holocene. *Quaternary Research*, 58(2), pp.205-  
1209 | 209.

1210 | Charnay, B., E. Barth, S. Rafkin, C. Narteau, S. Lebonnois, S. Rodriguez, S.C. Du Pont, A. Lucas, 2015.  
1211 | Methane storms as a driver of Titan's dune orientation. *Nature Geoscience*, 8(5), pp.362-366.

1212 | Cheng, A.F., C.M. Lisse, M. A'Hearn, 2013. Surface geomorphology of Jupiter Family Comets: A geologic  
1213 | process perspective. *Icarus* 222, 808-817, doi:10.1016/j.icarus.2012.10.004.

1214 | Christensen, P.R., 1983. Eolian intracrater deposits on Mars: Physical properties and global distribution.  
1215 | *Icarus* 56(3), 496-518.

1216 | Christensen, P.R., J.L. Bandfield, J.F. Bell III, N. Gorelick, V.E. Hamilton, A. Ivanov, B.M. Jakosky, H.H.  
1217 | Kieffer, M.D. Lane, M.C. Malin, T. McConnochie, 2003. Morphology and composition of the surface  
1218 | of Mars: Mars Odyssey THEMIS results. *Science* 300(5628), pp.2056-2061.

1219 | Chojnacki, M., D.M. Burr, J.E. Moersch, T.I. Michaels, 2011. Orbital observations of contemporary dune  
1220 | activity in Endeavour Crater, Meridiani Planum, Mars. *J. Geophys. Res.* 116, E00F19.

1221 | [Chojnacki, M., D. M. Burr, J. E. Moersch, J. J. Wray, 2014. Valles Marineris dune sediment provenance](#)  
1222 | [and pathways. \*Icarus\*, 232\(0\), 187–219, doi:10.1016/j.icarus.2014.01.011.](#)

1223 | [Chojnacki, M., J.R. Johnson, J.E. Moersch, L.K. Fenton, T.I. Michaels, J.F. Bell III, 2015. Persistent aeolian](#)  
1224 | [activity at Endeavour crater, Meridiani Planum, Mars; new observations from orbit and the surface.](#)  
1225 | [Icarus, 251\(0\), 275–290, doi:10.1016/j.icarus.2014.04.044.](#)

1226 | Claudin, P., B. Andreotti, 2006. A scaling law for aeolian dunes on Mars, Venus, Earth, and for  
1227 | subaqueous ripples. *Earth and Planetary Science Letters*, vol. 252, 30–44,  
1228 | doi:10.1016/j.epsl.2006.09.004.

1229 | Colinvaux, P.A., G. Irion, M.E. Räsänen, M.B. Bush, J.N. De Mello, 2001. A paradigm to be discarded:  
1230 | geological and paleoecological data falsify the Haffer & Prance refuge hypothesis of Amazonian  
1231 | speciation. *Amazoniana*, 16(3), pp.609-646.

1232 | [Courrech du Pont, S., C. Narteau, X. Gao, 2014. Two modes for dune orientation. \*Geology\*, 42\(9\),](#)  
1233 | [doi:10.1130/G35657.1.](#)

1234 | Cousin, A., O. Forni, P.-Y. Meslin, S. Schröder, O. Gasnault, N. Bridges, B. Elhmann, S. Maurice, R. Wiens,  
1235 | 2016. Chemical diversity among fine-grained soils at Gale (Mars): a chemical transition as the rover  
1236 | is approaching the Bagnold Dunes? 47<sup>th</sup> LPSC, Ab. 2044.

1237 | Cutts, J.A., R.S.U. Smith, 1973. Eolian deposit and Dunes on Mars. *J. Geophys. Res.* 78(20), 4139-4154.

1238 | Cutts, J.A., K.R. Blasius, G.A. Briggs, M.H. Carr, R. Greeley, H. Masursky, 1976. North Polar Region of  
1239 | Mars: Imaging Results from Viking 2. *Science*, vol. 194(4271), 1329-37.

1240 | Diniega, S., K. Glasner, S. Byrne, 2010a. Long scale evolution of aeolian sand dune fields: influences of  
1241 | initial conditions and dune collisions. *Geomorphology* (special edition: Planetary Dunes) 121, 55-68,  
1242 | doi:10.1016/j.geomorph.2009.02.010.

1243 | Diniega, S., S. Byrne, N.T. Bridges, C.M. Dundas, A.S. McEwen, 2010b. Seasonality of present-day Martian  
1244 | dune-gully activity. *Geology* 38, no. 11, 1047-1050, doi:10.1130/G31287.1.

1245 | Diniega, S., A. Allen, T. Perez, C.J. Hansen, 2016. Tracking gully activity within the north polar erg, Mars.  
1246 | 47<sup>th</sup> LPSC, Ab. 1740.

1247 | Doran, P.T., S.M. Clifford, S.L. Forman, L. Nyquist, D.A. Papanastassiou, B.W. Stewart, N.C. Sturchio, T.D.  
1248 | Swindle, T. Cerling, J. Kargel, G. McDonald, 2004. Mars chronology: assessing techniques for  
1249 | quantifying surficial processes. *Earth-Science Reviews*, 67(3), pp.313-337.

1250 | Dundas C.M., S. Diniega, C.J. Hansen, S. Byrne, A.S. McEwen, 2012. HiRISE observations of seasonal  
1251 | activity and morphological changes in Martian gullies. *Icarus* 220(1), 124-143,  
1252 | doi:10.1016/j.icarus.2012.04.005.

1253 | Dundas, C.M., S. Diniega, A.S. McEwen, 2015. Long-Term Monitoring of Martian Gully Formation on  
1254 | Mars and Evolution with HiRISE. *Icarus* (special issue: Dynamic Mars) 251, 244-263,  
1255 | doi:10.1016/j.icarus.2014.05.013.

1256 | Edgett, K.S., P.R. Christensen, 1991. The particle size of Martian aeolian dunes. *JGR* 96(E5), 22765-22776.

1257 | [Edgett, K. S., N. Lancaster \(1993\), Volcaniclastic aeolian dunes: terrestrial examples and application to](#)  
1258 | [martian sands, \*J. Arid Environ.\*, 25\(3\), 271–297, doi:10.1006/jare.1993.1061.](#)

1259 | Elhmann, B.L., N. Bridges, A.A. Fraeman, M.G.A. Lapotre, K. Edgett, J.R. Johnson, A. Cousin, A. Yen, P.  
1260 | Conrad, L. Thompson, J. Van Beek, D. Vaniman, S. Schröder, A. Vasavada, 2016. Chemistry and  
1261 | mineralogy in situ at the Bagnold sand dunes: Evidence for aeolian sorting and size-dependence in  
1262 | sand composition. 47<sup>th</sup> LPSC, Ab. 1536.

1263 Ewing, R.C., G. Kocurek, 2010. Aeolian dune-field pattern boundary conditions. *Geomorphology* 114,  
1264 175-187, doi:10.1016/j.geomorph.2009.06.015.

1265 Ewing, R.C. A. Peyret, G. Kocurek and M.C. Bourke, 2010. Dune-field pattern formation and recent  
1266 transporting winds in the Olympia Undae dune field, north polar region of Mars. *Journal of*  
1267 *Geophysical Research*, 115, E08005.

1268 Ewing, R.C., A.G. Hayes, A. Lucas, 2015. Sand dune patterns on Titan controlled by long-term climate  
1269 cycles. *Nature* 10.1038/ngeo2323.

1270 Ewing, R.C., N.T. Bridges, R. Sullivan, M.G.A. Lapotre, W.W. Fisher, M.P. Lamb, D. Rubin, K. Lewis, S.  
1271 Gupta, 2016. Aeolian Sedimentary Processes at the Bagnold Dunes, Mars: Implications for Modern  
1272 Dune Dynamics and Sedimentary Structures in the Aeolian Stratigraphic Record of Mars. 47<sup>th</sup> LPSC,  
1273 Ab. 2783.

1274 Fenton, L.K., 2005. Potential sand sources for the dune fields in Noachis Terra, Mars. *J. Geophys. Res.*,  
1275 vol. 110, E11, doi:10.1029/2005JE002436.

1276 Fenton, L. K., 2006. Dune migration and slip face advancement in the Rabe Crater dune field, Mars.  
1277 *Geophys. Res. Lett.*, 33(20), 1-5, doi:10.1029/2006GL027133.

1278 Fenton, L., 2016. Dunes on Pluto? Blog on SETI website: [http://www.seti.org/seti-institute/dunes-on-](http://www.seti.org/seti-institute/dunes-on-pluto)  
1279 [pluto](http://www.seti.org/seti-institute/dunes-on-pluto).

1280 Fenton, L.K., R.K. Hayward, 2010. Southern high latitude dune fields on Mars: Morphology, aeolian  
1281 inactivity, and climate change, *Geomorphology* 121, Issues 1-2, 98-121,  
1282 doi:10.1016/j.geomorph.2009.11.006.

1283 Fenton, L.K., A.D. Toigo, M.I. Richardson, 2005. Aeolian processes in Proctor Crater on Mars: Mesoscale  
1284 modeling of dune-forming winds. *J. Geophys. Res.* 110, E06005, doi:10.1029/2004JE002309.

1285 Florensky, C.P., A.T. Basilevsky, V.P. Krychkov, R.O. Kusmin, O.V. Nikolaeva, A.A. Pronin, I.M. Chernaya,  
1286 Yu.S. Tyufin, A.S. Selivanov, M.K. Naraeva, L.B. Ronca, 1983. Venera 13 and Venera 14: Sedimentary  
1287 rocks on Venus? *Science* 221, 57-589.

1288 Fryberger, S.G., G. Dean, 1979. Dune forms and wind regime, in, E. D. McKee, ed., *A study of global sand*  
1289 *seas*. U.S. Geol. Surv. Prof. Pap. 1052, 137-169.

1290 Gardin, E., P. Allemand, C. Quantin, S. Silvestro, C. Delacourt, 2012. Dune fields on Mars: Recorders of a  
1291 climate change? *Planet. Space Sci.* 60, 314-321, doi:10.1016/j.pss.2011.10.004.

1292 Garzanti, E., P. Vermeesch, S. Andò, G. Vezzoli, M. Valagussa, K. Allen, K.A. Kadi, A.I.A. Al-Juboury, 2013.  
1293 Provenance and recycling of Arabian desert sand. *Earth-Sci. Rev.* 120, 1-19,  
1294 doi:10.1016/j.earscirev.2013.01.005.

1295 Geissler, P.E., N.W. Stantzos, N.T. Bridges, M.C. Bourke, S. Silvestro, L.K. Fenton, 2013. Shifting sands on  
1296 Mars: Insights from tropical intra-crater dunes. *ESP&L*, vol. 38, no. 4, 407-412,  
1297 doi:10.1002/esp.3331.

1298 Golombek, M. P., J.A. Grant, L.S. Crumpler, R. Greeley, R.E. Arvidson, J.F. Bell III, C.M. Weitz, R. Sullivan,  
1299 P.R. Christensen, L.A. Soderblom, S.W. Squyres, 2006. Erosion rates at the Mars Exploration Rover  
1300 landing sites and long-term climate change on Mars. *J. Geophys. Res.*, 111, E12S10,  
1301 doi:10.1029/2006JE002754.

1302 Golombek, M.P., N.H. Warner, V. Ganti, M.P. Lamb, T.J. Parker, R.L. Fergason, R. Sullivan, 2014. Small  
1303 crater modification on Meridiani Planum and implications for erosion rates and climate change on  
1304 Mars. *J. Geophys. Res. Planets*, 119, doi:10.1002/2014JE004658.

1305 Goudie, A., 2002. Great warm deserts of the world: landscapes and evolution (Vol. 1). Oxford University  
1306 Press on Demand.

1307 Grant, J. A., R.E. Arvidson, L.S. Crumpler, M.P. Golombek, B. Hahn, A.F.C. Haldemann, R. Li, L.A.  
1308 Soderblom, S. W. Squyres, S.P. Wright, W.A. Watters, 2006. Crater gradation in Gusev crater and  
1309 Meridiani Planum, Mars. *J. Geophys. Res.*, 111, E02S08, doi:10.1029/2005JE002465.

1310 Grant, J. A., S. A. Wilson, B. A. Cohen, M. P. Golombek, P. E. Geissler, R. J. Sullivan, R. L. Kirk, and T. J.  
1311 Parker (2008), Degradation of Victoria crater, Mars, *J. Geophys. Res.*, 113, E11010,  
1312 doi:10.1029/2008JE003155.

1313 Grant, J. A., T. J. Parker, L. S. Crumpler, S. A. Wilson, M. P. Golombek, and D. W. Mittlefehldt, (2016), The  
1314 degradational history of Endeavour crater, Mars: *Icarus*, doi:10.1016/j.icarus.2015.08.019, in press.

1315 [Greeley, R., R.E. Arvidson, 1990. Aeolian processes on Venus. \*Earth, Moon, and Planets\*, 50/51, 127-157.](#)

1316 Greeley, R., J.D. Iversen, J.B. Pollack, N. Udovich, B. White, 1974. Wind tunnel studies of martian aeolian  
1317 processes. *Proc. Royal Soc. London*, 341, 331-360, doi: 10.1098/rspa.1974.0191.

1318 Greeley, R., R. Leach, B. White, J. Iversen, J. Pollack, 1980. Threshold windspeeds for sand on Mars: Wind  
1319 tunnel simulations. *Geophysical Research Letters*, vol. 7, no. 2, 121-124.

1320 Greeley, R., J. Iversen, R. Leach, J. Marshall, B. White, S. Williams, 1984a. Windblown sand on Venus:  
1321 Preliminary results of laboratory simulations. *Icarus* 57(1), 112-124.

1322 Greeley, R., J.R. Marshall, R.N. Leach, 1984b. Microdunes and other aeolian bedforms on Venus: Wind  
1323 tunnel experiments. *Icarus* 60, 152-160.

1324 [Greeley R., N. Lancaster, S. Lee, P. Thomas, 1992a. Martian aeolian processes, sediments and features,](#)  
1325 [in: Kieffer, H., Jakosky, B.M., Snyder, C.W., Matthews, M.S. \(Eds.\), \*Mars\*. University of Arizona Press,](#)  
1326 [Tucson, pp. 730-767.](#)

1327 Greeley, R., R.E. Arvidson, C. Elachi, M.A. Geringer, J.J. Plaut, R.S. Saunders, G. Schubert, E.R. Stofan,  
1328 E.J.P. Thouvenot, S.D. Wall, C.M Weitz, 1992b. Aeolian features on Venus: Preliminary Magellan  
1329 results. *J. Geophys. Res.* 97(E8), 13319-13345.

1330 [Greeley, R. K. Bender, P.E. Thomas, G. Schubert, D. Limonadi, C.M. Weitz, 1995. Wind-related features](#)  
1331 [and processes on Venus: Summary of Magellan results. \*Icarus\*, 115, 399-420.](#)

1332 Greeley, R., K.C. Bender, R.S. Saunders, G. Schubert, C. M. Weitz, 1997. Aeolian processes and features  
1333 on Venus, in *Venus II: Geology, Geophysics, Atmosphere, and Solar Wind Environment*, edited by S.  
1334 W. Bougher, D. M. Hunten, and R. J. Philips, pp. 547– 590, Univ. of Ariz. Press, Tucson.

1335 Greeley, R., R.E. Arvidson, P.W. Barlett, D. Blaney, N.A. Cabrol, P.R. Christensen, R.L. Fergason, M.P.  
1336 Golombek, G.A. Landis, M.T. Lemmon, S.M. McLennan, J.N. Maki, T. Michaels, J.E. Moersch, L.D.V.  
1337 Neakrase, S.C.R. Rafkin, L. Richter, S.W. Squyres, P.A. de Souza Jr., R.J. Sullivan, S.D. Thompson, P.L.  
1338 Whelley, 2006. Gusev crater: Wind-related features and processes observed by the Mars Exploration  
1339 Rover Spirit. *J. Geophys. Res.* 111, E02S09, doi:10.1029/2005JE002491.

1340 Hanna, S.R., 1969. The formation of longitudinal sand dunes by large helical eddies in the atmosphere.  
1341 *Journal of Applied Meteorology*, 8(6), pp.874-883.

1342 Hansen, C. J., M. Bourke, N.T. Bridges, S. Byrne, C. Colon, S. Diniega, C. Dundas, K. Herkenhoff, A.  
1343 McEwen, M. Mellon, G. Portyankina, N. Thomas, 2011. Seasonal erosion and restoration of Mars'  
1344 northern polar dunes. *Science* 331(6017), 575-578, doi:10.1126/science.1197636.



1345 Hansen, C.J., S. Diniega, N. Bridges, S. Byrne, C. Dundas, A. McEwen, G. Portyankina, 2015. Agents of  
1346 change on Mars' northern dunes: CO<sub>2</sub> ice and wind, *Icarus* (special issue: Dynamic Mars) 251, 264-  
1347 274. doi:10.1016/j.icarus.2014.11.015.

1348 Hayward, R.K., 2011. Mars Global Digital Dune Database (MGD3): north polar region (MC-1) distribution,  
1349 applications, and volume estimates. *Earth Surface Processes and Landforms*, doi:10.1002/esp.2219.

1350 Hayward, R.K., K.F. Mullins, L.K. Fenton, T.M. Hare, T.N. Titus, M.C. Bourke, A. Colaprete, P.R.  
1351 Christensen, 2007. Mars global digital dune database and initial science results. *J. Geophys. Res.*  
1352 *Planets* 112, E11007, doi:10.1029/2007JE002943.

1353 Hayward, R.K., T.N. Titus, T.I. Michaels, L.K. Fenton, A. Colaprete, P.R. Christensen, 2009. Aeolian dunes  
1354 as ground truth for atmospheric modeling on Mars. *J. Geophys. Res.*, 114, E11012,  
1355 doi:10.1029/2009JE003428.

1356 [Hayward, R.K., L.K. Fenton, T.N. Titus, 2014. Mars Global Digital Dune Database \(MGD3\): Global dune  
1357 distribution and wind pattern observations. \*Icarus\*, 230, 38-46, doi:10.1016/j.icarus.2013.04.011.](#)

1358 Hersen, P., 2004. On the crescentic shape of barchan dunes. *Eur. Phys. J. B.*, vol. 37, 507-514,  
1359 doi:10.1140/epjb/e2004-00087-y.

1360 Hesse, P., N. Lancaster, M.W. Telfer, 2015. Digital mapping of the extent of global dune systems.  
1361 *Geophysical Research Abstracts*, vol. 17, EGU2015-3638.

1362 Horgan, B.H.N., J.F. Bell, 2012. Seasonally active slipface avalanches in the north polar sand sea of Mars:  
1363 Evidence for a wind-related origin. *Geophys. Res. Lett.* 39, L09201, doi:10.1029/2012GL051329.

1364 Hugenholtz, C.H., T.E. Barchyn, 2010. Spatial analysis of sand dunes with a new global topographic  
1365 dataset: new approaches and opportunities. *Earth Surf. Process. Landforms* 35, 986-992,  
1366 doi:10.1002/esp.2013.

1367 Jackson, D.W.T., M.C. Bourke, T.A.G. Smyth, 2015. The dune effect on sand-transporting winds on Mars.  
1368 *Nature Communications*, DOI: 10.1038/ncomms9796.

1369 Jain, M., C.E. Andersen, L. Bøtter-Jensen, A.S. Murray, H. Haack, J.C. Bridges, 2006. Luminescence dating  
1370 on Mars: OSL characteristics of Martian analogue materials and GCR dosimetry. *Radiation*  
1371 *Measurements*, 41(7), pp.755-761.

1372 [Jerolmack, D.J., R.C. Ewing, F. Falcini, R.L. Martin, C. Masteller, C. Phillips, M.D. Reitz, I. Buynevich, 2012.  
1373 Internal boundary layer model for the evolution of desert dune fields. \*Nature Geoscience\*, 5, 3, 1-4,  
1374 DOI: 10.1038/ngeo1381.](#)

1375 JPL, Oct. 4, 2012. NASA Mars Curiosity Rover Prepares to Study Martian Soil, Press Release  
1376 <http://www.jpl.nasa.gov/news/news.php?feature=3540>.

1377 JPL, Feb. 3, 2014. Through the Gap: Curiosity Mars Rover Crosses Dune, Press Release  
1378 <http://www.jpl.nasa.gov/news/news.php?release=2014-035>.

1379 JPL, Dec. 10, 2015. NASA Mars Rover Curiosity Reaches Sand Dunes, Press Release  
1380 <http://www.jpl.nasa.gov/news/news.php?feature=4787>.

1381 Klammer, G., 1982. Die Paleoste des Pantanal von Mato Grosso und die pleistozene Klimageschichte der  
1382 brasilianischen Randtropen. *Zeitschrift. der Geomorphologie N.F.*, vol. 26, 393-416

1383 Kieffer, S.W., R. Lopes-Gautier, A. McEwen, W. Smythe, L. Keszthelyi, R. Carlson, 2000. Prometheus: Io's  
1384 wandering plume. *Science* 288, 1204-1208.

1385 Kocurek, G., R.C. Ewing, 2005. Aeolian dune field self-organization – implications for the formation of  
1386 simple versus complex dune-field patterns. *Geomorphology* 72, 94-105,  
1387 doi:10.1016/j.geomorph.2005.05.005.

1388 [Kocurek, G., N. Lancaster, 1999. Aeolian system sediment state: theory and Mojave Desert Kelso dune](#)  
1389 [field example. \*Sedimentology\* 46, 505 - 515.](#)

1390 Kocurek, G., R.C. Ewing, 2012. Source-to-sink: An Earth/Mars comparison of boundary conditions for  
1391 eolian dune systems. *Sedimentary Geology of Mars*, SEPM Special Publication No. 102, Print ISBN  
1392 978-1-56576-312-8, CD/DVD ISBN 978-1-56576-313-5, p. 151–168.

1393 Kocurek, G., M. Carr, R. Ewing, K.G. Havholm, Y.C. Nagar, A.K. Singhvi, 2007. White Sands Dune Field,  
1394 New Mexico: age, dune dynamics and recent accumulations. *Sedimentary Geology*, vol. 197, no. 3,  
1395 313-331.

1396 Kok, J. F., N.O. Renno, 2009. A comprehensive numerical model of steady state saltation (COMSALT). *J.*  
1397 *Geophys. Res.*, 114, D17204, doi:10.1029/2009JD011702.

1398 Kok, J.F., 2010. Difference in the wind speeds required for initiation versus continuation of sand  
1399 transport on Mars: Implications for dunes and dust Storms. *Phys. Rev. Lett.* 104, 074502,  
1400 doi:10.1103/PhysRevLett.104.074502.

1401 Kok J.F., E.J.R. Parteli, T.I. Michaels, D. Bou Karam, 2012. The physics of windblown sand and dust. *Rep.*  
1402 *Prog. Phys.* 75, 106901, doi:10.1088/0034-4885/75/10/106901.

1403 Kreslavsky, M.A., R.V. Vdovichenko, 1999. Anisotropy of Venus Surface Properties Inferred from  
1404 Magellan Side-Looking-Radar Data. *Solar System Research* 33, 110.

1405 Lancaster, N., 1995. *Geomorphology of Desert Dunes* (Routledge Physical Environment Series).  
1406 Routledge. ISBN-13: 978-0415060943.

1407 [Lancaster, N., S. Wolfe, D. Thomas, C. Bristow, O. Bubbenzer, S. Burrough, G. Duller, A. Halfen, P. Hesse, J.](#)  
1408 [Roskin, A. Singhvi, H. Tsoar, A. Tripaldi, X. Yang, M. Zárate, M., in press. The INQUA Dunes Atlas](#)  
1409 [chronologic database. \*Quaternary International\*, doi:10.1016/j.quaint.2015.10.044.](#)

1410 [Lapotre, M.G.A., R.C. Ewing, M.P. Lamb, W.W. Fischer, J.P. Grotzinger, D.M. Rubin, K.W. Lewis, M.](#)  
1411 [Ballard, M. Day, S. Gupta, S.G. Banham, N.T. Bridges, D.J. Des Marais, A.A. Fraeman, J.A. Grant, K.E.](#)  
1412 [Herkenhoff, D.W. Ming, M.A. Mischna, M.S. Rice, D.A. Summer, A.R. Vasavada, R.A. Yingst, 2016a.](#)  
1413 [Large wind ripples on Mars: A record of atmospheric evolution. \*Science\*, 353, 6294, 55-58, DOI:](#)  
1414 [10.1126/science.aaf3206.](#)

1415 ~~Lapotre, M.G.A., R.C. Ewing, M.P. Lamb, W.W. Fischer, K. Lewis, M. Ballard, M. Day, D. Rubin, J.P.~~  
1416 ~~Grotzinger (2016a) Orbital and in-situ observations in support of the existence of an unknown stable~~  
1417 ~~aeolian bedform regime on Mars, 47<sup>th</sup> LPSC, Ab. 1510. *[this has also been written up in a submitted*~~  
1418 ~~*paper]*~~

1419 Lapotre, M.G.A., B.L. Ehlmann, A.A. Fraeman, S.E. Minson, F. Ayoub, R.C. Ewing, R.E. Arvidson, N.T.  
1420 Bridges, MSL Science Team, 2016b. A quantitative assessment of aeolian fractionation at the  
1421 Bagnold dunes of Gale crater, Mars, from orbit to the ground. 47<sup>th</sup> LPSC, Ab. 1513.

1422 Le Gall, A., A.G. Hayes, R. Ewing, M.A. Janssen, J. Radebaugh, C. Savage, P. Encrenaz, 2012. Latitudinal  
1423 and altitudinal controls of Titan’s dune field morphometry. *Icarus*, vol. 217, issue 1, 231–242.

1424 Livingstone, I., G.F. Wiggs, C.M. Weaver, 2007. Geomorphology of desert sand dunes: a review of recent  
1425 progress. *Earth-Science Reviews*, 80(3), 239-257.

1426 Livingstone, I., C. Bristow, R.G. Bryant, J. Bullard, K. White, G.F. Wiggs, A.C. Baas, M.D. Bateman, D.S.  
1427 Thomas, 2010. The Namib Sand Sea digital database of aeolian dunes and key forcing variables.  
1428 *Aeolian Research*, 2(2), 93-104.

1429 Lorenz, R.D., 2015. Heights of Fortuna-Meshknet Dunes (Al-Uzza Undae), Venus, from Magellan  
1430 Radarclinometry. Fourth Annual International Planetary Dunes Workshop, abstract # 8004.

1431 Lorenz, R.D., J. Radebaugh, 2009. Global pattern of Titan's dunes: Radar survey from the Cassini prime  
1432 mission. *Geophys. Res. Lett.* 36, L03202, doi:10.1029/2008/GL036850.

1433 Lorenz, R.D., J.I. Lunine, J.A. Grier, M.A. Fisher, 1995. Prediction of aeolian features on planets:  
1434 Application to Titan paleoclimatology. *JGR* 100(E12), 26377-26386.

1435 Lorenz, R.D., S. Wall, J. Radebaugh, G. Boubin, E. Reffet, M. Janssen, E. Stofan, R. Lopes, R. Kirk, C. Elachi,  
1436 J. Lunine, K. Mitchell, F. Paganelli, L. Soderblom, C. Wood, L. Wye, H. Zebker, Y. Anderson, S. Ostro,  
1437 M. Allison, R. Boehmer, P. Callahan, P. Encrenaz, G.G. Ori, G. Francescetti, Y. Gim, G. Hamilton, S.  
1438 Hensley, W. Johnson, K. Kelleher, D. Muhleman, G. Picardi, F. Posa, L. Roth, R. Seu, S. Shaffer, B.  
1439 Stiles, S. Vetrella, E. Flamini, R. West, 2006. The sand seas of Titan: Cassini RADAR observations of  
1440 longitudinal dunes. *Science* 312, 724–727, doi: 10.1126/science.1123257.

1441 Lorenz, R. D., R.M. Lopes, F. Paganelli, J.I. Lunine, R.L. Kirk, K.L. Mitchell, L.A. Soderblom, E.R. Stofan, G.  
1442 Ori, M. Myers, H. Miyamoto, J. Radebaugh, B. Stiles, S.D. Wall, C.A. Wood and the Cassini RADAR  
1443 Team, 2008. Fluvial channels on Titan: Initial Cassini RADAR observations. *Planet. Space Sci.* 56,  
1444 1132-1144.

1445 Lucas, A., S. Rodriguez, C. Nartean, B. Charnay, S.C. Pont, T. Tokano, A. Garcia, M. Thiriet, A.G. Hayes,  
1446 R.D. Lorenz, O. Aharonson, 2014. Growth mechanisms and dune orientation on Titan.  
1447 *Geophysical Research Letters*, 41(17), 6093-6100.

1448 Madigan, C.T., 1936. The Australian sand-ridge deserts. *Geographical Review*, 26(2), 205-227.

1449 Malaska, M.J., R.M. Lopes, A.G. Hayes, J. Radebaugh, R.D. Lorenz and E.P. Turtle, 2016. Material  
1450 transport map of Titan: The fate of dunes. *Icarus*, 270, 183-196,  
1451 doi:10.1016/j.icarus.2015.09.029.

1452 Marsh, G.P., 1864. *Man and Nature*. (Reprint, 1965) Cambridge, MA: Harvard University Press.

1453 Marshall, J.R., R. Greeley, 1992. An experimental study of aeolian structures on Venus. *J. Geophys. Res.*  
1454 97(E1), 1007-1016.

1455 May, J.H., 2013. Dunes and dune fields in the Bolivian Chaco as potential records of environmental  
1456 change. *Aeolian Research*, vol. 10, 89-102.

1457 McCauley, J.F., M.H. Carr, J.A. Cutts, W.K. Hartmann, H. Masursky, D.J. Milton, R.P. Sharp, D.E.  
1458 Wilhelms, 1972. Preliminary Mariner 9 Report on the Geology of Mars, *Icarus* 17(2), 289-327,  
1459 doi:10.1016/0019-1035(72)90003-6.

1460 McEwen, A.S., M.J.S. Belton, H.H. Breneman, S.A. Fagents, P. Geissler, R. Greeley, J.W. Head, G. Hoppa,  
1461 W.L. Jaeger, T.V. Johnson, L. Keszthelyi, 2000. Galileo at Io: Results from high-resolution imaging.  
1462 *Science*, vol. 288, no. 5469, 1193-1198.

1463 McEwen, A.S., E.M. Eliason, J.W. Bergstrom, N.T. Bridges, C.J. Hansen, W.A. Delamere, J.A. Grant, V.C.  
1464 Gulick, K.E. Herkenhoff, L. Keszthelyi, R.L. Kirk, 2007. Mars Reconnaissance Orbiter's High Resolution  
1465 Imaging Science Experiment (HiRISE). *J. Geophys. Res.*, 112, issue E5, doi:10.1029/2005JE002605.

1466 McKeever, S.W.S., D. Banerjee, M. Blair, S.M. Clifford, M.S. Cloudsley, S.S. Kim, M. Lamothe, K. Lepper,  
1467 M. Leuschen, K.J. McKeever, M. Prather, 2003. Concepts and approaches to in situ luminescence  
1468 dating of martian sediments. *Radiation Measurements*, 37(4), pp.527-534.

1469 [Mellon, M.T., B.M. Jakovsky, H.H. Kieffer, P.R. Christensen, 2000. High-resolution thermal inertia](#)  
1470 [mapping from the Mars Global Surveyor Thermal Emissino Spectrometer. \*Icarus\*, 148, 437-455,](#)  
1471 [doi:10.1006/icar.2000.6503.](#)

1472 Milazzo, M.P., L.P. Keszthelyi, A.S. McEwen, 2001. Observations and initial modeling of lava-SO<sub>2</sub>  
1473 interactions at Prometheus, Io. *J. Geophys. Res.* 106, 33,121-33,127.

1474 Moore, J.M., A.D. Howard, P.M. Schenk, W.B. McKinnon, R.T. Pappalardo, R.C. Ewing, E.B. Bierhaus, V.J.  
1475 Bray, J.R. Spencer, R.P. Binzel, B. Buratti, 2015. Geology before Pluto: Pre-encounter considerations.  
1476 *Icarus*, 246, 65-81, doi:10.1016/j.icarus.2014.04.028.

1477 Moore, J.M., W.B. McKinnon, J.R. Spencer, A.D. Howard, P.M. Schenk, R.A. Beyer, F. Nimmo, K.N. Singer,  
1478 O.M. Umurhan, O.L. White, S.A. Stern, K. Ennico, C.B. Olkin, H.A. Weaver, L.A. Young, R.P. Binzel,  
1479 M.W. Buie, B.J. Buratti, A.F. Cheng, D.P. Cruikshank, W.M. Grundy, I.R. Linscott, H.J. Reitsema, D.C.  
1480 Reuter, M.R. Showalter, V.J. Bray, C.L. Chavez, C.J.A. Howett, T.R. Lauer, C.M. Lisse, A.H. Parker, S.B.  
1481 Porter, S.J. Robbins, K. Runyon, T. Stryk, H.B. Throop, C.C.C. Tsang, A.J. Verbiscer, A.M. Zangari, A.L.  
1482 Chaikin, D.E. Wilhelms, New Horizons Science Team, 2016. The geology of Pluto and Charon through  
1483 the eyes of New Horizons. *Science* v. 351, Is. 6279, 1284-93. doi:10.1126/science.aad7055.

1484 Mottola, S., Arnold, G., Grothues, H.-G., Jaumann, R., Michaelis, H., Neukum, G., Bibring, J.-P., Schröder,  
1485 S.E., Hamm, M., Otto, K.A., Pelivan, I., Proffe, G., Scholten, F., Tirsch, D., Kreslavsky, M., Remetean,  
1486 E., Souvannavong, F., Dolives, B., 2015. The structure of the regolith on 67P/Churyumov-  
1487 Gerasimenko from ROLIS descent imaging. *Science* 349, 10.1126/science.aab0232.

1488 [Neakrase, L.D.V. \(2015\), Terrestrial Analogues for Venusian dunes: Sub-aqueous, seafloor dune fields.](#)  
1489 [Fourth International Planetary Dunes Workshop. Ab. 8023.](#)

1490 New Horizons, 2015. [http://www.nasa.gov/mission\\_pages/newhorizons/images/index.html](http://www.nasa.gov/mission_pages/newhorizons/images/index.html).

1491 New Horizons, 2016. The Tartarus Dorsa Mountains Rise Up Along Pluto, <http://www.nasa.gov/image->  
1492 [feature/the-tartarus-dorsa-mountains-rise-up-along-pluto](http://www.nasa.gov/image-feature/the-tartarus-dorsa-mountains-rise-up-along-pluto).

1493 NRC, 2013. Vision and Voyages for Planetary Science in the Decade 2013-2022. Steering Group Chair:  
1494 Squyres, National Academies Press, Washington, D.C., ISBN: 0-309-20955-2.

1495 O'Connell-Cooper, C.D., L.M. Thompson, J.G. Spray, J.A. Berger, E.D. Desouza, M. Schmidt, A. Yen, N.I.  
1496 Boyd, R. Gellert, G. Perrett, S.J. VanBommel, the MSL-APXS Team, 2016. Preliminary comparison of  
1497 soils within Gale crater to those from Gusev Crater and Meridiani Planum. 47<sup>th</sup> LPSC, Ab. 2477.

1498 Parteli, E.J.R., O. Durán, H.J. Herrmann, 2007. Minimal size of a barchan dune. *Phys. Rev. E* 75, 011301,  
1499 doi:10.1103/PhysRevE.75.011301.

1500 Parteli, E.J.R., O. Durán, H. Tsoar, V. Schwämmle, H.J. Herrmann, 2009. Dune formation under bimodal  
1501 winds. *PNAS* 106(52), 22085-22089, doi:10.1073/pnas.0808646106.

1502 Pelletier, J.D., 2009. Controls on the height and spacing of eolian ripples and transverse dunes: A  
1503 numerical modeling investigation. *Geomorphology*, v. 105, 322-333,  
1504 doi:10.1016/j.geomorph.2008.10.010.

1505 [Pelletier, J.D., 2015. Controls on the large-scale spatial variations of dune field properties in the](#)  
1506 [barchanoid portion of White Sands dune field, New Mexico. \*J. Geophys. Res. Earth Surf.\*, 120, 453-](#)  
1507 [473, doi:10.1002/2014JF003314.](#)

1508 | [Portyankina, G., A. Pommerol, K.-M. Aye, C. J. Hansen, N. Thomas, 2012. Polygonal cracks in the](#)  
1509 | [seasonal semi-translucent CO<sub>2</sub> ice layer in Martian polar areas. \*J. Geophys. Res.\*, 117\(E2006\),](#)  
1510 | [doi:10.1029/2011JE003917.](#)

1511 | Radebaugh, J., 2013. Dunes on Saturn's moon Titan at the end of the Cassini Equinox Mission. *Aeolian*  
1512 | *Research* 11, 23-41.

1513 | Radebaugh, J., R. Lorenz, J. Lunine, S. Wall, G. Boubin, E. Reffet, R. Kirk, R. Lopes, E. Stofan, L. Soderblom,  
1514 | M. Allison, M. Janssen, P. Paillou, P. Callahan and the Cassini Radar Team, 2008. Dunes on Titan  
1515 | observed by Cassini Radar. *Icarus* 194, 690-703, doi:10.1016/j.icarus.2007.10.015.

1516 | Radebaugh, J., R. Lorenz, T. Farr, P. Paillou, C. Savage, C. Spencer, 2010. Linear dunes on Titan and earth:  
1517 | Initial remote sensing comparisons. *Geomorphology* 121, 122–132,  
1518 | doi:10.1016/j.geomorph.2009.02.022.

1519 | Radebaugh, J., R.D. Lorenz, S.D. Wall, R.L. Kirk, C.A. Wood, J.I. Lunine, E.R. Stofan, R.M.C. Lopes, P.  
1520 | Valora, T.G. Farr, A.G. Hayes, B. Stiles, G. Mitri, H. Zebker, M. Janssen, L. Wye, A. Le Gall, K.L.  
1521 | Mitchell, F. Paganelli, R.D. West, E.L. Schaller and the Cassini RADAR Team, 2011. Regional  
1522 | geomorphology and history of Titan's Xanadu province. *Icarus* 211, 672-685.

1523 | Radebaugh, J., D. Ventra, R.D. Lorenz, T. Farr, R. Kirk, A. Hayes, M.J. Malaska, S. Birch, Z. Y-C. Liu, J.  
1524 | Lunine, J. Barnes, A. Le Gall, R. Lopes, E. Stofan, S. Wall, P. Paillou, 2016. Alluvial and fluvial fans on  
1525 | Saturn's moon Titan reveal processes, materials and regional geology, in, Ventra, D. & Clarke, L. E.  
1526 | (eds) *Geology and Geomorphology of Alluvial and Fluvial Fans: Terrestrial and Planetary*  
1527 | *Perspectives*. Geological Society, London, Special Publications, 440, doi: 10.1144/SP440.6.

1528 | Rubin, D.M., 2012. A unifying model for planform straightness of ripples and dunes in air and water.  
1529 | *Earth-Science Reviews*, 113(3), 176-185, doi:10.1016/j.earscirev.2012.03.010.

1530 | Rubin, D.M., P.A. Hesp, 2009. Multiple origins of linear dunes on Earth and Titan. *Nature Geoscience*,  
1531 | vol. 2, 653-658, doi:10.1038/ngeo610.

1532 | [Rubin, D.M., R.E. Hunter, 1987. Bedform alignment in directionally varying flows. \*Science\*, 237\(4812\),](#)  
1533 | [276–8, doi:10.1126/science.237.4812.276.](#)

1534 | [Rubin, D. M., H. Ikeda, 1990. Flume experiments on the alignment of transverse, oblique, and](#)  
1535 | [longitudinal dunes in directionally varying flows. \*Sedimentology\*, 37, 673–684.](#)

1536 | Sagan, C., J. Veverka, P. Fox, R. Dubisch, J. Lederberg, E. Levinthal, L. Quam, R. Tucker, J.B. Pollack, B.A.  
1537 | Smith, 1972. Variable features on Mars: Preliminary Mariner 9 television results. *Icarus* 17(2), 346-  
1538 | 372, doi:10.1016/0019-1035(72)90005-X.

1539 | Sauermann, G., K. Kroy, H.J. Herrmann, 2001. Continuum saltation model for sand dunes, *Physical*  
1540 | *Review E*, v. 64, 031305, doi:10.1103/PhysRevE.64.031305.

1541 | Savage, C.J., J. Radebaugh, E.H. Christiansen, R.D. Lorenz, 2014. Implications of dune pattern analysis for  
1542 | Titan's surface history. *Icarus* 230, 180-190.

1543 | Silvestro, S., G. Di Achille, G.G. Ori, 2010a. Dune morphology, sand transport pathways and possible  
1544 | source areas in east Thaumasia Region (Mars), *Geomorphology*, vol. 121, 84-97,  
1545 | doi:10.1016/j.geomorph.2009.07.019.

1546 | Silvestro, S., L.K. Fenton, D.A. Vaz, N.T. Bridges, G.G. Ori, 2010b. Ripple migration and dune activity on  
1547 | Mars: Evidence for dynamic processes. *Geophys. Res. Lett.* 37, L20203 (2010).

1548 Silvestro, S., D.A. Vaz, L.K. Fenton and P.E. Geissler, 2011. Active Aeolian processes on Mars: a regional  
1549 study in Arabia and Meridiani Terrae, *Geophysical Research Letters*, 38, L20201,  
1550 doi:10.1029/2011GL048955.

1551 [Silvestro S., L.K. Fenton, T.I. Michaels, A. Valdez, G.G. Ori, 2012. Interpretation of the complex dune  
1552 morphology on Mars: dune activity, modeling and a terrestrial analogue. \*Earth Surface Processes  
1553 and Landforms\*, 37, 13, 1424-1436, DOI: 10.1002/esp.3286.](#)

1554 Silvestro, S., D.A. Vaz, R.C. Ewing, A.P. Rossi, L.K. Fenton, T.I. Michaels, J. Flahaut and P.E.  
1555 Geissler, 2013. Pervasive aeolian activity along Curiosity's traverse in Gale Crater, Mars.  
1556 *Geology*, 41 (4), 483–486, doi:10.1130/G34162.1.

1557 [Silvestro S., D.A. Vaz, H. Yizhaq, F. Esposito, 2016. Dune-like dynamic of martian aeolian large ripples.  
1558 \*Geophysical Research Letters\*, doi:10.1002/2016GL070014.](#)

1559 [Silvestro, S., D.A. Vaz, I.C. Popa and F. Esposito \(2015\), Longitudinal aeolian depositional features on  
1560 Mars? Fourth International Planetary Dunes Workshop, LPI Contribution No. 1843,  
1561 p.8049.](#)

1562 Singhvi, A.K., N. Porat, 2008. Impact of luminescence dating on geomorphological and palaeoclimate  
1563 research in drylands. *Boreas*, 37(4), 536-558.

1564 Soderblom, L., J. Anderson, K. Baines, J. Barnes, J. Barrett, R. Brown, B. Buratti, R. Clark, D. Cruikshank, C.  
1565 Elachi, M. Janssen, R. Jaumann, R. Kirk, E. Karkoschka, S. Lemouelic, R. Lopes, R. Lorenz, J. Lunine, T.  
1566 McCord, P. Nicholson, J. Radebaugh, B. Rizk, C. Sotin, E. Stofan, T. Sucharski, M. Tomasko, S. Wall,  
1567 2007. Correlations between Cassini VIMS spectra and RADAR SAR images: Implications for Titan's  
1568 surface composition and the character of the Huygens Probe landing site. *Planetary and Space  
1569 Science*, 55, 2025-2036.

1570 Sohbaty, R., A.S. Murray, J.P. Buylaert, M. Ortuño, P.P. Cunha, E. Masana, 2012. Luminescence dating of  
1571 Pleistocene alluvial sediments affected by the Alhama de Murcia fault (eastern Betics, Spain)—a  
1572 comparison between OSL, IRSL and post-IR IRSL ages. *Boreas*, 41(2), 250-262, doi:10.1111/j.1502-  
1573 3885.2011.00230.x

1574 Sokolów, N. A., 1894. *Die Dunen; Bildung, Entwicklung, and innerer Bau* (German transl. from Russian  
1575 4th ed., 1884, assisted by A. Arzruni). Berlin, Springer.

1576 [Sullivan, R., R. Arvidson, J.F. Bell III, R. Gellert, M. Golombek, R. Greeley, K. Herkenhoff, J. Johnson, S.  
1577 Thompson, P. Whelley, J. Wray, 2008. Wind-driven particle mobility on Mars: Insights from Mars  
1578 Exploration Rover observations at "El Dorado" and surroundings at Gusev Crater. \*J. Geophys. Res.\*,  
1579 113, E06S07, doi:10.1029/2008JE003101.](#)

1580 Stern and the New Horizons team, 2016. New Horizons: The Exploration of the Pluto System and the  
1581 Kuiper Belt beyond. 47<sup>th</sup> LPSC, Ab. 1317.

1582 Sullivan, R., D. Banfield, J. F. Bell III, W. Calvin, D. Fike, M. Golombek, R. Greeley, J. Grotzinger, K.  
1583 Herkenhoff, D. Jerolmack, M. Malin, D. Ming, L. A. Soderblom, S. W. Squyres, S. Thompson, W.A.  
1584 Watters, C.M. Weitz, A. Yen, 2005. Aeolian processes at the Mars Exploration Rover Meridiani  
1585 Planum landing site. *Nature* 436, 58-61, doi:10.1038/nature03641.

1586 Tanaka, K.L., Rodriguezb, J.A.P., Skinner Jr., J.A., Bourke, M.C., Fortezzo, C.M., Herkenhoff, K.E., Kolb, E.J.,  
1587 Okubo, C.H., 2008. North polar region of Mars: Advances in stratigraphy, structure, and erosional  
1588 modification. *Icarus* 192, 318-358, 10.1016/j.icarus.2008.01.021.

1589 Teeuw, R.M., E.J. Rhodes, 2004. Aeolian activity in northern Amazonia: optical dating of Late Pleistocene  
1590 and Holocene palaeodunes. *Journal of Quaternary Science*, 19(1), pp.49-54.

1591 Telfer, M.W., 2011. Growth by extension, and reworking, of a south-western Kalahari linear dune. *Earth*  
1592 *Surface Processes and Landforms*, 36(8), pp.1125-1135.

1593 Telfer, M.W. and Thomas, D.S.G., 2007. Late Quaternary linear dune accumulation and  
1594 chronostratigraphy of the southwestern Kalahari: implications for aeolian palaeoclimatic  
1595 reconstructions and predictions of future dynamics. *Quaternary Science Reviews*, 26(19), pp.2617-  
1596 2630.

1597 Tirsch, D., R. Jaumann, A. Pacifici, F. Poulet, 2011. Dark aeolian sediments in Martian craters:  
1598 Composition and sources. *J Geophys Res-Planet* 116, doi: 10.1029/2009je003562.

1599 [Thomas, P., 1981. North-south asymmetry of eolian features in martian polar regions: Analysis based on](#)  
1600 [crater-related wind markers. \*Icarus\*, 48, 76-90.](#)

1601 [Thomas, P., 1982. Present wind activity on Mars - Relation to large latitudinally zoned sediment](#)  
1602 [deposits. \*J. Geophys. Res.\*, 87\(B12\), 9999-10,008, doi:10.1029/JB087iB12p09999.](#)

1603 Thomas, N., H. Sierks, C. Barbieri, P.L. Lamy, R. Rodrigo, H. Rickman, D. Koschny, H.U. Keller, J. Agarwal,  
1604 M.F. A'Hearn, F. Angrilli, A.-T. Auger, M.A. Barucci, J.-L. Bertaux, I. Bertini, S. Besse, D. Bodewits, G.  
1605 Cremonese, V. Da Deppo, B. Davidsson, M. De Cecco, S. Debei, S., M.R. El-Maarry, F. Ferri, S.  
1606 Fornasier, M. Fulle, L. Giacomini, O. Groussin, P.J. Gutierrez, C. Güttler, S.F. Hviid, W.-H. Ip, L. Jorda,  
1607 J. Knollenberg, J.-R. Kramm, E. Kührt, M. Küppers, F. La Forgia, L.M. Lara, M. Lazzarin, J.J.L. Moreno,  
1608 S. Magrin, S. Marchi, F. Marzari, M. Massironi, H. Michalik, R. Moissl, S. Mottola, G. Naletto, N.  
1609 Ookay, M. Pajola, A. Pommerol, F. Preusker, L. Sabau, F. Scholten, C. Snodgrass, C. Tubiana, J.-B.  
1610 Vincent, K.-P. Wenzel, 2015a. The morphological diversity of comet 67P/Churyumov-Gerasimenko.  
1611 *Science* 347, doi:10.1126/science.aaa0440.

1612 [Thomas, N., B. Davidsson, M.R. El-Maarry, S. Fornasier, L. Giacomini, A.G. Gracia-Berná, S.F. Hviid, W.-H.](#)  
1613 [Ip, L. Jorda, H.U. Keller, J. Knollenberg, E. Kührt, F. La Forgia, I.L. Lai, Y. Liao, R. Marschall, M.](#)  
1614 [Massironi, S. Mottola, M. Pajola, O. Poch, A. Pommerol, F. Preusker, F. Scholten, C.C. Su, J.S. Wu, J.-](#)  
1615 [B. Vincent, H. Sierks, C. Barbieri, P.L. Lamy, R. Rodrigo, D. Koschny, H. Rickman, M.F. A'Hearn, M.A.](#)  
1616 [Barucci, J.-L. Bertaux, I. Bertini, G. Cremonese, V. Da Deppo, S. Debei, M. de Cecco, M. Fulle, O.](#)  
1617 [Groussin, P.J. Gutierrez, J.-R. Kramm, M. Küppers, L.M. Lara, M. Lazzarin, J.J. Lopez Moreno, F.](#)  
1618 [Marzari, H. Michalik, G. Naletto, J. Agarwal, C. Güttler, N. Ookay, C. Tubiana, 2015b. Redistribution of](#)  
1619 [particles across the nucleus of comet 67P/Churyumov-Gerasimenko. \*Astronomy and Astrophysics\*,](#)  
1620 [583, A17, doi:10.1051/0004-6361/201526049.](#)

1621 Treiman, A.H., 2003. Geologic settings of Martian gullies: Implications for their origins. *J. Geophys. Res.*  
1622 108 (E4), 8031, doi:10.1029/2002JE001900.

1623 Tricart, J., 1974. Existence de periodes seches au Quaternaire en Amazonie et dans les regions voisines.  
1624 *Révue de Géomorphologie Dynamique* 23, 145-158

1625 Trilling, D.E., 2016. The surface age of Sputnik Planum, Pluto, must be less than 10 million years. *PLOS*  
1626 *ONE*, doi:10.1371/journal.pone.0147386.

1627 Tripaldi, A., M.A. Zárate, in press. A review of Late Quaternary inland dune systems of South America  
1628 east of the Andes. *Quaternary International*, available online 24 July 2014,  
1629 doi:10.1016/j.quaint.2014.06.069.

1630 Tokano, T., 2008. Dune-forming winds on Titan and the influence of topography. *Icarus* 194, 243-262,

1631 doi:10.1016/j.icarus.2007.10.007.

1632 Tsoar, H., R. Greeley, A.R. Peterfreund, 1979. Mars: The north polar sand seas and related wind patterns.

1633 J. Geophys. Res., vol. 84, no. B14, 8167-8180.

1634 Tsoar, H., 1983. Dynamic processes acting on a longitudinal (seif) dune. *Sedimentology* 30, 567-578.

1635 Vaz, D.A., S. Silvestro, 2014. Mapping and characterization of small-scale aeolian structures on Mars: An

1636 example from the MSL landing site in Gale Crater. *Icarus*, vol. 230, 151-161.

1637 Vaz, D.A., S. Silvestro, P.T.K. Sarmiento, M. Cardinale, submitted. Migrating meter-scale bedforms on

1638 Martian dark dunes: are terrestrial aeolian ripples good analogs? *To Aeolian Research* (this issue).

1639 Ward, A.W., 1979. Yardangs on Mars: Evidence of recent wind erosion. *J. Geophys. Res* 84, 8147-8165.

1640 [Ward, A.W., K.B. Doyle, P.J. Helm, M.K. Weisman, N.E. Witbeck, 1985. Global map of eolian features on](#)

1641 [Mars. J. Geophys. Res., 90\(B2\), 2038-2056.](#)

1642 Weitz, C.M., J.J. Plaut, R. Greeley, R.S. Saunders, 1994. Dunes and microdunes on Venus: Why were so

1643 few found in the Magellan data? *Icarus* 112, 282-295.

1644 Werner, B.T., 1995. Eolian dunes: Computer simulations and attractor interpretation. *Geology* 23(12),

1645 1107-1110.

1646 Williams, S.H., R. Greeley, 1994. Windblown sand on Venus: The effect of high atmospheric density.

1647 *Geophys. Res. Lett.* 21(25), 2825-2828.

1648 Wilson, L., J.W. Head, 1994. Mars: Review and analysis of volcanic eruption theory and relationships to

1649 observed landforms. *Reviews of Geophysics*, vol. 32, no. 3, 221-263.

1650 [Zimbelman, J.R., 2000. Non-active dunes in the Acheron Fossae Region of Mars between the Viking and](#)

1651 [Mars Global Surveyor eras. Geophys. Res. Lett. 27, 1069-1072, doi:10.1029/1999GL008399.](#)



**Title: Our evolving understanding of aeolian bedforms, based on observation of dunes on different worlds**

© 2016. All rights reserved

Authors:

Serina Diniega<sup>\*,a</sup>,  
Mikhail Krevalevsky<sup>b</sup>,  
Jani Radebaugh<sup>c</sup>,  
Simone Silverstro<sup>d,e</sup>,  
Matt Telfer<sup>f</sup>,  
Daniela Tirsch<sup>g</sup>

\*corresponding author: serina.diniega@jpl.nasa.gov, +1 818-393-1487

<sup>a</sup>Jet Propulsion Laboratory, California Institute of Technology, 4800 Oak Grove Dr., Pasadena, CA 91109, USA

<sup>b</sup>Earth and Planetary Sciences, University of California - Santa Cruz, 1156 High Str. Santa Cruz, CA, 95064, USA

<sup>c</sup>Department of Geological Sciences, Brigham Young University, Provo, UT 84602, USA

<sup>d</sup>INAF Osservatorio Astronomico di Capodimonte, Via Moiariello 16, 80131, Napoli, Italy

<sup>e</sup>Carl Sagan Center, SETI Institute 189 N Bernardo Ave, Mountain View, CA 94043, USA

<sup>f</sup>SOGEES, University of Plymouth, Drake Circus, Plymouth, Devon. PL4 8AA. UK.

<sup>g</sup>Institute of Planetary Research, German Aerospace Center (DLR), Rutherfordstrasse 2, 12489 Berlin, Germany.

**Abstract (253 words)**

Dunes, dune fields, and ripples are unique and useful records of the interaction between wind and granular materials – finding such features on a planetary surface immediately suggests certain information about climate and surface conditions (at least during the dunes’ formation and evolution). Additionally, studies of dune characteristics under non-Earth conditions allow for “tests” of aeolian process models based primarily on observations of terrestrial features and dynamics, and refinement of the models to include consideration of a wider range of environmental and planetary conditions. To-date, the planetary aeolian community has found and studied dune fields on Mars, Venus, and the Saturnian moon Titan. Additionally, we have observed candidate “aeolian bedforms” on Comet 67P/Churyumov-Gerasimenko, the Jovian moon Io, and – most recently -- Pluto. In this paper, we hypothesize that the progression of investigations of aeolian bedforms and processes on a particular planetary body follows a consistent sequence – primarily set by the acquisition of data of particular types and resolutions, and by the maturation of knowledge about that planetary body. We define that sequence of generated knowledge and new questions (within seven investigation phases) and discuss examples from all of the studied bodies. The aim of such a sequence is to better define our past and current state of understanding about the aeolian bedforms of a particular body, to highlight the related assumptions that require re-analysis with data acquired during later investigations, and to use lessons learned from planetary and terrestrial aeolian studies to predict what types of investigations could be most fruitful in the future.

**Keywords** (at least 6): Planetary; terrestrial; aeolian bedforms; aeolian science; dunes; ripples

**Highlights** (max 85 characters, including spaces)

- Planetary dunes yield useful/unique information about climate & surface conditions.
- Aeolian bedform studies progress logically in questions/assumptions/new knowledge.
- Considering this progression exposes gaps/assumptions to be reviewed with new data.
- Comparing planetary progression with Earth aeolian studies yields lessons for each.

**Acknowledgements:** We thank Matt Chojnacki, Lori Fenton, and an anonymous reviewer for their very helpful comments, and in particular for the suggestions of additional references, which strengthened this manuscript. We also thank the space mission and instrument teams who collect the observations, start the analysis, and enable the described advancements in our understanding of aeolian bedform evolution and interpretation. SD's work was carried out at the Jet Propulsion Laboratory, California Institute of Technology, under a contract with the National Aeronautics and Space Administration. SS was supported by ASI through the ASI-CISAS agreement I/018/ 12/0: "DREAMS EDM Payload—ExoMars 2016."