

1 **Linking the biological impacts of ocean acidification on oysters to**
2 **changes in ecosystem services: A review**

3

4 Anaëlle J. Lemasson^a, Stephen Fletcher^{b,c}, Jason M Hall-Spencer^a, Antony M.
5 Knights^a

6

7 ^a Marine Biology and Ecology Research Centre, School of Marine Science and
8 Engineering, Plymouth University, Davy Building, Plymouth, PL4 8AA, UK

9

10 ^b UNEP World Conservation Monitoring Centre (UNEP-WCMC), Cambridge,
11 CB3 0DL, UK

12

13 ^c Centre for Marine and Coastal Policy Research (MarCoPol), School of Marine
14 Science and Engineering, Plymouth University, Plymouth, PL4 8AA, UK

15

16 **Abstract:** Continued anthropogenic carbon dioxide emissions are acidifying our
17 oceans, and hydrogen ion concentrations in surface oceans are predicted to
18 increase 150% by 2100. Ocean acidification (OA) is changing ocean carbonate
19 chemistry, including causing rapid reductions in calcium carbonate availability
20 with implications for many marine organisms, including biogenic reefs formed by
21 oysters. The impacts of OA are marked. Adult oysters display both decreased
22 growth and calcification rates, while larval oysters show stunted growth,
23 developmental abnormalities, and increased mortality. These physiological
24 impacts are affecting ecosystem functioning and the provision of ecosystem
25 services by oyster reefs. Oysters are ecologically and economically important,

26 providing a wide range of ecosystem services, such as improved water quality,
27 coastlines protection, and food provision. OA has the potential to alter the
28 delivery and the quality of the ecosystem services associated with oyster reefs,
29 with significant ecological and economic losses. This review provides a
30 summary of current knowledge of OA on oyster biology, but then links these
31 impacts to potential changes to the provision of ecosystem services associated
32 with healthy oyster reefs.

33

34 Keywords: climate change; shellfish; aquaculture; sustainability; ecosystem
35 approach.

36

37 **1. Introduction**

38

39 The risks arising from climate change are now widely acknowledged as a major
40 cause for concern, yet awareness of ocean acidification is far less prevalent
41 (Gattuso et al., 2015). Consequently, our understanding of the scope and
42 severity of ocean acidification (OA herein) and its impacts on the marine
43 environment remain relatively limited, and especially, the implications of OA to
44 the continued provision of valuable ecosystem services.

45

46 Since 1750, the oceans have absorbed approximately 30% of anthropogenic
47 CO₂, altering oceanic carbonate chemistry by reducing carbonate ion
48 concentrations (CO₃²⁻), and reducing the saturation states of calcite and
49 aragonite. The result – lower pH, or ‘ocean acidification’ (Gattuso et al., 2014).
50 Historic OA linked to the Permian-Triassic mass extinction led to the

51 disappearance of ~90% of marine species (Clarkson et al., 2015). Today,
52 without significant cuts in CO₂ emissions, a 150% increase in the concentration
53 of surface ocean H⁺ is predicted by 2100 (Stocker et al., 2013).

54

55 OA may be of benefit to some organisms, such as jellyfish and toxic species of
56 algae (Hall-Spencer and Allen, 2015; Uthicke et al., 2015), but for other species,
57 such as corals and molluscs that use calcium carbonate in their structures, OA
58 is expected and has been shown to cause considerable direct harm (Basso et
59 al., 2015; Comeau et al., 2015; Gazeau et al., 2014; Houlbrèque et al., 2015;
60 Kim et al., 2016; Milazzo et al., 2014; Sui et al., 2016; Tahil and Dy, 2016). It is
61 therefore unsurprising that it is the negative effects of OA on individual
62 organisms that have received the most attention in the literature to date (see
63 reviews by Albright, 2011; Brander et al., 2012; Fabricius et al., 2014; Gazeau
64 et al., 2013; Hoegh-Guldberg et al., 2007; Pandolfi et al., 2011; Parker et al.,
65 2013). However, the ecosystem effects and loss of ecosystem services
66 associated with OA remain conspicuously absent, despite the increased
67 prevalence of ecosystem-based approaches in environmental legislation and
68 management. Here, we address that gap and introduce the current state of
69 knowledge required to underpin a multidisciplinary evaluation (Knights et al.,
70 2014), that considers the ecological, social and economic consequences of OA.

71

72 We have focused our review on an important ecosystem engineer (*sensu* Jones
73 et al., 1996) and commercially valuable species, the oyster, although much of
74 the discussion will also be relevant to other reef forming species. Oysters
75 provide a number of ecosystem services (ESs herein) to society, including the

76 formation of extensive reef structures that not only improve water quality, but
77 are also an important food source (see Section 3). Worldwide, oyster reefs have
78 dramatically declined in the past century and are now at the centre of many
79 conservation measures and restoration strategies (Beck et al., 2011; Grabowski
80 and Peterson, 2007), but these efforts are being undermined by OA. A plethora
81 of recent reviews and meta-analyses have highlighted the threat of OA to
82 marine fauna (see references above), but are often restricted to the description
83 of biological effects on a range of taxa, and do not focus on specific species or
84 groups of organisms (but see Albright, 2011; Gazeau et al., 2013; Hoegh-
85 Guldberg et al., 2007; Parker et al., 2013, for reviews on corals and shelled
86 molluscs). To date, there have been no reviews focused on oysters, despite
87 their ecological, economic and societal value.

88

89 This review is in two parts: firstly, we undertake a brief review of the biological
90 and ecological impacts of OA on oysters. This includes an assessment of the
91 effects of OA on individual life history stages (planktic larvae and sessile
92 juveniles and adults), populations and ecosystem-level responses. We then
93 review the range of ecosystem services that are provided by oysters, including
94 an assessment of their economic value and associated metrics. We conclude
95 by considering how impacts at the organismal-level can affect the provision of
96 ecosystem services.

97

98 **2. The biological impacts of ocean acidification on oysters**

99

100 2.1. Effects of OA on reproduction and planktic life-history stages

101

102 The planktic larval stage is a crucial life-history stage for many benthic
103 organisms and changes in development, performance or survival of this stage
104 can critically influence juvenile-adult population dynamics and ecosystem
105 functioning (Bachelet, 1990; Green et al., 2004; Rumrill, 1990). The early
106 development stages of marine calcifiers were quickly identified as particularly
107 vulnerable to OA, with the potential to alter population size and dynamics, and
108 community structure (Kurihara, 2008). As such, there has been a burgeoning
109 literature describing larval responses to OA (reviewed in Albright, 2011; Byrne,
110 2011; Przeslawski et al., 2014; Ross et al., 2011).

111

112 OA has been shown to induce narcotic effects on motile life-history stages,
113 reducing fertilisation success (Byrne, 2011). In a number of instances, OA
114 effects include reduced sperm motility, reductions in fertilisation success and
115 hatching rates of embryos (Barros et al., 2013; Parker et al., 2009; Parker et al.,
116 2010), although in the case of Parker et al. (2009), changes could not be solely
117 attributed to OA due to the effects being conflated with suboptimal culture
118 temperatures. However, OA-induced narcosis has not been consistently shown,
119 with disparity between studies of the same species (e.g. Kurihara et al., 2007;
120 Parker et al., 2012). Parker et al. (2012) suggest this disparity may be the result
121 of intraspecific phenotypic plasticity, whereas Byrne (2011) argues that the
122 fertilisation process in marine invertebrates can be resilient to fluctuations in pH
123 and may not be a reliable end-point. Neither Parker et al. (2012) or Byrne's
124 (2011) theories have been tested, but the inconsistencies shown highlight the

125 need for comparative studies using discrete populations to determine if OA has
126 consistent and repeatable effects, irrespective of scale or location.

127

128 In contrast to the fertilisation process, embryos and larvae are considered less
129 tolerant to the effects of OA (Kroeker et al., 2010; Parker et al., 2012), in part
130 because molluscs and other calcareous shell-forming species commonly lack
131 the specialised ion-regulatory epithelium used to maintain acid-base status
132 (reviewed in Lannig et al., 2010). The process of shell mineralisation begins at
133 the trochophore (prodissoconch I) stage (reviewed in Gazeau et al., 2013).

134 Larvae use two types of calcium carbonate, firstly mineralising highly soluble
135 amorphous calcium carbonate (ACC) (Brečević and Nielsen, 1989) before
136 switching to aragonite (Weiner and Addadi, 2002; Weiss et al., 2002). In
137 juvenile and adult stages, this again changes to the use of low solubility calcite
138 instead (Lee et al., 2006; Stenzel, 1964). Because the calcium carbonate
139 structures formed in these early life-history stages play a crucial role in
140 protection, feeding, buoyancy and pH regulation, disruption of calcification from
141 OA could have significant consequences for survival (Barros et al., 2013;
142 Simkiss and Wilbur, 2012). In other taxa, OA has been shown to greatly alter
143 the structure of the important larval shell of calcifying organisms, including
144 affecting dissolution rates and causing shell malformation, stunted growth,
145 altered mineral content, and weaker skeletons (reviewed in Byrne, 2011; and
146 Kurihara, 2008).

147

148 OA can also affect development rates. Multi-stressor experiments manipulating
149 $p\text{CO}_2$, pH, total alkalinity, and Ω_{arag} in order to simulate future acidification

150 scenarios have shown that oyster larvae are highly sensitive to predicted future
151 conditions. Responses include lower survivorship, abnormal development,
152 smaller body size, and altered shell properties (Gazeau et al., 2013; Guo et al.,
153 2015; Hettinger et al., 2013a; Hettinger et al., 2012; Parker et al., 2009; Parker
154 et al., 2013; Talmage and Gobler, 2009; Timmins-Schiffman et al., 2012;
155 Watson et al., 2009). However, the response remains inconsistent, with
156 differences between species and regions apparent (see Gazeau et al., 2011;
157 Kurihara et al., 2007; Parker et al., 2010 for a regional comparison of
158 *Crassostrea gigas* performance), with the differences within species suggestive
159 of pre-adaptation determined by exposure in their respective natural
160 environment (*sensu* environmental filtering, Kraft et al., 2015).

161

162 OA places individuals under stress as they try to regulate or maintain
163 physiological function. Processes including shell mineralisation, maintenance of
164 internal acid-base balance, somatic growth, swimming and feeding are
165 energetically expensive (Pörtner et al., 2004), and require additional energy for
166 maintenance under OA (Pörtner, 2008). As such, the planktic larval stage can
167 be extended to allow individuals to compensate for inefficient feeding and
168 delayed development, but doing so may subsequently affect the fitness,
169 competitive ability and survivorship of the individual at later life-history stages
170 (Anil et al., 2001; Gazeau et al., 2010; Rumrill, 1990; Talmage and Gobler,
171 2009). Trade-offs between calcification and other physiological aspects are
172 expected to occur, but the extent to which these occur and their impact, will
173 depend on an individuals' ability to obtain sufficient energy from their

174 environment to counteract any negative effects of acidification (Hettinger et al.,
175 2013a).

176

177 2.2. Carry-over or latent effects: metamorphosis to juvenile-adult stages

178

179 Changes in larval fitness are expected to impact adult population success
180 through a combination of latent/carry-over effects (see Pechenik, 2006) and
181 bottleneck effects (Schneider et al., 2003). These effects may only be transient,
182 for instance, in cases where larval development is slower and the increased
183 risks associated with extended larval duration (e.g. mortality from predation,
184 starvation) enhance bottleneck effects. However, if larval development is
185 unchanged and larval duration does not vary, the full suite of consequences will
186 be transferred to the juvenile (carry-over effects). Consequences may include
187 reduced environmental tolerance, decreased predation resistance, and
188 increased mortality, which can introduce an additional bottleneck for the adult
189 population (see Gaylord et al., 2011).

190

191 The negative impacts of OA on both pre- and post-settlement processes in
192 oysters are clear. These include: reduced metamorphosis success (Hettinger et
193 al., 2013a); greater mortality of juveniles (Beniash et al., 2010; Dickinson et al.,
194 2012); shell weakening (Dickinson et al., 2012) and greater prevalence of
195 micro-fractures; a reduction in shell dry mass, soft-tissue mass (Beniash et al.,
196 2010; Dickinson et al., 2012) and growth (Hettinger et al., 2013b; Hettinger et
197 al., 2012; Parker et al., 2011).

198

199 Metamorphosis is a crucial step in population development and growth and
200 mortality is often high due to the high energetic cost (Gosselin and Qian, 1997;
201 Videla et al., 1998). OA can lead to the depletion of energy reserves (e.g. lipids),
202 impair larval fitness and decrease the likelihood of successful metamorphosis
203 by up to 30% (e.g. Talmage and Gobler, 2009). A delay in metamorphosis can
204 reduce energy reserves and lead to settlement in suboptimal habitat, such that
205 post-settlement competence is impaired and mortality rates increased (see the
206 extensive work by Jan Pechenik, including Pechenik, 2006; Pechenik et al.,
207 1998). Given that post-settlement mortality often exceeds 90% under natural
208 environmental conditions (Thorson, 1950), any additional impact on juveniles
209 fitness or survival associated with OA impacts is likely to lead to significant
210 consequences in terms of adult population density.

211

212 It has been suggested that negative consequences of OA on juvenile and adult
213 oysters are carried-over as a result of energetic deficits experienced during the
214 early (planktonic) life stage (Hettinger et al., 2013b); a less fit larva will likely
215 become a less fit (e.g. smaller) juvenile/adult (Hettinger et al., 2012). These
216 consequences may, in general, be negative but individuals from specific regions
217 or some species appear to have developed coping mechanisms. Ko et al.
218 (2013) recently provided evidence of compensatory mechanisms in *C. gigas*
219 juveniles raised under acidified conditions, in which individuals displayed more
220 rapid calcification (without a reduction in shell thickness or change in
221 microcrystalline structure). Parker et al. (2011) demonstrated that some bred
222 populations of *Saccostrea glomerata* were more resistant to OA than wild
223 populations. While both bred and wild-caught individuals displayed significant

224 reduction in shell growth at pH 7.8, wild populations were more susceptible to
225 OA conditions than the bred population. The authors argue that these
226 differences could be due to phenotypic plasticity emanating from different
227 parental history, and/or differences in enzymatic activity of carbonic anhydrase -
228 an enzyme linked with acid-base regulation and shell formation. Irrespective of
229 the mechanism(s) of resilience, some species may be better suited for future
230 OA conditions, and identification of resilient lineages may provide important
231 insights for future food biosecurity and decision-making (see part 3.3).

232

233 2.3 OA impacts on adult oysters

234

235 Early studies indicated adult oysters were relatively robust to OA, leading to the
236 rapid focus on early life history stages. Negative responses of adults were
237 largely overlooked (see review by Gazeau et al., 2013; Parker et al., 2013),
238 despite evidence of reduced calcification and shell growth (Beniash et al., 2010;
239 Gazeau et al., 2007; Parker et al., 2009; Ries et al., 2009; Waldbusser et al.,
240 2011b; Wright et al., 2014), decreased shell density, weight, and strength
241 (Bamber, 1990; Welladsen et al., 2010), increased shell dissolution
242 (Waldbusser et al., 2011a), and increased mortality (Beniash et al., 2010; Dove
243 and Sammut, 2007). Metabolic activities have also been shown to be impaired,
244 but have received little attention to date, with additional studies needed. In the
245 few studies available, a consistent response across species has been apparent,
246 notably impaired feeding activity (Dove and Sammut, 2007). Impaired filtration
247 and feeding has the potential to effect energy supply and metabolic

248 maintenance in adult oysters, and may affect resilience and persistence to OA
249 in the long term.

250

251 Lower pH has been shown to affect the immune response of several taxa
252 (bivalves and echinoderms, Asplund et al., 2014; Beesley et al., 2008; Bibby et
253 al., 2008; Dupont and Thorndyke, 2012; Matozzo et al., 2012), but not others
254 (decapods, Small et al., 2010). In oysters, several studies have shown negative
255 effects. Li et al. (2009) noted impediment to metabolic activities of *C. gigas*
256 under food deprivation and extended recovery time post-spawning, which they
257 linked to an impaired immunological response. Wang et al. (2016) recently
258 demonstrated that reduced pH (≤ 7.8) negatively impacted the immune system
259 of *C. gigas* by increasing haemocyte apoptosis and reactive oxygen species
260 production, inhibiting the activity of antioxidant enzymes, and influencing the
261 mRNA expression pattern of immune related genes.

262

263 In *Crassostrea virginica*, *Crassostrea angulata*, and *C. gigas*, suppression of
264 immune-related functions including haemocyte production and antioxidant
265 defence was compromised leading to greater sensitivity to metal pollutants
266 (Ivanina et al., 2014; Moreira et al., 2016). Li et al. (2015) showed short-term
267 OA and warming significantly altered immune parameters of the Pearl oyster,
268 *Pinctada fucata*, impacting acid-base regulation capacity, immune system
269 functioning and bio-mineralization. Altered immune functions of bivalves from
270 acidification can potentially lead to higher susceptibility to pathogens (Asplund
271 et al., 2014; Beesley et al., 2008; Bibby et al., 2008; Ellis, 2013). For example,
272 *Vibrio tubiashii*, a well-known pathogen of oysters that causes significant losses

273 to the aquaculture sector (Dorfmeier, 2012; Elston et al., 2008; Richards et al.,
274 2015a) grew more quickly, increasing the likelihood of outbreaks, although the
275 susceptibility of the oysters to pathogens did not change.

276

277 2.4 From individual to ecosystem-level effects: consequences for the oyster reef

278

279 It is apparent that oysters, like many other marine invertebrates, are vulnerable
280 to OA, yet consequences at the population and ecosystem level remain largely
281 unknown. Many of the effects of OA are non-lethal, but substantial ecosystem
282 changes can be expected as systems become restructured, with 'winners' and
283 'losers', through environmental filtering and niche partitioning (Barry et al.,
284 2011; Fulton, 2011; Kraft et al., 2015; Somero, 2010). Such changes have
285 already been seen in other taxa based on laboratory, mesocosm and field-
286 based experiments, which all show that acidification favours some species,
287 such as macroalgae and invasive invertebrates (Hall-Spencer and Allen, 2015),
288 over others (Brodie et al., 2014). Ecosystem changes can be dramatic. For
289 example, Christen et al. (2013) observed a phase shift from a calcareous-
290 dominated system to one dominated by non-calcareous species. In Australia,
291 the introduced Pacific oyster, *C. gigas*, is more resilient to acidification than the
292 native *S. glomerata* (Parker et al., 2010) and may dominate interactions in the
293 future. Should there be a shift in dominance toward the non-native species, the
294 question remains as to whether *C. gigas* can provide functional redundancy and
295 continue to provide the suite of ESs currently derived from *S. glomerata* or be
296 lost altogether?

297

298 OA may affect community composition by altering interspecific interactions,
299 between different trophic levels. For oysters, OA is predicted to lead to greater
300 vulnerability to predation due to the negative effects on shell dissolution and
301 microstructure (see references above). In some instances, the predator
302 themselves may also be affected. For example, Sanford et al. (2014) found that
303 *Urosalpinx cinerea*, a major gastropod predator of *Ostrea lurida*, consumed
304 significantly more oysters in acidified treatments than in control treatments.
305 Suggested reasons for this were reduced energetic value of the prey species,
306 reduced prey handling time, increased energetic requirements of the predator
307 or a combination of these (Kroeker et al., 2014).

308

309 Sites with naturally acidified conditions, such as volcanic seeps, lagoons and
310 upwelling areas, can provide insights into long-term community responses to
311 OA, particularly in areas subjected to anthropogenic stress (see studies by
312 Basso et al., 2015; Range et al., 2012; Thomsen et al., 2012; Thomsen et al.,
313 2010; Tunnicliffe et al., 2009). Field studies of mussel and vermetid reefs have
314 shown that, in oligotrophic conditions, reduced pH levels benefit non-calcified
315 algae, but impairs mollusc larval recruitment and dissolves carbonate habitat
316 (Cigliano et al., 2010; Comeau et al., 2015; Kroeker et al., 2012; Milazzo et al.,
317 2014; Rodolfo-Metalpa et al., 2011). Studies of the impacts of shallow-water
318 oyster reef degradation show that loss or damage to these habitats can trigger
319 cascading effects, including loss of biodiversity, reductions in biofiltration, and
320 loss of coastal habitat protection (Rossoll et al., 2012).

321

322 2.5 Potential for adaptation

323

324 There are concerns over whether species will be able to adapt to the current
325 unprecedented rates of environmental change (Somero, 2010; Sunday et al.,
326 2014; Visser, 2008). Fast-generating species such as the pond algae,
327 *Chlamydomonas*, did not show evidence of adaptation to acidified conditions
328 after 1000 generations (Collins and Bell, 2004), although recently, the marine
329 polychaete, *Ophryotrocha labronica*, demonstrated acclimation after just two
330 generations (Rodríguez-Romero et al., 2015) indicating vastly different
331 propensity for adaptation. It might be reasonable to assume that long-lived,
332 slow generation time species, such as oysters, are even less likely to evolve
333 rapidly (Barry et al., 2011; Byrne, 2011). However, Parker et al. (2015; 2012)
334 demonstrated conferred tolerance from adults exposed to elevated CO₂ to their
335 offspring (also dependent on if the oysters originated from wild or aquaculture-
336 reared populations), suggesting hereditary traits as a mechanism of resilience.
337 But as in fast generation time species, the potential for adaptation may be
338 species-specific, as well as dependent on the heritage of the individual (Parker
339 et al., 2010; Parker et al., 2011 respectively; Thompson et al., 2015).

340

341 A possible mechanism through which resilience to environmental stress can be
342 achieved may be linked to the 'quality' of the environment. In the Baltic Sea,
343 seasonal upwelling of eutrophic water leads to very high CO₂ levels and
344 phytoplankton blooms, nevertheless, blue mussel reefs are able maintain their
345 structure and function. Thomsen et al. (2012) argued that the high food supply
346 allows the mussels to offset the metabolic costs of hypercapnia (Thomsen et al.,
347 2012); a hypothesis that would support why aquaculture-reared individuals may

348 be more tolerant to OA than wild individuals. However, what is not yet clear is
349 the extent to which resilience and survival is achieved at the expense of ES
350 provision?

351

352 **3. The impacts of ocean acidification on oyster-reef ecosystem services**

353

354 3.1 Using ESs to assess the impacts of OA

355

356 3.1.1 Assessing and valuing ESs

357

358 ESs allow society to evaluate and estimate the social and economic impacts of
359 changes in resource availability (Beaumont et al., 2007; Cooley, 2012). They
360 provide a tangible link between ecosystem health and human use (Figure 1)
361 that help to inform decision-making of how to sustainably use and protect
362 ecosystems (e.g. deciding whether to provide financial incentives, Knights et al.,
363 2014). This is especially important as human well-being is linked to the
364 sustained provision of resources including food, fuel, shelter, and water (Díaz et
365 al., 2006). However, clear context and definition of ESs is critical to their
366 perceived 'value' (Friedrich et al., 2015).

367

368 There are a number of ways to value ESs. The most obvious - '*monetary value*'
369 - can be useful in conjunction with frameworks, such as the *Total Economic*
370 *Valuation* (TEV) framework, which provides a classification of the different types
371 of economic values associated with ESs (DEFRA, 2013; Herbert et al., 2012). In
372 TEV, ESs are classified as either 'use values' (those derived from human

373 interactions with a particular resource) or 'non-use values' (derived solely from
374 the knowledge that the resource exists currently and will continue to exist in the
375 future (Figure 2)).

376

377 3.1.2 Valuing oyster-derived ESs

378

379 The value of a service can be relatively easily estimated when a particular ES
380 has a market price. For example, in 2009, Pacific oysters bought for the food
381 industry were valued at £1,815/tonne (Herbert et al., 2012). Other services,
382 such as carbon sequestration (carbon credit values), or raw materials (shell
383 cultch) can be valued in a similar way. But when no market price exists or the
384 ES value is subjective, estimating value can be a difficult task. In these
385 instances, approaches such as 'Willingness-to-Pay' (WTP. e.g. De Groot et al.,
386 2002; Fletcher et al., 2014) can provide a 'value' based on what people are
387 willing to pay to avoid or counteract the adverse effects of the loss of the ES.
388 The WTP approach is less precise than using market values, but can still be
389 used to infer ecosystem or ES value.

390

391 3.1.3 ESs and OA

392

393 Future OA conditions are predicted to negatively affect the availability and
394 quality of ESs through direct or indirect effects on species and the ecosystem
395 (Frommel et al., 2012) with major social and economic consequences (Figure 3)
396 (Cooley et al., 2009). Studies have typically focused on particular habitats, such
397 as coral reefs (Brander et al., 2012), locations of significant value (Bosello et al.,

398 2015; Hilmi et al., 2014; Lacoue-Labarthe et al., 2016; Rodrigues et al., 2013),
399 or an economic sector (Cooley and Doney, 2009; Moore, 2011; Narita and
400 Rehdanz, 2016; Narita et al., 2012). Although an increasing number of studies
401 have tried to quantify the economic implications of OA on ES provision (Brander
402 et al., 2012; Cooley and Doney, 2009; Moore, 2011; Narita et al., 2012), they
403 are often qualitative in nature due to a lack of quantitative data (Cooley, 2012;
404 Hilmi et al., 2014; Hilmi et al., 2013). While qualitative assessments are
405 valuable for indicating directionality of change in ES provision, it is challenging
406 to include the findings in management decisions and investment prioritization
407 without clear quantitative estimates of change. Therefore, future studies
408 providing more quantitative estimates are critically needed in order to bridge
409 those gaps.

410

411 3.2 Oyster-reefs ESs, their value, and impacts of OA

412

413 3.2.1 Oyster reef ecosystem services

414

415 Biogenic reefs provide a wide range of ecosystem services including supporting,
416 provisioning, regulating, and cultural services (MEA, 2005; Teagle et al. this
417 issue) (Figure 1). Oysters are ecosystem engineers (Jones et al., 1996; Jones
418 et al., 1997), in that they form biogenic reefs providing habitat for a range of
419 other species and contribute a number of ecosystem functions and services
420 (Fletcher et al., 2012; Herbert et al., 2012) (Figure 4, Table 1). Importantly,
421 oysters are both allogenic and autogenic engineers, and ESs originate from
422 both individual oysters and the wider reef structure (Wallis et al., 2015).

423 Allogenic ESs include water filtration, benthic-pelagic coupling, nutrient cycling,
424 carbon sequestration, and food provision (from oyster harvesting), while
425 autogenic ESs include habitat formation, food provision, erosion protection and
426 shoreline stabilization (Figure 4, Table 1). Additionally, cultural services
427 associated with oyster reefs include recreational harvesting, educational use
428 (research) and cultural heritage (Paolisso and Dery, 2010; Scyphers et al.,
429 2014).

430

431 The perceived and relative importance of oyster-derived ESs is context-specific
432 (Scyphers et al., 2014). Local or regional characteristics, such as environmental
433 conditions, but also local economy and communities, can influence the
434 significance of each service, but also their susceptibility to OA. For instance,
435 biofiltration is a particularly valued ES in areas that are: polluted or susceptible
436 to eutrophication; intensely used for recreation; or located near to seagrass
437 beds (Cerco and Noel, 2007). Alternatively, shoreline stabilisation and
438 protection from erosion by oysters is of importance in coastal areas under threat
439 of climate change and extreme weather events (Brumbaugh et al., 2010; La
440 Peyre et al., 2014), and in other areas, seafood is considered most important
441 (Cooley et al., 2012).

442

443 Determining the importance of ESs to society and the economy can be
444 achieved using one of the valuation methods described in brief above. Several
445 studies have estimated the value of ESs association with oyster reefs (Beseres-
446 Pollack et al., 2013; Grabowski et al., 2012; Grabowski and Peterson, 2007;
447 Henderson and O'Neil, 2003; Kroeger, 2012; Volety et al., 2014) but estimates

448 are wide-ranging (e.g. Grabowski et al. (2012) valued oyster reefs at between
449 \$5500-\$99000 ha⁻¹ yr⁻¹, excluding the economic value from harvesting). Given
450 local/regional priorities may vary, such varied estimates are unsurprising, but
451 makes predicting the economic impact of OA on oyster reefs challenging and
452 perhaps explains why no study has attempted to do so to date. In lieu of such
453 an analysis, studies that estimate the economic losses emanating from
454 damaged oyster reefs may be a useful proxy. Here, we use these studies to
455 provide a first assessment of OA impacts on ES provision from oyster reefs.
456 While it can be argued that drawing conclusions on the effects of OA from
457 damaged reefs is unrealistic or inaccurate, it has the merit to reinforce the high
458 value of oyster reef ESs, allow an initial estimate of losses to be undertaken,
459 and provides direction for future studies.

460

461 3.2.2 Current state of oyster reefs

462

463 Despite the numerous ESs provided by healthy oyster reefs, reef conservation
464 and health is rarely considered unless populations are in danger of collapsing or
465 are threatened (Kirby, 2004). It is only in these instances that the value of
466 oyster reefs is assessed; the outcomes used to direct restoration efforts (Beck
467 et al., 2011; Coen et al., 2007; Grabowski et al., 2012; Grabowski and Peterson,
468 2007; La Peyre et al., 2014; Volety et al., 2014). A recent study estimated that
469 85% of native oyster reefs had been lost globally and that in many bays and
470 ecoregions, reefs were less than 10% of their historical abundance and
471 'functionally extinct' (Beck et al., 2011). As such, there is an urgent need for an
472 assessment of ESs provided by oyster reefs.

473

474 There is, however, on-going debate over what is considered an acceptable level
475 of 'ecological health' for oyster reefs (Alleway and Connell, 2015). While it is
476 recognised that reef degradation can lead to a decrease in, or even loss of,
477 provision of many ESs (Coen et al., 2007; Table 1; Jackson et al., 2001; Paerl
478 et al., 1998; zu Ermgassen et al., 2013), the impact of OA on ESs derived from
479 oyster reefs is less clear. In fact, the potential consequences of OA are largely
480 speculative, and based either on the physiological and ecological impacts
481 observed during laboratory experiments, or using *in situ* field experiments in
482 naturally acidified sites (see Section 2.4). These studies suggest that population
483 sizes may decrease below the minimum threshold required for the desired level
484 of ESs (Figure 4) and indicate an overall negative effect of OA on oysters' early
485 life stages, with direct consequences on juvenile and adult physiology as well
486 as recruitment and population dynamics (see discussion in Section 2.4).
487 However, while there have been studies examining the effects of OA on
488 filtration rates, growth, and survival of oysters, there has been no attempt to
489 date to link those effects to ES provision.

490

491 3.2.3 Ecosystem services associated with oyster reefs

492

493 Oyster reefs provide a number of important provisioning, regulating, habitat and
494 cultural ESs (Table 1). Below we describe each of the ESs in turn, grouped
495 using the MEA (MEA, 2005) and more recent ODEMM assessment (White et al.,
496 2013), and consider how OA is likely to affect the provision of those services in
497 the future.

498

499 3.2.3.1 Habitat/Structural Services

500

501 Oysters create large complex 3-dimensional structures (Knights and Walters,
502 2010), providing a unique habitat for many organisms, assuring life-cycle
503 maintenance, and securing a diverse gene pool (Table 1). The maintenance of
504 the reef structure relies notably on the successful recruitment of oyster larvae
505 and juveniles into the adult population. By creating a mismatch between
506 environmental conditions and larval/juvenile performance (see Section 2), OA
507 can have direct negative impacts on the formation and replenishment of the reef
508 structure (Table 1). Moreover, it can hold further negative consequences for the
509 reef, by lowering gene pool diversity and creating additional bottlenecks. OA-
510 induced reef deterioration is likely to alter the available niches for other species,
511 and restructure the overall habitat, which can hold critical consequences for the
512 provision of other ESs.

513

514 3.2.3.2 Provisioning Services

515

516 *Biodiversity and Seafood (other than oysters)*

517 Biogenic reefs are important to food webs and fisheries (Peterson and Lipcius,
518 2003) and many organisms use reefs as refugia from predators, as nests, or
519 feeding grounds. Oyster reefs create critical habitat for a number of species,
520 including other economically important molluscs, crustaceans, and various
521 species of fish (reviewed in Coen et al., 2007; Coen et al., 1999; La Peyre et al.,
522 2014; Scyphers et al., 2011; Tolley and Volety, 2005; Volety et al., 2014). In a

523 recent analysis, Grabowski et al. (2012) estimated that one ha. of healthy oyster
524 reef increases biodiversity and enhances commercial fish value by up to
525 ~\$4123 yr⁻¹ (see also Grabowski and Peterson, 2007; Table 1), although that
526 value-added benefit may only be apparent when the oyster reef is not located
527 near to other biogenic habitats that provides a similar function (Geraldi et al.,
528 2009). The value of oyster reefs can be used to justify a management action
529 (Knights et al., 2014), for example, the decision to restore an oyster reef
530 following degradation (an effective method for increasing fish and large
531 crustaceans production, Peterson et al., 2003). At the time of writing, there is no
532 indication or means of disentangling the 'value' of oyster reefs to biodiversity
533 and seafood in differing states of health, nor a clear understanding of the likely
534 state of oyster reefs under OA beyond an expectation that some reefs will be
535 more susceptible to damage and lead to a reduction in ES provision. Current
536 research is on going to determine if other/non-native species, appearing more
537 robust to OA, can provide redundancy for the loss of biogenic reef species
538 threatened by OA.

539

540 *Ornamental and raw materials*

541 Oyster shell is valuable to a number of sectors (Yao et al., 2014), including
542 construction, agriculture, and wastewater treatment. The cost of oyster cultch is
543 relatively low (~\$126/tonne, Kwon et al., 2004; Table 1) and the calcium
544 carbonate from the shells is used as grit for the rearing of poultry (Çath et al.,
545 2012; Scott et al., 1971); as a construction material and substitute for aggregate
546 in concrete or mortar (Yang et al., 2010; Yang et al., 2005; Yoon et al., 2003;
547 Yoon, 2004); as a liming material and soil stabilizer (Lee et al., 2008; Ok et al.,

548 2010); and to treat discharged wastewaters to remove phosphates and traces
549 of toxic heavy metals such as cadmium, copper and nickel (Hsu, 2009; Kwon et
550 al., 2004). As shell dissolution rates increase with OA, the availability of oyster
551 shell for use as a raw material may decline in the future (in terms of abundance
552 and/or 'quality'), increasing the cost of the material to industry and raising the
553 cost to the consumer.

554

555 The use of oysters for ornamental purposes has decreased in modern times,
556 but there are historical references of the use of oyster shells as raw material for
557 the creation of glass (Wedepohl and Baumann, 2000). Shells are still available
558 commercially, with values ranging from £5-15 a shell (Lemasson et al.
559 *unpublished*). OA is likely to impact on the provision of raw materials, by
560 negatively impacting on the calcification process of oysters, promoting adult
561 shell dissolution, and may well reduce the value of shells as a raw material. The
562 economic costs that would be incurred because of the loss of this service are
563 unclear, and no estimates are found in the literature to date.

564

565 3.2.3.3 Regulating Services

566

567 *Climate regulation and carbon sequestration*

568 A number of studies have suggested that oysters provide climate change
569 mitigation as a result of carbon sequestration during the calcification process
570 (Dehon, 2010; Grabowski and Peterson, 2007; Lee et al., 2010; Peterson and
571 Lipcius, 2003; Wingard and Lorenz, 2014). Although it can be argued that
572 oysters are net CO₂ producers on their own, this service is particularly valid

573 when oysters are present in association with algae (Hall et al., 2011). While the
574 calcification process in oysters has been extensively researched, the extent to
575 which climate can be mitigated is not easily quantifiable, and the impact on the
576 carbon cycle not easily determined (Hickey, 2008). Under OA, shell calcification
577 is negatively affected as shell dissolution increases (Welladsen et al., 2010) as
578 a result of corrosive waters. The carbon sequestration process and efficiency of
579 the climate regulation service is therefore at risk. At the time of writing, there
580 was no data assessing the value of such service, although estimates could be
581 derived from the price of carbon credits, or be related to bequest value of the
582 bio-sequestration of CO₂ (Herbert et al., 2012).

583

584 *Disturbance prevention and coastal erosion protection*

585 The 3-dimensional structure of oyster reefs provides protection from erosion
586 and stabilising shorelines. Adjacent critical habitats, such as seagrass beds or
587 saltmarshes, positively benefit from the attenuation of wave action and the
588 modification of the local water flow (Coen et al., 2007; Scyphers et al., 2011). It
589 is argued that shore stabilisation and erosion protection are the most valuable
590 ESs provided by oyster reefs (Grabowski et al., 2012). How OA will affect the
591 maintenance of the 3-dimensional reef structure is unclear (see also
592 provisioning services above), but the impacts on early life history, juvenile and
593 adult forms (discussed in Section 2) are predicted to lead to a reduction in reef
594 size, and therefore, a reduction or loss of coastal protection and shoreline
595 stabilisation properties. The 'value' of this ES is likely to continue to erode under
596 climate change as the number and intensity of extreme weather events

597 increases, such that the ability of reefs to attenuate wave action may be
598 reduced (Michener et al., 1997).

599

600 The cost of using engineered shoreline stabilization solutions as an alternative
601 to reefs can provide some insights into the value of this ES (termed 'avoidance
602 costs'), but as Grabowski et al. (2012) points out, this value is context-specific
603 and highly dependent on factors including the location, infrastructure types and
604 prices, local economy, and the level of exposure. Nevertheless, costs of coastal
605 defence structures can be in the a few dollars to millions of dollars depending
606 on the setting and requirements (Table 1, see Firth et al., 2014 for a review)

607

608 *Water treatment and quality*

609 Oysters are important bio-filters that improve water quality (Grizzle et al., 2008;
610 Hoellein et al., 2015; La Peyre et al., 2014; Nelson et al., 2004; Newell, 2004)
611 and affect nutrient cycling (Beseres-Pollack et al., 2013; Coen et al., 2007;
612 Hoellein et al., 2015; Kellogg et al., 2013; Newell et al., 2005; Piehler and
613 Smyth, 2011). A number of approaches have been used to attribute value to
614 this ES. Grabowski et al. (2012) used nitrogen permits as a proxy for market
615 price, estimating that one ha. of oyster reef removed a quantity of nitrogen
616 valued to \$1385-\$6716 yr⁻¹. Beseres-Pollack et al. (2013) used the costs of
617 adding a biological nutrient removal system to a wastewater treatment plant,
618 valuing the nitrogen removal service at \$293,993 yr⁻¹ in the Mission-Aransas
619 Estuary (NB this is equivalent to between 44 and 212 hectares of oyster reef
620 based on Grabowski et al., 2012 estimates). Oyster-mediated nitrogen removal
621 was valued at \$314,836 yr⁻¹ in the Choptank estuary, using an average

622 monetary value of ~\$24 kg⁻¹ of nitrogen removed specifically applied to their
623 study site (Newell et al., 2005), \$18,136 per million oysters in Chesapeake Bay
624 (Kasperski and Wieland, 2009), and \$3000 ac⁻¹ yr⁻¹ in Bogue Sound, using
625 values derived from the North Carolina nutrient offset trading program of \$13
626 per kg of nitrogen removed (Piehler and Smyth, 2011). These values are,
627 however, difficult to compare as they are based on multiple assumptions of
628 what constitutes 'value' and the bodies of water are of different sizes and
629 properties, but nevertheless highlight the important economic value of this
630 ecosystem service provided by oyster reefs.

631

632 The continued provision of this ES has important indirect benefits to other ES
633 provision that are often not considered. For example, improved water quality
634 affects the provision of ESs by other healthy and functioning species and
635 habitats, such as economically important seagrass beds (Cerco and Noel,
636 2007; Coen et al., 2007; Dennison et al., 1993; Grabowski et al., 2012; Kahn
637 and Kemp, 1985; Meyer et al., 1997). Increased seagrass bed coverage that
638 comes with improved water quality can also be incorporated into the economic
639 valuation of oyster reefs. Grabowski et al. (2012) estimated that one ha. of
640 oyster reef promoted 0.005 hectares of additional seagrass bed, valued at
641 \$2584 ha⁻¹. Therefore, under OA, further economic losses can originate from
642 adversely impacted adjacent habitats.

643

644 3.2.3.4 Cultural Services

645

646 *Recreation and leisure*

647 Improved water quality associated with health oyster reefs increases human
648 well-being by reducing the likelihood of eutrophication (Lipton, 2004), and
649 increasing recreational use of the environment. The value of recreation and
650 tourism can therefore be used as an indirect estimate of the value of oyster
651 reefs. Oyster reefs are socially recognised as a valuable resource; the U.S.
652 National Research Council (2004) valued reefs using willingness-to-pay
653 estimates at ~\$222 million (Grabowski and Peterson, 2007; see also Volety et
654 al., 2014). Reduced bio-filtration rates under OA will likely lead to increased
655 eutrophication and reduced water quality, diminishing public appeal and
656 generating financial losses from lower recreational use.

657

658 *Spiritual experience and cultural heritage*

659 Oysters hold a significant place in local culture, traditions and history. Many
660 countries have a long history of oyster harvest and consumption (Kirby, 2004),
661 including the USA (Dyer and Leard, 1994), France (Heral, 1989), and the UK
662 (Humphreys et al., 2014; Mac Con Iomaire, 2006), which is celebrated during
663 oyster festivals, such as the 'Bluff Oyster' in New Zealand (Panelli et al., 2008;
664 Rusher, 2003), 'Oysterfest' in Australia (Lee and Arcodia, 2011), and the
665 'Whitstable Oyster Festival' and the 'Falmouth Oyster Festival' in the UK.
666 Traditional oyster harvesting can be an important part of the local economy and
667 creates a sense of community and heritage, with the desire for this activity to be
668 sustainable and prosper (Dyer and Leard, 1994; Paolisso and Dery, 2010;
669 Scyphers et al., 2014). The impact of OA on the provision of cultural services is
670 difficult to assess, but a reduction in the persistence of oyster reefs (and the
671 number of harvestable oysters) will likely impact on the sense of heritage and

672 affect local economies and communities that rely on a long tradition of oyster
673 harvesting. In the worst case, OA could lead to the disappearance of oyster
674 festivals, leading to the loss of sense of tradition and community well-being, as
675 well as negative economic impacts locally through reductions of tourism.

676

677 3.3 Economic Impacts of OA on oyster harvest and aquaculture

678

679 The vulnerability of shellfisheries to OA and the likely economic consequences
680 is of growing concern (Ekstrom et al., 2015; Haigh et al., 2015; Seijo et al.,
681 2016). Although the 'value' of reefs is highly variable and context-dependent in
682 terms of harvest yield, OA is likely to negatively affect the resilience,
683 persistence, and sustainable use of wild oyster reefs into the future. In response,
684 wild harvests now represent only a small proportion of oyster production
685 worldwide, and increasingly replaced by aquaculture (130,754 tonnes of wild
686 harvest compared to 5,155,257 tonnes of aquaculture production; valued at
687 \$ 4,174,258,000 in 2014 - FAO data¹).

688

689 Aquaculture is the fastest growing food sector, and production reached an all-
690 time high of 90.4 million tonnes in 2012 (FAO, 2014). In the UK, the decline of
691 the native oyster, *O. edulis*, led to the introduction of the Pacific oyster, *C. gigas*,
692 which now represents over 90% of the country's oyster production (Humphreys
693 et al., 2014). Approximately 1200 tonnes of Pacific oysters are estimated to be
694 produced each year in the UK (Herbert et al., 2012) worth an estimated £10.14
695 million (Humphreys et al., 2014). Given the demand and value of shellfish

¹ <http://www.fao.org/fishery/statistics>

696 aquaculture, determining how OA will affect food security in the future is a
697 crucial question that remains unanswered, although in 2010, the United Nations
698 Environment Programme (UNEP) cited OA as a major threat to food security
699 (UNEP, 2010). At locations where the effects of OA are already being felt,
700 damages to the oyster aquaculture industry have been disastrous. In the US,
701 the Pacific North-west region hosts an oyster industry worth over \$72 million,
702 but since 2007, several oyster hatcheries have suffered from mass mortalities of
703 oyster larvae of up to 70-80% due to the upwelling of waters that were acidified,
704 highly saline and rich in *V. tubiashii* (Barton et al., 2015; Elston et al., 2008;
705 Feely et al., 2008). Similar impacts are expected in UK shellfisheries and
706 elsewhere under combined scenarios of acidification and warming (Callaway et
707 al., 2012).

708

709 Concerns have been expressed regarding the likely imbalance in the social and
710 economic consequences of OA experienced by different communities (Ciuriak,
711 2012; Cooley et al., 2012). Islands and coastal communities that rely heavily on
712 seafood as a source of protein and for their livelihood (e.g. tourism), are
713 expected to suffer most, particularly as their potential for adaptation and
714 mitigation is restricted due to lower financial means and limited access to
715 technologies (Cooley et al., 2009; Hilmi et al., 2014; Hilmi et al., 2013; UNEP,
716 2010). Seafood aquaculture holds the potential to provide food security in a
717 future where growing population and growing income are expected to increase
718 the demand for food. However, for aquaculture to be sustainable, there is a
719 need to recognise that the sector is nested in a sensitive system
720 interconnecting economic, social and ecological spheres, whereby impacts on

721 one sphere will likely disrupt the others (Bailey, 2008; Schmitt and Brugere,
722 2013; Soto et al., 2008).

723

724 Some argue that shellfish aquaculture will be impacted less by OA than wild
725 harvests (Rodrigues et al., 2013), for instance, the rearing of larval stages in
726 tanks may mitigate the impacts of OA. Further, the use of informed 'climate-
727 proof' management measures, designed to buffer the effect(s) of OA, such as
728 quality control and close monitoring of water quality may minimise any potential
729 effects (Hilmi et al., 2014; see also the case study on the mitigation of the
730 effects of ocean acidification on prawn and scallop fisheries in Australia by
731 Richards et al., 2015b). In the Pacific North-west, oyster hatcheries have
732 benefitted from close monitoring of seawater quality, which has greatly
733 improved yield and reduced mortality (Barton et al., 2015). Relocation of
734 hatcheries and farms to areas of higher water quality and environmental
735 conditions more suited to the rearing of larvae is an option already considered
736 by shellfish producers (Barton et al., 2015), although this may represent a costly
737 alternative. Aquaculture farms could focus on more resistant species, such as
738 *C. virginica* and *C. gigas* (Gobler and Talmage, 2014; Guo et al., 2015; Parker
739 et al., 2010), and hatcheries could select for lines that produce the highest
740 survival rates, such as selectively bred lines (Parker et al., 2011; Thompson et
741 al., 2015). However, such scenarios must rely on robust scientific knowledge of
742 the response of the different life stages of different species of oysters to OA.

743 Moore (2011) claims that damage to individuals, and especially their shell,
744 which appear to be susceptible to low pH (Welladsen et al., 2010), are unlikely
745 to hold significant economic impacts as long as the organism survives the

746 culture process. However, OA has the potential to impact on the final quality
747 and value of the product by affecting the wet tissue mass, the shell appearance
748 (due to corrosive waters), tissue texture and taste (Dupont et al., 2014), and
749 potentially their nutritional value.

750

751 OA is likely to increase production costs due to the necessary buffering
752 measures taken during husbandry procedures, such as pH and Ω_{ar}
753 manipulations and increased feeding, as well as the costs associated with the
754 longer developmental time of the early life stages, unless the production is
755 focused on OA-resistant species or lines. New models should be investigated in
756 order to link future levels of OA with variations in the production of oysters, and
757 to try and predict to what extent those variations in production are likely to
758 indirectly impact the economy due to welfare losses, and welfare effects of price
759 increase due to the reduced supply (Narita et al., 2012), changes in job
760 opportunities, and general reverberation into the wider economy. As Richards et
761 al. (2015b) stated “OA (*sic*) itself cannot be mitigated through fisheries
762 management, however management can be used to reduce the negative
763 effects and take advantage of positive effects associated with this
764 phenomenon”.

765

766 **4. Conclusion**

767

768 OA is likely to have negative effects throughout the life cycle of oysters,
769 although the effects may be difficult to decipher due to other stressors acting on
770 multiple life-history stages (Byrne and Przeslawski, 2013; Knights and Walters,

771 2010). Acclimation, parental (hereditary) traits, or adaptation may well reduce
772 the risk of negative consequences from OA, although the extent to which
773 individuals can respond to the threat remains to be seen.

774

775 Oysters are a key ecological species that provide a plethora of ES to humans,
776 but the provision, quantity, and quality of ESs in the future under climate
777 change and OA remains uncertain. OA is occurring alongside other
778 environmental and anthropogenic stressors, such as warming, hypoxia,
779 variations in salinity, eutrophication, or metal contamination, that are likely to
780 affect the organisms' responses. The outcomes of multi-stressor interactions
781 are difficult to predict and seem highly context-specific (see the reviews on the
782 impacts of ocean warming and acidification by Byrne, 2011; Byrne and
783 Przeslawski, 2013; and Harvey et al., 2013). Future studies should consider the
784 combination of multiple stressors, but should also focus on adult oysters in the
785 aim to link individual and population responses with the provision of associated
786 ESs. The consequences of OA on oysters are already being felt in parts of the
787 world (Cooley et al., 2015), where natural populations and hatcheries have
788 been negatively impacted upon; therefore ocean acidification will likely put
789 increased pressure on food security in the future, by reducing harvest and
790 aquaculture productions. Predicted levels of OA are likely to hold a number of
791 other significant social and economic consequences that goes beyond oyster
792 production from harvest and aquaculture, such as impoverished water quality,
793 reduced shoreline protection, or altered well-being.

794

795 The few studies to date that estimate the value of oyster-derived ESs give an
796 insight into the importance and value of oysters to the environment and society.
797 Whilst there is some variation in the economic value of oyster-derived ESs due
798 to local and/or regional differences in societal value placed on those services,
799 the role of oysters in supporting a healthy and functioning ecosystem is clear,
800 but which is under threat from OA. Further assessments of the social and
801 economic impacts of OA on oysters and oyster reefs are needed to emphasise
802 the 'value' of oysters to society, such that necessary steps are taken to ensure
803 their long-term future.

804

805 **5. Acknowledgements**

806

807 This research is supported by a grant awarded to AMK, JHS, and SF by the
808 School of Marine Science and Engineering, Plymouth University, as part of the
809 PhD research of AJL. Thank you also to Bayden Russell (editor) and two
810 anonymous reviewers whose comments greatly helped to improve this
811 manuscript.

812

813 **6. References**

814

- 815 Albright, R., 2011. Reviewing the effects of ocean acidification on sexual reproduction and early
816 life history stages of reef-building corals. *Journal of Marine Biology*, 1-14.
- 817 Alleway, H.K., Connell, S.D., 2015. Loss of an ecological baseline through the eradication of
818 oyster reefs from coastal ecosystems and human memory. *Conservation Biology*.

819 Anil, A.C., Desai, D., Khandeparker, L., 2001. Larval development and metamorphosis in *Balanus*
820 *amphitrite* Darwin (Cirripedia; Thoracica): significance of food concentration, temperature and
821 nucleic acids. *Journal of Experimental Marine Biology and Ecology* 263, 125-141.

822 Asplund, M.E., Baden, S.P., Russ, S., Ellis, R.P., Gong, N., Hernroth, B.E., 2014. Ocean acidification
823 and host-pathogen interactions: blue mussels, *Mytilus edulis*, encountering *Vibrio tubiashii*.
824 *Environmental microbiology* 16, 1029-1039.

825 Bachelet, G., 1990. Recruitment of soft-sediment infaunal invertebrates: the importance of
826 juvenile benthic stages. *La Mer* 28, 199-210.

827 Bailey, C., 2008. Human dimension of an ecosystem approach to aquaculture, in: Soto D, A.-M.J.,
828 Hishamunda N (Ed.), *Building an ecosystem approach to aquaculture*. FAO Fisheries and
829 Aquaculture Proceedings. No. 14. Rome, Italy: Food and Agriculture Organization,
830 FAO/Universitat de les Illes Balears Expert Workshop, 7–11 May 2007, Palma de Mallorca,
831 Spain, pp. 37–42.

832 Bamber, R.N., 1990. The effects of acidic seawater on three species of lamellibranch mollusc.
833 *Journal of Experimental Marine Biology and Ecology* 143, 181-191.

834 Barros, P., Sobral, P., Range, P., Chícharo, L., Matias, D., 2013. Effects of sea-water acidification
835 on fertilization and larval development of the oyster *Crassostrea gigas*. *Journal of*
836 *Experimental Marine Biology and Ecology* 440, 200-206.

837 Barry, J.P., Widdicombe, S., Hall-Spencer, J.M., 2011. Effects of ocean acidification on marine
838 biodiversity and ecosystem function. *Ocean Acidification*, 192-209.

839 Barton, A., Waldbusser, G., Feely, R., Weisberg, S., Newton, J., Hales, B., Cudd, S., Eudeline, B.,
840 Langdon, C., Jefferds, I., King, T., Suhrbier, A., McLaughlin, K., 2015. Impacts of coastal
841 acidification on the Pacific Northwest shellfish industry and adaptation strategies implemented
842 in response. *Oceanography* 25, 146-159.

843 Basso, L., Hendriks, I.E., Rodríguez-Navarro, A.B., Gambi, M.C., Duarte, C.M., 2015. Extreme pH
844 conditions at a natural CO₂ vent system (Italy) affect growth, and survival of juvenile Pen shells
845 (*Pinna nobilis*). Estuaries and Coasts.

846 Beaumont, N.J., Austen, M.C., Atkins, J.P., Burdon, D., Degraer, S., Dentinho, T.P., Deros, S.,
847 Holm, P., Horton, T., van Ierland, E., Marboe, A.H., Starkey, D.J., Townsend, M., Zarzycki, T.,
848 2007. Identification, definition and quantification of goods and services provided by marine
849 biodiversity: implications for the ecosystem approach. Mar Pollut Bull 54, 253-265.

850 Beck, M.W., Brumbaugh, R.D., Airoidi, L., Carranza, A., Coen, L.D., Crawford, C., Defeo, O., Edgar,
851 G.J., Hancock, B., Kay, M.C., Lenihan, H.S., Luckenbach, M.W., Toropova, C.L., Zhang, G., Guo,
852 X., 2011. Oyster reefs at risk and recommendations for conservation, restoration, and
853 management. BioScience 61, 107-116.

854 Beesley, A., Lowe, D.M., Pascoe, C.K., Widdicombe, S., 2008. Effects of CO₂-induced seawater
855 acidification on the health of *Mytilus edulis*. Climate Res 37, 215-225.

856 Beniash, E., Ivanina, A., Lieb, N.S., Kurochkin, I., Sokolova, I.M., 2010. Elevated level of carbon
857 dioxide affects metabolism and shell formation in oysters *Crassostrea virginica* (Gmelin).
858 Marine Ecology Progress Series 419, 95-108.

859 Beseres-Pollack, J., Yoskowitz, D., Kim, H.-C., Montagna, P.A., 2013. Role and value of nitrogen
860 regulation provided by oysters (*Crassostrea virginica*) in the Mission-Aransas Estuary, Texas,
861 USA. PloS one 8, e65314.

862 Bibby, R., Widdicombe, S., Parry, H., Spicer, J., Pipe, R., 2008. Effects of ocean acidification on
863 the immune response of the blue mussel *Mytilus edulis*. Aquatic Biology 2, 67-74.

864 Bosello, F., Delpiazzo, E., Eboli, F., 2015. Acidification in the Mediterranean sea: impacts and
865 adaptation strategies. Review of Environment, Energy and Economics (Re3), Forthcoming.

866 Brander, L.M., Rehdanz, K., Tol, R.S., Van Beukering, P.J., 2012. The economic impact of ocean
867 acidification on coral reefs. Climate Change Economics 3.

868 Brečević, L., Nielsen, A.E., 1989. Solubility of amorphous calcium carbonate. *J Cryst Growth* 98,
869 504-510.

870 Brodie, J., Williamson, C.J., Smale, D.A., Kamenos, N.A., Mieszkowska, N., Santos, R., Cunliffe, M.,
871 Steinke, M., Yesson, C., Anderson, K.M., Asnaghi, V., Brownlee, C., Burdett, H.L., Burrows, M.T.,
872 Collins, S., Donohue, P.J., Harvey, B., Foggo, A., Noisette, F., Nunes, J., Ragazzola, F., Raven, J.A.,
873 Schmidt, D.N., Suggett, D., Teichberg, M., Hall-Spencer, J.M., 2014. The future of the northeast
874 Atlantic benthic flora in a high CO₂ world. *Ecology and Evolution* 4, 2787-2798.

875 Brumbaugh, R.D., Beck, M.W., Hancock, B., Meadows, A.W., Spalding, M., zu Ermgassen, P.,
876 2010. Changing a management paradigm and rescuing a globally imperiled habitat. *National*
877 *Wetlands Newsletter*, 16-20.

878 Byrne, M., 2011. Impact of ocean warming and ocean acidification on marine invertebrate life
879 history stages: vulnerabilities and potential for persistence in a changing ocean. *Oceanography*
880 *and Marine Biology: An Annual Review* 49, 1-42.

881 Byrne, M., Przeslawski, R., 2013. Multistressor impacts of warming and acidification of the ocean
882 on marine invertebrates' life histories. *Integrative and comparative biology* 53, 582-596.

883 Callaway, R., Shinn, A.P., Grenfell, S.E., Bron, J.E., Burnell, G., Cook, E.J., Crumlish, M., Culloty, S.,
884 Davidson, K., Ellis, R.P., Flynn, K.J., Fox, C., Green, D.M., Hays, G.C., Hughes, A.D., Johnston, E.,
885 Lowe, C.D., Lupatsch, I., Malham, S., Mendzil, A.F., Nickell, T., Pickerell, T., Rowley, A.F., Stanley,
886 M.S., Tocher, D.R., Turnbull, J.F., Webb, G., Wootton, E., Shields, R.J., 2012. Review of climate
887 change impacts on marine aquaculture in the UK and Ireland. *Aquatic Conservation: Marine*
888 *and Freshwater Ecosystems* 22, 389-421.

889 Çath, A., Bozkurt, M., Küçükyılmaz, K., Çınar, M., Bintaş, E., Çöven, F., Atik, H., 2012.
890 Performance and egg quality of aged laying hens fed diets supplemented with meat and bone
891 meal or oyster shell meal. *South African Journal of Animal Science* 42, 74-82.

892 Cerco, C.F., Noel, M.R., 2007. Can oyster restoration reverse cultural eutrophication in
893 Chesapeake Bay? *Estuaries and Coasts* 30, 331-343.

894 Christen, N., Calosi, P., McNeill, C.L., Widdicombe, S., 2013. Structural and functional
895 vulnerability to elevated $p\text{CO}_2$ in marine benthic communities. *Marine Biology* 160, 2113-2128.

896 Cigliano, M., Gambi, M.C., Rodolfo-Metalpa, R., Patti, F.P., Hall-Spencer, J.M., 2010. Effects of
897 ocean acidification on invertebrate settlement at volcanic CO_2 vents. *Marine Biology* 157,
898 2489-2502.

899 Ciuriak, N., 2012. The quiet tsunamis: the ecological, economic, social, and political consequences
900 of ocean acidification. *Paterson Review of International Affairs* 12, 123-143.

901 Clarkson, M., Kasemann, S., Wood, R., Lenton, T., Daines, S., Richoz, S., Ohnemüller, F., Meixner,
902 A., Poulton, S., Tipper, E., 2015. Ocean acidification and the Permo-Triassic mass extinction.
903 *Science* 348, 229-232.

904 Coen, L.D., Brumbaugh, R.D., Bushek, D., Grizzle, R., Luckenbach, M.W., Posey, M.H., Powers,
905 S.P., Tolley, S., 2007. Ecosystem services related to oyster restoration. *Marine Ecology Progress*
906 *Series* 341, 303-307.

907 Coen, L.D., Luckenbach, M.W., Breitburg, D.L., 1999. The role of oyster reefs as essential fish
908 habitat: a review of current knowledge and some new perspectives, *American Fisheries Society*
909 *Symposium*, pp. 438-454.

910 Collins, S., Bell, G., 2004. Phenotypic consequences of 1,000 generations of selection at elevated
911 CO_2 in a green alga. *Nature* 431, 566-569.

912 Comeau, S., Carpenter, R.C., Lantz, C.A., Edmunds, P.J., 2015. Ocean acidification accelerates
913 dissolution of experimental coral reef communities. *Biogeosciences* 12, 365-372.

914 Cooley, S.R., 2012. How human communities could 'feel' changing ocean biogeochemistry.
915 *Current Opinion in Environmental Sustainability* 4, 258-263.

916 Cooley, S.R., Doney, S.C., 2009. Anticipating ocean acidification's economic consequences for
917 commercial fisheries. *Environmental Research Letters* 4, 024007.

918 Cooley, S.R., Kite-Powell, H.L., Doney, S.C., 2009. Ocean acidification's potential to alter global
919 marine ecosystem services. *Oceanography* 22.

920 Cooley, S.R., Lucey, N., Kite-Powell, H., Doney, S.C., 2012. Nutrition and income from molluscs
921 today imply vulnerability to ocean acidification tomorrow. *Fish and Fisheries* 13, 182-215.

922 Cooley, S.R., Ono, C.R., Melcer, S., Roberson, J., 2015. Community-level actions that can address
923 ocean acidification. *Frontiers in Marine Science* 2.

924 De Groot, R.S., Wilson, M.A., Boumans, R.M., 2002. A typology for the classification, description
925 and valuation of ecosystem functions, goods and services. *Ecological Economics* 41, 393-408.

926 DEFRA, 2013. An introductory guide to valuing ecosystem services. Department for Environment,
927 Food and Rural Affairs.

928 Dehon, D.D., 2010. Investigating the use of bioengineered oyster reefs as a method of shoreline
929 protection and carbon storage. MSc Thesis. Louisiana State University.

930 Dennison, W.C., Orth, R.J., Moore, K.A., Stevenson, J.C., Carter, V., Kollar, S., Bergstrom, P.W.,
931 Batiuk, R.A., 1993. Assessing water quality with submersed aquatic vegetation. *BioScience*, 86-
932 94.

933 Díaz, S., Fargione, J., Chapin, F.S., III, Tilman, D., 2006. Biodiversity loss threatens human well-
934 being. *PLoS Biol* 4, e277.

935 Dickinson, G.H., Ivanina, A.V., Matoo, O.B., Portner, H.O., Lannig, G., Bock, C., Beniash, E.,
936 Sokolova, I.M., 2012. Interactive effects of salinity and elevated CO₂ levels on juvenile eastern
937 oysters, *Crassostrea virginica*. *The Journal of experimental biology* 215, 29-43.

938 Dorfmeier, E.M., 2012. Ocean acidification and disease: How will a changing climate impact
939 *Vibrio tubiashii* growth and pathogenicity to Pacific oyster larvae?, School of Aquatic and
940 Fishery Sciences. University of Washington.

941 Dove, M.C., Sammut, J., 2007. Impacts of estuarine acidification on survival and growth of
942 Sydney rock oysters *Saccostrea glomerata* (Gould 1850). *Journal of Shellfish Research* 26, 519-
943 527.

944 Dupont, S., Hall, E., Calosi, P., Lundve, B., 2014. First evidence of altered sensory quality in a
945 shellfish exposed to decreased pH relevant to ocean acidification. *Journal of Shellfish Research*
946 33, 857-861.

947 Dupont, S., Thorndyke, M., 2012. Relationship between CO₂-driven changes in extracellular
948 acid–base balance and cellular immune response in two polar echinoderm species. *Journal of*
949 *Experimental Marine Biology and Ecology* 424–425, 32-37.

950 Dyer, C.L., Leard, R.L., 1994. Folk management in the oyster fishery of the US Gulf of Mexico.
951 *Folk Management in the World's Fisheries*, 55-89.

952 Ekstrom, J.A., Suatoni, L., Cooley, S.R., Pendleton, L.H., Waldbusser, G.G., Cinner, J.E., Ritter, J.,
953 Langdon, C., van Hooidek, R., Gledhill, D., Wellman, K., Beck, M.W., Brander, L.M., Rittschof,
954 D., Doherty, C., Edwards, P.E.T., Portela, R., 2015. Vulnerability and adaptation of US
955 shellfisheries to ocean acidification. *Nature Clim. Change* 5, 207-214.

956 Ellis, R.P., 2013. PhD Thesis: The impact of ocean acidification, increased seawater temperature
957 and a bacterial challenge on the immune response and physiology of the blue mussel, *Mytilus*
958 *edulis*. Plymouth University.

959 Elston, R.A., Hasegawa, H., Humphrey, K.L., Polyak, I.K., Hase, C.C., 2008. Re-emergence of *Vibrio*
960 *tubiaschii* in bivalve shellfish aquaculture: severity, environmental drivers, geographic extent
961 and management. *Dis Aquat Organ* 82, 119-134.

962 Fabricius, K.E., De'ath, G., Noonan, S., Uthicke, S., 2014. Ecological effects of ocean acidification
963 and habitat complexity on reef-associated macroinvertebrate communities. *P Roy Soc B-Biol*
964 *Sci* 281, 20132479.

965 FAO, 2014 *The State of World Fisheries and Aquaculture 2014*. Rome. 223 pp.

966 Feely, R.A., Sabine, C.L., Hernandez-Ayon, J.M., Janson, D., Hales, B., 2008. Evidence for
967 upwelling of corrosive "acidified" water onto the continental shelf. *Science* 320, 1490-1492.

968 Firth, L.B., Thompson, R.C., Bohn, K., Abbiati, M., Airoidi, L., Bouma, T.J., Bozzeda, F., Ceccherelli,
969 V.U., Colangelo, M.A., Evans, A., Ferrario, F., Hanley, M.E., Hinz, H., Hoggart, S.P.G., Jackson,

970 J.E., Moore, P., Morgan, E.H., Perkol-Finkel, S., Skov, M.W., Strain, E.M., van Belzen, J., Hawkins,
971 S.J., 2014. Between a rock and a hard place: Environmental and engineering considerations
972 when designing coastal defence structures. *Coastal Engineering* 87, 122-135.

973 Fletcher, S., Rees, S., Gall, S., Shellock, R., Dodds, W., Rodwell, L., 2014. Assessing the socio-
974 economic benefits of marine protected areas. A report for Natural Resources Wales by the
975 Centre for Marine and Coastal Policy Research, Plymouth University.

976 Fletcher, S., Saunders, J., Herbert, R., Roberts, C., Dawson, K., 2012. Description of the
977 ecosystem services provided by broad-scale habitats and features of conservation importance
978 that are likely to be protected by Marine Protected Areas in the Marine Conservation Zone
979 Project area., Natural England Commissioned Reports 088.

980 Friedrich, L.A., Dodds, W., Philippe, M., Glegg, G., Fletcher, S., Bailly, D., 2015. Improving
981 stakeholder engagement in marine management through ecosystem service assessment. A
982 guide for practitioners based on experience from the VALMER project.

983 Frommel, A.Y., Maneja, R., Lowe, D., Malzahn, A.M., Geffen, A.J., Folkvord, A., Piatkowski, U.,
984 Reusch, T.B.H., Clemmesen, C., 2012. Severe tissue damage in Atlantic cod larvae under
985 increasing ocean acidification. *Nature Climate Change* 2, 42-46.

986 Fulton, E.A., 2011. Interesting times: winners, losers, and system shifts under climate change
987 around Australia. *ICES Journal of Marine Science* 68, 1329-1342.

988 Gattuso, J.P., Hoegh-Guldberg, O., Pörtner, H.O., 2014. Cross-chapter box on ocean acidification,
989 in: Field, C.B., Barros, V.R., Dokken, D.J., Mach, K.J., Mastrandrea, M.D., Bilir, T.E., Chatterjee,
990 M., Ebi, K.L., Estrada, Y.O., Genova, R.C., Girma, B., Kissel, E.S., Levy, A.N., MacCracken, S.,
991 Mastrandrea, P.R., White, L.L. (Eds.), *Climate Change 2014: Impacts, Adaptation, and*
992 *Vulnerability. Part A: Global and Sectoral Aspects. Contribution of Working Group II to the Fifth*
993 *Assessment Report of the Intergovernmental Panel of Climate Change. Cambridge University*
994 *Press, Cambridge, United Kingdom and New York, NY, USA, pp. 97-100.*

995 Gattuso, J.P., Magnan, A., Bille, R., Cheung, W.W., Howes, E.L., Joos, F., Allemand, D., Bopp, L.,
996 Cooley, S.R., Eakin, C.M., Hoegh-Guldberg, O., Kelly, R.P., Portner, H.O., Rogers, A.D., Baxter,
997 J.M., Laffoley, D., Osborn, D., Rankovic, A., Rochette, J., Sumaila, U.R., Treyer, S., Turley, C.,
998 2015. Contrasting futures for ocean and society from different anthropogenic CO₂ emissions
999 scenarios. *Science* 349, aac4722.

1000 Gaylord, B., Hill, T.M., Sanford, E., Lenz, E.A., Jacobs, L.A., Sato, K.N., Russell, A.D., Hettinger, A.,
1001 2011. Functional impacts of ocean acidification in an ecologically critical foundation species.
1002 *The Journal of Experimental Biology* 214, 2586-2594.

1003 Gazeau, F., Alliouane, S., Correa, M.L., Gentile, M., Bock, C., Hirse, T., Bramanti, L., Pörtner, H.-O.,
1004 Ziveri, P., 2014. Impact of ocean acidification and warming on the Mediterranean mussel
1005 (*Mytilus galloprovincialis*). *Frontiers in Marine Science* 1.

1006 Gazeau, F., Gattuso, J.P., Dawber, C., Pronker, A.E., Peene, F., Peene, J., Heip, C.H.R., Middelburg,
1007 J.J., 2010. Effect of ocean acidification on the early life stages of the blue mussel *Mytilus edulis*.
1008 *Biogeosciences* 7, 2051-2060.

1009 Gazeau, F., Gattuso, J.P., Greaves, M., Elderfield, H., Peene, J., Heip, C.H., Middelburg, J.J., 2011.
1010 Effect of carbonate chemistry alteration on the early embryonic development of the Pacific
1011 oyster (*Crassostrea gigas*). *PloS one* 6, e23010.

1012 Gazeau, F., Parker, L.M., Comeau, S., Gattuso, J.-P., O'Connor, W.A., Martin, S., Pörtner, H.-O.,
1013 Ross, P.M., 2013. Impacts of ocean acidification on marine shelled molluscs. *Marine Biology*
1014 160, 2207-2245.

1015 Gazeau, F., Quiblier, C., Jansen, J.M., Gattuso, J.-P., Middelburg, J.J., Heip, C.H.R., 2007. Impact
1016 of elevated CO₂ on shellfish calcification. *Geophysical Research Letters* 34.

1017 Geraldi, N.R., Powers, S.P., Heck, K.L., Cebrian, J., 2009. Can habitat restoration be redundant?
1018 Response of mobile fishes and crustaceans to oyster reef restoration in marsh tidal creeks.
1019 *Marine Ecology Progress Series* 389, 171-180.

1020 Gobler, C.J., Talmage, S.C., 2014. Physiological response and resilience of early life-stage Eastern
1021 oysters (*Crassostrea virginica*) to past, present and future ocean acidification. Conservation
1022 Physiology 2.

1023 Gosselin, L.A., Qian, P.-Y., 1997. Juvenile mortality in benthic marine invertebrates. Marine
1024 Ecology Progress Series 146, 265-282.

1025 Grabowski, J.H., Brumbaugh, R.D., Conrad, R.F., Keeler, A.G., Opaluch, J.J., Peterson, C.H., Piehler,
1026 M.F., Powers, S.P., Smyth, A.R., 2012. Economic valuation of ecosystem services provided by
1027 oyster reefs. BioScience 62, 900-909.

1028 Grabowski, J.H., Peterson, C.H., 2007. Restoring oyster reefs to recover ecosystem services, in:
1029 Kim Cuddington, J.E.B.W.G.W., Alan, H. (Eds.), Theoretical Ecology Series. Academic Press, pp.
1030 281-298.

1031 Green, M.A., Jones, M.E., Boudreau, C.L., Moore, R.L., Westman, B.A., 2004. Dissolution
1032 mortality of juvenile bivalves in coastal marine deposits. Limnology and Oceanography 49, 727-
1033 734.

1034 Grizzle, R.E., Greene, J.K., Coen, L.D., 2008. Seston removal by natural and constructed intertidal
1035 Eastern oyster (*Crassostrea virginica*) reefs: a comparison with previous laboratory studies,
1036 and the value of in situ methods. Estuaries and Coasts 31, 1208-1220.

1037 Guo, X., Huang, M., Pu, F., You, W., Ke, C., 2015. Effects of ocean acidification caused by rising
1038 CO₂ on the early development of three mollusks. Aquatic Biology 23, 147-157.

1039 Haigh, R., Ianson, D., Holt, C.A., Neate, H.E., Edwards, A.M., 2015. Effects of ocean acidification
1040 on temperate coastal marine ecosystems and fisheries in the northeast Pacific. PloS one 10,
1041 e0117533.

1042 Hall-Spencer, J., Allen, R., 2015. The impact of CO₂ emissions on 'nuisance' marine species.
1043 Research and Reports in Biodiversity Studies, 33.

1044 Hall, S.G., Risinger, J.D., Farlow, J., 2011. Ecological engineering of artificial oyster reefs to
1045 enhance carbon sequestration via the algae-oyster complex. Conference proceedings.
1046 American Society of Agricultural and Biological Engineers, p. 1.

1047 Harvey, B.P., Gwynn-Jones, D., Moore, P.J., 2013. Meta-analysis reveals complex marine
1048 biological responses to the interactive effects of ocean acidification and warming. *Ecology and*
1049 *Evolution* 3, 1016-1030.

1050 Henderson, J., O'Neil, J., 2003. Economic values associate with construction of oyster reefs.
1051 Corps of Engineers, EMRRP Technical Notes Collection (ERDC TN-EMRRP-ER-01), US Army
1052 Engineer Research and Development Center, Vicksburg, MS.

1053 Heral, M., 1989. Traditional oyster culture in France, Barnabe. *Aquaculture*, pp. 342-387.

1054 Herbert, R.J.H., Robert, C., Humphreys, J., Fletcher, S., 2012. The Pacific Oyster (*Crassostrea*
1055 *gigas*) in the UK: economic, legal and environmental issues associated with its cultivation, wild
1056 establishment and exploitation. Report for the Shellfish Association of Great Britain.

1057 Hettinger, A., Sanford, E., Hill, T., Hosfelt, J., Russell, A., Gaylord, B., 2013a. The influence of food
1058 supply on the response of *Olympia* oyster larvae to ocean acidification. *Biogeosciences* 10,
1059 6629-6638.

1060 Hettinger, A., Sanford, E., Hill, T.M., Lenz, E.A., Russell, A.D., Gaylord, B., 2013b. Larval carry-over
1061 effects from ocean acidification persist in the natural environment. *Global Change Biology* 19,
1062 3317-3326.

1063 Hettinger, A., Sanford, E., Hill, T.M., Russell, A.D., Sato, K.N.S., Hoey, J., Forsch, M., Page, H.N.,
1064 Gaylord, B., 2012. Persistent carry-over effects of planktonic exposure to ocean acidification in
1065 the *Olympia* oyster. *Ecology* 93, 2758-2768.

1066 Hickey, 2008. Carbon sequestration potential of shellfish. In *Seminars in Sustainability*.
1067 University of South Australia.

1068 Hilmi, N., Allemand, D., Cinar, M., Cooley, S., Hall-Spencer, J., Haraldsson, G., Hattam, C., Jeffree,
1069 R., Orr, J., Rehdanz, K., Reynaud, S., Safa, A., Dupont, S., 2014. Exposure of Mediterranean
1070 countries to ocean acidification. *Water* 6, 1719-1744.

1071 Hilmi, N., Allemand, D., Dupont, S., Safa, A., Haraldsson, G., Nunes, P.A., Moore, C., Hattam, C.,
1072 Reynaud, S., Hall-Spencer, J.M., Fine, M., Turley, C., Jeffree, R., Orr, J., Munday, P.L., Cooley,
1073 S.R., 2013. Towards improved socio-economic assessments of ocean acidification's impacts.
1074 *Marine Biology* 160, 1773-1787.

1075 Hoegh-Guldberg, O., Mumby, P.J., Hooten, A.J., Steneck, R.S., Greenfield, P., Gomez, E., Harvell,
1076 C.D., Sale, P.F., Edwards, A.J., Caldeira, K., Knowlton, N., Eakin, C.M., Iglesias-Prieto, R.,
1077 Muthiga, N., Bradbury, R.H., Dubi, A., Hatzioiols, M.E., 2007. Coral reefs under rapid climate
1078 change and ocean acidification. *Science* 318, 1737-1742.

1079 Hoellein, T.J., Zarnoch, C.B., Grizzle, R.E., 2015. Eastern oyster (*Crassostrea virginica*) filtration,
1080 biodeposition, and sediment nitrogen cycling at two oyster reefs with contrasting water
1081 quality in Great Bay Estuary (New Hampshire, USA). *Biogeochemistry* 122, 113-129.

1082 Houlbrèque, F., Reynaud, S., Godinot, C., Oberhänsli, F., Rodolfo-Metalpa, R., Ferrier-Pagès, C.,
1083 2015. Ocean acidification reduces feeding rates in the scleractinian coral *Stylophora pistillata*.
1084 *Limnology and Oceanography* 60, 89-99.

1085 Hsu, T.C., 2009. Experimental assessment of adsorption of Cu^{2+} and Ni^{2+} from aqueous solution
1086 by oyster shell powder. *Journal of Hazardous Materials* 171, 995-1000.

1087 Humphreys, J., Herbert, R.J.H., Roberts, C., Fletcher, S., 2014. A reappraisal of the history and
1088 economics of the Pacific oyster in Britain. *Aquaculture* 428-429, 117-124.

1089 Ivanina, A.V., Hawkins, C., Sokolova, I.M., 2014. Immunomodulation by the interactive effects of
1090 cadmium and hypercapnia in marine bivalves *Crassostrea virginica* and *Mercenaria mercenaria*.
1091 *Fish and Shellfish Immunology* 37, 299-312.

1092 Jackson, J.B., Kirby, M.X., Berger, W.H., Bjorndal, K.A., Botsford, L.W., Bourque, B.J., Bradbury,
1093 R.H., Cooke, R., Erlandson, J., Estes, J.A., 2001. Historical overfishing and the recent collapse of
1094 coastal ecosystems. *Science* 293, 629-637.

1095 Jones, C., Lawton, J., Shachak, M., 1996. Organisms as Ecosystem Engineers, *Ecosystem*
1096 *Management*. Springer New York, pp. 130-147.

1097 Jones, C.G., Lawton, J.H., Shachak, M., 1997. Positive and negative effects of organisms as
1098 physical ecosystem engineers. *Ecology* 78, 1946-1957.

1099 Kahn, J.R., Kemp, W.M., 1985. Economic losses associated with the degradation of an
1100 ecosystem: The case of submerged aquatic vegetation in Chesapeake Bay. *Journal of*
1101 *Environmental Economics and Management* 12, 246-263.

1102 Kasperski, S., Wieland, R., 2009. When is it optimal to delay harvesting? The role of ecological
1103 services in the Northern Chesapeake Bay oyster fishery. *Mar Resour Econ* 24, 361-385.

1104 Kellogg, M.L., Cornwell, J.C., Owens, M.S., Paynter, K.T., 2013. Denitrification and nutrient
1105 assimilation on a restored oyster reef. *Marine Ecology Progress Series* 480, 1-19.

1106 Kim, J.-H., Yu, O.H., Yang, E.J., Kang, S.-H., Kim, W., Choy, E.J., 2016. Effects of ocean acidification
1107 driven by elevated CO₂ on larval shell growth and abnormal rates of the venerid clam, *Mactra*
1108 *veneriformis*. *Chin. J. Ocean. Limnol.*

1109 Kirby, M.X., 2004. Fishing down the coast: Historical expansion and collapse of oyster fisheries
1110 along continental margins. *Proceedings of the National Academy of Sciences of the United*
1111 *States of America* 101, 13096-13099.

1112 Knights, A.M., Culhane, F., Hussain, S.S., Papadopoulou, K.N., Piet, G.J., Raakær, J., Rogers, S.I.,
1113 Robinson, L.A., 2014. A step-wise process of decision-making under uncertainty when
1114 implementing environmental policy. *Environmental Science & Policy* 39, 56-64.

1115 Knights, A.M., Walters, K., 2010. Recruit-recruit interactions, density-dependent processes and
1116 population persistence in the eastern oyster *Crassostrea virginica*. *Marine Ecology Progress*
1117 *Series* 404, 79-90.

1118 Ko, W.K.G., Chan, V.B., Dineshram, R., Choi, K.S.D., Li, J.A., Ziniu, Y., Thiyagarajan, V., 2013. Larval
1119 and post-larval stages of Pacific oyster (*Crassostrea gigas*) are resistant to elevated CO₂. PLoS
1120 One 8, e64147.

1121 Kraft, N.J.B., Adler, P.B., Godoy, O., James, E.C., Fuller, S., Levine, J.M., Fox, J., 2015. Community
1122 assembly, coexistence and the environmental filtering metaphor. *Funct Ecol* 29, 592-599.

1123 Kroeger, T., 2012. Dollars and Sense: Economic benefits and impacts from two oyster reef
1124 restoration projects in the Northern Gulf of Mexico., The Nature Conservancy.

1125 Kroeker, K.J., Kordas, R.L., Crim, R.N., Singh, G.G., 2010. Meta-analysis reveals negative yet
1126 variable effects of ocean acidification on marine organisms. *Ecology Letters* 13, 1419-1434.

1127 Kroeker, K.J., Micheli, F., Gambi, M.C., 2012. Ocean acidification causes ecosystem shifts via
1128 altered competitive interactions. *Nature Climate Change* 3, 156-159.

1129 Kroeker, K.J., Sanford, E., Jellison, B.M., Gaylord, B., 2014. Predicting the effects of ocean
1130 acidification on predator-prey interactions: a conceptual framework based on coastal molluscs.
1131 *The Biological Bulletin* 226, 211-222.

1132 Kurihara, H., 2008. Effects of CO₂-driven ocean acidification on the early developmental stages
1133 of invertebrates. *Marine Ecology Progress Series* 373, 275-284.

1134 Kurihara, H., Kato, S., Ishimatsu, A., 2007. Effects of increased seawater pCO₂ on early
1135 development of the oyster *Crassostrea gigas*. *Aquatic Biology* 1, 91-98.

1136 Kwon, H.-B., Lee, C.-W., Jun, B.-S., Yun, J.-d., Weon, S.-Y., Koopman, B., 2004. Recycling waste
1137 oyster shells for eutrophication control. *Resources, Conservation and Recycling* 41, 75-82.

1138 La Peyre, M.K., Humphries, A.T., Casas, S.M., La Peyre, J.F., 2014. Temporal variation in
1139 development of ecosystem services from oyster reef restoration. *Ecological Engineering* 63,
1140 34-44.

1141 Lacoue-Labarthe, T., Nunes, P.A.L.D., Ziveri, P., Cinar, M., Gazeau, F., Hall-Spencer, J.M., Hilmi, N.,
1142 Moschella, P., Safa, A., Sauzade, D., Turley, C., 2016. Impacts of ocean acidification in a
1143 warming Mediterranean Sea: An overview. *Regional Studies in Marine Science* 5, 1-11.

1144 Lannig, G., Eilers, S., Portner, H.O., Sokolova, I.M., Bock, C., 2010. Impact of ocean acidification
1145 on energy metabolism of oyster, *Crassostrea gigas*-changes in metabolic pathways and
1146 thermal response. *Marine Drugs* 8, 2318-2339.

1147 LDWF, 2004. Louisiana's oyster shell recovery pilot project. Louisiana Department of Wildlife
1148 and Fisheries. Socioeconomics Research and Development Section and Marine Fisheries
1149 Division.

1150 Lee, C.H., Lee do, K., Ali, M.A., Kim, P.J., 2008. Effects of oyster shell on soil chemical and
1151 biological properties and cabbage productivity as a liming materials. *Waste Management* 28,
1152 2702-2708.

1153 Lee, I., Arcodia, C., 2011. The role of regional food festivals for destination branding.
1154 *International Journal of Tourism Research* 13, 355-367.

1155 Lee, S. W., Park, S.-B., Jeong, S.-K., Lim, K.-S., Lee, S.-H., Trachtenberg, M.C., 2010. On carbon
1156 dioxide storage based on biomineralization strategies. *Micron* 41, 273-282.

1157 Lee, S.W., Hong, S.M., Choi, C.S., 2006. Characteristics of calcification processes in embryos and
1158 larvae of the Pacific oyster, *Crassostrea gigas*. *Bulletin of Marine Science* 78, 309-317.

1159 Li, S., Liu, Y., Liu, C., Huang, J., Zheng, G., Xie, L., Zhang, R., 2015. Morphology and classification
1160 of hemocytes in *Pinctada fucata* and their responses to ocean acidification and warming. *Fish*
1161 *and Shellfish Immunology*.

1162 Li, Y., Qin, J.G., Li, X., Benkendorff, K., 2009. Spawning-dependent stress response to food
1163 deprivation in Pacific oyster *Crassostrea gigas*. *Aquaculture* 286, 309-317.

1164 Lipton, D., 2004. The value of improved water quality to Chesapeake Bay boaters. *Mar Resour*
1165 *Econ* 19, 265-270.

1166 Mac Con Iomaire, M., 2006. A history of seafood in Irish cuisine and culture, *Wild Food:*
1167 *Proceedings of the Oxford Symposium on Food and Cookery, 2004. Oxford Symposium.*

1168 Matozzo, V., Chinellato, A., Munari, M., Finos, L., Bressan, M., Marin, M.G., 2012. First evidence
1169 of immunomodulation in bivalves under seawater acidification and increased temperature.
1170 PloS One 7, e33820.

1171 MEA, 2005. Millenium Ecosystem Assessment. Ecosystems and human well-being: wetlands and
1172 water. World Resources Institute, Washington, DC.

1173 Meyer, D.L., Townsend, E.C., Thayer, G.W., 1997. Stabilization and erosion control value of
1174 oyster cultch for intertidal marsh. Restoration Ecology 5, 93-99.

1175 Michener, W.K., Blood, E.R., Bildstein, K.L., Brinson, M.M., Gardner, L.R., 1997. Climate change,
1176 hurricanes and tropical storms, and rising sea level in coastal wetlands. Ecological Applications
1177 7, 770-801.

1178 Milazzo, M., Rodolfo-Metalpa, R., Chan, V.B., Fine, M., Alessi, C., Thiyagarajan, V., Hall-Spencer,
1179 J.M., Chemello, R., 2014. Ocean acidification impairs vermetid reef recruitment. Science
1180 Report 4, 4189.

1181 Moore, C.C., 2011. Welfare impacts of ocean acidification: an integrated assessment model of
1182 the US mollusk fishery. National Center for Environmental Economics Working Paper, 11-06.

1183 Moreira, A., Figueira, E., Soares, A.M.V.M., Freitas, R., 2016. The effects of arsenic and seawater
1184 acidification on antioxidant and biomineralization responses in two closely related *Crassostrea*
1185 species. Science of The Total Environment 545–546, 569-581.

1186 Narita, D., Rehdanz, K., 2016. Economic impact of ocean acidification on shellfish production in
1187 Europe. Journal of Environmental Planning and Management, 1-19.

1188 Narita, D., Rehdanz, K., Tol, R.S., 2012. Economic costs of ocean acidification: a look into the
1189 impacts on global shellfish production. Climatic Change 113, 1049-1063.

1190 National Research Council, 2004. Non-native oysters in the Chesapeake Bay. Committee on non-
1191 native oysters in the Chesapeake Bay, Ocean Studies Board, Division on Earth and Life Studies.

1192 Nelson, K.A., Leonard, L.A., Posey, M.H., Alphin, T.D., Mallin, M.A., 2004. Using transplanted
1193 oyster (*Crassostrea virginica*) beds to improve water quality in small tidal creeks: a pilot study.
1194 Journal of Experimental Marine Biology and Ecology 298, 347-368.

1195 Newell, R.I., 2004. Ecosystem influences of natural and cultivated populations of suspension-
1196 feeding bivalve molluscs: a review. Journal of Shellfish Research 23, 51-62.

1197 Newell, R.I.E., Fisher, T.R., Holyoke, R.R., Cornwell, J.C., 2005. Influence of Eastern oysters on
1198 nitrogen and phosphorus regeneration in Chesapeake Bay, USA, in: Dame, R., Olenin, S. (Eds.),
1199 The Comparative Roles of Suspension-Feeders in Ecosystems. Springer Netherlands, pp. 93-120.

1200 Ok, Y., Oh, S.-E., Ahmad, M., Hyun, S., Kim, K.-R., Moon, D., Lee, S., Lim, K., Jeon, W.-T., Yang, J.,
1201 2010. Effects of natural and calcined oyster shells on Cd and Pb immobilization in
1202 contaminated soils. Environ Earth Sci 61, 1301-1308.

1203 Paerl, H.W., Pinckney, J.L., Fear, J.M., Peierls, B.L., 1998. Ecosystem responses to internal and
1204 watershed organic matter loading: consequences for hypoxia in the eutrophying Neuse River
1205 Estuary, North Carolina, USA. Marine Ecology Progress Series 166, 17-25.

1206 Pandolfi, J.M., Connolly, S.R., Marshall, D.J., Cohen, A.L., 2011. Projecting coral reef futures
1207 under global warming and ocean acidification. Science 333, 418-422.

1208 Panelli, R., Allen, D., Ellison, B., Kelly, A., John, A., Tipa, G., 2008. Beyond Bluff oysters? Place
1209 identity and ethnicity in a peripheral coastal setting. Journal of Rural Studies 24, 41-55.

1210 Paolisso, M., Dery, N., 2010. A cultural model assessment of oyster restoration alternatives for
1211 the Chesapeake Bay. Human Organization 69, 169-179.

1212 Parker, L.M., O'Connor, W.A., Raftos, D.A., Portner, H.O., Ross, P.M., 2015. Persistence of
1213 positive carryover effects in the oyster, *Saccostrea glomerata*, following transgenerational
1214 exposure to ocean acidification. PloS One 10, e0132276.

1215 Parker, L.M., Ross, P.M., O'Connor, W.A., 2009. The effect of ocean acidification and
1216 temperature on the fertilization and embryonic development of the Sydney rock oyster
1217 *Saccostrea glomerata* (Gould 1850). Global Change Biology 15, 2123-2136.

1218 Parker, L.M., Ross, P.M., O'Connor, W.A., Borysko, L., Raftos, D.A., Pörtner, H.-O., 2012. Adult
1219 exposure influences offspring response to ocean acidification in oysters. *Global Change Biology*
1220 18, 82-92.

1221 Parker, L.M., Ross, P.M., O'Connor, W.A., Portner, H.O., Scanes, E., Wright, J.M., 2013. Predicting
1222 the response of molluscs to the impact of ocean acidification. *Biology 2*, 651-692.

1223 Parker, L.M., Ross, P.M., O'Connor, W.A., 2010. Comparing the effect of elevated $p\text{CO}_2$ and
1224 temperature on the fertilization and early development of two species of oysters. *Marine*
1225 *Biology* 157, 2435-2452.

1226 Parker, L.M., Ross, P.M., O'Connor, W.A., 2011. Populations of the Sydney rock oyster,
1227 *Saccostrea glomerata*, vary in response to ocean acidification. *Marine Biology* 158, 689-697.

1228 Pechenik, J.A., 2006. Larval experience and latent effects-metamorphosis is not a new beginning.
1229 *Integrative and Comparative Biology* 46, 323-333.

1230 Pechenik, J.A., Wendt, D.E., Jarrett, J.N., 1998. Metamorphosis is not a new beginning.
1231 *Bioscience*, 901-910.

1232 Peterson, C.H., Grabowski, J.H., Powers, S.P., 2003. Estimated enhancement of fish production
1233 resulting from restoring oyster reef habitat: quantitative valuation. *Marine Ecology Progress*
1234 *Series* 264, 249-264.

1235 Peterson, C.H., Lipcius, R.N., 2003. Conceptual progress towards predicting quantitative
1236 ecosystem benefits of ecological restorations. *Marine Ecology Progress Series* 264, 297-307.

1237 Piehler, M.F., Smyth, A.R., 2011. Habitat-specific distinctions in estuarine denitrification affect
1238 both ecosystem function and services. *Ecosphere* 2, art12.

1239 Pörtner, H., 2008. Ecosystem effects of ocean acidification in times of ocean warming: a
1240 physiologist's view. *Marine Ecology Progress Series* 373, 203-217.

1241 Pörtner, H., Langenbuch, M., Reipschläger, A., 2004. Biological impact of elevated ocean CO_2
1242 concentrations: lessons from animal physiology and Earth history. *Journal of Oceanography* 60,
1243 705-718.

1244 Przeslawski, R., Byrne, M., Mellin, C., 2014. A review and meta-analysis of the effects of multiple
1245 abiotic stressors on marine embryos and larvae. *Global Change Biology*.

1246 Range, P., Piló, D., Ben-Hamadou, R., Chícharo, M.A., Matias, D., Joaquim, S., Oliveira, A.P.,
1247 Chícharo, L., 2012. Seawater acidification by CO₂ in a coastal lagoon environment: Effects on
1248 life history traits of juvenile mussels *Mytilus galloprovincialis*. *Journal of Experimental Marine*
1249 *Biology and Ecology* 424-425, 89-98.

1250 Richards, G.P., Watson, M.A., Needleman, D.S., Church, K.M., Hase, C.C., 2015a. Mortalities of
1251 Eastern and Pacific oyster larvae caused by the pathogens *Vibrio coralliilyticus* and *Vibrio*
1252 *tubishii*. *Appl Environ Microb* 81, 292-297.

1253 Richards, R.G., Davidson, A.T., Meynecke, J.-O., Beattie, K., Hernaman, V., Lynam, T., van Putten,
1254 I.E., 2015b. Effects and mitigations of ocean acidification on wild and aquaculture scallop and
1255 prawn fisheries in Queensland, Australia. *Fisheries Research* 161, 42-56.

1256 Ries, J.B., Cohen, A.L., McCorkle, D.C., 2009. Marine calcifiers exhibit mixed responses to CO₂-
1257 induced ocean acidification. *Geology* 37, 1131-1134.

1258 Rodolfo-Metalpa, R., Houlbreque, F., Tambutte, E., Boisson, F., Baggini, C., Patti, F.P., Jeffree, R.,
1259 Fine, M., Foggo, A., Gattuso, J.P., Hall-Spencer, J.M., 2011. Coral and mollusc resistance to
1260 ocean acidification adversely affected by warming. *Nature Climate Change* 1, 308-312.

1261 Rodrigues, L.C., van den Bergh, J.C.J.M., Ghermandi, A., 2013. Socio-economic impacts of ocean
1262 acidification in the Mediterranean Sea. *Marine Policy* 38, 447-456.

1263 Rodríguez-Romero, A., Jarrold, M.D., Massamba-N'Siala, G., Spicer, J.I., Calosi, P., 2015. Multi-
1264 generational responses of a marine polychaete to a rapid change in seawater pCO₂.
1265 *Evolutionary Applications*.

1266 Ross, P.M., Parker, L., O'Connor, W.A., Bailey, E.A., 2011. The impact of ocean acidification on
1267 reproduction, early development and settlement of marine organisms. *Water* 3, 1005-1030.

1268 Rossoll, D., Bermudez, R., Hauss, H., Schulz, K.G., Riebesell, U., Sommer, U., Winder, M., 2012.
1269 Ocean acidification-induced food quality deterioration constrains trophic transfer. *PLoS One* 7,
1270 e34737.

1271 Rumrill, S.S., 1990. Natural mortality of marine invertebrate larvae. *Ophelia* 32, 163-198.

1272 Rusher, K., 2003. The Bluff Oyster Festival and regional economic development: festivals as
1273 culture commodified. *Food tourism around the world: Development, management and*
1274 *markets*, 192-205.

1275 Sanford, E., Gaylord, B., Hettinger, A., Lenz, E.A., Meyer, K., Hill, T.M., 2014. Ocean acidification
1276 increases the vulnerability of native oysters to predation by invasive snails. *PLoS Biol* 12,
1277 e100181.

1278 Schmitt, L.H., Brugere, C., 2013. Capturing ecosystem services, stakeholders' preferences and
1279 trade-offs in coastal aquaculture decisions: a Bayesian belief network application. *PLoS One* 8,
1280 e75956.

1281 Schneider, D.W., Stoeckel, J.A., Rehmann, C.R., Douglas Blodgett, K., Sparks, R.E., Padilla, D.K.,
1282 2003. A developmental bottleneck in dispersing larvae: implications for spatial population
1283 dynamics. *Ecology Letters* 6, 352-360.

1284 Scott, M., Hull, S., Mullenhoff, P., 1971. The calcium requirements of laying hens and effects of
1285 dietary oyster shell upon egg shell quality. *Poultry Science* 50, 1055-1063.

1286 Scyphers, S.B., Picou, J.S., Brumbaugh, R.D., Powers, S.P., 2014. Integrating societal perspectives
1287 and values for improved stewardship of a coastal ecosystem engineer. *Ecology and Society* 19.

1288 Scyphers, S.B., Powers, S.P., Heck, K.L., Jr., Byron, D., 2011. Oyster reefs as natural breakwaters
1289 mitigate shoreline loss and facilitate fisheries. *PLoS One* 6, e22396.

1290 Seijo, J.C., Villanueva-Poot, R., Charles, A., 2016. Bioeconomics of ocean acidification effects on
1291 fisheries targeting calcifier species: A decision theory approach. *Fisheries Research* 176, 1-14.

1292 Simkiss, K., Wilbur, K.M., 2012. *Biom mineralization*. Elsevier.

1293 Small, D., Calosi, P., White, D., Spicer, J.I., Widdicombe, S., 2010. Impact of medium-term
1294 exposure to CO₂ enriched seawater on the physiological functions of the velvet swimming crab
1295 *Necora puber*. *Aquatic Biology* 10, 11-21.

1296 Somero, G.N., 2010. The physiology of climate change: how potentials for acclimatization and
1297 genetic adaptation will determine 'winners' and 'losers'. *The Journal of Experimental Biology*
1298 213, 912-920.

1299 Soto, D., Aguilar-Manjarrez, J., Brugère, C., Angel, D., Bailey, C., Black, K., Edwards, P., Costa-
1300 Pierce, B., Chopin, T., Deudero, S., 2008. Applying an ecosystem-based approach to
1301 aquaculture: principles, scales and some management measures, in: Soto D, A.-M.J.,
1302 Hishamunda N (Ed.), *Building an ecosystem approach to aquaculture*. FAO Fisheries and
1303 *Aquaculture Proceedings No. 14*. Rome, Italy: Food and Agriculture Organization, pp. 15–35.

1304 Stenzel, H.B., 1964. Oysters: Composition of the larval shell. *Science* 145, 155-156.

1305 Stocker, T., Qin, D., Plattner, G., Tignor, M., Allen, S., Boschung, J., Nauels, A., Xia, Y., Bex, B.,
1306 Midgley, B., 2013. IPCC, 2013: Climate change 2013: the physical science basis. Contribution of
1307 working group I to the fifth assessment report of the intergovernmental panel on climate
1308 change.

1309 Sui, Y., Kong, H., Huang, X., Dupont, S., Hu, M., Storch, D., Portner, H.O., Lu, W., Wang, Y., 2016.
1310 Combined effects of short-term exposure to elevated CO₂ and decreased O₂ on the physiology
1311 and energy budget of the thick shell mussel *Mytilus coruscus*. *Chemosphere* 155, 207-216.

1312 Sunday, J.M., Calosi, P., Dupont, S., Munday, P.L., Stillman, J.H., Reusch, T.B., 2014. Evolution in
1313 an acidifying ocean. *Trends in Ecology and Evolution* 29, 117-125.

1314 Tahil, A.S., Dy, D.T., 2016. Effects of reduced pH on the early larval development of hatchery-
1315 reared Donkey's ear abalone, *Haliotis asinina* (Linnaeus 1758). *Aquaculture*.

1316 Talmage, S.C., Gobler, C.J., 2009. The effects of elevated carbon dioxide concentrations on the
1317 metamorphosis, size, and survival of larval hard clams (*Mercenaria mercenaria*), bay scallops

1318 (*Argopecten irradians*), and Eastern oysters (*Crassostrea virginica*). Limnology and
1319 Oceanography 54, 2072-2080.

1320 Teagle, H., Hawkins, S.J., Moore, P.J., Smale, D.A., This issue. The role of kelp species as biogenic
1321 habitat formers in coastal marine ecosystems. J Exp Mar Biol Ecol.

1322 Thompson, E.L., O'Connor, W., Parker, L., Ross, P., Raftos, D.A., 2015. Differential proteomic
1323 responses of selectively bred and wild Sydney rock oyster populations exposed to elevated CO₂.
1324 Molecular Ecology 24(6), pp.1248-1262.

1325 Thomsen, J., Casties, I., Pansch, C., Kortzinger, A., Melzner, F., 2012. Food availability outweighs
1326 ocean acidification effects in juvenile *Mytilus edulis*: laboratory and field experiments. Global
1327 change biology 19, 1017-1027.

1328 Thomsen, J., Gutowska, M.A., Saphörster, J., Heinemann, A., Trübenbach, K., Fietzke, J.,
1329 Hiebenthal, C., Eisenhauer, A., Körtzinger, A., Wahl, M., Melzner, F., 2010. Calcifying
1330 invertebrates succeed in a naturally CO₂-rich coastal habitat but are threatened by high levels
1331 of future acidification. Biogeosciences 7, 3879-3891.

1332 Thorson, G., 1950. Reproductive and larval ecology of marine bottom invertebrates. Biological
1333 Reviews 25, 1-45.

1334 Timmins-Schiffman, E., O'Donnell, M.J., Friedman, C.S., Roberts, S.B., 2012. Elevated pCO₂
1335 causes developmental delay in early larval Pacific oysters, *Crassostrea gigas*. Marine Biology
1336 160, 1973-1982.

1337 Tolley, S.G., Volety, A.K., 2005. The role of oysters in habitat use of oyster reefs by resident
1338 fishes and decapod crustaceans. Journal of Shellfish Research 24, 1007-1012.

1339 Tunnicliffe, V., Davies, K.T.A., Butterfield, D.A., Embley, R.W., Rose, J.M., Chadwick Jr, W.W.,
1340 2009. Survival of mussels in extremely acidic waters on a submarine volcano. Nature
1341 Geoscience 2, 344-348.

1342 UNEP, 2010. UNEP Emerging Issues: Environmental Consequences of Ocean Acidification: A
1343 Threat to Food Security. UNEP.

1344 Uthicke, S., Logan, M., Liddy, M., Francis, D., Hardy, N., Lamare, M., 2015. Climate change as an
1345 unexpected co-factor promoting coral eating seastar (*Acanthaster planci*) outbreaks. Scientific
1346 Reports 5, 8402.

1347 Videla, J., Chaparro, O., Thompson, R., Concha, I., 1998. Role of biochemical energy reserves in
1348 the metamorphosis and early juvenile development of the oyster *Ostrea chilensis*. Marine
1349 Biology 132, 635-640.

1350 Visser, M.E., 2008. Keeping up with a warming world; assessing the rate of adaptation to climate
1351 change. P Roy Soc B-Biol Sci 275, 649-659.

1352 Volety, A.K., Haynes, L., Goodman, P., Gorman, P., 2014. Ecological condition and value of oyster
1353 reefs of the Southwest Florida shelf ecosystem. Ecological Indicators 44, 108-119.

1354 Waldbusser, G.G., Steenson, R.A., Green, M.A., 2011a. Oyster shell dissolution rates in estuarine
1355 waters: effects of pH and shell legacy. Journal of Shellfish Research 30, 659-669.

1356 Waldbusser, G.G., Voigt, E.P., Bergschneider, H., Green, M.A., Newell, R.I.E., 2011b.
1357 Biocalcification in the Eastern oyster (*Crassostrea virginica*) in relation to long-term trends in
1358 Chesapeake Bay pH. Estuaries and Coasts 34, 221-231.

1359 Walles, B., Mann, R., Ysebaert, T., Troost, K., Herman, P.M.J., Smaal, A.C., 2015. Demography of
1360 the ecosystem engineer *Crassostrea gigas*, related to vertical reef accretion and reef
1361 persistence. Estuarine, Coastal and Shelf Science 154, 224-233.

1362 Wang, Q., Cao, R., Ning, X., You, L., Mu, C., Wang, C., Wei, L., Cong, M., Wu, H., Zhao, J., 2016.
1363 Effects of ocean acidification on immune responses of the Pacific oyster *Crassostrea gigas*. Fish
1364 & Shellfish Immunology 49, 24-33.

1365 Watson, S.-A., Southgate, P.C., Tyler, P.A., Peck, L.S., 2009. Early larval development of the
1366 Sydney rock oyster *Saccostrea glomerata* under near-future predictions of CO₂-driven ocean
1367 acidification. Journal of Shellfish Research 28, 431-437.

1368 Wedepohl, K.H., Baumann, A., 2000. The use of marine molluskan shells for roman glass and
1369 local raw glass production in the Eifel Area (Western Germany). *Naturwissenschaften* 87, 129-
1370 132.

1371 Weiner, S., Addadi, L., 2002. Calcium carbonate formation in biology: the involvement of an
1372 amorphous calcium carbonate precursor phase. *Geochimica et Cosmochimica Acta* 66, A827.

1373 Weiss, I.M., Tuross, N., Addadi, L., Weiner, S., 2002. Mollusc larval shell formation: amorphous
1374 calcium carbonate is a precursor phase for aragonite. *Journal of Experimental Zoology* 293,
1375 478-491.

1376 Welladsen, H.M., Southgate, P.C., Heimann, K., 2010. The effects of exposure to near-future
1377 levels of ocean acidification on shell characteristics of *Pinctada fucata* (Bivalvia: Pteriidae).
1378 *Molluscan Research* 30, 125.

1379 White, L.J., Koss, R.S., Knights, A.M., Eriksson, A., Robinson, L.A., 2013. ODEMM Linkage
1380 Framework Userguide (Version 2). ODEMM Guidance Document Series No.3. EC FP7 project
1381 (244273) 'Options for Delivering Ecosystem-based Marine Management'. University of
1382 Liverpool. ISBN: 978-0- 906370-87-6: 14 pp.

1383 Wingard, G.L., Lorenz, J.J., 2014. Integrated conceptual ecological model and habitat indices for
1384 the southwest Florida coastal wetlands. *Ecological Indicators* 44, 92-107.

1385 Wright, J.M., Parker, L.M., O'Connor, W.A., Williams, M., Kube, P., Ross, P.M., 2014. Populations
1386 of Pacific oysters *Crassostrea gigas* respond variably to elevated CO₂ and predation by *Morula*
1387 *marginalba*. *The Biological Bulletin* 226, 269-281.

1388 Yang, E.-I., Kim, M.-Y., Park, H.-G., Yi, S.-T., 2010. Effect of partial replacement of sand with dry
1389 oyster shell on the long-term performance of concrete. *Construction and Building Materials* 24,
1390 758-765.

1391 Yang, E.-I., Yi, S.-T., Leem, Y.-M., 2005. Effect of oyster shell substituted for fine aggregate on
1392 concrete characteristics: Part I. Fundamental properties. *Cement Concrete Res* 35, 2175-2182.

1393 Yao, Z., Xia, M., Li, H., Chen, T., Ye, Y., Zheng, H., 2014. Bivalve shell: not an abundant useless
1394 waste but a functional and versatile biomaterial. *Critical Reviews in Environmental Science and*
1395 *Technology* 44, 2502-2530.

1396 Yoon, G.-L., Kim, B.-T., Kim, B.-O., Han, S.-H., 2003. Chemical–mechanical characteristics of
1397 crushed oyster-shell. *Waste Management* 23, 825-834.

1398 Yoon, H., 2004. Oyster shell as substitute for aggregate in mortar. *Waste Management Research*
1399 22, 158-170.

1400 zu Ermgassen, P.S.E., Spalding, M.D., Grizzle, R.E., Brumbaugh, R.D., 2013. Quantifying the loss
1401 of a marine ecosystem service: filtration by the Eastern oyster in US estuaries. *Estuaries and*
1402 *Coasts* 36, 36-43.

1403

1404 Table 1 (web version (colour); print (grayscale)): Ecosystem goods and services provided by oysters and oyster reefs, following the
 1405 ODEMM Linkage Framework description (White et al., 2013), and the potential impacts of ocean acidification (OA). '√' indicates that
 1406 the service is provided by oysters, 'X' indicates the service is not provided (according to the available literature). Direction of arrows
 1407 indicates expected change in the ecosystem service (i.e. ↑ indicates increase in ES provision, ↓ a decrease in ES provision). '~'
 1408 indicates that no consensus is reached or the change is context-specific.

	Marine Ecosystem Goods and Services	Provided by Oysters or Oyster Reefs?	Estimated Value?	Affected by OA?
Habitat	Lifecycle maintenance	√ (assumed from 3-D structure)	Unknown	Yes ↓~
	Gene pool protection	√ (assumed from 3-D structure)	Unknown	Yes ↓~
Provisioning	Sea Food	√ (harvest, aquaculture, extended fisheries)	\$20 850-\$52 224/hectare of reef (oyster harvest value- Grabowski et al. (2012)) \$4123/year/hectare (extended fisheries- Grabowski et al. (2012)) 809.7\$/tonne (aquaculture production (FAO data [†]))	Yes ↓ or ↑ ~
	Sea Water	X	X	X
	Raw materials	√ (clutch material, construction material)	\$7-10 lb ⁻¹ (Baywater Oyster Seeds LLC) \$17 yd ⁻³ (Pontchartrain Materials Corp., New Orleans) \$17 yd ⁻³ (LDWF, 2004) ~\$126/ton (Kwon et al., 2004)	Yes ↓~
	Genetic resources	√ (aquaculture bred lines/triploid)	Unknown	Unknown

	Medicinal resources	X	X	X
	Ornamental resources	√ (<i>shell collection</i>)	~\$13m ⁻³ (Lemasson, unpublished)	Yes ↓~
Regulating	Air purification	X	X	X
	Climate regulation (incl. carbon sequestration)	√ (<i>carbon sequestration</i>)	Unknown	Yes ↓~
	Disturbance prevention or moderation	√	Upward of \$6 million (<i>for coastal defence structures- Firth et al., 2014</i>)	Yes ↓~
	Regulation of water flows	X	X	X
	Waste treatment (esp. water purification)	√ (<i>water purification, Chl-a removal, nutrient cycling</i>)	\$1385-\$6716/yr/ha. (Grabowski et al., 2012) \$314 836/yr (<i>Newell et al., 2005</i>) \$18 135/million oysters (<i>Kasperski and Wieland, 2009</i>) \$3000/ac./yr (<i>Piehler and Smyth, 2011</i>) \$293 993/yr (<i>Beseres-Pollack et al., 2013</i>)	Yes ↓~
	Coastal erosion prevention	√ (<i>shoreline stabilization, erosion prevention</i>)	Upward of \$6 million for coastal defence structures (<i>Firth et al. 2014</i>)	Yes ↓~
	Biological control	√ (<i>facilitate submerged aquatic vegetation</i>)	Unknown	Yes ↓~
Cultural	Aesthetic information	X	X	X
	Recreation & leisure	√	\$222 million (<i>U.S. National Research Council, 2004</i>)	Yes ↓~
	Inspiration for culture, art and design	X	X	X
	Spiritual experience	√ (<i>assumed</i>)	Unknown	Unknown

1409

	Information for cognitive development	√ (<i>assumed</i>)	<i>Unknown</i>	<i>Unknown</i>
	Cultural Heritage and Identity	√ (<i>sense of tradition, oyster festivals</i>)	<i>Unknown</i>	<i>Unknown</i>

1410 Figure 1: Categories of ecosystem goods and services, as described (left) in the
1411 Millennium Ecosystem Assessment (MEA, 2005), and additional examples
1412 (right) from the ODEMM Linkage Framework (White et al., 2013).

1413

1414 Figure 2: Total Economic Value framework (modified from Herbert et al., 2012).


1415

1416 Figure 3: Relationships between ocean acidification effects, various levels of
1417 biological complexity, the provision of ecosystem services, and the associated
1418 social and economic effects (adapted from Le Quesne and Pinnegar, 2012).

1419


1420 Figure 4: Conceptual diagram depicting some of the ecosystem services
1421 provided by oyster reefs (right) and the potential effects of ocean acidification
1422 on their provision (left). SAV=Submerged Aquatic Vegetation (Figure created by
1423 the authors, symbols courtesy of the Integration and Application Network,
1424 University of Maryland Center for Environmental Science
1425 (ian.umces.edu/symbols/)).

1426



1429 Figure 2: (web version)


1430



1431


1432

1433 Figure 2: (print version)




1434

1435 Figure 3: (web version)




1436

1437 Figure 3: (print version)




1438

Present day



Future ocean acidification scenario



1440

1441

1442

1443

1444 **Highlights:**

1445 • Ocean acidification (OA) threatens all stages of the life-cycle of oysters.

1446 • Oyster reefs provide numerous ecosystem services (ES)

1447 • The delivery of ESs associated with healthy oyster reefs may be threatened
1448 by OA.

1449 • Further studies linking OA and ES are urgently needed.