

Slice Fourier transform and convolutions

Lander Cnudde, Hendrik De Bie

Abstract

Recently the construction of various integral transforms for slice monogenic functions has gained a lot of attention. In line with these developments, the article at hand introduces the slice Fourier transform. In the first part, the kernel function of this integral transform is constructed using the Mehler formula. An explicit expression for the integral transform is obtained and allows for the study of its properties. In the second part, two kinds of corresponding convolutions are examined: Mustard convolutions and convolutions based on generalised translation operators. The paper finishes by demonstrating the connection between both.

Keywords: Clifford-Hermite functions, slice Fourier transform, Mustard convolution, generalised convolution

MSC: 30G35, 33C45

1 Introduction

In recent years a lot of work has been put into the study of various integral transforms in a slice Clifford analytic setting. In line with earlier research on slice monogenic functions (see e.g. [4,8,9]), recent developments comprise the generalisation of various well-known integral transforms to this particular setting. So far the cases of the Bergman-Sce transform [5], the Cauchy transform [9] and the (dual) Radon transform [7] have been treated. However, an integral transform of major importance has not yet been studied. The paper at hand introduces the classical Fourier transform to the slice monogenic framework, giving rise to the slice Fourier transform.

In various papers (see e.g. [1, 2, 10, 11, 13]) the Fourier transform has been generalised to the general hypercomplex setting. Depending on the parity of the dimension of the underlying real Clifford algebra, these efforts even lead to closed forms for the corresponding kernel functions. The study of these new Fourier transforms has also been carried out from an application point of view, examining the possibility to use them for image processing (see e.g. [14, 15, 18, 20]).

Up to now it has not been possible to generalise the Fourier transform to the slice Clifford analytic setting because a solid basis was missing. Owing to recent work on Clifford-Hermite functions and their properties (see [4]), however, the insight in the corresponding function space has been growing. Hence it is now a natural question to address the construction of a slice Fourier transform based

on these findings. In order to retrieve a realisation of the $\mathfrak{osp}(1|2)$ Lie super-algebra, [4] manipulates the slice Cauchy-Riemann operator of [6] such that a slice Dirac operator is obtained. This in turn allowed for the construction of Clifford-Hermite functions which exhibit interesting differential properties. For this reason they qualify extremely well for an eigenfunction basis of the slice Fourier transform to be obtained.

In a first part of this paper, the slice Fourier transform is constructed and its main properties are studied. The classical Fourier transform being an integral transform, the search for a generalisation to the above mentioned slice monogenic setting comes down to finding an appropriate kernel function. Next, the explicit expression of the integral transform allows for a closer study of its basic properties. It is also proved that the slice Fourier transform is well-defined on a subspace of a Clifford analogue of the Hilbert space L^2 .

In this paper the construction of the kernel is achieved using the Mehler formula. Based on a set of eigenfunctions that are orthogonal with respect to a well-chosen inner product and their corresponding eigenvalues, this formula allows for a formal expression of the kernel function. Because the aforementioned Clifford-Hermite functions show the desired properties, they are declared eigenfunctions of the slice Fourier transform and assigned corresponding eigenvalues based on the scalar differential equation. Using their explicit expressions and several technical manipulations, we ultimately find a closed form for the integral transform.

Though this method merely starts from eigenfunctions and eigenvalues analogous to those of the classical Fourier transform, the resulting slice Fourier transform shows even more similarities. Not only does the final kernel expression comprise of familiar exponential functions, it is also proven that this kernel function obeys a system of Clifford algebra-valued partial differential equations analogous to the classical transform. At this point, a natural question to ask is whether it also yields a nice convolution property.

In the second part of this article, two interpretations of the classical convolution are explored to generalise the classical convolution property to the case of the slice Fourier transform. Based on [3] both a Mustard convolution and a generalised translation operator are defined and proven to show the desired property. Given the non-commutative nature of the Clifford setting, the commutativity of the classical convolution can of course not be retrieved. One thus has to distinguish between left and right Mustard convolutions and generalised translation operators.

In the first approach, the (left) Mustard convolution \star_S is defined such that its slice Fourier transform \mathcal{F}_S equals the product of the slice Fourier transforms of the constituting functions, or, for certain Clifford-valued functions f and g ,

$$\mathcal{F}_S(f \star_S g) = \mathcal{F}_S(f)\mathcal{F}_S(g).$$

In the second approach, the focus is on the translation operator t_y in the definition of the classical convolution

$$(f \star g)(x) = \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} t_y(f)(x)g(y)dy, \quad (1.1)$$

where $t_y(f)(x) = f(x - y)$. The operator t_y is such that its classical Fourier transform generates an extra factor identical to the classical kernel function, thus giving rise to the above convolution property. Based on these observations, a (left) generalised translation operator T_y is defined which behaves analogously under the slice Fourier transform and leads to a slice convolution property as well. Given that both approaches give rise to the same behaviour in the slice Fourier domain, we conclude by pinpointing the connection between them.

The paper is organised as follows. Section 2 summarises, besides some general preliminary results with respect to the slice Clifford context, the main results concerning the Clifford-Hermite functions studied in [4]. In section 3, the explicit form of the slice Fourier transform is obtained using the Mehler formula and several of its properties are studied. Section 4 addresses two ways to define an appropriate convolution with respect to this transform. Apart from a Mustard-type convolution, the problem is also approached using generalised translation operators. Conclusions can be found in section 5.

2 Preliminaries

This preliminary section contains some general background on the slice approach in \mathbb{R}^{m+1} and briefly summarizes the definitions and properties of the Clifford-Hermite functions as defined and proved in [4].

2.1 Slice approach in \mathbb{R}^{m+1}

The $(m + 1)$ -dimensional real Clifford algebra Cl_{m+1} has $m + 1$ basis vectors $e_i, i = 0, \dots, m$, which satisfy the anti-commutation relations

$$e_i e_j + e_j e_i = -2\delta_{ij}, \quad i, j = 0, \dots, m.$$

A k -vector (where $k \leq m + 1$) is an element e_A of Cl_{m+1} such that

$$e_A = e_{i_1} \dots e_{i_k}$$

where $i_j \in \{0, \dots, m\}$ for all $j \in \{1, \dots, k\}$ and with $i_1 < \dots < i_k$. The variable $\mathbf{x} \in Cl_{m+1}$ is defined as the 1-vector which corresponds to the $(m + 1)$ -tuple $(x_0, \dots, x_m) \in \mathbb{R}^{m+1}$ by

$$\mathbf{x} = x_0 e_0 + x_1 e_1 + \dots + x_m e_m.$$

Using spherical coordinates to describe the Cl_m -part \underline{x} of \mathbf{x} , one can also write

$$\begin{aligned} \mathbf{x} &= x_0 e_0 + \underline{x} \\ &= x_0 e_0 + r \underline{\omega}, \end{aligned}$$

where $r = \sqrt{x_1^2 + \dots + x_m^2}$ and $\underline{\omega} = \underline{x}/r$. A general element \mathbf{x} is thus defined by the triplet $(x_0, r, \underline{\omega}) \in \mathbb{R} \times \mathbb{R}^+ \times \mathbb{S}^{m-1}$, where \mathbb{S}^{m-1} denotes the $(m-1)$ -dimensional sphere in \mathbb{R}^m . The variable \mathbf{x} therefore lives in the subspace spanned by the fixed basis vector e_0 and the unit 1-vector $\underline{\omega}$. This subspace is called a slice, referring to the slice concept used in literature (see e.g. [8] and the book [9]). The slice at hand, however, is not directly a subalgebra of the ambient algebra as $1, e_0, \underline{\omega}$ and $e_0\underline{\omega}$ generate, for fixed $\underline{\omega}$, the quaternions \mathbb{H} . It should therefore rather be interpreted as a ‘rotated’ slice.

As a consequence, a general function f of \mathbf{x} will depend on x_0, r and $\underline{\omega}$. Throughout this article such functions f will be written both as $f(\mathbf{x})$ and as $f(x_0, r, \underline{\omega})$ because the former is more compact and the latter shows its dependences explicitly. Based on considerations in [6], the following definition was proposed in [4].

Definition 2.1 (Slice Dirac operator). *The slice Dirac operator D_0 is the partial differential operator defined as*

$$D_0 = e_0 \partial_{x_0} + \underline{\omega} \partial_r.$$

Given that $-e_0 D_0$ corresponds to the Cauchy-Riemann operator in Cl_{m+1} , null-solutions of D_0 correspond to slice monogenic functions as studied in e.g. [8, 16, 17]. Together with the multiplication operator \mathbf{x} and the slice Dirac operator D_0 , the Euler operator $\mathbb{E} = \sum_{i=0}^m x_i \partial_{x_i}$ establishes a realisation of the $\mathfrak{osp}(1|2)$ -superalgebra, see [12].

Theorem 2.2. *The operators \mathbf{x} , D_0 and \mathbb{E} constitute a Lie superalgebra, isomorphic to $\mathfrak{osp}(1|2)$, with relations*

$$\begin{array}{ll} \text{(i)} & \{\mathbf{x}, \mathbf{x}\} = -2|\mathbf{x}|^2 \\ \text{(iii)} & \{\mathbf{x}, D_0\} = -2(\mathbb{E} + 1) \\ \text{(v)} & [|\mathbf{x}|^2, D_0] = -2\mathbf{x} \\ \text{(vii)} & [\partial_{x_0}^2 + \partial_r^2, \mathbf{x}] = 2D_0 \\ \text{(ix)} & [\partial_{x_0}^2 + \partial_r^2, |\mathbf{x}|^2] = 4(\mathbb{E} + 1) \end{array} \quad \begin{array}{ll} \text{(ii)} & \{D_0, D_0\} = -2(\partial_{x_0}^2 + \partial_r^2) \\ \text{(iv)} & [\mathbb{E} + 1, D_0] = -D_0 \\ \text{(vi)} & [\mathbb{E} + 1, \mathbf{x}] = \mathbf{x} \\ \text{(viii)} & [\mathbb{E} + 1, \partial_{x_0}^2 + \partial_r^2] = -2(\partial_{x_0}^2 + \partial_r^2) \\ \text{(x)} & [\mathbb{E} + 1, |\mathbf{x}|^2] = 2|\mathbf{x}|^2. \end{array}$$

Definition 2.3. *The Clifford conjugation $\bar{\cdot}$ is defined as*

$$\begin{array}{ll} \bar{\lambda} = \lambda^* & \lambda \in \mathbb{C} \\ \bar{e}_i = -e_i & i = 0, \dots, m. \\ \overline{ab} = \bar{b} \bar{a} & a, b \in Cl_{m+1}. \end{array}$$

where $*$ denotes the standard complex conjugation.

Definition 2.4. *The vector space \mathcal{L}^2 is defined as*

$$\begin{aligned} \mathcal{L}^2 &= L^2(\mathbb{R}^{m+1}, r^{1-m} d\mathbf{x}) \otimes Cl_{m+1} \\ &= \left\{ f : \mathbb{R}^{m+1} \rightarrow Cl_{m+1} \mid \left[\int_{\mathbb{R}^{m+1}} \overline{f(\mathbf{x})} f(\mathbf{x}) r^{1-m} d\mathbf{x} \right]_0 < +\infty \right\} \end{aligned}$$

where $[\cdot]_0$ denotes the scalar part of the expression between the brackets.

On \mathcal{L}^2 an inner product was defined to be self-adjoint with respect to D_0 .

Definition 2.5. *The vector space \mathcal{L}^2 is given the structure of a right Hilbert module by defining the inner product of two functions $f, g : \mathbb{R}^{m+1} \rightarrow Cl_{m+1}$ as*

$$\langle f, g \rangle = \int_{\mathbb{R}^{m+1}} \overline{f(\mathbf{x})} g(\mathbf{x}) r^{1-m} d\mathbf{x} = \int_{\mathbb{R}^{m+1}} \overline{f(D\mathbf{x})} g(\mathbf{x}) dx_0 d\mathbf{r} d\sigma_{\underline{x}}$$

where $d\sigma_{\underline{x}}$ denotes the measure on the unit sphere \mathbb{S}^{m-1} corresponding to the \underline{x} -part of \mathbf{x} .

The inner product has some interesting properties.

Proposition 2.6. *The inner product $\langle f, g \rangle = \int_{\mathbb{R}^{m+1}} \overline{f} g dx_0 d\mathbf{r} d\omega$ on the right Cl_{m+1} -module \mathcal{L}^2 obeys the relations*

$$\begin{aligned} \langle D_0 f, g \rangle &= \langle f, D_0 g \rangle, \\ \langle \mathbf{x} f, g \rangle &= -\langle f, \mathbf{x} g \rangle \end{aligned}$$

on a dense subset of \mathcal{L}^2 .

2.2 The Clifford-Hermite functions

Based on the classical definitions, the Clifford-Hermite polynomials and functions are defined using the kernel of the differential operator D_0 and the $\mathfrak{osp}(1|2)$ -relations.

In [4] it was shown that the polynomial kernel of D_0 is a right Cl_{m+1} -module which is spanned by the homogeneous polynomials $m_k(\mathbf{x}) = (e_0 - 1)(x_0 + \underline{x})^k$ of degree $k \in \mathbb{N}$.

Definition 2.7 (Clifford-Hermite polynomials). *The Clifford-Hermite polynomials $h_{j,k}$ of degree j and order k are defined as*

$$h_{j,k}(\mathbf{x}) m_k(\mathbf{x}) = (\mathbf{x} - cD_0)^j m_k(\mathbf{x})$$

where $c \in \mathbb{R}_0^+$ and $j \in \mathbb{N}$.

The parameter c adds some freedom to the definition. In order not to overload notation, however, its presence will not be denoted explicitly. We briefly summarise the most important properties of these polynomials (see [4]).

Theorem 2.8. *The polynomials $H_j(m_k)(\mathbf{x}) = h_{j,k}(\mathbf{x}) m_k(\mathbf{x})$ are solutions of the differential equation*

$$cD_0^2 H_j(m_k)(\mathbf{x}) - \mathbf{x} D_0 H_j(m_k)(\mathbf{x}) + C(j, k) H_j(m_k)(\mathbf{x}) = 0$$

with $C(j, k) = -2t$ if $j = 2t$ and $C(j, k) = -2(k + t + 1)$ if $j = 2t + 1$.

Theorem 2.9. *The Hermite polynomials $h_{j,k}$ can be expressed as*

$$\begin{aligned} h_{2t,k}(\mathbf{x}) &= (2c)^t t! L_t^k \left(\frac{|\mathbf{x}|^2}{2c} \right) \\ h_{2t+1,k}(\mathbf{x}) &= (2c)^t t! \mathbf{x} L_t^{k+1} \left(\frac{|\mathbf{x}|^2}{2c} \right) \end{aligned}$$

where L_t^k are the generalised Laguerre polynomials of degree t and order k on the real line.

Based on the Clifford-Hermite polynomials $h_{j,k}$, Clifford-Hermite functions $\psi_{j,k}$ are defined and their main properties are summarised as well.

Definition 2.10 (Clifford-Hermite functions). *The Clifford-Hermite functions $\psi_{j,k}$ are defined as*

$$\psi_{j,k}(\mathbf{x}) = h_{j,k}(\mathbf{x}) m_k(\mathbf{x}) \exp(-|\mathbf{x}|^2/4c)$$

where $c \in \mathbb{R}_0^+$ is the same parameter as in Definition 2.7.

Proposition 2.11. *The Clifford-Hermite functions $\psi_{j,k}$ satisfy the relations*

$$\begin{cases} \psi_{j,k} &= \tilde{D}_c \psi_{j-1,k} \\ \tilde{D}_c^\dagger \psi_{j,k} &= -c C(j,k) \psi_{j-1,k} \end{cases} \quad (2.2)$$

with $\tilde{D}_c = \frac{\mathbf{x}}{2} - cD_0$, $C(j,k)$ as in Theorem 2.8 and where \cdot^\dagger denotes the adjoint with respect to the inner product.

Theorem 2.12. *Let $\psi_{j_i, k_i} = [(\mathbf{x} - cD_0)^{j_i} m_{k_i}(\mathbf{x})] \exp(-|\mathbf{x}|^2/4c)$ with $m_{k_i}(\mathbf{x}) = (e_0 - 1)(x_0 + \underline{x})^{k_i}$ for $i = 1, 2$. The inner product of these two Clifford-Hermite functions ψ_{j_1, k_1} and ψ_{j_2, k_2} is given by*

$$\langle \psi_{j_1, k_1}, \psi_{j_2, k_2} \rangle = A(j_1, k_1) \delta_{j_1 j_2} \delta_{k_1 k_2}$$

with

$$A(j_1, k_1) = \begin{cases} 2(2c)^{2t_1+k_1+1} t_1! (k_1 + t_1)! \frac{\pi^{m/2+1}}{\Gamma(m/2)} & j_1 = 2t_1, \\ 2(2c)^{2t_1+k_1+2} t_1! (k_1 + t_1 + 1)! \frac{\pi^{m/2+1}}{\Gamma(m/2)} & j_1 = 2t_1 + 1. \end{cases}$$

Theorem 2.13. *The Clifford-Hermite functions $\psi_{j,k}$ are solutions of the scalar differential equation*

$$\left(cD_0^2 + \frac{|\mathbf{x}|^2}{4c} \right) \psi_{j,k}(\mathbf{x}) = (j + k + 1) \psi_{j,k}(\mathbf{x}). \quad (2.3)$$

Definition 2.14. *The set of right finite linear combinations of Clifford-Hermite functions over Cl_{m+1} will be denoted as \mathcal{V} , so $\mathcal{V} = \text{span}_{Cl} \{\psi_{j,k}\}$ and $\mathcal{V} \subset \mathcal{L}^2$.*

Remark 2.15. *As a consequence any function $f \in \mathcal{V}$ can be written as $f(\mathbf{x}) = f_1(x_0, r) + \varpi f_2(x_0, r)$, where f_1 and f_2 are Cl_{m+1} -valued functions defined on $\mathbb{R} \times \mathbb{R}^+$. Moreover these functions f_1 and f_2 obey the relations $f_1(x_0, -r) = f_1(x_0, r)$ and $f_2(x_0, -r) = -f_2(x_0, r)$ so all functions $f \in \mathcal{V}$ are slice functions.*

3 Slice Fourier transform

Based on Theorem 2.13, a formal definition of the slice Fourier transform (see [19]) on \mathcal{V} is obtained by taking the exponential of the scalar differential equation (2.3):

$$e^{-i\frac{\pi}{2}\left(cD_0^2 + \frac{|\mathbf{x}|^2}{4c}\right)}\psi_{j,k}(\mathbf{x}) = (-i)^{(j+k+1)}\psi_{j,k}(\mathbf{x}) \quad (3.4)$$

where the choice for the sign of $-i$ is arbitrary. The aim of this section is however to construct an explicit integral expression corresponding to the above formal exponential operator. To this end functions f will be restricted to $f \in \mathcal{V}$. Given its eigenvalues and corresponding eigenfunctions, an appropriate kernel function \mathcal{K}^M will thus be constructed such that

$$\mathcal{F}_S(f)(\mathbf{y}) = \int_{\mathbb{R}^{m+1}} \mathcal{K}^M(\mathbf{x}, \mathbf{y}) f(\mathbf{x}) r^{1-m} d\mathbf{x} \quad (3.5)$$

where the integration is performed over the first argument of \mathcal{K}^M . Though the slice Fourier transform is defined using the eigenfunction approach, we will demonstrate that it generalises other properties of the classical Fourier transform as well.

3.1 The Mehler construction

For the functions $\psi_{j,k}$ to be eigenfunctions of the slice Fourier transform, the action of the transform on these functions should be to multiply them with proportionality factors given by their corresponding eigenvalues $(-i)^{j+k+1}$ (see expression (3.4)). Together with the orthogonality of the Clifford-Hermite functions proved in Theorem 2.12, this yields the following formal series expansion for the kernel function \mathcal{K}^M :

$$\mathcal{K}^M(\mathbf{x}, \mathbf{y}) = \sum_{j,k=0}^{+\infty} \frac{\psi_{j,k}(\mathbf{y})(-i)^{j+k+1}\overline{\psi_{j,k}(\mathbf{x})}}{\langle \psi_{j,k}, \psi_{j,k} \rangle}, \quad (3.6)$$

where the denominator accounts for the normalisation of the eigenfunctions. Indeed, after substituting this expression in equation (3.5) and changing the order of integration and summation, what appears is the sum of the projections of f onto all $\psi_{j,k}$'s times the slice Fourier transforms of these $\psi_{j,k}$'s. By definition this exactly yields the image of f under the slice Fourier transform.

3.1.1 Differential and symmetry properties of \mathcal{K}^M

Before focussing on the explicit calculation of the kernel function, we show that the formal Mehler formula also generalises the differential equations of the classical Fourier transform.

Theorem 3.1. *The Mehler formula (3.6) obeys the following system of Clifford-valued partial differential equations:*

$$\begin{cases} D_0^{\mathbf{y}} \mathcal{K}^M(\mathbf{x}, \mathbf{y}) &= -\frac{i}{2c} \mathcal{K}^M(\mathbf{x}, \mathbf{y}) \mathbf{x} \\ i\mathbf{y} \mathcal{K}^M(\mathbf{x}, \mathbf{y}) &= -2c [\mathcal{K}^M(\mathbf{x}, \mathbf{y}) D_0^{\mathbf{x}}] \end{cases} \quad (3.7)$$

where $[\cdot D_0^{\mathbf{x}}]$ denotes the differential operator $D_0^{\mathbf{x}}$ acting from the right.

Proof. Both expressions can be proved similarly so we only treat the first relation in full detail. From the definition of the Clifford-Hermite functions $\psi_{j,k}$ and the differential properties of the Clifford-Hermite polynomials $h_{j,k}$, one derives that

$$\begin{aligned} \psi_{j+1,k}(\mathbf{x}) &= \left[(\mathbf{x} - cD_0^{\mathbf{x}}) h_{j,k}(\mathbf{x}) m_k(\mathbf{x}) \right] \exp\left(-\frac{|\mathbf{x}|^2}{4c}\right) \\ &= \mathbf{x} \psi_{j,k}(\mathbf{x}) - c C(j, k) \psi_{j-1,k}(\mathbf{x}) \end{aligned}$$

where the real-valued factor $C(j, k)$ is as in Theorem 2.8. In Proposition 6.2 of [4] the following relation for the Clifford-Hermite functions was obtained:

$$\psi_{j,k}(\mathbf{x}) = \left(\frac{\mathbf{x}}{2} - cD_0 \right) \psi_{j-1,k}(\mathbf{x}), \quad j = 1, 2, 3 \dots$$

Therefore the action of $D_0^{\mathbf{y}}$ on \mathcal{K}^M can be written as

$$\begin{aligned} D_0^{\mathbf{y}} \mathcal{K}^M(\mathbf{x}, \mathbf{y}) &= \sum_{j,k=0}^{\infty} \frac{[D_0^{\mathbf{y}} \psi_{j,k}(\mathbf{y})] (-i)^{j+k+1} \overline{\psi_{j,k}(\mathbf{x})}}{\langle \psi_{j,k}, \psi_{j,k} \rangle} \\ &= \sum_{j,k=0}^{\infty} \frac{[\frac{\mathbf{y}}{2c} \psi_{j,k}(\mathbf{y}) - \frac{1}{c} \psi_{j+1,k}(\mathbf{y})] (-i)^{j+k+1} \overline{\psi_{j,k}(\mathbf{x})}}{\langle \psi_{j,k}, \psi_{j,k} \rangle} \\ &= \frac{1}{2c} \sum_{j,k=0}^{\infty} \frac{[\psi_{j+1,k}(\mathbf{y}) + c C(j, k) \psi_{j-1,k}(\mathbf{y}) - 2\psi_{j+1,k}(\mathbf{y})] (-i)^{j+k+1} \overline{\psi_{j,k}(\mathbf{x})}}{\langle \psi_{j,k}, \psi_{j,k} \rangle} \\ &= \frac{1}{2c} \sum_{j,k=0}^{\infty} \frac{[c C(j, k) \psi_{j-1,k}(\mathbf{y}) - \psi_{j+1,k}(\mathbf{y})] (-i)^{j+k+1} \overline{\psi_{j,k}(\mathbf{x})}}{\langle \psi_{j,k}, \psi_{j,k} \rangle}. \end{aligned}$$

Changing the summation indices yields

$$\begin{aligned} D_0^{\mathbf{y}} \mathcal{K}^M(\mathbf{x}, \mathbf{y}) &= \frac{1}{2} \sum_{j,k=0}^{\infty} C(j+1, k) \frac{\psi_{j,k}(\mathbf{y}) (-i)^{j+k+2} \overline{\psi_{j+1,k}(\mathbf{x})}}{\langle \psi_{j+1,k}, \psi_{j+1,k} \rangle} \\ &\quad - \frac{1}{2c} \sum_{j=1, k=0}^{\infty} \frac{\psi_{j,k}(\mathbf{y}) (-i)^{j+k} \overline{\psi_{j-1,k}(\mathbf{x})}}{\langle \psi_{j-1,k}, \psi_{j-1,k} \rangle}. \end{aligned}$$

Based on Theorem 2.12, one has the following relation for all $j, k \in \mathbb{N}$:

$$\langle \psi_{j,k}, \psi_{j,k} \rangle = -c C(j, k) \langle \psi_{j-1,k}, \psi_{j-1,k} \rangle$$

so $D_0^{\mathbf{y}} \mathcal{K}^M(\mathbf{x}, \mathbf{y})$ can finally be written as

$$\begin{aligned} & -\frac{1}{2c} \sum_{j,k=0}^{\infty} \frac{\psi_{j,k}(\mathbf{y})(-i)^{j+k+2} \overline{\psi_{j+1,k}(\mathbf{x})}}{\langle \psi_{j,k}, \psi_{j,k} \rangle} + \frac{1}{2} \sum_{j,k=0}^{\infty} C(j, k) \frac{\psi_{j,k}(\mathbf{y})(-i)^{j+k} \overline{\psi_{j-1,k}(\mathbf{x})}}{\langle \psi_{j,k}, \psi_{j,k} \rangle} \\ &= \frac{i}{2c} \sum_{j,k=0}^{\infty} \frac{\psi_{j,k}(\mathbf{y})(-i)^{j+k+1} [\overline{\psi_{j+1,k}(\mathbf{x})} + c C(j, k) \overline{\psi_{j-1,k}(\mathbf{x})}]}{\langle \psi_{j,k}, \psi_{j,k} \rangle} \\ &= \frac{i}{2c} \sum_{j,k=0}^{\infty} \frac{\psi_{j,k}(\mathbf{y})(-i)^{j+k+1} \mathbf{x} \overline{\psi_{j,k}(\mathbf{x})}}{\langle \psi_{j,k}, \psi_{j,k} \rangle} \\ &= -\frac{i}{2c} \mathcal{K}^M(\mathbf{x}, \mathbf{y}) \mathbf{x}, \end{aligned}$$

which proves the first relation. The proof of the second relation is completely analogous when starting from the right-hand side. \square

Furthermore, equation (3.6) exhibits an interesting symmetry property. By the noncommutativity of \mathbf{x} and \mathbf{y} , however, it will only show up with respect to an anti-involution. Indeed, taking the Clifford conjugation of (3.6) yields

$$\begin{aligned} \overline{\mathcal{K}^M(\mathbf{x}, \mathbf{y})} &= \sum_{j,k=0}^{\infty} \frac{\overline{\psi_{j,k}(\mathbf{y})(-i)^{j+k+1} \overline{\psi_{j,k}(\mathbf{x})}}}{\langle \psi_{j,k}, \psi_{j,k} \rangle} \\ &= \sum_{j,k=0}^{\infty} \frac{\psi_{j,k}(\mathbf{x})(-i)^{j+k+1} \overline{\psi_{j,k}(\mathbf{y})}}{\langle \psi_{j,k}, \psi_{j,k} \rangle} = (\mathcal{K}^M(\mathbf{y}, \mathbf{x}))^*. \end{aligned}$$

Remark 3.2. *The symmetry of \mathcal{K}^M allows to demonstrate the equivalence of the two partial differential equations. Indeed, from $\overline{\mathcal{K}^M(\mathbf{x}, \mathbf{y})} = (\mathcal{K}^M(\mathbf{y}, \mathbf{x}))^*$ it follows that*

$$\begin{aligned} \overline{D_0^{\mathbf{y}} \mathcal{K}^M(\mathbf{x}, \mathbf{y})} &= \overline{\frac{-i}{2c} \mathcal{K}^M(\mathbf{x}, \mathbf{y}) \mathbf{x}} \\ \Leftrightarrow \left[\overline{\mathcal{K}^M(\mathbf{x}, \mathbf{y})} \quad \overline{D_0^{\mathbf{y}}} \right] &= \frac{i}{2c} \overline{\mathbf{x} \mathcal{K}^M(\mathbf{x}, \mathbf{y})} \\ \Leftrightarrow \left[(\mathcal{K}^M(\mathbf{y}, \mathbf{x}))^* \quad D_0^{\mathbf{y}} \right] &= \frac{i}{2c} \mathbf{x} (\mathcal{K}^M(\mathbf{y}, \mathbf{x}))^* \\ \Leftrightarrow \left[\mathcal{K}^M(\mathbf{x}, \mathbf{y}) \quad D_0^{\mathbf{x}} \right] &= \frac{-i}{2c} \mathbf{y} \mathcal{K}^M(\mathbf{x}, \mathbf{y}), \end{aligned}$$

which equals the second equation. In the last step the complex conjugate is taken and the variables \mathbf{x} and \mathbf{y} , regarded as dummy variables, are interchanged.

3.1.2 Closed form of the kernel function \mathcal{K}^M

Because all entities in (3.6) are known, their explicit expressions can be substituted and a closed expression for \mathcal{K}^M can be obtained. The explicit form of the Clifford-Hermite functions $\psi_{j,k}$ depends on the parity of their degree j . Using the Laguerre form (see Theorem 2.9) for the Clifford-Hermite polynomials, one has

$$\begin{aligned}\psi_{2t,k}(\mathbf{x}) &= (2c)^t t! L_t^k \left(\frac{|\mathbf{x}|^2}{2c} \right) (e_0 - 1)(x_0 + \underline{x})^k \exp \left(-\frac{|\mathbf{x}|^2}{4c} \right) \\ \psi_{2t+1,k}(\mathbf{x}) &= (2c)^t t! \mathbf{x} L_t^{k+1} \left(\frac{|\mathbf{x}|^2}{2c} \right) (e_0 - 1)(x_0 + \underline{x})^k \exp \left(-\frac{|\mathbf{x}|^2}{4c} \right)\end{aligned}$$

and their norms, who show up in the denominator of (3.6), are given in Theorem 2.12. After direct substitution of these expressions, one obtains

$$\begin{aligned}\mathcal{K}^M(\mathbf{x}, \mathbf{y}) &= \frac{-i\Gamma\left(\frac{m}{2}\right)}{2\pi^{m/2+1}} \exp \left(-\frac{|\mathbf{x}|^2 + |\mathbf{y}|^2}{4c} \right) \\ &\times \left[\sum_{k=0}^{+\infty} \frac{(1 - e_0)(y_0 + \underline{y})^k (-i)^k (x_0 - \underline{x})^k (e_0 + 1)}{(2c)^{k+1}} H_t^k(\mathbf{x}, \mathbf{y}) \right. \\ &\left. + i \sum_{k=0}^{+\infty} \frac{\mathbf{y}(1 - e_0)(y_0 + \underline{y})^k (-i)^k (x_0 - \underline{x})^k (e_0 + 1)\mathbf{x}}{(2c)^{k+2}} H_t^{k+1}(\mathbf{x}, \mathbf{y}) \right].\end{aligned}$$

In each summation, all t -dependencies are real-valued and could therefore be grouped in a separate factor H_t^k which reads

$$H_t^k(\mathbf{x}, \mathbf{y}) = \sum_{t=0}^{+\infty} \frac{(-1)^t t! L_t^k \left(\frac{|\mathbf{y}|^2}{2c} \right) L_t^k \left(\frac{|\mathbf{x}|^2}{2c} \right)}{(k+t)!}$$

and is equal for both summations, apart from interchanging $k \leftrightarrow k+1$. The latter expression constitutes a special case of the so-called Hille-Hardy formula [22], which reads

$$\sum_{t=0}^{+\infty} \frac{t!}{\Gamma(k+t+1)} L_t^k(x) L_t^k(y) z^t = \frac{(xyz)^{-\frac{k}{2}}}{(1-z)} \exp \left(-\frac{(x+y)z}{1-z} \right) I_k \left(\frac{2\sqrt{xyz}}{1-z} \right)$$

for $|z| < 1$ and where the modified Bessel function I_k obeys $I_k(x) = i^{-k} J_k(ix)$. The fact that H_t^k is just outside the general domain of convergence is compensated for by the heuristic character of this reasoning. The validity of the final expression for \mathcal{K}^M will be verified explicitly afterwards in Theorem 3.5.

Moreover, recalling that $\mathbf{x} = x_0 e_0 + \underline{x}$ and $\mathbf{y} = y_0 e_0 + \underline{y}$ these variables obey the identities $(e_0 + 1)\mathbf{x} = (x_0 - \underline{x})(e_0 - 1)$ and $\mathbf{y}(1 - e_0) = (1 + e_0)(y_0 + \underline{y})$. Thereby

\mathcal{K}^M is proportional to

$$(1 - e_0) \sum_{k=0}^{+\infty} \frac{(y_0 + \underline{y})^k (-i)^k (x_0 - \underline{x})^k}{(|\mathbf{x}||\mathbf{y}|)^k} J_k \left(\frac{|\mathbf{x}||\mathbf{y}|}{2c} \right) (e_0 + 1) \\ - (1 + e_0) \sum_{k=0}^{+\infty} \frac{(y_0 + \underline{y})^{k+1} (-i)^{k+1} (x_0 - \underline{x})^{k+1}}{(|\mathbf{x}||\mathbf{y}|)^{k+1}} J_{k+1} \left(\frac{|\mathbf{x}||\mathbf{y}|}{2c} \right) (e_0 - 1).$$

Terms corresponding to Bessel functions of the same order can now be gathered by changing the summation index in the second sum, yielding

$$2J_0 \left(\frac{|\mathbf{x}||\mathbf{y}|}{2c} \right) \\ + 2 \sum_{k=1}^{+\infty} (-i)^k \frac{(y_0 + \underline{y})^k (x_0 - \underline{x})^k + (y_0 - \underline{y})^k (x_0 + \underline{x})^k}{(|\mathbf{x}||\mathbf{y}|)^k} J_k \left(\frac{|\mathbf{x}||\mathbf{y}|}{2c} \right). \quad (3.8)$$

At this point, we wish to use the following property of an infinite sum of products of Bessel and cosine functions [23]:

$$e^{-iz \cos(\phi)} = J_0(z) + 2 \sum_{n=1}^{+\infty} (-i)^n J_n(z) \cos(n\phi). \quad (3.9)$$

In order to retrieve this structure in our series expression, we introduce polar coordinates for \mathbf{x} and \mathbf{y} , respectively in the (x_0, r) - and the (y_0, g) -plane, using the respective angles χ and ϕ :

$$\frac{x_0 + \underline{x}}{\sqrt{x_0^2 + r^2}} = \cos(\chi) + \underline{\omega} \sin(\chi) \\ \frac{y_0 + \underline{y}}{\sqrt{y_0^2 + g^2}} = \cos(\phi) + \underline{\eta} \sin(\phi).$$

Regarding $\underline{\omega}$ and $\underline{\eta}$ as counterparts of the classical complex unit, these definitions can be split in their real and imaginary parts. Substituting the latter in (3.8), \mathcal{K}^M turns out to be proportional to

$$2J_0 \left(\frac{|\mathbf{x}||\mathbf{y}|}{2c} \right) + 2 \sum_{k=1}^{+\infty} (-i)^k (\cos(k(\phi + \chi)) + \cos(k(\phi - \chi))) J_k \left(\frac{|\mathbf{x}||\mathbf{y}|}{2c} \right) \\ - 2 \sum_{k=1}^{+\infty} (-i)^k (\cos(k(\phi - \chi)) - \cos(k(\phi + \chi))) J_k \left(\frac{|\mathbf{x}||\mathbf{y}|}{2c} \right) \underline{\eta} \underline{\omega}$$

where the appropriate trigonometric formulas have been used. Recalling equation (3.9), the previous expression can still be rewritten as

$$\mathcal{K}^M(\mathbf{x}, \mathbf{y}) = \frac{-i\Gamma\left(\frac{m}{2}\right)}{8c\pi^{m/2+1}} \left[\left(e^{-\frac{i}{2c}|\mathbf{x}||\mathbf{y}| \cos(\phi+\chi)} + e^{-\frac{i}{2c}|\mathbf{x}||\mathbf{y}| \cos(\phi-\chi)} \right) \right. \\ \left. - \left(e^{-\frac{i}{2c}|\mathbf{x}||\mathbf{y}| \cos(\phi-\chi)} - e^{-\frac{i}{2c}|\mathbf{x}||\mathbf{y}| \cos(\phi+\chi)} \right) \underline{\eta} \underline{\omega} \right].$$

Finally, eliminating the angular variables and grouping corresponding terms, we obtain

$$\mathcal{K}^M(\mathbf{x}, \mathbf{y}) = \frac{-i\Gamma\left(\frac{m}{2}\right)}{8c\pi^{m/2+1}} \left[(1 + \underline{\eta}\underline{\omega})e^{-\frac{i}{2c}(x_0y_0 - rg)} + (1 - \underline{\eta}\underline{\omega})e^{-\frac{i}{2c}(x_0y_0 + rg)} \right]. \quad (3.10)$$

A straightforward verification shows that (3.10) is indeed a solution of the Clifford-valued partial differential equations

$$\begin{cases} D_0^y \mathcal{K}^M(\mathbf{x}, \mathbf{y}) &= -\frac{i}{2c} \mathcal{K}^M(\mathbf{x}, \mathbf{y}) \mathbf{x} \\ i\mathbf{y} \mathcal{K}^M(\mathbf{x}, \mathbf{y}) &= -2c [\mathcal{K}^M(\mathbf{x}, \mathbf{y}) D_0^x] \end{cases}$$

and also obeys $\overline{\mathcal{K}^M(\mathbf{x}, \mathbf{y})} = (\mathcal{K}^M(\mathbf{y}, \mathbf{x}))^*$. Having obtained this closed expression for the kernel function, we can conclude this subsection with the explicit definition of the slice Fourier transform.

Definition 3.3 (Slice Fourier transform). *The slice Fourier transform of a function $f \in \mathcal{V}$ is given by*

$$\mathcal{F}_S(f)(\mathbf{y}) = \frac{-i\Gamma\left(\frac{m}{2}\right)}{8c\pi^{m/2+1}} \times \int_{\mathbb{R}^{m+1}} \left[(1 + \underline{\eta}\underline{\omega})e^{-\frac{i}{2c}(x_0y_0 - rg)} + (1 - \underline{\eta}\underline{\omega})e^{-\frac{i}{2c}(x_0y_0 + rg)} \right] f(\mathbf{x}) \, dx_0 dr d\sigma_{\underline{x}}$$

with $\mathbf{x} = x_0e_0 + r\underline{\omega}$ and $\mathbf{y} = y_0e_0 + g\underline{\eta}$.

Remark 3.4. *Using Eulers formula, the following equivalent expression for the slice Fourier transform is obtained:*

$$\mathcal{F}_S(f)(\mathbf{y}) = \frac{-i\Gamma\left(\frac{m}{2}\right)}{4c\pi^{m/2+1}} \int_{\mathbb{R}^{m+1}} e^{-\frac{i}{2c}(x_0y_0)} \left[\cos\left(\frac{rg}{2c}\right) + i\underline{\eta}\underline{\omega} \sin\left(\frac{rg}{2c}\right) \right] f(\mathbf{x}) dx_0 dr d\sigma_{\underline{x}}.$$

3.2 Verification of the eigenfunctions

Now the integral expression for the slice Fourier transform has been obtained explicitly, we can further investigate its properties. In this first subsection we demonstrate that definition 3.3 is indeed an integral transform whose eigenfunctions are the Clifford-Hermite functions.

Theorem 3.5. *The functions $\psi_{j,k}$ are eigenfunctions of the integral transform \mathcal{F}_S with eigenvalues $(-i)^{j+k+1}$.*

Proof. We already proved that the kernel function \mathcal{K}^M obeys the system of Clifford-valued partial differential equations

$$\begin{cases} D_0^y \mathcal{K}^M(\mathbf{x}, \mathbf{y}) = -\frac{i}{2c} \mathcal{K}^M(\mathbf{x}, \mathbf{y}) \mathbf{x} & (3.11) \\ i\mathbf{y} \mathcal{K}^M(\mathbf{x}, \mathbf{y}) = -[2c \mathcal{K}^M(\mathbf{x}, \mathbf{y}) D_0^x] & (3.12) \end{cases}$$

where $[\cdot D_0^{\mathbf{x}}]$ denotes the differential operator $D_0^{\mathbf{x}}$ acting from the right. Writing $\psi_{j,k}(\mathbf{x}) = \left(\frac{\mathbf{x}}{2} - cD_0^{\mathbf{x}}\right) \psi_{j-1,k}(\mathbf{x})$ for $j > 0$ and performing partial integration on D_0 yields

$$\begin{aligned} \mathcal{F}_S(\psi_{j,k})(\mathbf{y}) &= \int_{\mathbb{R}^{m+1}} \mathcal{K}^M(\mathbf{x}, \mathbf{y}) \left(\frac{\mathbf{x}}{2} - cD_0^{\mathbf{x}}\right) \psi_{j-1,k}(\mathbf{x}) \, dx_0 \, dr \, d\sigma_{\underline{x}} \\ &= \int_{\mathbb{R}^{m+1}} \left(\mathcal{K}^M(\mathbf{x}, \mathbf{y}) \frac{\mathbf{x}}{2} + c[\mathcal{K}^M(\mathbf{x}, \mathbf{y}) D_0^{\mathbf{x}}]\right) \psi_{j-1,k}(\mathbf{x}) \, dx_0 \, dr \, d\sigma_{\underline{x}}. \end{aligned}$$

Using (3.11) and (3.12) and performing this procedure j times, we get

$$\mathcal{F}_S(\psi_{j,k})(\mathbf{y}) = (-i)^j \left(\frac{\mathbf{y}}{2} - cD_0^{\mathbf{y}}\right)^j \int_{\mathbb{R}^{m+1}} \mathcal{K}^M(\mathbf{x}, \mathbf{y}) \psi_{0,k}(\mathbf{x}) \, dx_0 \, dr \, d\sigma_{\underline{x}}$$

so we only have to prove that the functions $\psi_{0,k}$ are eigenfunctions of \mathcal{F}_S with corresponding eigenvalues $(-i)^{k+1}$. Using the equality

$$(x_0 + \underline{x}) = \left(\frac{1-e_0}{\sqrt{2}}\right) \mathbf{x} \left(\frac{1-e_0}{\sqrt{2}}\right),$$

the expression for the Hermite functions $\psi_{0,k}$ can be rewritten as

$$\psi_{0,k}(\mathbf{x}) = (e_0 - 1) \left[\left(\frac{1-e_0}{\sqrt{2}}\right) \mathbf{x} \left(\frac{1-e_0}{\sqrt{2}}\right) \right]^k \exp\left(-\frac{|\mathbf{x}|^2}{4c}\right).$$

Given that both $\underline{\omega}$ and $\underline{\eta}$ in \mathcal{K}^M anticommute with e_0 , one has $\mathcal{K}^M e_0 = e_0 \mathcal{K}^M$. Using (3.11), the slice Fourier transform of $\psi_{0,k}$ is given by

$$\begin{aligned} &\int_{\mathbb{R}^{m+1}} \mathcal{K}^M(\mathbf{x}, \mathbf{y}) (e_0 - 1) \left[\left(\frac{1-e_0}{\sqrt{2}}\right) \mathbf{x} \left(\frac{1-e_0}{\sqrt{2}}\right) \right]^k \exp\left(-\frac{|\mathbf{x}|^2}{4c}\right) \, dx_0 \, dr \, d\sigma_{\underline{x}} \\ &= (e_0 - 1) (2ci)^k \left[\left(\frac{1-e_0}{\sqrt{2}}\right) D_0^{\mathbf{y}} \left(\frac{1-e_0}{\sqrt{2}}\right) \right]^k \\ &\quad \times \int_{\mathbb{R}^{m+1}} \mathcal{K}^M(\mathbf{x}, \mathbf{y}) \exp\left(-\frac{|\mathbf{x}|^2}{4c}\right) \, dx_0 \, dr \, d\sigma_{\underline{x}}. \end{aligned}$$

Using the fact that $\int_{\mathbb{S}^{m-1}} \underline{\omega} \, d\sigma_{\underline{x}} = 0$ together with the identities

$$\int_{-\infty}^{+\infty} e^{\pm i \frac{x_0 y_0}{2c}} e^{-\frac{x_0^2}{4c}} \, dx_0 = 2\sqrt{\pi c} e^{-\frac{y_0^2}{4c}} \quad \text{and} \quad \int_0^{+\infty} \cos\left(\frac{rg}{2c}\right) e^{-\frac{r^2}{4c}} \, dr = \sqrt{\pi c} e^{-\frac{g^2}{4c}},$$

the integral reduces to

$$\frac{2}{4\pi c} \int_{-\infty}^{+\infty} \int_0^{+\infty} e^{-\frac{i}{2c} x_0 y_0} \cos\left(\frac{rg}{2}\right) \exp\left(-\frac{|\mathbf{x}|^2}{4c}\right) \, dx_0 \, dr = \exp\left(-\frac{|\mathbf{y}|^2}{4c}\right)$$

and we finally obtain

$$\begin{aligned}
\mathcal{F}_S(\psi_{0,k})(\mathbf{y}) &= (2ci)^k (e_0 - 1) \left[\left(\frac{1 - e_0}{\sqrt{2}} \right) D_0^{\mathbf{y}} \left(\frac{1 - e_0}{\sqrt{2}} \right) \right]^k \exp \left(-\frac{|\mathbf{y}|^2}{4c} \right) \\
&= -i(2ci)^k (e_0 - 1) (\partial_{y_0} + \underline{\eta} \partial_g)^k \exp \left(-\frac{|\mathbf{y}|^2}{4c} \right) \\
&= -i \frac{(2ci)^k}{(-2c)^k} (e_0 - 1) (y_0 + \underline{y})^k \exp \left(-\frac{|\mathbf{y}|^2}{4c} \right) \\
&= (-i)^{k+1} \psi_{0,k}(\mathbf{y}).
\end{aligned}$$

As was noticed above, the degree j yields another factor $(-i)^j$ so we have proven that the functions $\psi_{j,k}$ are eigenfunctions of \mathcal{F}_S with eigenvalues $(-i)^{j+k+1}$. \square

3.3 Verification of the action of the inverse

In the derivation of the integral kernel, we required the eigenvalues of the functions $\psi_{j,k}$ to be $(-i)^{j+k+1}$. Now the action of the inverse transform is to undo the action of the slice Fourier transform, so to transform $(-i)^{j+k+1} \psi_{j,k}$ into the original Clifford-Hermite function $\psi_{j,k}$. Therefore we require the eigenvalues of the inverse transform with respect to $\psi_{j,k}$ to be i^{j+k+1} . Given the formal series expression (3.6) of \mathcal{K}^M , one quickly observes that by this reasoning the kernel of the inverse transform will be the complex conjugate of the kernel of the slice Fourier transform. We thus get the following theorem.

Theorem 3.6. *With $\mathbf{x} = x_0 e_0 + r\underline{\omega}$, $\mathbf{y} = y_0 e_0 + g\underline{\eta}$ and $F \in \mathcal{V}$, the functions $\psi_{j,k}$ are eigenfunctions of the integral transform*

$$\begin{aligned}
\mathcal{F}_S^{-1}(F)(\mathbf{y}) &= \frac{i\Gamma\left(\frac{m}{2}\right)}{8c\pi^{m/2+1}} \\
&\int_{\mathbb{R}^{m+1}} \left[(1 + \underline{\eta}\underline{\omega}) e^{\frac{i}{2c}(x_0 y_0 - rg)} + (1 - \underline{\eta}\underline{\omega}) e^{\frac{i}{2c}(x_0 y_0 + rg)} \right] F(\mathbf{x}) dx_0 dr d\sigma_{\underline{x}}
\end{aligned}$$

with eigenvalues given by i^{j+k+1} .

Proof. The kernel function of the inverse slice Fourier transform, denoted as $\mathcal{K}^{M,-1}$, is the complex conjugate of \mathcal{K}^M . Therefore we obtain from equation (3.7) that it obeys the following system of Clifford-valued partial differential equations:

$$\begin{cases} D_0^{\mathbf{y}} \mathcal{K}^{M,-1}(\mathbf{x}, \mathbf{y}) &= \frac{i}{2c} \mathcal{K}^{M,-1}(\mathbf{x}, \mathbf{y}) \mathbf{x} \\ i\mathbf{y} \mathcal{K}^{M,-1}(\mathbf{x}, \mathbf{y}) &= [2c \mathcal{K}^{M,-1}(\mathbf{x}, \mathbf{y}) D_0^{\mathbf{x}}]. \end{cases}$$

The rest of the proof is completely analogous to the proof of Theorem 3.5. \square

Theorem 3.7. *The integral transform \mathcal{F}_S^{-1} in the above Theorem 3.6 is the inverse slice Fourier transform of a function $F \in \mathcal{V}$.*

Proof. Using Theorem 3.5 and Theorem 3.6 we get for all Clifford-Hermite functions $\psi_{j,k}$ that

$$(\mathcal{F}_S^{-1} \circ \mathcal{F}_S)(\psi_{j,k}) = (\mathcal{F}_S^{-1})((-i)^{j+k+1}\psi_{j,k}) = (i)^{j+k+1}(-i)^{j+k+1}\psi_{j,k} = \psi_{j,k}$$

and analogously for $(\mathcal{F}_S \circ \mathcal{F}_S^{-1})(\psi_{j,k})$. Because $F \in \text{span}\{\psi_{j,k}\}$ this proves the theorem. \square

3.4 Basic properties of the slice Fourier transform

Within this subsection the constant prefactor of the trigonometric expression for \mathcal{K}^M will be denoted as

$$C_m = \frac{-i\Gamma\left(\frac{m}{2}\right)}{4c\pi^{m/2+1}}.$$

As announced, the explicit expression for the integral kernel allows for the study of various properties of the slice Fourier transform by direct calculation. Here some of these basic properties are summarised.

3.4.1 Translation property

Denoting a translation in the e_0 -direction as $t_a f(x_0, r, \underline{\omega}) = f(x_0 - a, r, \underline{\omega})$, one has

$$\begin{aligned} \mathcal{F}_S(t_a f)(\mathbf{y}) &= \int_{\mathbb{R}^{m+1}} \mathcal{K}(\mathbf{x}, \mathbf{y}) f(x_0 - a, r, \underline{\omega}) dx_0 dr d\sigma_{\underline{x}} \\ &= e^{-\frac{ia y_0}{2c}} \mathcal{F}_S(f)(\mathbf{y}). \end{aligned}$$

There is no analogous property for translations in the $\underline{\omega}$ -direction because of its spherical nature.

3.4.2 Reflection property

Denoting a reflection with respect to the origin as $sf(x_0, r, \underline{\omega}) = f(-x_0, r, -\underline{\omega})$, one has

$$\begin{aligned} \mathcal{F}_S(sf)(\mathbf{y}) &= C_m \int_{\mathbb{R}^{m+1}} e^{\frac{ix_0 y_0}{2c}} \left[\cos\left(\frac{rg}{2c}\right) - \underline{\eta}\underline{\omega}i \sin\left(\frac{rg}{2c}\right) \right] f(x_0, r, \underline{\omega}) dx_0 dr d\sigma_{\underline{x}} \\ &= \mathcal{F}_S(f)(-y_0, g, -\underline{\eta}) \\ &= s\mathcal{F}_S(f)(\mathbf{y}). \end{aligned}$$

3.4.3 Complex conjugation property

Denoting the complex conjugate of f as before as f^* , one has

$$\begin{aligned}\mathcal{F}_S(f^*)(\mathbf{y}) &= C_m \int_{\mathbb{R}^{m+1}} e^{-\frac{ix_0y_0}{2c}} \left[\cos\left(\frac{rg}{2c}\right) + \underline{\eta}\underline{\omega}i \sin\left(\frac{rg}{2c}\right) \right] f^* dx_0 dr d\sigma_{\underline{x}} \\ &= -C_m^* \int_{\mathbb{R}^{m+1}} \left(e^{-\frac{ix_0(-y_0)}{2c}} \left[\cos\left(\frac{rg}{2c}\right) + (-\underline{\eta})\underline{\omega}i \sin\left(\frac{rg}{2c}\right) \right] f(\mathbf{x}) \right)^* dx_0 dr d\sigma_{\underline{x}} \\ &= -\mathcal{F}_S^*(f)(-y_0, g, -\underline{\eta}).\end{aligned}$$

3.4.4 Commutation with e_0

Given that both $\underline{\omega}$ and $\underline{\eta}$ anticommute with e_0 , one has

$$\mathcal{F}_S(e_0 f)(\mathbf{y}) = e_0 \mathcal{F}_S(f)(\mathbf{y}).$$

3.4.5 Twofold transform

Applying two consecutive slice Fourier transforms to a function $f \in \text{span}\{\psi_{j,k}\}$ yields

$$\mathcal{F}_S(\mathcal{F}_S(f))(\mathbf{x}) = -f(-\mathbf{x})$$

because $\mathcal{F}_S(\mathcal{F}_S(\psi_{j,k}))(\mathbf{x}) = (-i)^{2+2j+2k} \psi_{j,k}(\mathbf{x}) = (-1)^{j+k+1} \psi_{j,k}(\mathbf{x})$, which equals $-\psi_{j,k}(-\mathbf{x})$ because $\psi_{j,k}(-\mathbf{x}) = (-1)^{j+k} \psi_{j,k}(\mathbf{x})$.

3.5 Explicit calculation

In this last subsection, we take a closer look at the computational load of the slice Fourier transform. To this end we consider the computation of the slice Fourier transform of a function $f \in \mathcal{V}$ explicitly. Using Remark 2.15 and performing the spherical integration in the definition of the transform, we obtain

$$\begin{aligned}\mathcal{F}_S(f)(\mathbf{y}) &= \frac{-i}{4\pi c} \int_{-\infty}^{+\infty} \int_0^{+\infty} \left[\left(e^{-\frac{i}{2c}(x_0y_0-rg)} + e^{-\frac{i}{2c}(x_0y_0+rg)} \right) f_1(x_0, r) \right. \\ &\quad \left. - \left(e^{-\frac{i}{2c}(x_0y_0-rg)} - e^{-\frac{i}{2c}(x_0y_0+rg)} \right) \underline{\eta} f_2(x_0, r) \right] dx_0 dr,\end{aligned}$$

which equals

$$\begin{aligned}\mathcal{F}_S(f)(\mathbf{y}) &= \frac{-i}{2\pi c} \int_{-\infty}^{+\infty} \int_0^{+\infty} e^{-\frac{i}{2c}x_0y_0} \cos\left(\frac{rg}{2c}\right) f_1(x_0, r) dx_0 dr \\ &\quad - \underline{\eta} \frac{1}{2\pi c} \int_{-\infty}^{+\infty} \int_0^{+\infty} e^{-\frac{i}{2c}x_0y_0} \sin\left(\frac{rg}{2c}\right) f_2(x_0, r) dx_0 dr.\end{aligned}$$

Extending f_1 to the function $f_1^+ : \mathbb{R}^2 \rightarrow Cl^{m+1} : (x_0, r) \mapsto f_1(x_0, |r|)$ which is even in its second argument and analogously extending f_2 to the function

$$f_2^- : \mathbb{R}^2 \rightarrow Cl^{m+1} : (x_0, r) \mapsto \begin{cases} f_2(x_0, r) & r > 0 \\ -f_2(x_0, -r) & r < 0 \end{cases}$$

which is odd in its second argument, the integrals can be rewritten as

$$\mathcal{F}_S(f)(\mathbf{y}) = \frac{-i}{4\pi c} \int_{-\infty}^{+\infty} \int_{-\infty}^{+\infty} e^{-\frac{i}{2c}(x_0 y_0 + r g)} [f_1^+(x_0, r) + \underline{\eta} f_2^-(x_0, r)] dx_0 dr.$$

Therefore we can conclude that the slice Fourier transform $\mathcal{F}_S(f)$ of a real Clifford-valued function f can be calculated solely using the classical two-dimensional Fourier transform of a single two-dimensional function $F(x_0, r) = f_1^+(x_0, r) - i f_2^-(x_0, r)$:

$$\mathcal{F}_S(f)(\mathbf{y}) = \frac{-i}{2c} (FT(F)^+(y_0, g) + \underline{\eta} FT(F)^-(y_0, g)).$$

Again the superscripts $+$ and $-$ denote, respectively, the even and odd parts of the functions in their second argument and FT is the classical two-dimensional Fourier transform of a complex-valued function F defined as

$$FT(F)(y_0, g) = \frac{1}{2\pi} \int_{-\infty}^{+\infty} \int_{-\infty}^{+\infty} e^{-\frac{i}{2c}(x_0 y_0 + r g)} F(x_0, r) dx_0 dr.$$

4 Convolutions

In this section two different approaches to the convolution operator associated with the slice Fourier transform are treated. Both of them satisfy the demand that the slice Fourier transform of them equals the product of the separate Fourier transforms. In a first approach the classical definition of the convolution is generalised using the Mustard convolution (see [21]). In a second approach the translation operator in the integrand of the classical convolution is generalised, based on [3]. Finally, given that both approaches give rise to the same behaviour in the Fourier domain, we pinpoint the connection between both in the last subsection.

A major drawback in constructing convolutions in a Clifford setting, however, is the lack of commutativity. Any Clifford counterpart to the classical convolution \star will therefore automatically fail to generalise one of its primary properties, namely $f \star g = g \star f$.

Remark 4.1. *In this section all generalisations are constructed such that in the Fourier domain the function f is right multiplied with g . Analogous constructions can be done for left multiplication.*

4.1 Technical prerequisites

Let us first recall the following lemma which computes the integral of a squared component of a unit vector over the unit sphere.

Lemma 4.2. *With $\underline{\omega} = \omega_1 e_1 + \omega_2 e_2 + \dots + \omega_m e_m$ a unit vector in Cl_m , one has*

$$\int_{\mathbb{S}^{m-1}} \omega_i^2 d\sigma_{\underline{\omega}} = \frac{2\pi^{m/2}}{m\Gamma(\frac{m}{2})}.$$

Proof. Given that

$$\int_{\mathbb{S}^{m-1}} \underline{\omega}^2 d\sigma_{\underline{\omega}} = - \int_{\mathbb{S}^{m-1}} 1 d\sigma_{\underline{\omega}} = - \sum_{i=1}^m \int_{\mathbb{S}^{m-1}} \omega_i^2 d\sigma_{\underline{\omega}},$$

the above expression follows from the symmetry of the problem and the expression for the area of an m -dimensional sphere, $\frac{2\pi^{m/2}}{\Gamma(\frac{m}{2})}$. \square

Another useful lemma addresses the sum of a k -vector squeezed between all basis elements of Cl_m .

Lemma 4.3. *With $\underline{a}^{(k)}$ a k -vector in Cl_m (so $k \leq m$), one has*

$$\sum_{i=1}^m e_i \underline{a}^{(k)} e_i = (-1)^k (2k - m) \underline{a}^{(k)}.$$

Proof. Given that $\underline{a}^{(k)}$ is a k -vector, possible values for k range from 0 to m . We prove this lemma using induction.

- **$k = 0$** When $\underline{a}^{(k)}$ is a scalar the expression reads

$$\sum_{i=1}^m e_i \underline{a}^{(0)} e_i = -m \underline{a}^{(0)},$$

which is identically true.

- **$k = k' + 1$ ($\leq m$)** Assuming the validity for $k = k'$ ($k' < m$), one has $\sum_{i=1}^m e_i \underline{a}^{(k')} e_i = (-1)^{k'} (2k' - m) \underline{a}^{(k')}$. Without loss of generality we assume $\underline{a}^{(k')}$ to consist of one single term. Now a $(k' + 1)$ -vector can be constructed as $\underline{a}^{(k'+1)} = \underline{a}^{(k')} e_\ell$ where the basis vector e_ℓ is not contained

in $\underline{a}^{(k')}$. We get

$$\begin{aligned}
\sum_{i=1}^m e_i \underline{a}^{(k'+1)} e_i &= \sum_{i=1}^m e_i \underline{a}^{(k')} e_\ell e_i \\
&= \sum_{i=1}^m e_i \underline{a}^{(k')} (-2\delta_{i\ell} - e_i e_\ell) \\
&= -2e_\ell \underline{a}^{(k')} - \left(\sum_{i=1}^m e_i \underline{a}^{(k')} e_i \right) e_\ell \\
&= -2(-1)^{k'} \underline{a}^{(k')} e_\ell - (-1)^{k'} (2k' - m) \underline{a}^{(k')} e_\ell \\
&= (-1)^{k'+1} [2 + (2k' - m)] \underline{a}^{(k')} e_\ell \\
&= (-1)^{k'+1} (2(k' + 1) - m) \underline{a}^{(k'+1)},
\end{aligned}$$

which proves the expression for $k = k' + 1$.

By induction, this proves the lemma. \square

Now we can state the main technical theorem. The integral treated by this theorem will be met several times in the following subsections.

Theorem 4.4. *With $\underline{a} \in Cl_m$, one has*

$$\int_{\mathbb{S}^{m-1}} \underline{\omega} \underline{a} \underline{\omega} \, d\omega = \frac{2\pi^{m/2}}{m\Gamma\left(\frac{m}{2}\right)} \sum_{k=0}^m (-1)^k (2k - m) \underline{a}^{(k)}.$$

Proof. Writing

$$\underline{\omega} = \sum_{i=1}^m \omega_i e_i,$$

the integral can be rewritten as

$$\int_{\mathbb{S}^{m-1}} \underline{\omega} \underline{a} \underline{\omega} \, d\omega = \sum_{i,j=1}^m \left(\int_{\mathbb{S}^{m-1}} \omega_i \omega_j \, d\sigma_\omega \right) e_i \underline{a} e_j = \sum_{i=1}^m \left(\int_{\mathbb{S}^{m-1}} \omega_i^2 \, d\sigma_\omega \right) e_i \underline{a} e_i$$

where the last equality holds because the integral over the sphere vanishes when $i \neq j$. Using lemmas 4.2 and 4.3, the theorem follows. \square

4.2 Mustard convolution

As pointed out in the introduction, a Mustard convolution is defined such that its Fourier transform equals the product of the Fourier transforms of both functions separately. The Mustard convolution corresponding to the slice Fourier transform is therefore defined as

$$f \star_S g = \mathcal{F}_S^{-1} (\mathcal{F}_S(f) \mathcal{F}_S(g)), \tag{4.13}$$

where f and g are Cl_{m+1} -valued functions belonging to \mathcal{V} . Using $\mathbf{y} = y_0 e_0 + g\eta$, $\mathbf{z} = z_0 e_0 + n\underline{\zeta}$ and $\mathbf{u} = u_0 e_0 + h\underline{\nu}$ as integration variables, the right-hand side reads in full

$$\begin{aligned} \mathcal{F}_S^{-1}(\mathcal{F}_S(f)\mathcal{F}_S(g))(\mathbf{x}) &= -i \left(\frac{\Gamma(\frac{m}{2})}{8c\pi^{m/2+1}} \right)^3 \int_{\mathbb{R}^{m+1}} \int_{\mathbb{R}^{m+1}} \int_{\mathbb{R}^{m+1}} \\ &\quad \left[(1 + \underline{\omega}\eta) e^{+\frac{i}{2c}(x_0 y_0 - rg)} + (1 - \underline{\omega}\eta) e^{+\frac{i}{2c}(x_0 y_0 + rg)} \right] \times \\ &\quad \left[(1 + \underline{\eta}\zeta) e^{-\frac{i}{2c}(y_0 z_0 - gn)} + (1 - \underline{\eta}\zeta) e^{-\frac{i}{2c}(y_0 z_0 + gn)} \right] f(\mathbf{z}) \times \\ &\quad \left[(1 + \underline{\eta}\nu) e^{-\frac{i}{2c}(y_0 u_0 - gh)} + (1 - \underline{\eta}\nu) e^{-\frac{i}{2c}(y_0 u_0 + gh)} \right] g(\mathbf{u}) \times \\ &\quad dy_0 dg d\sigma_{\underline{y}} dz_0 dn d\sigma_{\underline{z}} du_0 dh d\sigma_{\underline{u}}. \end{aligned}$$

Performing the integration over y_0 formally, a factor $4\pi c \delta(x_0 - z_0 - u_0)$ shows up in the integrand. Rearranging terms, keeping in mind the integration over $\underline{\eta}$ and using distributivity, the remaining part of the integrand reads

$$\begin{aligned} &(1 - \underline{\omega}\zeta) f(\mathbf{z}) \left(e^{-\frac{i}{2c}(r-n-h)g} + e^{-\frac{i}{2c}(r-n+h)g} + e^{\frac{i}{2c}(r-n+h)g} + e^{\frac{i}{2c}(r-n-h)g} \right) \\ &+ (1 + \underline{\omega}\zeta) f(\mathbf{z}) \left(e^{-\frac{i}{2c}(r+n-h)g} + e^{-\frac{i}{2c}(r+n+h)g} + e^{\frac{i}{2c}(r+n+h)g} + e^{\frac{i}{2c}(r+n-h)g} \right) \\ &+ (\underline{\omega}\eta + \underline{\eta}\zeta) f(\mathbf{z}) \underline{\eta}\nu \left(e^{-\frac{i}{2c}(r-n-h)g} - e^{-\frac{i}{2c}(r-n+h)g} - e^{\frac{i}{2c}(r-n+h)g} + e^{\frac{i}{2c}(r-n-h)g} \right) \\ &+ (\underline{\omega}\eta - \underline{\eta}\zeta) f(\mathbf{z}) \underline{\eta}\nu \left(e^{-\frac{i}{2c}(r+n-h)g} - e^{-\frac{i}{2c}(r+n+h)g} - e^{\frac{i}{2c}(r+n+h)g} + e^{\frac{i}{2c}(r+n-h)g} \right) \end{aligned}$$

times $g(\mathbf{u})$. Using the formal equality

$$\int_0^{+\infty} e^{-\frac{i}{2c}(r-n-h)g} dg = \frac{1}{2} \int_0^{+\infty} e^{-\frac{i}{2c}(r-n-h)g} dg + \frac{1}{2} \int_{-\infty}^0 e^{\frac{i}{2c}(r-n-h)g} dg$$

to dissolve all terms in the previous expression and recombining terms with respect to the integrands, formal delta distributions are obtained in r , n and h too. Writing $f, g \in \mathcal{V} = \text{span}_{Cl}\{\psi_{j,k}\}$ as $f(\mathbf{z}) = f_1(z_0, n) + \underline{\zeta}f_2(z_0, n)$ and $g(\mathbf{u}) = g_1(u_0, h) + \underline{\nu}g_2(u_0, h)$, we get

$$\begin{aligned} \mathcal{F}_S^{-1}(\mathcal{F}_S(f)\mathcal{F}_S(g))(\mathbf{x}) &= -i \left(\frac{\Gamma(\frac{m}{2})}{8c\pi^{m/2+1}} \right) \int_{\mathbb{S}^{m-1}} \int_{-\infty}^{+\infty} \int_0^{+\infty} \int_{-\infty}^{+\infty} \int_0^{+\infty} \delta(x_0 - z_0 - u_0) \times \\ &\quad \left[(f_1(z_0, n) + \underline{\omega} f_2(z_0, n)) g_1(u_0, h) (\delta(r - n - h) + \delta(r - n + h)) \right. \\ &\quad - \underline{\omega} (\underline{\eta}f_1(z_0, n) + \underline{\omega}\eta f_2(z_0, n)) \underline{\eta}g_2(u_0, h) (\delta(r - n - h) - \delta(r - n + h)) \\ &\quad + (f_1(z_0, n) - \underline{\omega} f_2(z_0, n)) g_1(u_0, h) (\delta(r + n - h) + \delta(r + n + h)) \\ &\quad \left. - \underline{\omega} (\underline{\eta}f_1(z_0, n) - \underline{\omega}\eta f_2(z_0, n)) \underline{\eta}g_2(u_0, h) (\delta(r + n - h) - \delta(r + n + h)) \right] \\ &\quad \times d\sigma_{\underline{y}} dz_0 dn du_0 dh, \end{aligned}$$

where we have integrated over $\underline{\zeta}$ and $\underline{\nu}$. Given that r, n and h can only take positive values, terms containing $\delta(r + n + h)$ do not contribute to the final result. Using Theorem 4.4, the integration over $\underline{\eta}$ yields

$$\begin{aligned}
\mathcal{F}_S^{-1}(\mathcal{F}_S(f)\mathcal{F}_S(g))(\mathbf{x}) &= \frac{-i}{4\pi c} \int_{-\infty}^{+\infty} \int_0^{+\infty} \int_{-\infty}^{+\infty} \int_0^{+\infty} \delta(x_0 - z_0 - u_0) \times \\
&\left\{ \left[f(z_0, n, \underline{\omega})g_1(u_0, h) - \underline{\omega} \sum_{k=0}^m (-1)^k \left(\frac{2k}{m} - 1 \right) f^{(k)}(z_0, n, \underline{\omega})g_2(u_0, h) \right] \right. \\
&\quad \times \delta(r - n - h) \\
&+ \left[f(z_0, n, \underline{\omega})g_1(u_0, h) + \underline{\omega} \sum_{k=0}^m (-1)^k \left(\frac{2k}{m} - 1 \right) f^{(k)}(z_0, n, \underline{\omega})g_2(u_0, h) \right] \\
&\quad \times \delta(r - n + h) \\
&+ \left[f(z_0, n, -\underline{\omega})g_1(u_0, h) - \underline{\omega} \sum_{k=0}^m (-1)^k \left(\frac{2k}{m} - 1 \right) f^{(k)}(z_0, n, -\underline{\omega})g_2(u_0, h) \right] \\
&\quad \left. \times \delta(r + n - h) \right\} dz_0 dn du_0 dh,
\end{aligned}$$

where the k -vector parts of f_1 and f_2 have been denoted as $f_1^{(k)}$ and $f_2^{(k)}$, respectively. Performing the integrations over u_0 and h , the expression for the

Mustard convolution reads

$$\begin{aligned}
\mathcal{F}_S^{-1}(\mathcal{F}_S(f)\mathcal{F}_S(g))(\mathbf{x}) &= \frac{-i}{4\pi c} \int_{-\infty}^{+\infty} \\
&\left\{ \int_0^r \left[f_1(z_0, n)g_1(x_0 - z_0, r - n) + \sum_{k=0}^m (-1)^k \left(\frac{2k}{m} - 1 \right) f_2^{(k)}(z_0, n)g_2(x_0 - z_0, r - n) \right] dn \right. \\
&+ \int_r^{+\infty} \left[f_1(z_0, n)g_1(x_0 - z_0, n - r) - \sum_{k=0}^m (-1)^k \left(\frac{2k}{m} - 1 \right) f_2^{(k)}(z_0, n)g_2(x_0 - z_0, n - r) \right] dn \\
&+ \int_0^{+\infty} \left[f_1(z_0, n)g_1(x_0 - z_0, r + n) - \sum_{k=0}^m (-1)^k \left(\frac{2k}{m} - 1 \right) f_2^{(k)}(z_0, n)g_2(x_0 - z_0, r + n) \right] dn \\
&+ \underline{\omega} \int_0^r \left[f_2(z_0, n)g_1(x_0 - z_0, r - n) - \sum_{k=0}^m (-1)^k \left(\frac{2k}{m} - 1 \right) f_1^{(k)}(z_0, n)g_2(x_0 - z_0, r - n) \right] dn \\
&+ \underline{\omega} \int_r^{+\infty} \left[f_2(z_0, n)g_1(x_0 - z_0, n - r) + \sum_{k=0}^m (-1)^k \left(\frac{2k}{m} - 1 \right) f_1^{(k)}(z_0, n)g_2(x_0 - z_0, n - r) \right] dn \\
&\left. - \underline{\omega} \int_0^{+\infty} \left[f_2(z_0, n)g_1(x_0 - z_0, r + n) + \sum_{k=0}^m (-1)^k \left(\frac{2k}{m} - 1 \right) f_1^{(k)}(z_0, n)g_2(x_0 - z_0, r + n) \right] dn \right\} \\
&dz_0. \tag{4.14}
\end{aligned}$$

Given the complexity of the resulting expression, we have implemented and thoroughly verified this result using the computersoftware Maple, which confirmed our calculations. Nonetheless, a careful study of the different terms suggests some hidden symmetry underneath. Indeed, defining even and odd extensions in the second argument as

$$f_1^{even}(z_0, n) = \begin{cases} f_1(z_0, n) & n > 0 \\ f_1(z_0, -n) & n < 0 \end{cases}$$

and

$$f_1^{odd}(z_0, n) = \begin{cases} f_1(z_0, n) & n > 0 \\ -f_1(z_0, -n) & n < 0 \end{cases}$$

we finally obtain

$$\begin{aligned}
& \mathcal{F}_S^{-1}(\mathcal{F}_S(f)\mathcal{F}_S(g))(\mathbf{x}) \\
&= \frac{-i}{4\pi c} \int_{-\infty}^{+\infty} \left\{ \int_{-\infty}^{+\infty} \left[f_1^{even}(z_0, n) g_1^{even}(x_0 - z_0, r - n) \right. \right. \\
&\quad \left. \left. + \sum_{k=0}^m (-1)^k \left(\frac{2k}{m} - 1 \right) f_2^{odd, (k)}(z_0, n) g_2^{odd}(x_0 - z_0, r - n) \right] dn \right. \\
&\quad \left. + \underline{\omega} \int_{-\infty}^{+\infty} \left[f_2^{odd}(z_0, n) g_1^{even}(x_0 - z_0, r - n) \right. \right. \\
&\quad \left. \left. - \sum_{k=0}^m (-1)^k \left(\frac{2k}{m} - 1 \right) f_1^{even, (k)}(z_0, n) g_2^{odd}(x_0 - z_0, r - n) \right] dn \right\} dz_0.
\end{aligned}$$

We can thus end this subsection with the following definition and theorem.

Definition 4.5 (Mustard convolution). *The Mustard convolution \star_S corresponding to the slice Fourier transform is defined as*

$$\begin{aligned}
& (f \star_S g)(\mathbf{x}) \\
&= \frac{-i}{4\pi c} \left\{ f_1^{even} \star g_1^{even}(x_0, r) + \sum_{k=0}^m (-1)^k \left(\frac{2k}{m} - 1 \right) f_2^{odd, (k)} \star g_2^{odd}(x_0, r) \right. \\
&\quad \left. + \underline{\omega} \left(f_2^{odd} \star g_1^{even}(x_0, r) - \sum_{k=0}^m (-1)^k \left(\frac{2k}{m} - 1 \right) f_1^{even, (k)} \star g_2^{odd}(x_0, r) \right) \right\},
\end{aligned}$$

where $f, g \in \mathcal{V}$ were written as $f(\mathbf{x}) = f_1(x_0, r) + \underline{\omega}f_2(x_0, r)$ and $g(\mathbf{x}) = g_1(x_0, r) + \underline{\omega}g_2(x_0, r)$ and the superscripts even and odd denote the respective extensions in the second argument. The symbol \star denotes the classical two-dimensional convolution given by

$$(f \star g)(x_1, x_2) = \int_{\mathbb{R}} \int_{\mathbb{R}} f(x_1 - y_1, x_2 - y_2) g(y_1, y_2) dy_1 dy_2.$$

Theorem 4.6. *Under the action of the slice Fourier transform, the Mustard convolution \star_S obeys the classical convolution property*

$$\mathcal{F}_S(f \star_S g) = \mathcal{F}_S(f)\mathcal{F}_S(g).$$

Proof. Given that the above definition for the Mustard convolution \star_S is a mere elaboration of expression (4.13), taking the slice Fourier transform of the latter proves the theorem. \square

4.3 The generalised translation

Another way to generalise the convolution property is by adapting the translation operator t_y in the definition of the classical convolution stated in (1.1). Using $T_{\mathbf{y}}$ to denote this generalised translation operator, the Fubini theorem requires

$$\mathcal{F}_S \left(\int_{\mathbb{R}^{m+1}} T_{\mathbf{y}}(f)(\mathbf{x})g(\mathbf{y})d\mathbf{y} \right) (\mathbf{z}) = \int_{\mathbb{R}^{m+1}} \mathcal{F}_S(T_{\mathbf{y}}(f))(\mathbf{z}) g(\mathbf{y}) d\mathbf{y}$$

to be equal to

$$(\mathcal{F}_S(f)\mathcal{F}_S(g))(\mathbf{z}) = \int_{\mathbb{R}^{m+1}} \mathcal{F}_S(f)(\mathbf{z}) \mathcal{K}^M(\mathbf{y}, \mathbf{z}) g(\mathbf{y}) d\mathbf{y},$$

which is achieved when the integrands are equal. The generalised translation operator $T_{\mathbf{y}}$ is thus defined as

$$T_{\mathbf{y}}f(\mathbf{x}) = \mathcal{F}_S^{-1}(\mathcal{F}_S(f)(\mathbf{z})\mathcal{K}^M(\mathbf{y}, \mathbf{z})), \quad (4.15)$$

so its slice Fourier transform generates an extra factor $\mathcal{K}^M(\mathbf{y}, \mathbf{z})$ to the right of the slice Fourier transform of the function it is acting upon.

Remark 4.7. *Note the crucial difference between the above defined generalised translation $T_{\mathbf{y}}$ and the translation operator studied in [13]. As the slice Fourier transform of $T_{\mathbf{y}}(f)$ adds the factor \mathcal{K}^M to the right of $\mathcal{F}_S(f)$, the kernel expression \mathcal{K}^M can subsequently be right-multiplied with a neighbouring factor.*

Denoting the kernel function of the inverse slice Fourier transform as $\mathcal{K}^{M,-1}$, the following expression is shown to equal the inverse Fourier transform of the product of the slice Fourier transforms of f and g :

$$\begin{aligned} & \int_{\mathbb{R}^{m+1}} T_{\mathbf{y}}(f)(\mathbf{x})g(\mathbf{y})d\mathbf{y} \\ &= \int_{\mathbb{R}^{m+1}} \left(\int_{\mathbb{R}^{m+1}} \mathcal{K}^{M,-1}(\mathbf{z}, \mathbf{x})\mathcal{F}_S(f)(\mathbf{z})\mathcal{K}^M(\mathbf{y}, \mathbf{z})g(\mathbf{y}) dz_0 d\mathbf{n} d\sigma_{\underline{z}} \right) dy_0 dg d\sigma_{\underline{y}} \\ &= \int_{\mathbb{R}^{m+1}} \mathcal{K}^{M,-1}(\mathbf{z}, \mathbf{x})\mathcal{F}_S(f)(\mathbf{z}) \left(\int_{\mathbb{R}^{m+1}} \mathcal{K}^M(\mathbf{y}, \mathbf{z}) g(\mathbf{y}) dy_0 dg d\sigma_{\underline{y}} \right) dz_0 d\mathbf{n} d\sigma_{\underline{z}} \\ &= \int_{\mathbb{R}^{m+1}} \mathcal{K}^{M,-1}(\mathbf{z}, \mathbf{x}) (\mathcal{F}_S(f)(\mathbf{z})\mathcal{F}_S(g)(\mathbf{z})) dz_0 d\mathbf{n} d\sigma_{\underline{z}} \\ &= \mathcal{F}_S^{-1}(\mathcal{F}_S(f)\mathcal{F}_S(g))(\mathbf{x}). \end{aligned}$$

As the above defined generalised translation operator $T_{\mathbf{y}}$ shows the desired behaviour, we will now calculate its explicit expression. Substituting all known

kernel functions and writing $\mathbf{x} = x_0 + r\underline{\omega}$, $\mathbf{y} = y_0 + g\underline{\eta}$, $\mathbf{z} = z_0 + h\underline{\zeta}$ and $\mathbf{u} = u_0 + n\underline{\nu}$ yields

$$\begin{aligned}
T_{\mathbf{y}}(f)(\mathbf{x}) &= -i \left(\frac{\Gamma\left(\frac{m}{2}\right)}{8c\pi^{m/2+1}} \right)^3 \int_{\mathbb{R}^{m+1}} \int_{\mathbb{R}^{m+1}} \\
&\quad \left[(1 + \underline{\omega}\underline{\zeta})e^{+\frac{i}{2c}(x_0z_0-rn)} + (1 - \underline{\omega}\underline{\zeta})e^{+\frac{i}{2c}(x_0z_0+rn)} \right] \times \\
&\quad \left[(1 + \underline{\zeta}\underline{\nu})e^{-\frac{i}{2c}(z_0u_0-nh)} + (1 - \underline{\zeta}\underline{\nu})e^{-\frac{i}{2c}(z_0u_0+nh)} \right] f(\mathbf{u}) \times \\
&\quad \left[(1 + \underline{\zeta}\underline{\eta})e^{-\frac{i}{2c}(y_0z_0-gn)} + (1 - \underline{\zeta}\underline{\eta})e^{-\frac{i}{2c}(y_0z_0+gn)} \right] dz_0 \, dn \, d\sigma_{\underline{z}} \, du_0 \, dh \, d\sigma_{\underline{u}}.
\end{aligned}$$

As usual, integrating over z_0 formally gives a delta distribution. Rearranging the remaining terms we get

$$\begin{aligned}
T_{\mathbf{y}}(f)(\mathbf{x}) &= -i \left(\frac{\Gamma\left(\frac{m}{2}\right)}{8c\pi^{m/2+1}} \right)^3 \int_0^{+\infty} \int_{\mathbb{S}^{m-1}} \int_{\mathbb{R}^{m+1}} 4\pi c \, \delta(x_0 - y_0 - u_0) \times \\
&\quad \left\{ \left[(1 + \underline{\omega}\underline{\zeta} + \underline{\zeta}\underline{\nu} - \underline{\omega}\underline{\nu})e^{-\frac{i}{2c}(r-h-g)n} + (1 + \underline{\omega}\underline{\zeta} - \underline{\zeta}\underline{\nu} + \underline{\omega}\underline{\nu})e^{-\frac{i}{2c}(r+h-g)n} \right. \right. \\
&\quad \left. \left. + (1 - \underline{\omega}\underline{\zeta} + \underline{\zeta}\underline{\nu} + \underline{\omega}\underline{\nu})e^{+\frac{i}{2c}(r+h+g)n} + (1 - \underline{\omega}\underline{\zeta} - \underline{\zeta}\underline{\nu} - \underline{\omega}\underline{\nu})e^{+\frac{i}{2c}(r-h+g)n} \right] \right. \\
&\quad \left. \times f(\mathbf{u})(1 + \underline{\zeta}\underline{\eta}) \right. \\
&\quad \left. + \left[(1 + \underline{\omega}\underline{\zeta} + \underline{\zeta}\underline{\nu} - \underline{\omega}\underline{\nu})e^{-\frac{i}{2c}(r-h+g)n} + (1 + \underline{\omega}\underline{\zeta} - \underline{\zeta}\underline{\nu} + \underline{\omega}\underline{\nu})e^{-\frac{i}{2c}(r+h+g)n} \right. \right. \\
&\quad \left. \left. + (1 - \underline{\omega}\underline{\zeta} + \underline{\zeta}\underline{\nu} + \underline{\omega}\underline{\nu})e^{+\frac{i}{2c}(r+h-g)n} + (1 - \underline{\omega}\underline{\zeta} - \underline{\zeta}\underline{\nu} - \underline{\omega}\underline{\nu})e^{+\frac{i}{2c}(r-h-g)n} \right] \right. \\
&\quad \left. \times f(\mathbf{u})(1 - \underline{\zeta}\underline{\eta}) \right\} dn \, d\sigma_{\underline{z}} \, du_0 \, dh \, d\sigma_{\underline{u}}.
\end{aligned}$$

Because of symmetry reasons, all terms containing $\underline{\zeta}$ will disappear after integration over \mathbb{S}^{m-1} . As before, only functions in the span of $\{\psi_{j,k}\}$ are considered so f can be written as $f(\mathbf{u}) = f_1(u_0, h) + \underline{\nu}f_2(u_0, h)$. Performing the same manipulations as before (see section 4.2) on the integrals over $[0, +\infty[$ and integrating

over $n, \underline{\zeta}$ and $\underline{\nu}$ yields

$$\begin{aligned}
T_{\mathbf{y}}(f)(\mathbf{x}) &= \frac{-i\Gamma\left(\frac{m}{2}\right)}{8c\pi^{m/2+1}} \int_{-\infty}^{+\infty} \int_0^{+\infty} \delta(x_0 - y_0 - u_0) \times \\
&\quad \left\{ \left[f(u_0, h, \underline{\omega}) + \underline{\omega} \sum_{k=0}^m (-1)^k \left(\frac{2k}{m} - 1\right) \left(f_1^{(k)} + \underline{\omega} f_2^{(k)}\right)(u_0, h) \underline{\eta} \right] \right. \\
&\quad \times \delta(r - h - g) \\
&\quad + \left[f(u_0, h, \underline{\omega}) - \underline{\omega} \sum_{k=0}^m (-1)^k \left(\frac{2k}{m} - 1\right) \left(f_1^{(k)} + \underline{\omega} f_2^{(k)}\right)(u_0, h) \underline{\eta} \right] \\
&\quad \times \delta(r - h + g) \\
&\quad + \left[f(u_0, h, -\underline{\omega}) + \underline{\omega} \sum_{k=0}^m (-1)^k \left(\frac{2k}{m} - 1\right) \left(f_1^{(k)} - \underline{\omega} f_2^{(k)}\right)(u_0, h) \underline{\eta} \right] \\
&\quad \left. \times \delta(r + h - g) \right\} du_0 dh.
\end{aligned}$$

Performing the last two integrations, we finally obtain the explicit action of the generalised translation $T_{\mathbf{y}}$ on a function f :

$$\begin{aligned}
T_{\mathbf{y}}(f)(\mathbf{x}) &= \frac{-i\Gamma\left(\frac{m}{2}\right)}{8c\pi^{m/2+1}} \times \\
&\quad \left\{ f(x_0 - y_0, r - g, \underline{\omega}) + \underline{\omega} \sum_{k=0}^m (-1)^k \left(\frac{2k}{m} - 1\right) f^{(k)}(x_0 - y_0, r - g, \underline{\omega}) \underline{\eta} \right. \\
&\quad + f(x_0 - y_0, r + g, \underline{\omega}) - \underline{\omega} \sum_{k=0}^m (-1)^k \left(\frac{2k}{m} - 1\right) f^{(k)}(x_0 - y_0, r + g, \underline{\omega}) \underline{\eta} \\
&\quad \left. + f(x_0 - y_0, g - r, -\underline{\omega}) + \underline{\omega} \sum_{k=0}^m (-1)^k \left(\frac{2k}{m} - 1\right) f^{(k)}(x_0 - y_0, g - r, -\underline{\omega}) \underline{\eta} \right\}
\end{aligned}$$

where terms in f and $f^{(k)}(\mathbf{x}) = f_1^{(k)}(x_0, r) + \underline{\omega} f_2^{(k)}(x_0, r)$ are only included when

their second arguments are positive. Making the $\underline{\omega}$ -dependence explicit yields

$$\begin{aligned}
T_{\mathbf{y}}(f)(\mathbf{x}) &= \frac{-i\Gamma\left(\frac{m}{2}\right)}{8c\pi^{m/2+1}} \times \\
&\left\{ \left[f_1(x_0 - y_0, r - g) - \sum_{k=0}^m (-1)^k \left(\frac{2k}{m} - 1\right) f_2^{(k)}(x_0 - y_0, r - g)\underline{\eta} \right] \right. \\
&+ \left[f_1(x_0 - y_0, r + g) + \sum_{k=0}^m (-1)^k \left(\frac{2k}{m} - 1\right) f_2^{(k)}(x_0 - y_0, r + g)\underline{\eta} \right] \\
&+ \left[f_1(x_0 - y_0, g - r) + \sum_{k=0}^m (-1)^k \left(\frac{2k}{m} - 1\right) f_2^{(k)}(x_0 - y_0, g - r)\underline{\eta} \right] \\
&+ \underline{\omega} \left[f_2(x_0 - y_0, r - g) + \sum_{k=0}^m (-1)^k \left(\frac{2k}{m} - 1\right) f_1^{(k)}(x_0 - y_0, r - g)\underline{\eta} \right] \\
&+ \underline{\omega} \left[f_2(x_0 - y_0, r + g) - \sum_{k=0}^m (-1)^k \left(\frac{2k}{m} - 1\right) f_1^{(k)}(x_0 - y_0, r + g)\underline{\eta} \right] \\
&+ \underline{\omega} \left[-f_2(x_0 - y_0, g - r) + \sum_{k=0}^m (-1)^k \left(\frac{2k}{m} - 1\right) f_1^{(k)}(x_0 - y_0, g - r)\underline{\eta} \right] \left. \right\}.
\end{aligned}$$

where again terms in f_1 and f_2 are only included when their second argument is positive (so depending on $r \in [0, +\infty[$ being smaller or bigger than $g \in [0, +\infty[$).

4.4 Connection between both approaches

Proposition 4.8. *The Mustard convolution \star_S and the generalised translation $T_{\mathbf{y}}$ satisfy*

$$(f \star_S g)(\mathbf{x}) = \int_{\mathbb{R}^{m+1}} T_{\mathbf{y}}(f)(\mathbf{x})g(\mathbf{y}) \, d\mathbf{y}. \quad (4.16)$$

Proof. Taking the slice Fourier transform of both sides of (4.16) gives the same expression: $\mathcal{F}(f)\mathcal{F}(g)$. In section 3.5 it was shown how the slice Fourier transform (and therefore its inverse as well) can be calculated using the classical two-dimensional Fourier transform. Given that the latter is a bijective transform, the above equality follows immediately. \square

We can also verify this proposition explicitly:

$$\begin{aligned}
\int_{\mathbb{R}^{m+1}} T_{\mathbf{y}}(f)(\mathbf{x})g(\mathbf{y})d\mathbf{y} &= \frac{-i\Gamma\left(\frac{m}{2}\right)}{8c\pi^{m/2+1}} \int_{-\infty}^{+\infty} \int_0^{+\infty} \int_{\mathbb{S}^{m-1}} \\
&\left\{ f(x_0 - y_0, r - g, \underline{\omega}) + \underline{\omega} \sum_{k=0}^m (-1)^k \left(\frac{2k}{m} - 1\right) f^{(k)}(x_0 - y_0, r - g, \underline{\omega}) \underline{\eta} \right. \\
&+ f(x_0 - y_0, r + g, \underline{\omega}) - \underline{\omega} \sum_{k=0}^m (-1)^k \left(\frac{2k}{m} - 1\right) f^{(k)}(x_0 - y_0, r + g, \underline{\omega}) \underline{\eta} \\
&+ \left. f(x_0 - y_0, g - r, -\underline{\omega}) + \underline{\omega} \sum_{k=0}^m (-1)^k \left(\frac{2k}{m} - 1\right) f^{(k)}(x_0 - y_0, g - r, -\underline{\omega}) \underline{\eta} \right\} \\
&\times g(\mathbf{y}) dy_0 dg d\sigma_{\underline{y}}
\end{aligned}$$

where the different terms are only taken into account when the second argument of f or $f^{(k)}$ is positive. Writing $g(\mathbf{y}) = g_1(y_0, g) + \underline{\eta}g_2(y_0, g)$ and performing the integration over $\underline{\eta}$, we obtain

$$\begin{aligned}
\int_{\mathbb{R}^{m+1}} T_{\mathbf{y}}(f)(\mathbf{x})g(\mathbf{y})d\mathbf{y} &= \frac{-i}{4\pi c} \int_{-\infty}^{+\infty} \int_0^{+\infty} \int_{\mathbb{S}^{m-1}} \\
&\left\{ f(x_0 - y_0, r - g, \underline{\omega})g_1(y_0, g) - \underline{\omega} \sum_{k=0}^m (-1)^k \left(\frac{2k}{m} - 1\right) f^{(k)}(x_0 - y_0, r - g, \underline{\omega})g_2(y_0, g) \right. \\
&+ f(x_0 - y_0, r + g, \underline{\omega})g_1(y_0, g) + \underline{\omega} \sum_{k=0}^m (-1)^k \left(\frac{2k}{m} - 1\right) f^{(k)}(x_0 - y_0, r + g, \underline{\omega})g_2(y_0, g) \\
&+ \left. f(x_0 - y_0, g - r, -\underline{\omega})g_1(y_0, g) - \underline{\omega} \sum_{k=0}^m (-1)^k \left(\frac{2k}{m} - 1\right) f^{(k)}(x_0 - y_0, g - r, -\underline{\omega})g_2(y_0, g) \right\} \\
&dy_0 dg.
\end{aligned}$$

Performing the substitutions $z_0 = x_0 - y_0$ and $n = \pm r \pm g$, where the signs depend on the integrand, we indeed retrieve the expression for $\mathcal{F}_S^{-1}(\mathcal{F}_S(f)\mathcal{F}_S(g))$ from (4.14).

5 Conclusion

The article at hand introduces the classical Fourier transform to the slice Clifford setting. Using the Clifford-Hermite functions defined in [4] as a basis of eigenfunctions, corresponding eigenvalues are obtained from the scalar differential equation they satisfy. The combination of these data allowed for a formal definition of the slice Fourier transform. Its explicit expression was obtained using the Mehler and Hille-Hardy formulas. The closed integral expression evoked a

closer study of its main characteristics including its differential and symmetry properties and the construction of the inverse transform. Also the computational load of the slice Fourier transform has been briefly addressed.

The second part of the paper investigates how to construct an appropriate counterpart of the classical convolution. For its slice Fourier transform to equal the product of the slice Fourier transforms of the functions it is acting upon, two possibilities are presented. First the Mustard convolution is studied and its explicit expression is obtained. In a second approach, a generalised translation operator is introduced to the integrand of the classical convolution expression. Given that both approaches show the same extension of the behaviour of the classical convolution under the classical Fourier transform, the paper finishes by highlighting their connection.

References

- [1] F. Brackx, N. De Schepper, and F. Sommen. The Clifford-Fourier transform. *J. Fourier Anal. Appl.* **11** (2005), 669–681.
- [2] F. Brackx, N. De Schepper, and F. Sommen. The two-dimensional Clifford-Fourier transform. *J. Math. Imaging Vision* **26** (2006), 5–18.
- [3] R. Bujack, H. De Bie, N. De Schepper, and G. Scheuermann. Convolution products for hypercomplex fourier transforms. *J. Math. Imaging Vision* **48** (2014), 606–624.
- [4] L. Cnudde, H. De Bie, and G. Ren. Algebraic approach to slice monogenic functions. *Complex Anal. Oper. Theory* **9** (2015), 1065–1087.
- [5] F. Colombo, J. O. Gonzàles Cervantes, and I. Sabadini. The Bergman-Sce transform for slice monogenic functions. *Math. Methods Appl. Sci.* **34** (2011), 1896–1909.
- [6] F. Colombo, J. O. González-Cervantes, and I. Sabadini. A nonconstant coefficients differential operator associated to slice monogenic functions. *Trans. Amer. Math. Soc.* **365** (2013), 303–318.
- [7] F. Colombo, R. Lávička, I. Sabadini, and V. Souček. The Radon transform between monogenic and generalized slice monogenic functions. *Math. Ann.* **363** (2015), 733–752.
- [8] F. Colombo, I. Sabadini, and D. C. Struppa. Slice monogenic functions. *Israel J. Math.* **171** (2009), 385–403.
- [9] F. Colombo, I. Sabadini, and D. C. Struppa. *Noncommutative functional calculus*, volume 289 of *Progress in Mathematics*. Birkhäuser/Springer Basel AG, Basel (2011). Theory and applications of slice hyperholomorphic functions.

- [10] H. De Bie. Clifford algebras, Fourier transforms, and quantum mechanics. *Math. Methods Appl. Sci.* **35** (2012), 2198–2228.
- [11] H. De Bie, N. De Schepper, and F. Sommen. The class of Clifford-Fourier transforms. *J. Fourier Anal. Appl.* **17** (2011), 1198–1231.
- [12] H. De Bie, B. Ørsted, P. Somberg, and V. Souček. Dunkl operators and a family of realizations of $\mathfrak{osp}(1|2)$. *Trans. Amer. Math. Soc.* **364** (2012), 3875–3902.
- [13] H. De Bie and Y. Xu. On the Clifford-Fourier transform. *Int. Math. Res. Not. IMRN* (2011), no. 22, pages 5123–5163.
- [14] P. Denis, P. Carre, and C. Fernandez-Maloigne. Spatial and spectral quaternionic approaches for colour images. *Computer Vision and Image Understanding* **107** (2007), 74 – 87.
- [15] T. A. Ell and S. J. Sangwine. Hypercomplex fourier transforms of color images. *IEEE Trans. Image Process.* **16** (2007), 22–35.
- [16] G. Gentili and D. C. Struppa. A new theory of regular functions of a quaternionic variable. *Adv. Math.* **216** (2007), 279 – 301.
- [17] R. Ghiloni and A. Perotti. Slice regular functions on real alternative algebras. *Adv. Math.* **226** (2011), 1662–1691.
- [18] C. Guo and L. Zhang. A novel multiresolution spatiotemporal saliency detection model and its applications in image and video compression. *IEEE Trans. Image Process.* **19** (2010), 185–198.
- [19] R. Howe. The oscillator semigroup. The mathematical heritage of Hermann Weyl, Proc. Symp., Durham/NC 1987, Proc. Symp. Pure Math. 48, 61-132 (1988). (1988).
- [20] C. E. Moxey, S. J. Sangwine, and T. A. Ell. Hypercomplex correlation techniques for vector images. *IEEE Trans. Signal Process.* **51** (2003), 1941–1953.
- [21] D. Mustard. Fractional convolution. *J. Aust. Math. Soc. Series B. Applied Mathematics* **40** (1998), 257–265.
- [22] G. Szegő. *Orthogonal polynomials*. American Mathematical Society, Providence, R.I., fourth edition (1975). American Mathematical Society, Colloquium Publications, Vol. XXIII.
- [23] G. N. Watson. *A Treatise on the Theory of Bessel Functions*. Cambridge University Press, Cambridge, England; The Macmillan Company, New York (1944).

6 Acknowledgements

L. Cnudde and H. De Bie are supported by the UGent BOF starting grant 01N01513.