



គម្រោងថ្នាក់តំបន់ស្តីពី ការវិវឌ្ឍន៍ជីវិតតំបន់កម្ពុជា និងអាស៊ីអាគ្នេយ៍
The Regional Fisheries Livelihoods Programme for South and Southeast Asia (RFLP) - Cambodia

CATCH MONITORING



2012



REGIONAL FISHERIES LIVELIHOODS PROGRAMME FOR SOUTH AND SOUTHEAST ASIA (RFLP)

Activity 1.5 (2011):

Systems and procedures for participatory monitoring of management measures developed, introduced and implemented – catch monitoring

**For the Regional Fisheries Livelihoods Programme for the South and Southeast Asia
Cambodia Component**

Prepared by

Peng Bun NGOR and Kong HENG
Monitoring and Evaluation Division
Planning, Finance and International Cooperation Department
Fisheries Administration

January 2012

Acknowledgements

The authors are very grateful to His Excellency Dr. Nao Thuok, Director General of the Fisheries Administration (FiA) and Mrs. Kaing Khim, Deputy Director General of the Fisheries Administration, and Mr. Thor Sensereivorth, Acting Director of Planning, Finance and International Cooperation Department of the Fisheries Administration for their support of this study.

Our sincere thanks are also given to Mr. Hun Kimtek, Fisheries Officer who kindly assisted in data entry and checking for errors before final analysis for reporting could be made. The team would like to express our thanks to Mr. Ngor Pengby, database manager, for his kind assistance in commenting on the database design, creating queries to check for errors and data analysis for final reporting.

We would like to profoundly thank FAO Cambodia - Regional Fisheries Livelihoods Programme for South and Southeast Asia (RFLP), which is funded by Spain for its financial support. Without this support, this study would have been impossible. Sincere thanks are also given to the RFLP regional and national staff and particularly Mr. Jose Parajua, RFLP Regional Programme Manager, Mr. Don Griffiths, RFLP Chief Technical Advisor, Mr. Yos Chanthana, Fisheries Co-management Consultant and Mr. Thay Somony, RFLP Cambodia National Project Coordinator for their comments on the report and support during the study period.

We would also like to thank the Fisheries Administration Cantonments of Kampot, Kampong Som and Koh Kong for their support. Specifically, we thank Mr. Ly Seyha (Deputy Chief of Prek Chak Division), Chak Sinead (Deputy Chief of Kep Division), Mr. Phan Sopha (Kampot FiA fisheries officer), Mr. Pheun Phalla (Chief of Kep FiA Division), Em Phea (Deputy Chief of Prey Nub FiA Division), Mr. Chan Sophy (Chief of Sre Ambel FiA Unit) and Mr. Nen Chanroeun (Deputy Head of Kampong Som FiA Cantonment) for their kind assistance and support during this field study.

Finally the authors would like to thank all the 26 fishers from the five selected Community Fisheries (CFi's) who kindly collected and provided the data on which this report is based.

Table of contents

Acknowledgements	i
Table of contents	ii
Table of Tables	ivv
Table of Figures	v
Summary	1
1. Background.....	3
2. Objective of the study.....	3
3. Scope of the study.....	4
4. Methods.....	4
4.1 Study areas	4
4.2 Fishing geared monitored.....	4
4.3 Survey design	5
5. Results and discussion.....	7
5.1 Fisher profiles.....	7
5.2 Average catch and value per fishing trip	7
5.3. Monthly catch of top ten fishers.....	10
5.4. Monthly value of top ten fishers.....	11
5.5. Key important species caught by fishers	11
5.5.1 Fisher ID 1: fish gillnet.....	12
5.5.2 Fisher ID 2: Crab gillnet.....	12
5.5.3 Fisher ID 3: Crab trap	13
5.5.4 Fisher ID 4: Hand pushnet.....	13
5.5.5 Fisher ID 5: Wedge-shaped scoop basket.....	14
5.5.6 Fisher ID 6: Fish gillnet.....	14
5.5.7 Fisher ID 7: Crab trap	15
5.5.8 Fisher ID 8: Hand pushnet.....	15
5.5.9 Fisher ID 9: Karav (Threadfin) gillnet	16
5.5.10 Fisher ID 10: Kabork (Mullet) gillnet	16
5.5.11 Fisher ID 11: Fish trap.....	17
5.5.12 Fisher ID 12: Squid hook and line.....	18
5.5.13 Fisher ID 13: Squid hook and line.....	18
5.5.14 Fisher ID 14: Fish trap.....	18
5.5.15 Fisher ID 15: Squid hook and line.....	19
5.5.16 Fisher ID 16: Gastropod long line	19

5.5.17 Fisher ID 17: Crab trap	20
5.5.18 Fisher ID 18: Crab trap	20
5.5.19 Fisher ID 19: Spade and hand collection of cockles	21
5.5.20 Fisher ID 20: Hand collection plus spade (targeting crabs)	21
5.5.21 Fisher ID 21: Hand collection plus spade (targeting crabs)	22
5.5.22 Fisher ID 22: Crab trap	22
5.5.23 Fisher ID 23: Fish gillnet.....	23
5.5.24 Fisher ID 24: Crab trap	23
5.5.25 Fisher ID 25: Crab gillnet.....	24
5.5.26 Fisher ID 26: Hand collection of blood cockles	24
5.6. Monthly average prices at landing sites by species by fisher	25
5.7. Frequencies versus average maximum length by fisher by species	25
6. Conclusions	25
7. Suggestions for further studies	27
Annexes	29
Annex 1: Fisher logbook	29
Annex 2: Name of fishers and their fishing gears used in five target community fisheries	31
Annex 3: Average weight (kg) and value (Riel) per fisher per trip (ranking according to value at the landing site).....	33
Annex 4: Monthly catch (kg) and value (Riel) by fisher (Ranking according to value) .	34
Annex 5: Monthly catch (kg) by species and by fishers, ranking according to total catch over the period from April to November 2011	35
Annex 6: Monthly landing value (Riel) by species and by fishers over the period from April to November 2011 (ranked according to total value)	51
Annex 7: Average price (Riel/kg) of catch by species by fisher (April-November 2011)	66
Annex 8: Average maximum length (cm) versus frequencies by fisher measured over the period from April to November 2011	80

Table of Tables

Table 1: List of fishing gears monitored in this study.....	5
Table 2: Average catch (kg) and value (Riel) of crab traps per fishing trip (hour) from April to November 2011, ranked according to the total catch.....	8

Table of Figures

Figure 1: Map showing the Community Fisheries covered in this study	4
Figure 2: Average catch per fishing trip by fisher (April-November 2011)	9
Figure 3: Average value (value at fish landing site) obtained per fishing trip by fisher....	9
Figure 4: Monthly catch of top ten fishers ranking according to the total catch over the study period from April to November 2011.	11
Figure 5: Monthly value of top ten fishers ranking according to the total value over the study period from April to November 2011.	11
Figure 6: Fisher ID 1: Catch (kg) (April-November 2011).....	12
Figure 7: Fisher ID 2: Catch (kg) (April-November 2011).....	13
Figure 8: Fisher ID 3: Catch (kg) (April-November 2011).....	13
Figure 9: Fisher ID 4: Catch (kg) (April-November 2011).....	14
Figure 10: Fisher ID 5: Catch (kg) (April-November 2011).....	14
Figure 11: Fisher ID 6: Catch (kg) (April-November 2011).....	15
Figure 12: Fisher ID 7: Catch (kg) (April-November 2011).....	15
Figure 13: Fisher ID 8: Catch (kg) (April-November 2011).....	16
Figure 14: Fisher ID 9: Catch (kg) (April-November 2011).....	16
Figure 15: Fisher ID 10: Catch (kg) (April-November 2011)	17
Figure 16: Fisher ID 11: Catch (kg) (April-November 2011)	17
Figure 17: Fisher ID 12: Catch (kg) (April-November 2011)	18
Figure 18: Fisher ID 13: Catch (kg) (April-November 2011)	18
Figure 19: Fisher ID 14: Catch (kg) (April-November 2011)	19
Figure 20: Fisher ID 15: Catch (kg) (April-November 2011)	19
Figure 21: Fisher ID 16: Catch (kg) (April-November 2011)	20
Figure 22: Fisher ID 17: Catch (kg) (April-November 2011)	20
Figure 23: Fisher ID 18: Catch (kg) (April-November 2011)	21
Figure 24: Fisher ID 19: Catch (kg) (April-November 2011)	21
Figure 25: Fisher ID 20: Catch (kg) (April-November 2011)	22
Figure 26: Fisher ID 21: Catch (kg) (April-November 2011)	22
Figure 27: Fisher ID 22: Catch (kg) (April-November 2011)	23
Figure 28: Fisher ID 23: Catch (kg) (April-November 2011)	23
Figure 29: Fisher ID 24: Catch (kg) (April-November 2011)	24
Figure 30: Fisher ID 25: Catch (kg) (April-November 2011)	24
Figure 31: Fisher ID 29: Catch (kg) (April-November 2011)	25

Summary

Coastal fisheries are very essential for supporting the livelihoods of many rural poor in the coastal areas, particularly coastal community fisheries members. They serve as sources of food, employment and income generation for many coastal families. “Coastal Community Fisheries Catch Monitoring” Project which was conducted from April to November 2011 provides some data which indicates the importance of small-scale fisheries. The project was financially supported by the Regional Fisheries Livelihoods Programme, Cambodian component (RFLP/CMB) and was activity 1.5 of the approved RFLP CMB 2011 activity work plan and budget.

For this catch monitoring study, 26 small-scale subsistence fishers, including 05 women, from five community fisheries (CFi's) in the RFLP CMB area of geographic coverage from the four coastal provinces of Cambodia were selected and following appropriate training collected specific catch data and recorded it in fisher's logbook on a daily basis for the purpose of getting a better understanding of catch per unit of effort (CEPU), the health of inshore fish stock and the contribution of aquatic products to small-scale fisher households along the coast of Cambodia. The key data items recorded included total catch weight, catch weight by species, total catch sale price, fish price of the main species and total lengths of some key species. The study involved designing logbooks, training the selected 26 fishers as data collectors on data collection methods, collecting data from all the selected fishers, designing a database and entering all the collected data into the database, checking for errors and analyzing the collected data for final reporting and preparing report.

The main findings from this eight month study are:

- More than 100 different marine species were recorded by the 26 participating fishers involved in the study. Species caught varied according to different fishing gear types. Data on catch, value, prices as well as maximum length of all species are made available in annexes of this report.
- Some fishing gears were highly selective. These included crab traps, gastropod (octopus) long line, and squid hook and line, as well as hand collection of blood cockles.
- A few fishing gears such as crab gillnet, fish traps and hand collection with supplementary equipment are fairly selective, capturing about 10 different aquatic species and usually have only limited by-catch or non-target species low value species and juveniles of commercial species.
- Other fishing gears such as different types of fish gillnet and hand pushnet capture up to more than 30 different aquatic species and usually have a high proportion of by-catch.
- Overall, fishers spent an average of 8.9 hours actually fishing, with some variations, per fishing trip. Average catch and value at landing sites per fishing trip were around 18 kg of aquatic product with an average value of Riel 80,000 (approximately US\$¹ 20), though there was considerable variation over time in both catch quantity and value.

¹US\$ 1 = approximately Riels 4,000.

- Fish trap fishers left their traps in the water an average of about 40 hours per fishing trip. Fish traps were only operated in one of the five sampled CFI's at Koh Rounsanlem and targeted high value species such as groupers. The catch per fishing trip (10-20 kg) was relatively low compared to other fishers. However, the monetary value from selling the catch was comparatively higher at between Riel 200,000 and 300,000 (US\$ 50 – 75) per fishing trip.
- Gillnetfishers in Trapiang Ropov CFI had their nets in the water on average for 07 hours per fishing trip and caught between 100 and 130 kg of aquatic species with the total value of ranging typically from Riel 150,000 - 400,000.
- Crab trap fishers had around 10 actual fishing hours, with an average of catch of around 8 kg which is equivalent to a total value of about Riel 100,000 at fish landing sites (first sale value).
- A gastropod (octopus) long line fisher (1,000 gastropod shells) in Koh Rounsanlem had, on average, around 2 actual fishing hours, capturing around 10 kg of *Sepiolo sp.* (*Meuk dong*) with a total value of around 110,000 Riel.
- Hand pushnet are practiced especially in Kep and Kampot Provinces. Hand pushnet fishers had an average of around 8 actual fishing hours with a catch varied between 2 and 5 kilogram and value of around Riel 11,000 to Riel 20,000 per fishing trip.
- Hand collection fishers with supplementary equipment such as spade, sliding board, and basket are the least efficient fishers, having the least catch. They spent around 6 hours per trip in search for marine aquatic animals. On average, they could collect 3-4 kg of cockles, the cheapest species, with a total of around Riel 10,000.

For further study to better understand catch per unit of effort, it is suggested that:

- A standardized fishing gear should be applied for a standard sampling in order to understand CPUE which is based on a gear type. Moreover, in order to generalize the findings from the study, sampling sites should be geographically represented the coastal areas and different fish habitat types and ecosystems. For instance, fishers expect different fish species in mangrove forests, in sea grass and in coral reef areas.
- Biological, socio-economical and environmental parameters should be properly included in order to obtain a broad view point about the size of the catch, value and the species occurrence as well as their explanatory factors from socio-economic and environmental view points.
- Findings from this study should be presented to the 5 piloted CFI members especially those directly involved in recording and reporting the data in order to give them the opportunity to provide feedback and validate the findings as well as to make use these data to plan, monitor and understand the catch status in their CFI waters.
- Last but not least, a field guide containing a majority of marine species with proper local Khmer names, English names, scientific names and a brief description on the identification of the species should be made available as this type of field guide is very useful for this type of research. Through literature, marine field guide for Cambodia is available for only key main marine species. A large number of marine species has not been properly documented.

1. Background

As the world's marine fish stocks have become more heavily fished and particularly inshore small-scale fisheries, so fisheries data collection at the community fisheries level has become increasingly important, especially for understanding fish stock health, species abundance and diversity as well as the current catch rate being enjoyed by fishers. Community-based catch monitoring, especially with regards to the monitoring of Catch per Unit of Effort (CPUE), fish catch composition and prices of fish at landing sites, are also essential for the establishment of community fisheries baseline data. More specifically, community-based monitoring will provide information on current catches to fishers and local authorities, allowing them to understand changes in catch and species composition and prices as well as catch rates per fisher and per fishing gear for a particular time and place in order that actions or interventions can be designed or modified, using a factual basis. Apart from this, data and information generated by a community-based monitoring system will be of crucial importance for planning, monitoring and evaluation of community fisheries management and development which has been practiced by the Fisheries Administration (FiA) as part of its decentralized fisheries management during over the last ten years. Moreover, community-based fisheries monitoring may provide vital inputs for the formulation of a rural food security policy.

Therefore, the establishment of “systems and procedures for participatory monitoring of management measures” was one of the prioritized activities under Output 1 of the Regional Fisheries Livelihoods Programme RFLP/CMB Work Plan for implementation in 2011. The RFLP (GCP/RAS/237/SPA), which is funded by Spain and executed by the Food and Agriculture Organization of the United Nations in collaboration with the national agencies of fisheries has activities in Cambodia, Indonesia, the Philippines, Sri Lanka, Timor-Leste, and Viet Nam. The four year programme began on 1st September 2009 and will run until 31st August 2013.

More specifically, this activity contributes to achieving the RFLP intermediate outcome on “Co-management mechanisms for sustainable utilization of fishery resources” and the RFLP overall outcome which is ‘Strengthened capacity among participating small-scale fishing communities and their supporting institutions towards improved livelihoods and sustainable fisheries resources management’.

2. Objective of the study

The main objective of this activity was to establish a community-based catch monitoring programme and to collect baseline data on catch rate or catch per unit of effort (CEPU) of important fishing gears in five selected Community Fisheries currently being targeted by RFLP/CMB. The specific objectives of this activity were to:

- Set up a team of field data collectors in five community fisheries (CFis) to record details of their catch by species and effort by gears.
- Design a logbook for selected CFi fishers (26 fishers) to record details of their catch by species and effort by gears.
- Train (50) CFi members from the selected CFis under the RFLP/CAM target on how to use the logbooks for the recording of their daily catches.
- Design a database for storing and using the collected data;

- Technically support field data collection.
- Generate a catch monitoring report (with baseline data on catches from each selected CFI).

3. Scope of the study

The study targeted five (5) coastal community fisheries. The study period was undertaken over approximately eight month period between April and November 2011. Following logbook design, participant fisher selection and training actual field actual data collection commenced in week 4 of April and ran for just over seven months.

4. Methods

4.1 Study areas

The study area covered the four coastal provinces of Cambodia namely Kep, Kampot, Preah Sihanouk and Koh Kong Provinces where RFLP/CAM (2009-13) is conducting activities in collaboration with FiA. Five coastal Community Fisheries (CFI's), namely Okrasa in Kep Province, Trapiang Ropov in Kampot Province, Chumpou Khmao and Koh Rungsanlem in Preah Sihanouk Province and Koh Khyorang in Koh Kong Province were selected for this study, as being geographically representative of the coastal conditions and inshore fishing areas found along the 400 km coastline of Cambodia. The selected locations are shown in Figure 1 below.



Figure 1: Map showing the Community Fisheries covered in this study

4.2 Fishing geared monitored

A total of 11 key fishing gears were selected for monitoring in the five selected coastal community fisheries. These selected fishing gears were chosen as being typical and commonly used small-scale fishing gears which are commonly used in in-shore fisheries by subsistence or family-scale fishers. They were selected through a participatory consultation meeting with each of the selected CFs. Amongst these fishing gears, hand collection or gleaning was one of the most common practices used by the community fisheries members. Often villagers (with a spade, basket and a sliding board) spend approximately half a day to one day in the coastal areas, mangrove forests or sea grass searching for marine aquatic animals such as blood cockle, crabs and shrimp.

It is worth noting that the legal status of the fishing gears selected for this monitoring project is still unclear. Chapter 9 (Article 45) of the current fisheries law of Cambodia briefly states that “all types of fishery exploitations in the marine fishery domain, except subsistence fishing, shall be allowed only in the possession of license and these exploitations shall follow the conditions and obligation in the fishing logbook”. Article 45 also indicates that “the model of the fishing logbook shall be determined by the proclamation of the Minister of Agriculture, Forestry and Fisheries”. However, there is no current official proclamation which clearly categorizes ‘in detail’ the dimensions for different types of fishing gears used in the marine waters of Cambodia. Therefore, in principle, it is difficult to clearly categorise whether the fishing gears selected for this monitoring project were legal or illegal and whether they should be considered as subsistence fishing gears or otherwise – even though, in practice, it was claimed that they all were subsistence fishing gears. The full list of the 11 fishing gears monitored in this study is given in Table 1 below.

Table 1: List of fishing gears monitored in this study

No.	Khmer name	English name
1	Lop kdam	Crab trap
2	Lob trey	Fish trap
3	Mong trey	Fish gillnet
4	Mong kdam	Crab gillnet
5	Mong Karav	Karav (threadfin) gillnet
7	Mong Kabork	Kabork (mullet) gillnet
6	Santouch Os Meuk	Squid hook and line
7	Ronorng Khyorng	Gastropod (octopus) long line
8	Chheb runh dai	Hand pushnet
9	Chhnieng chunhchoat	Wedge-shaped scoop basket
10	Chorb chhik neg dai	Spade and hand collection
11	Riev Krieng Chhiem	Hand collection of blood cockle

4.3 Survey design

Data on coastal fisheries from 26 subsistence fishers, including 05 women, in the selected community fisheries was collected on a daily basis for the purpose of better understanding catch (catch per unit of effort) and corresponding value that subsistence fishers obtain for their catches if it was sold. The main data items recorded were catch weight (total and by key species), prices (total and by key species), and total lengths of key species, plus data on gear types, and fishing duration. The main steps in the research process were: (1) designing logbooks, (2) training selected fishers as data collectors on data collection methods, (3)

collecting data from all the selected fishers, (4) designing a database and (5) entering all the collected data into the database, checking for errors and analysing the collected data for final reporting.

- (1) To begin with, a logbook was designed in order to facilitate the data collection process. The logbook consisted of cover page and data recording pages. The cover page details the name of the project, the involved institutions (Fisheries Administration and RFLP/CAM) and the fisher's contact details as well as the community fisheries to which he/she belongs. The data recording pages were designed to capture data concerning the dimension of fishing gears used by the selected fishers, total catch per fishing trip, species composition and their prices per kilogram and total lengths of each species with the largest size etc. Logbooks were always made available to all fishers for recording their daily catch on a monthly basis. Each logbook was usually composed of a cover page, 30-31 pages of data recording sheets and diagrams showing how samples and sub-samples should be taken when recording data into the data recording sheets, as well as how the length of marine aquatic species should be measured. The logbooks were produced using simple, easy to understand Khmer language. An example logbook is given as Annex 1 of this report.
- (2) Next, before the actual data collection was started, a team of data collectors (fishers) in each of the five selected community fisheries was established. The full list of the data collection team names and gear types used in each community fisheries are given as Annex 2. All the data collectors were trained in data collection methods and on the identification of different aquatic species. Data collectors were first trained on how to fill in the logbook. Then, the training was started with data collection methods which included how to sample the catch and sort the sub-sample by species. The training also included how to measure the total length of fish and how to measure other marine aquatic animals such as crabs, shrimp, blood cockles and squid. Data collectors were also trained on how to identify fish species before recording the information onto the data sheets in the logbook. Fishers were advised to put in X-others or *trey cham ros* (by-catch or trash fish) if they had any difficulties sampling and identifying species which were of small size.
- (3) The third step was the actual data collection process by the 26 fishers using 11 different fishing gear types from the five selected coastal community fisheries. Logbooks were given out to all the selected fishers in advance. To properly record the catch per unit of effort (CPUE), the time when fishing gear was in the water and the time when the catch was removed from the sea was recorded. Logbooks were designed and explained in a way which allowed fishers to properly record the actual fishing period in hours for their corresponding catches. To record the catch by species for each fishing gear, samples were taken for analysis of species composition. A sub-sample of fish was sorted by species and each species was weighed on calibrated balances. The price of each species was also recorded, based on information from fishers who, in some cases, sell their catches to a middle-person or in a local market. For some common species, total fish lengths of the largest species were measured using a measuring ruler accurate to 1 mm.
- (4) The fourth survey step was to design a database. The database was designed in Microsoft Access 2007 in accordance with the agreed logbook. Data concerning

fishers' profiles, fishing gears used and marine fish and other aquatic animals were entered, stored and analyzed in the database. The database was first tested with actual field data collected by fishers to make sure that it functioned well. Thereafter one key identified staff from the Department of Planning, Finance and International Cooperation of the Fisheries Administration was given training on operation of the database. The concerned staff member was responsible for data entry for the whole period of the study.

- (5) The last step was data entry, data checking for errors and analysis. All the logbooks (with data) were collected from the 26 selected fishers on a monthly basis. The logbook data were sorted and entered into the database according to the community fisheries name, the selected fisher's name and fisher codes to facilitate data entry. Data cleaning and checking for errors was an essential next step that was conducted to ensure that there were no data anomalies and validated before final analysis for reporting. Data storage, processing, cleaning and some data analysis were undertaken in Microsoft Access 2007. Later, the data was exported to Electronic Spreadsheet for final analysis.

In addition to the above, the local Khmer names of aquatic species reported by fishers were verified using existing field guides in order to check for corresponding English and scientific names against Fishbase. Unknown/unidentified species were preserved in a plastic bottle in alcohol and were taken back to a laboratory for further identification. Samples of any unidentified species have been kept at the Fisheries Administration office in Phnom Penh.

5. Results and discussion

5.1 Fisher profiles

A total of 26 coastal fishers from five coastal community fisheries in the four coastal six provinces were involved in this study. All the selected fishers practice only small-scale fishing using gears to fish in support of their livelihoods. An identification number or fisher ID (1-26) was given to each fisher for the purpose of data analysis. The details of each fisher including their name, CF name, and province to which they belong, the fishing gear used, fishing gear dimensions and the number of fishing gear used (if applicable) are given in Annex 2 of this report.

5.2 Average catch and value per fishing trip

Table 2 presents the average catch per crab fishing trip reported by each fisher monitored between late April and November 2011 from the 05 piloted CFI's.

Overall for the 26 selected fishers, the average time of gear operation was 8.9 hours, though there was a lot of variation between different fishing gears, between fishers and fishing trips. The average catch and value at landing sites per fishing trip was around 18 kg and Riel 80,000. Detailed average catch and value against average actual fishing time in hours for each fisher, which were reported for their catch are given in Annex 3 of this report.

It is also interesting to note that fishers using fish traps (fisher ID: 11 and 14) from Koh Rounsanglem community fisheries left their fish traps² in the water for around 40 hours before retrieving them again. These fish trap types were only used in Koh Rounsanglem (amongst the five piloted CFIs). They often capture economically high value species such as groupers. Although their catch were only of between 10-20 kg per fishing trip) which was lower than other fishers, their income from selling their catch was higher. The two fish trap fishers - Fisher ID: 11 and 14 were earning approximately Riel 200,000 and 300,000 (US\$ 50 and US\$ 75) each respectively per fishing trip.

Similarly, fish gillnet (fisher ID: 6³) and *karav* (threadfin) gillnet (fisher ID: 9⁴) set their nets for an average of 7 hours per fishing trip. However, the average catch per fishing trip reported by the fishers was about 100 and 130 kg respectively over the study period. The average value obtained per fishing trip was around Riel 150,000 for fisher ID: 9 and almost Riel 400,000 for fisher ID: 6.

Also, from Table 2, it appears that trap crab fishers (Fisher ID: 7, 24, 22, 3, 18 and 17) earn significantly more from their small-scale fishing than most other fishers except those targeting high value fish like groupers and crab gillnet fishers. This is perhaps because of the high market price of crab. Average catch and average value per fishing trip from crab trap fishers is given in Table 2 below. Detailed information on prices (Riel/kg) at the landing site for all species caught by each fisher is given at Annex 7 of this report.

Table 2: Average catch (kg) and value (Riel) of crab traps per fishing trip from April to November 2011, ranking according to the total catch

Fisher ID	Main crab species caught	N of Traps	Average catch per trip (kg)	Average catch per 100 traps (kg)	Average Value per trip (Riels)	Average Fishing hour per trip
7	<i>Portunus pelagicus</i>	820	15.44	1.88	160,703.48	9.06
24	<i>Scylla serrata</i>	300	14.48	4.83	133,904.76	11.53
22	<i>Portunus pelagicus</i> and <i>Scylla serrata</i> (32%)	300	12.39	4.13	115,681.08	12.25
3	<i>Portunus pelagicus</i>	150	4.01	2.67	70,936.42	7.31
18	<i>Scylla serrata</i>	50	1.93	3.87	22,716.59	8.45
17	<i>Scylla serrata</i>	50	1.84	3.69	29,177.98	7.02
Average		278	8.35	3.00	100,788.46	9.72

The crab fishers listed in Table 2 above use similar crab traps, but crab trap fishers ID 17 and 18 caught far less than fishers ID 3, 7, 22 and 24 because they had non-motorized boats and had far less traps namely only 50 traps while the others had motorized boats and between 150 - 850 traps. The average crab catch per 100 traps ranged from 1.88 – 4.83 kg, with both fisher ID 17 and 18 having quite high catch rates per 100 traps at 3.69 and 3.87 kg compared to crab

²Fisher ID 11 operated 40 traps; each trap was 1 m long and 0.6 m deep.

Fisher ID 14 operated 20 traps; each trap was 0.7 m long and 0.6 m deep.

³ Fisher ID 6 operated a fish gillnet of 2,300 m long, 3 m deep, with a stretched mesh size of 4.5 cm, operated from a boat with a 13 HP engine.

⁴ Fisher ID 9 operated a *karav* gillnet of 1,500 m long, 6 m deep and with a stretched mesh size of 0.35 cm, operated from a boat with a 13 HP engine.

fishers with motorized boats. The mud crab (*Scylla serrata*) is taken closer to the coast and often in streams and tributaries within the CFi fishing area, while the amount of swimming crab (*Portunus pelagicus*) caught increases further from the coast.

It was also observed that hand collection fishers (Fisher ID: 5, 19, 20, 21 and 26) with some supplementary equipment such as spades, baskets and sliding boards were the least efficient fishers, collecting 02-03 kg of marine fish and other aquatic marine animals such as crabs, cockles and shrimp in mangrove forests and sea grass areas. These fishers spent around 3 to 6 hours per session searching for other marine aquatic animals (non-fish species) and typically only earned around Riel 10,000 to 20,000 per fishing expedition.




Figure 2: Average catch per fishing trip by fisher (April-November 2011)




Figure 3: Average value (value at fish landing site) obtained per fishing trip by fisher

5.3. Monthly catch of top ten fishers

Figure 3 shows the variation in monthly catchweight (kg) of top ten fishing gears (in terms reported total catch) from April to November 2011. Overall, it is seen that monthly catches reported by top ten fishers show no clear trends, but varied around a mean of approximately 800 kg. Figure 3, also shows that monthly catches from fish gillnets and fourfinger threadfin (*Eleutheronema tetradactylum*) karav gillnets peaked in July at 4,096 and 7,712 kg respectively and were the highest monthly total catches reported among the 26 selected fishers. These gillnets were used by fisher ID 6 and 9 from Trapiang Ropov CFI, Kampot province.

Interestingly, hand collection with spade (fisher ID 19 from Chum Pou Khmao CFI, Preah Sihanouk Province) ranked number 9 in terms of the total catch over the period from April to November. This particular fisher spent an average of 8 hours daily collecting various kinds of bivalve mollusks like cockles from mangrove forests and coastal areas. The reported monthly gleaning (hand-collection) catch varied between 170 kg and 290 kg per month with an average of 223 kg. While cockles were freely available in the community, they fetch a very low market price. Fisher ID 19 earned the least income from his catch of all the 26 fishers monitored between April-November 2011 (for additional details see Annex 3).

More details on the monthly catch reported by all fishers are given in Annex 4.




Figure 4: Monthly catch of top the ten fishers ranking according to the total catch over the study period from April to November 2011.

5.4. Monthly value of top ten fishers

Figure 5 shows the variation in monthly catch value obtained by top ten fishers over the period from April to November 2011. The top ten fishers (in terms of total value), earned between Riel 2,000,000 to 4,000,000 (US\$ 500 – 1,000) a month from fishing. Exceptionally fisher ID: 6 from Trapiang Ropov CFi, who used a fish gillnet of around 2,300 meters long, 3 meters deep and a mesh size of 4.5 cm, reportedly earned around Riel 13 million (US\$ 3,250) in August and up to Riel 14.5 million in November (US\$ 3,625). This spike in income by this fish gillnetter was reported to be due to the fish price from about Riel 2,200/kg in July to around Riel 3,800/kg in August, and Riel 3,500 in November.

More details about the monthly value reported by all fishers are given in Annex 4.




Figure 5: Monthly total monetary value of top ten ranked fishers over the study period from April to November 2011.

5.5. Key important species caught by fishers

Overall, more than 100 different aquatic fish, shrimp, gastropod and bivalve mollusk species were reported the all 26 studied fishers over the seven month plus survey period. Gillnets and hand pushnet were reported as having captured a greater bio-diversity of species than other fishing gear types. However, some gillnet types that were designed to target only a small

number of fish species and crabs such as the threadfin *orkarav* gillnet, the mullet or *kabork* gillnet and the crab gillnet were more selective. In addition, some fishing gears were highly selective, targeting and catching only one or two species and these included crab traps, squid hook and line gear and gastropod long lining.

In this section, an attempt has been made to provide detailed information on the key species caught by each of the 26 selected fishers who were involved in recording their catches over the study period from late April to November 2011. More detailed information about the catch and value by species by each fisher are given in Annexes 5 and Annex 6 of this report.

5.5.1 Fisher ID 1: fish gillnet

Overall, 32 species and many other species which were grouped as “X-others” were recorded by this fisher. More than 70 percent of the catch was made up of the 10 species shown in Figure 6. The short mackerel (*Rastrelliger brachysoma*) comprised 28% of the total catch during the survey period, while deep bodied mojarra (*Gerres abbreviatus*) known in Khmer as *Trey do Angkor*, comprised 24 percent of the total catch of fisher ID 1.




Figure 6: Fisher ID 1: Catch (kg) (April-November 2011)

5.5.2 Fisher ID 2: Crab gillnet

Fisher ID 2 reported only catching 8 species over the study period. Figure 7 below shows the main six species caught by fisher ID 2 during the survey period. 95 percent of the catch was composed of only three species namely blue swimming crab (*Portunus pelagicus*) (*Kdam Seih*), the gastropod mollusk *Conus vexillum* (*Khyorng thnot*) and the gastropod mollusk (*Melo melo*) (*Khyorng dong*) comprised 54, 28 and 13 percent of the catch. Interestingly, although this fishing gear specifically targets swimming crab, it also captures several other economically important species and gastropod mollusks in particular.




Figure 7: Fisher ID 2: Catch (kg) (April-November 2011)

5.5.3 Fisher ID 3: Crab trap

Fisher ID 3 reported a total of 16 different species in the catch. However, only one crab species, *Portunus pelagicus* (locally known *Kdam Seih*), formed a large share (59 percent) of the total catch. Other important species included in the catch were the stomatopod crustacean (*Clorida decorata*) (*Bangkorng kandob*), the crab locally named *Kdam krahorm*, the crab (*Matuta victor*) (*Kdam skor krorb/Kdam sor*), the mangrove mud crab (*Scylla serrata*) (*Kdam Thmor*), see Figure 8.




Figure 8: Fisher ID 3: Catch (kg) (April-November 2011)

5.5.4 Fisher ID 4: Hand pushnet

Fisher ID 4 reported only 10 species taken using a hand pushnet, but excluding many other small-sized or trash fish or by-catch which was recorded in “X-others”. Hand pushnets are often operated in inshore areas where there are sea grassbeds. The main species reported by the hand pushnetter included 54 percent banana prawns (*Penaeus merguensis*) (locally known as *Bangkea kor gneub*) as shown in Figure 9.




Figure 9: Fisher ID 4: Catch (kg) (April-November 2011)

5.5.5 Fisher ID 5: Wedge-shaped scoop basket

Fisher ID 5 reported a total of 19 identified species in the catch, plus by-catch (X-others) which was probably made up of several other species. The wedge-shaped scoop basket is also primarily operated in sea grassbeds areas. Apart from this, the same fisher also reported collecting/gleaning marine aquatic animals by hand in the coastal areas and mangrove forests. Seven species taken by the wedge-shaped scoop basket contributed around 70 percent of the catch including *Trey mrech*, *Trey prorkong*, *Bangkea sor*, *Bangkea phourk*, *Trey damlong*, *Trey kanchanh chras*, *Trey kranh*. These these species were only reported and recorded under the local Khmer name. As yet the team has been able to identify the scientific names of any of the species captured by fisher ID 5.




Figure 10: Fisher ID 5: Catch (kg) (April-November 2011)

5.5.6 Fisher ID 6: Fish gillnet

15 species were recorded by this fisher. By-catch which composed of several other species was also reported. As shown in Figure 11, the gillnet targeted mainly mackerel species (*Rastrelliger* spp.), known in Khmer as *trey kamong* comprised 77 percent of the total catch

and included the short mackerel (*Rastrelliger brachysoma*) (55 percent) and the Indian mackerel (*Rastrelliger kanagurta*) (22 percent) respectively.




Figure 11: Fisher ID 6: Catch (kg) (April-November 2011)

5.5.7 Fisher ID 7: Crab trap

Crab traps are a highly selective type of fishing gear. Fisher ID 7 captured 100 percent blue swimming crab (*Portunus pelagicus*) (*Kdam Seih*) and no by-catch, as shown in Figure 10 below.




Figure 12: Fisher ID 7: Catch (kg) (April-November 2011)

5.5.8 Fisher ID 8: Hand pushnet

Fisher ID 8 reported a total of 25 different identified aquatic species in the catch from a hand pushnet. However as Figure 13 shows, seven species contribute 75 percent of the total catch. These species are *Bangka sor* (local name), bluespot mullet (*Moolgarda seheli*) (*Trey kbork*), the banana shrimp (*Penaeus merguensis*) (*Bangka kor gneub*), the black-barred halfbeak, (*Hemiramphus far*) (*Trey phtong phka*), the white-spotted spinefoot, (*Siganus*

canaliculatus) (*Trey kantang kra-ob*), the barhead spinefoot (*Siganus virgatus*) (*Trey kantang thmor*), and the keel-jawed needlefish (*Tylosurus acus melanotus*) (*Trey pthong prort*).




Figure 13: Fisher ID 8: Catch (kg) (April-November 2011)

5.5.9 Fisher ID 9: Karav (Threadfin) gillnet

Karavor trey karav is a local name of the fourfinger threadfin fish species. This fisher designed his own gillnet to target *trey kavav* or fourfinger threadfin (*Eleutheronema tetradactylum*) and that is the reason why this gillnet was referred to as a *karav* gillnet. However, according to the reported catch composition shown in Figure 14, over 93 percent of catch was the shortmackerel (*Rastrelliger brachysoma*) (*Trey kamong khoun khley*). *Trey karav* was the second most caught species, but it contributed only 7 percent of the total catch.




Figure 14: Fisher ID 9: Catch (kg) (April-November 2011)

5.5.10 Fisher ID 10: Kabork (Mullet) gillnet

Likewise the *Kabork* gillnet targets the bluespot mullet (*Moolgarda seheli*) or *trey kabork* in Khmer. Reportedly, 33 species and by-catch “X-others” were recorded as captured by the selected fisher ID 10 during the study period. As illustrated in Figure 15, the top ten 10 species contributed 72 percent of the catch from the *kabork* gillnet during the study period; bluespot mullet (*Moolgarda seheli*) or *trey kabork* ranked top and contributed around 23 percent of the total catch. Banana shrimp (*Penaeus merguensis*) was ranked second and

contributed 14% of the total catch. However the proportion of by-catch was high at 28 percent.




Figure 15: Fisher ID 10: Catch (kg) (April-November 2011)

5.5.11 Fisher ID 11: Fish trap

Fish traps were only operated at Koh Rounsalem CFI, Preah Sihanouk Province. The fish traps captured 13 different identified species and some by-catch of less valuable species. More than 80 percent of catch from this fishing gear was made up of the 9 key marine species shown in Figure 16 below.

The main species caught by fish traps was parrotfish (*Scarus* spp.) which constituted 26% of the catch. Fish traps also captured high value fish species such as groupers, with the highfin coral grouper (*Plectropomus oligacanthus*) (*Trey teukei ouch kiev*) ranked third and contributing 13 percent of the total catch weight for fisher ID 11. This species sells at Riel 100,000 per kg (US\$ 25 per kg).




Figure 16: Fisher ID 11: Catch (kg) (April-November 2011)

5.5.12 Fisher ID 12: Squid hook and line

Squid hook and line is a type of fishing gear that uses artificial bait (with hooks) to attract squid. It was used by fishers in Koh Rounsalem CFI. When in operation, the boat tows the hooks and line with the artificial bait behind to attract squid. As shown in Figure 17, the gear is highly selective with 97 percent of the reported catch being the squid (*Sepioteuthis lessoniana*) (Meuk huysay).




Figure 17: Fisher ID 12: Catch (kg) (April-November 2011)

5.5.13 Fisher ID 13: Squid hook and line

Similarly, Fisher ID 13 operated squid hook and line gear. He reported catching only two species of squids, namely *Sepioteuthis lessoniana* (Meuk huysay) and *Uroteuthis duvaucelii* (Meuk bompong/meit) which constituted 94 and 6 percent of the catch respectively, see Figure 18 below.




Figure 18: Fisher ID 13: Catch (kg) (April-November 2011)

5.5.14 Fisher ID 14: Fish trap

Similar to Fisher ID 11, Fisher ID 14 operated fish traps in Koh Rounsalem CFI, and recorded 11 species in his catch. Parrotfish (*Scarus* spp.) or *Trey Seik* constituted 26 percent of

the catch, and the high value highfin coral grouper (*Plectropomus oligacanthus*) (*Trey teukei ouch kiev*) was ranked second, with 23 percent of the catch, see Figure 19 below.




Figure 19: Fisher ID 14: Catch (kg) (April-November 2011)

5.5.15 Fisher ID 15: Squid hook and line

Like Fisher ID 12 and 13, Fisher ID 15 operated squid hook and line gear and reported catching the same two squid species namely 92 percent *Sepioteuthis lessoniana* (*Meuk huysay*) and 8 percent *Uroteuthis duvaucelii* (*Meuk bompong/meit*), see Figure 20 below.




Figure 20: Fisher ID 15: Catch (kg) (April-November 2011)

5.5.16 Fisher ID 16: Gastropod long line

Gastropod long lining gear was used by fishers in Koh Rounsanlem. Shells from dead gastropods were used to provide shelter and hiding places for octopus species. This fisher used 1,000 shells, connecting by a line with an interval of one meter between shells. The gear was extremely selective catching only octopus species, *Sepioloa* sp. (known in Khmer as *Meuk dong*), see Figure 21 below.




Figure 21: Fisher ID 16: Catch (kg) (April-November 2011)

5.5.17 Fisher ID 17: Crab trap

All crab trap fishers reported only catching mainly the blue swimming crab (*Portunus pelagicus*) (*Kdam Seih*). Most of the crab trapfishers fished in deeper water using motorized boats.

However, fisher ID 17 from Chum PouKhmao CFi could not afford a motorized boat and so used a non-motorized paddle boat to set his 50 crab traps in canals and streams and nearby coastal areas in shallow waters. This fisher reported only catching the mangrove mud crab (*Scylla serrata*) (*Kdam Thmor*) which fetched a high market price.




Figure 22: Fisher ID 17: Catch (kg) (April-November 2011)

5.5.18 Fisher ID 18: Crab trap

Like fisher ID 17, fisher ID 18 was also from Chum PouKhmao CFi. He also used a non-motorized paddle boat to set his 50 crab traps in canals and streams and nearby coastal areas in shallow waters. His catch was also 100 percent mangrove mud crab (*Scylla serrata*), see Figure 23 below.




Figure 23: Fisher ID 18: Catch (kg) (April-November 2011)

5.5.19 Fisher ID 19: Spade and hand collection of cockles

Fisher ID 19 regularly searched for marine aquatic animals in coastal areas especially in mangrove forests. He used his hands and a spade to glean almost 100 percent Princess Venus clams (*Periglypta sp.*) (*Ngveav sor*), which only fetch 600 Riel/kg at market). As shown in Figure 22, he occasionally gets lucky and may catch a mangrove mud crab (*Scylla serrata*) (*Kdam Thmor*).




Figure 24: Fisher ID 19: Catch (kg) (April-November 2011)

5.5.20 Fisher ID 20: Hand collection plus spade (targeting crabs)

This fisher was from Chum Pou Khmao CFi and he used his hands and spade to search for marine aquatic animals. He targeted the tree-climbing vinegar crab (*Episesarma singaporensis*) (*Kdam Chor*). As shown in Figure 23, 79 percent of the catch was of this particular species. *Kdam Chor* were sold at around Riel 4,000 per kilogram (US\$ 1). 9 percent of his catch was mangrove mud crab *Scylla serrata* (*Kdam Thmor*) see Figure 25 below.




Figure 25: Fisher ID 20: Catch (kg) (April-November 2011)

5.5.21 Fisher ID 21: Hand collection plus spade (targeting crabs)

Fisher ID 21 belonged to the same CFi as fisher ID 20 and practiced the same fishing method, i.e. – hand collection and using a spade in search of crabs. 89 percent of the reported catch was the tree climbing vinegar crab (*Episesarma singaporens*) (*Kdam Chor*), while the remainder included 10 percent mangrove mud crab (*Scylla serrata*) (*Kdam Thmor*) and some mud clams (*Polymesoda erosa*) (*Ngeav phourk*), see Figure 26 below.




Figure 26: Fisher ID 21: Catch (kg) (April-November 2011)

5.5.22 Fisher ID 22: Crab trap

As shown in Figure 27, 98 percent of the catch of crab trap reported by Fisher ID 22 was composed of only four crab species namely blue swimming crab *Portunus pelagicus* (*Kdam Seih*) 51 percent, mangrove mud crab (*Scylla serrata*) (*Kdam Thmor*) 32 percent, the swimming crab (*Charybdis anisodon*) (*Kdam Sor/Dangkeab Sor*) 12 percent and (*Matuta victor*) (*Kdam skor krorb/Kdam sor*) 3 percent.




Figure 27: Fisher ID 22: Catch (kg) (April-November 2011)

5.5.23 Fisher ID 23: Fish gillnet

Fisher ID 23 reported 20 different identified aquatic species. However, as shown in Figure 28, 80 percent of the catch consisted of only three species which included the frigate tuna (*Auxis thazard*) (Trey beika) 46 percent, the gizzard shad (*Anodontostoma chacunda*) (Trey kamoy) 21 percent and the shortmackerel (*Rastrelliger brachysoma*) (Trey kamong khloy) 13 percent.




Figure 28: Fisher ID 23: Catch (kg) (April-November 2011)

5.5.24 Fisher ID 24: Crab trap

Fisher ID 24 was from Koh Khyornng CFi. As shown in Figure 29, only crab species were captured. 95 percent of the catch were from three main crab species the mangrove mud crab (*Scylla serrata*) (Kdam Thmor) 39 percent, the blue swimming crab (*Portunus pelagicus*) (Kdam Seih) 37 percent and the swimming crab (*Charybdis anisodon*) (Kdam Sor/Dangkeab Sor) 19 percent.




Figure 29: Fisher ID 24: Catch (kg) (April-November 2011)

5.5.25 Fisher ID 25: Crab gillnet

As Figure 30 shows, the crab gillnet of fisher ID 25 caught not only crabs, but also fish species. Around 50 percent of the catch was made of two species of crabs namely the blue swimming crab (*Portunus pelagicus*) (*Kdam Seih*) and the mangrove mud crab (*Scylla serrata*) (*Kdam Thmor*) which comprised 28 and 22 percent of the catch respectively. The balance of the catch consisted of fish species which included 17 percent scaly whipray (*Himantura imbricata*) (*Trey borbel moan*), 7 percent of the frigate tuna (*Auxis thazard*) (*Trey beika*) and 4 percent of island mackerel (*Rastrelliger faughni*) (*Trey balang*).




Figure 30: Fisher ID 25: Catch (kg) (April-November 2011)

5.5.26 Fisher ID 26: Hand collection of blood cockles

Fisher ID 26 was from Koh Khyorng CFi. He collected only blood cockles in the coastal area during the low tide. He used only a wooden sliding board to sit on and to move across the mud flats from place to place in search of blood cockles in the mud along the CFi coastline. As Figure 31 shows, 97 percent of his reported catch was blood cockles (*Anadara granosa*) (*Kreing chheam*).




Figure 31: Fisher ID 29: Catch (kg) (April-November 2011)

5.6. Monthly average prices at landing sites by species by fisher

Daily prices per kilogram of all species captured by all fishers at the landing sites were recorded. Average monthly prices per kilogram of each aquatic species were then calculated. The average monthly prices by aquatic species varied slightly between months over the study period. The detailed monthly average prices of all aquatic species reported by the 26 selected fishers at fish landing sites are given in Annex 7 of this report.

5.7. Frequencies versus average maximum length by fisher by species

Length frequency is an indicator that is used by fisheries biologists to better understand about the health status of aquatic stocks. Therefore, for this study, each fisher was asked to also record the maximum length of each species caught on a daily basis when they reported their catches. Average maximum length of each species captured by each of the 26 selected fishers involved in this study was then calculated against their frequencies (number of marine fish or other aquatic marine animal). These data are provided in Annex 8. They serve as an important baseline data which can be referred to see if fish lengths of the aquatic species being caught decreases, which would be an indication of declining stock health.

It is also worth noting that under the current fisheries law of Cambodia, the sizes of aquatic animals to be harvested are determined by the proclamation of the Minister of Agriculture, Forestry and Fisheries as stated in Chapter 5, article 23. However, there appears to be no proclamation which details the minimum size restrictions on species to be caught despite the fact there is a subdecree (signed on August 12, 2009) which identifies 58 endangered aquatic animals including 29 marine fish, reptile and mammal species and spells out which freshwater and marine animals are banned from being transported or traded unless they are being farmed or are in compliance with the domestic fisheries law and the Convention on International Trade in Endangered Species of Wild Fauna and Flora (CITES) which Cambodia ratified in 1997.

6. Conclusions

This seven month study provides some essential baseline data provided by 26 coastal small-scale fishers using 11 different gear types in the coastal inshore waters of Cambodia. Fishing was the main livelihood and income source of the 26 selected fishers which were from five coastal CFi's of Kep, Kampot, Preah Sihanouk and Koh Kong Provinces. Data on species diversity, gear selectivity, total catch weight and value, catch weight and value by species,

average actual fishing time in hours per fishing trip, prices of marine aquatic fish and other aquatic species caught, as well as the maximum length frequencies of species captured were recorded from late April to November 2011, i.e. just over a seven month study period. The major survey findings were:

- More than 100 marine aquatic species were reported by the 26 selected fishers involved in this study. The species caught varied according to the different types of fishing gears used, location, and season. Data on the catch weight and value for total catch and by species, and prices, as well as the maximum length of all species were collected, entered onto a data, checked for accuracy and analysed and are available in Annexes 1 – 8 of this report.
- Some fishing gears were highly selective and had resulted in virtually no by-catch. These included crab traps, the gastropod shell long line system for octopus, and squid hook and line gear, as well as hand collection of blood cockles.
- Some fishing gears such as crab gillnets, fish traps and hand collection with supplementary equipment were moderately selective, capturing around 10 aquatic species or less and usually with only limited by-catch.
- Other fishing gears such as different types of fish gillnet and the hand pushnet used particularly in sea grass beds were highly unselective and destructive, capturing up to more than 30 different aquatic species and usually with significant proportions of by-catch.
- The overall average time that fishing gear was in the water was 8.9 hours though there was significant variation between fishing gears, fishers and trips.
- The overall average catch and value at landing sites per fishing trip were around 18 kg of aquatic products worth Riel 80,000 (US\$ 4) though again there was significant variation across fishing gears, locations, fishers and season.
- Fish trap fishers left their traps in the water for around 40 hours each time before retrieval. Fish traps were however only operated and sampled in Koh Rounsanglem (one of the five piloted CFIs) and targeted high value species such as groupers. While the catch per fishing trip at typically 10-20 kg was relatively low compared to other fishers, the monetary value of the catch was comparatively higher at around between Riel 200,000 - 300,000 (US\$ 50 – 75) per fishing trip.
- Fish gillnets in Trapiang Ropov CFI were set for an average of 07 hours per fishing trip and caught between 100 and 130 kg of aquatic products with a total value of around Riel 150,000 and Riel 400,000 (US\$ 37.5 - 100).
- Crab trap fishers set their traps for an average of 10 hours, and caught an average of 8 kg of crabs, with a total average value of about Riel 100,000 (US\$ 25).
- A gastropod shell long line fisher (with 1,000 gastropod shells) who fished for octopus around Koh Rounsanglem set his long line for on average of around 2 hours, and typically captured around 10 kg of *Sepiolo* sp. (Meuk dong) with a total value of around Riel 110,000 (US\$ 27.5).
- Hand push-netting is practiced especially in Kep and Kampot Provinces. Hand pushnet fishers typically worked for 8 hours per expedition and caught between 2 and 5 kg of aquatic products with a value of ranging between Riel 11,000 to Riel 20,000 (US\$ 2.75 – 5) each fishing trip.
- Hand collection fishers with supplementary equipment such as a spade, a sliding board, and basket were the least efficient fishers, having the least catch. They typically spent around 6 hours per trip in search for marine aquatic animals. On

average, they were able to collect 3-4 kg of cockles which are the lowest value species, with a total sale value of around Riel 10,000 (US\$ 2.5).

7. Suggestions for further studies

While this study provides useful baseline data and information about catch and value of the fisheries as well as species diversities being enjoyed by a small selection (n = 26) of small-scale individual fishers in five pilot coastal community fisheries of the four coastal provinces of Cambodia, a number of suggestions are made for further studies in order to obtain more necessary data and information for a better understanding on Catch per Unit of Effort (CPUE) and biological and socio-economic aspects of the coastal fisheries and their contribution to the livelihoods of coastal fishers.

This study totally depends on catches reported by fishers. Better off fishers often change their gear types for aquatic marine species in accordance with seasons and the migration behavior of different target species. In addition, the number of fishing gears used often fluctuates because of various reasons such as thieves, loss due to heavy storms, seasonality of species occurrence at specific locations, the availability of workers and many other factors. All the aforementioned factors make CPUE calculation difficult to allow calculation across gear types. For this reason throughout this study, calculations were therefore based on a per fishing trip basis rather than on a per unit of fishing gear used. For further study, therefore, standardized fishing gear units should be used to provide standard sampling that will give a better understanding of CPUE per gear type. Moreover, in order to be able to extrapolate the study findings to other areas of Cambodia, the sampling sites should be more geographically representative of the coastal areas of Cambodia and cover a greater range of different fish habitat types and ecosystems. For instance, fishers expect to catch different aquatic and fish species in mangrove forests, in sea grass beds and in coral reef areas.

This study focuses only on the reported catch and value by species of individual fishers. None data was collected on the socio-economic aspects of fishers and the fishers' families as well as on the coastal communities to which they belong. Very little environmental data was collected during this survey, though environmental parameters would have been relevant and would have impacted on this study.

In reality the size of catch and its value, as well as species occurrence reported by the selected fishers was probably affected by many external factors including the number of other fishing gears in the area during the survey period, the presence of foreign fishing vessels and changing environmental conditions, etc. For future studies comparison this and other biological, socio-economical and environmental parameters should also be properly considered and studied.

The data and information reported in this study should be presented back to the 5 pilot CFI members and especially to those directly involved in recording and reporting the data. This is important as it will give the research team and the fishers an opportunity to provide feedback and validate the findings. Also, it is important that data from this study be used by the pilot CFI committees to better plan and monitor and understand the catch status in their CFI waters.

Last but not least, a field guide which contains the majority of marine aquatic species with proper local Khmer names, English names, and scientific names and a brief description on the identification of the species should be produced and made available, as this type of field guide would be very useful for this type of research. Currently marine field guides only exist in Cambodia for the most common key main marine aquatic species. A large number of marine species still remain to be properly documented.

Annexes

Annex 1: Fisher logbook

Cover Page



គម្រោងរៀនសូត្រស្តីពីការចិញ្ចឹមត្រីតំបន់អាស៊ីអាគ្នេយ៍និងអាស៊ីបូព៌ា

The Regional Fisheries Livelihoods Programme for South and Southeast Asia (RFLP) - Cambodia

Fishing Logbook

(Local Language)

Fisher name: Village: Commune:
District: Province:
Name of Community Fisheries:

Fisheries Administration, # 185 Preah Norodom Blvd., Khan Cham P.O.Box 582, Phnom Penh Cambodia

Annex 2: Name of fishers and their fishing gears used in five target community fisheries

FisherID	Fisher name	Community Fisheries	Province	Fishing gears used	Fishing gear dimensions	Gear unit
1	Prum Phorn	Okrasa	Kep	Fish gillnet	Length: 500 m, Depth: 2.5 m	
2	Choey Chan Thorn	Okrasa	Kep	Crab gillnet	Length: 800 m, Depth: 0.8 m Mesh size: 9 cm.	
3	Liet Chhoeun	Okrasa	Kep	Crab trap	Length: 0.4 m, Depth: 0.2 m	150 traps
4	Thuok Sokhon	Okrasa	Kep	Hand pushnet	Length: 3.5 m, Width: 5.5 m, Depth: 1 m	
5	Chea Ranut	Okrasa	Kep	Wedge-shaped scoop basket plus hand collection of marine aquatic animals (targeting shrimp)	Length: 1 m, Depth: 0.8 m	
6	Sos Seth	Trapiang Ropov	Kampot	Fish gillnet	Length: 2,300 m, Depth: 3m, mesh size: 4.5 cm	
7	Chan Sok	Trapiang Ropov	Kampot	Crab trap	Length: 0.5 m, Width: 0.25m, Depth 0.15 m	820 traps
8	Lan Mork	Trapiang Ropov	Kampot	Hand pushnet	Length: 4 m, Width: 5 m	
9	Rem Mi	Trapiang Ropov	Kampot	Karav (Threadfin)gillnet	Length: 1,500 m, Depth: 6 m, Mesh size: 3.5 cm	
10	Ni Phas	Trapiang Ropov	Kampot	Kabork (Mullet) gillnet	Length: 150 m, Depth: 3 m, Mesh size: 3.5 cm	
11	Chet San	Koh Rounsanlem	Preah Sihanouk	Fish trap	Length: 1 m, Depth: 0.6 m	40
12	Chiv Mut (Bela)	Koh Rounsanlem	Preah Sihanouk	Squid hook and line	Length: 4.5 m	16 hooks
13	Kong Thorn	Koh Rounsanlem	Preah Sihanouk	Squid hook and line	Length: 4.5 m	35 hooks
14	Ma Thorn	Koh Rounsanlem	Preah Sihanouk	Fish trap	Length: 0.7 m, Depth: 0.6 m	20 traps
15	Moul Raksmeay	Koh Rounsanlem	Preah Sihanouk	Squid hook and line	Length: 4.5 m	18 hooks

						(0.35 cm)
16	Kong Eang	Koh Rounsanlem	Preah Sihanouk	Gastropod long line	Length: 1,000 m	1000 gastropods
17	Uon Aem	Chum Pou Khmao	Preah Sihanouk	Crab trap	Length: 0.4 m, Width: 0.25m, Depth: 0.15 m	50 traps
18	Chhoeun Phal	Chum Pou Khmao	Preah Sihanouk	Crab trap	Length: 0.4 m, Width: 0.25m, Depth: 0.15 m	50 traps
19	Kun Sarun	Chum Pou Khmao	Preah Sihanouk	Spade and hand collection of cockle	-	-
20	Norn Ri	Chum Pou Khmao	Preah Sihanouk	Hand collection of crab (targeting mud crab) plus spade	-	-
21	Khan Ratha	Chum Pou Khmao	Preah Sihanouk	Hand collection of crab (targeting mud crab) plus spade	-	-
22	Touch Vorn	Koh Khyorng	Koh Kong	Crab trap	Length: 0.45 m, Width: 0.29 m, Depth: 0.19 m	300
23	Chuon Kea	Koh Khyorng	Koh Kong	Fish gillnet	Length: 560 m, Depth: 0.3 m, Mesh size: 0.38 cm	
24	Ly Cheak	Koh Khyorng	Koh Kong	Crab trap	Length: 0.45 m, Width: 0.25m, Depth: 0.15 m	300
25	Hang Mom	Koh Khyorng	Koh Kong	Crab gillnet	Length: 900 m, Depth: 08 m, Mesh size=8 cm	
26	Kea Hin	Koh Khyorng	Koh Kong	Hand collection of blood cockle plus sliding board and basket	-	-

**Annex 3: Average weight (Kg) and value (Riel) per fisher per trip
(ranking according to value at the landing site)**

Fisher ID	Fishing gears used	Average weight (Kg)	Average value (Riel)	Average time spent per fishing trip (hour)
6	Fish gillnet	130.42	394,966.10	6.51
11	Fish trap	21.98	296,802.31	39.08
14	Fish trap	8.17	202,617.46	42.85
2	Crab gillnet	13.96	167,019.71	12.13
7	Crab trap	15.44	160,703.48	9.06
9	Karav gillnet	99.16	143,365.21	6.00
24	Crab trap	14.48	133,904.76	11.53
1	Fish gillnet	28.08	128,624.32	5.76
22	Crab trap	12.39	115,681.08	12.25
16	Gastropod (octopus) long line	10.32	110,150.54	2.15
15	Squid hook and line	6.31	74,727.78	6.41
3	Crab trap	4.01	70,936.42	7.31
12	Squid hook and line	4.36	58,837.06	7.51
25	Crab gillnet	4.48	37,513.94	23.91
23	Fish gillnet	19.78	33,940.83	6.22
17	Crab trap	1.84	29,177.98	7.02
13	Squid hook and line	1.93	25,373.60	3.90
10	<i>Kabork</i> gillnet	5.72	24,175.72	7.84
18	Crab trap	1.93	22,716.59	8.45
8	Hand pushnet	5.39	20,409.13	7.75
5	Wedge-shaped scoop basket plus hand collection	3.87	17,076.18	4.21
20	Hand collection plus spade (crabs)	2.36	12,232.42	6.96
21	Hand collection plus spade (crabs)	2.35	12,124.70	4.95
4	Hand pushnet	1.92	11,601.64	7.72
26	Hand collection of blood cockle plus sliding board	2.07	10,582.91	3.68
19	Hand collection plus spade (cockles)	9.03	5,597.66	7.88
Grand Total		18.08	82,304.34	8.88

Annex 4: Monthly catch (Kg) and value (Riel) by fisher (Ranking according to value)

Fisher ID	Catch (Kg)									Value (Riel)								
	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Total	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Total
6	460	1,289	2,909	4,096	3,395	3,367	3,308	4,143	22,973	920,550	4,089,150	5,467,400	9,246,400	13,046,000	10,449,000	11,747,000	14,506,500	69,472,000
9	508	990	2,123	7,712	2,346	2,802	2,672	2,366	21,527	1,092,250	2,140,900	3,281,800	9,571,300	2,918,000	4,302,400	3,486,000	4,317,600	31,110,250
24	204	806	378	403	311	249	255	348	2,978	2,569,900	7,246,000	3,203,000	3,570,500	2,845,100	2,403,100	2,298,475	3,194,550	27,330,625
7	307	293	246	327	421	363	181	302	2,447	2,962,650	2,848,350	2,324,200	3,698,750	5,139,900	4,168,800	1,601,600	2,646,900	25,391,150
2	119	260	182	269	317	325	284	171	1,929	1,394,450	2,957,700	2,272,360	3,405,960	3,739,200	3,747,050	3,386,600	2,145,400	23,048,720
22		454	441	323	396	240	276	285	2,437		3,837,350	3,894,981	3,058,100	3,671,300	2,266,080	3,202,150	2,627,850	22,557,811
1	396	348	448	442	624	635	933	695	4,522	1,349,550	1,251,300	2,055,100	2,253,935	3,893,490	4,057,500	3,312,340	2,535,300	20,708,515
11	12	21	93	137	175	232	318	442	1,440	129,500	328,600	1,420,550	2,280,500	2,631,000	2,870,000	5,006,000	4,626,000	19,292,150
3	73	112	105	52	83	84	136	158	804	1,154,020	1,393,620	1,657,619	442,515	1,019,240	1,415,685	2,282,970	4,821,615	14,187,283
14	3	39	36	58	79	91	99	109	529	56,000	1,505,800	780,900	1,375,000	1,836,300	2,042,700	2,224,000	2,944,200	12,764,900
16		9	55	38	38	200	320	280	956		85,500	522,500	380,000	456,000	2,400,000	3,400,000	2,800,000	10,044,000
12	48	116	72	93	96	65	127	119	749	581,500	1,506,200	941,700	1,227,200	1,257,700	882,400	1,647,600	1,896,000	9,940,300
15	18	193	58	143	130	51	33	170	810	240,800	2,234,400	627,900	1,568,500	1,571,600	624,900	440,800	2,106,800	9,415,700
17	24	38	43	51	60	66	62	59	419	246,600	492,800	625,100	776,300	922,500	1,176,300	1,071,500	1,049,700	6,360,800
23	15	64	212	239	250	523	824	1,212	3,363	83,425	409,000	431,050	504,700	691,550	1,049,900	1,391,850	1,263,230	5,824,705
25	15	91	82	65	70	77	111	75	612	139,500	660,126	725,850	473,500	451,860	638,000	1,056,150	769,340	4,914,326
18	19	39	63	63	59	58	51	57	426	195,000	465,000	730,600	759,300	724,100	671,000	598,400	649,800	4,793,200
10	84	143	169	194	115	115	130	121	1,080	407,610	503,950	711,000	826,700	469,600	492,900	590,800	518,300	4,520,860
8	76	98	103	172	111	106	135	159	968	305,700	319,700	350,550	548,950	365,025	469,800	602,000	671,100	3,632,825
13	21	52	32	32	20	15	34	35	254	271,600	652,800	397,100	405,900	257,400	199,500	466,400	521,000	3,171,700
5	21	144	202	88	51	50	25	109	694	175,250	659,300	594,900	266,780	164,150	172,150	190,300	816,730	3,039,560
20	42	57	54	69	25	53	37	54	410	173,300	279,150	275,350	402,150	198,400	273,100	163,800	253,100	2,018,350
21	39	55	54	74	57	52	28	25	406	178,200	239,750	270,650	509,050	304,350	241,600	122,000	117,850	1,983,450
4	42	11	40	32	53	42	27	33	285	264,590	53,260	355,850	158,500	321,450	231,890	149,210	159,090	1,693,840
26	20	35	41	36	38	27	21	26	269	80,000	144,100	164,700	295,100	174,050	119,250	141,750	119,250	1,238,200
19	172	273	203	215	176	236	287	-	1,581	103,200	163,500	121,800	132,300	108,200	147,600	191,796	-	968,396

Annex 5: Monthly catch (Kg) by species and by fishers, ranking according to total catch over the period from April to November 2011

Fisher ID: 1

Local Name	Scientific Name	Catch (kg)								
		Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Total
<i>Trey do angkor</i>	<i>Gerres abbreviatus</i>			164.1	124.5	192.4	212.6	52.5	312.2	1,058.3
<i>Trey prorleuk</i>	<i>Leiognathus splendens</i>						59.2	245.4	73.9	378.5
<i>Trey kamoy</i>	<i>Anodontostoma chacunda</i>	150.5	184.0	7.3	0.3		4.5			346.6
<i>Trey Krorb khnol</i>	<i>Lethrinus nebulosus</i>			4.0	71.0	99.3	93.7	35.6	23.0	326.6
<i>Trey krorhorm srakakras</i>	<i>Scolopis taeniopterus</i>						121.4	85.0	80.0	286.4
<i>Trey krorhorm kbalkor</i>	(blank)				60.5	145.2		10.4	22.7	238.8
<i>Trey kamong khoun khley</i>	<i>Rastrelliger brachysoma</i>	89.0	73.0	8.5						170.5
<i>Trey prorlos</i>	<i>Sillago sihama</i>			37.4	38.0	78.4				153.8
<i>Trey Kor/Romeas</i>	<i>Monacanthus chinensis</i>				33.0	28.4	39.5	15.0	33.8	149.7
<i>Trey chek</i>	<i>Seriolina nigrofasciata</i>	38.0	15.0	8.0	33.0	15.1	11.0	1.0	25.0	146.1
<i>Trey koun</i>	<i>Hilsa keele</i>	26.0	18.0	3.0				32.9	9.5	89.4
<i>Trey krorhorm</i>	<i>Lutjanus malabaricus</i>	1.0		43.6	11.9	10.3				66.8
<i>Trey phneik Lan</i>	<i>Priacanthus hamrur</i>			49.6	7.0				3.5	60.1
<i>Trey udouy</i>	<i>Selaroides leptolepis</i>	5.0		1.7	2.5	11.7	29.3	5.7	3.0	58.9
<i>Trey angrei</i>	<i>Sphyraena putnamae</i>					8.9	36.3	6.0	4.6	55.8
<i>Kdam skor krorb/Kdam sor</i>	<i>Matuta victor</i>			10.5	25.8	6.3				42.6
<i>Trey kantang kra-ob</i>	<i>Siganus canaliculatus</i>			13.6	6.0	8.0	3.0	1.0	2.8	34.4
<i>Trey chhlam chhkout</i>	<i>Chiloscyllium griseum</i>			4.5	5.6	9.7	2.0		1.9	23.7
<i>Trey prorkong</i>	(blank)				18.5					18.5
<i>Kdam Seih</i>	<i>Portunus pelagicus</i>	0.5	0.7	3.2	1.9	2.9	1.8	0.2	1.3	12.4
<i>Trey andeng</i>	(blank)			11.5						11.5
<i>Trey andeng poy</i>	(blank)					7.0				7.0
<i>Kdam neak</i>	<i>Charybdis natator</i>	5.0	1.0	1.0						7.0

<i>Trey trorsork</i>	<i>Terapon theraps</i>							7.0		7.0
<i>Meuk huysay</i>	<i>Sepioteuthis lessoniana</i>	1.0	3.3							4.3
<i>Trey deik koul</i>	(blank)							4.0		4.0
<i>Kdam krahorm</i>	(blank)		3.0							3.0
<i>Trey sach</i>	(blank)	1.0								1.0
<i>Trey kbork</i>	<i>Moolgarda seheli</i>								0.7	0.7
<i>Trey kalang</i>	<i>Scomberoides commersonianus</i>								0.6	0.6
<i>Trey balang</i>	<i>Rastrelliger faughni</i>								0.5	0.5
<i>Bangkorng kandob</i>	<i>Clorida decorata</i>			0.2						0.2
<i>X-others</i>	<i>X-others</i>	79.0	49.0	76.5	3.0		20.9	430.8	96.1	755.3
Total		396.0	347.0	448.2	442.5	623.6	635.2	932.5	695.1	4,520

Fisher ID: 2

Local Name	Scientific Name	Catch (kg)								
		Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Total
<i>Kdam Seih</i>	<i>Portunus pelagicus</i>	78	157	111	137	172	173	134	85	1,046.1
<i>Khyorng thnot</i>	<i>Conus vexillum</i>	29	51	41	78	85	109	93	55	540.6
<i>Khyorng dong</i>	<i>Melo melo</i>	7	21	27	46	45	24	47	25	241.7
<i>Trey chhlam chhkout</i>	<i>Chiloscyllium griseum</i>	3	14	2	3	10	2	5	2	40.4
<i>Trey borbel moan</i>	<i>Himantura imbricata</i>				5	6	16	6	3	35.2
<i>Trey borbel ork</i>	<i>Aetobatus narinari</i>	3	13	2						16.5
<i>Trey kam kourch</i>	<i>Carangoides ferdau</i>		5							4.5
<i>Trey chenchas/pakchiv</i>	<i>Alectis ciliaris</i>								2	2.2
Total		119	260	182	269	317	325	284	171	1,927

Fisher ID: 3

Local Name	Scientific Name	Catch (kg)								
		Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Total
<i>Kdam Seih</i>	<i>Portunus pelagicus</i>	52.0	62.2	63.0	25.0	39.3	41.6	90.5	102.5	476.1
<i>Bangkorng kandob</i>	<i>Clorida decorata</i>			1.6	6.3	17.1	19.3	25.0	14.3	83.5
<i>Kdam krahorm</i>	(blank)	8.5	32.7	20.1	5.9	2.8		5.0	1.6	76.5
<i>Kdam skor krorb/Kdam sor</i>	<i>Matuta victor</i>		4.0		10.3	12.7	9.5	11.7	8.5	56.6
<i>Kdam Thmor</i>	<i>Scylla serrata</i>	5.8	7.9	6.4	1.6	1.5	1.1	1.2		25.4
<i>Bangkea sor</i>	(blank)	6.1	3.1	4.0		1.5	1.7		5.3	21.7
<i>Trey antanou</i>	<i>Pentapodus setosus</i>								19.4	19.4
<i>Bangkorng pak</i>	(blank)	0.4	1.0		1.2	3.9	8.6	1.3	0.3	16.6
<i>Kdam Sor/Dangkeab Sor</i>	<i>Charybdis anisodon</i>			9.0						9.0
<i>Kdam neak</i>	<i>Charybdis natator</i>			0.5	0.4	3.7		0.7	2.5	7.8
<i>Bangkea kor gneub</i>	<i>Penaeus merguensis</i>		1.3				2.3			3.6
<i>Trey Kor/Romeas</i>	<i>Monacanthus chinensis</i>								2.6	2.6
<i>Kdam pingpeang</i>	<i>Arcania undecimspinosa</i>			0.4	1.0					1.4
<i>Meuk bompong/meit</i>	<i>Uroteuthis duvaucelii</i>							0.5	0.7	1.2
<i>Khyorng thnot</i>	<i>Conus vexillum</i>					0.3				0.3
Total		73	112	105	52	83	84	136	158	801

Fisher ID: 4

Local Name	Scientific Name	Catch (kg)								
		Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Total
<i>Bangkea kor gneub</i>	<i>Penaeus merguensis</i>	20.5	6.9	26.1	18.0	26.6	21.6	13.3	15.5	148.5
<i>Bangkea sor</i>	(blank)	10.1	1.5	0.5	1.9	7.9	4.3	4.3	8.0	38.5
<i>Trey prorkong</i>	(blank)					5.4	3.0			8.4
<i>Trey mrech</i>	(blank)					6.9	0.5			7.4
<i>Bangkea para/kloy</i>	<i>Penaeus semisulcatus</i>	0.5		3.9	0.7	0.2				5.3

<i>Trey phtong prortkheat</i>	<i>Ablennes hians</i>				0.7	1.0	3.1			4.8
<i>Kdam Thmor</i>	<i>Scylla serrata</i>	1.0	0.2					0.4	2.0	3.6
<i>Trey angkeuy</i>	<i>Lutjanus johnii</i>			1.0						1.0
<i>Meuk bompong/meit</i>	<i>Uroteuthis duvaucelii</i>								0.4	0.4
<i>X-others</i>	<i>X-others</i>	9.9	2.6	8.8	10.8	5.2	9.9	8.9	6.7	62.8
Total		42	11	40	32	53	42	27	33	281

Fisher ID: 5

Local Name	Scientific Name	Catch (kg)								
		Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Total
<i>Trey mrech</i>	(blank)	3.5	24.9	107	30.2	15.9	15.7	5.1	17.9	220.2
<i>Trey prorkong</i>	(blank)		3	42.1	11.3	7.7	6.3	1	12.1	83.5
<i>Bangkea sor</i>	(blank)	8.5	16	4.2	3.8	1.4	3.6	6.8	19.9	64.2
<i>Bangkea phourk</i>	(blank)	2	17.3	3.6	4.1	3.5	4.3	1	8.8	44.6
<i>Trey damlong</i>	(blank)			0.6	7.3	5.2	5	5.9	14.5	38.5
<i>Trey kanchanh chras</i>	(blank)				16.8	4.8	6.1		4.5	32.2
<i>Trey kranh</i>	<i>Lobotes surinamensis</i>		3.5	0.7	4.4	3.2	3.5	1	2	18.3
<i>Kdam Thmor</i>	<i>Scylla serrata</i>	2.4	2.2					2.5	7.6	14.7
<i>Bangkea kor gneub</i>	<i>Penaeus merguensis</i>							1.5	7.4	8.9
<i>Trey prorlos</i>	<i>Sillago sihama</i>		6.7	2.2						8.9
<i>Trey chek</i>	<i>Seriolina nigrofasciata</i>					0.5	0.4		7.8	8.7
<i>Kdam Seih</i>	<i>Portunus pelagicus</i>		3.5							3.5
<i>Trey kantong kra-ob</i>	<i>Siganus canaliculatus</i>					0.8			1.8	2.6
<i>Trey do angkor</i>	<i>Gerres abbreviatus</i>		2							2.0
<i>Trey trorsork</i>	<i>Terapon theraps</i>								2	2.0
<i>Trey andeng</i>	(blank)		0.8			0.3	0.8			1.9
<i>Trey andeng poy</i>	(blank)					0.8				0.8
<i>Bangkea para/kloy</i>	<i>Penaeus semisulcatus</i>							0.5		0.5

<i>Trey kbork</i>	<i>Moolgarda seheli</i>					0.4				0.4
<i>X-others</i>	<i>X-others</i>	4.2	63.8	41.6	9.9	6.2	4.2		3	132.9
Total		21	144	202	88	51	50	25	109	689

Fisher ID: 6

Local Name	Scientific Name	Catch (kg)								
		Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Total
<i>Trey kamong khoun khley</i>	<i>Rastrelliger brachysoma</i>	35	340	2330	2855	625	2325	2025	2172	12,707.0
<i>Trey kamong khoun veng</i>	<i>Rastrelliger kanagurta</i>	312	487	80	610	1580	425	528	935	4,957.0
<i>Trey krorthorm kbalkor</i>	(blank)		260	155				489	790	1,694.0
<i>Trey krorthorm srakakras</i>	<i>Scolopsis taeniopterus</i>				215	290	155	70	115	845.0
<i>Trey krorthorm</i>	<i>Lutjanus malabaricus</i>		20		230	640				890.0
<i>Trey beika</i>	<i>Auxis thazard</i>	4.2	35	19.3	1	22	3.5	35	3	123.0
<i>Trey krahorm phneak thom</i>	<i>Priacanthus macracanthus</i>		77	150			5	30		262.0
<i>Trey Kor/Romeas</i>	<i>Monacanthus chinensis</i>						257			257.0
<i>Trey krorthorm phneklien</i>	<i>Priacanthus tayenus</i>		5		45	80				130.0
<i>Bangkea sor</i>	(blank)	2.5	3	15.5						21.0
<i>Trey karav</i>	<i>Eleutheronema tetradactylum</i>	23		2						25.0
<i>Bangkea para/kloy</i>	<i>Penaeus semisulcatus</i>			5.2						5.2
<i>Trey borbel ork</i>	<i>Aetobatus narinari</i>		5							5.0
<i>Kdam Seih</i>	<i>Portunus pelagicus</i>	1	0.5							1.5
<i>Trey kamoy</i>	<i>Anodontostoma chacunda</i>	1.5								1.5
<i>X-others</i>	<i>X-others</i>	81	56.5	152	140	158	196	131	128	1,042.5
Total		460	1,289	2,909	4,096	3,395	3,367	3,308	4,143	22,967

Fisher ID: 7

Local Name	Scientific Name	Catch (kg)								
		Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Total
<i>Kdam Seih</i>	<i>Portunus pelagicus</i>	307.4	292.9	246.1	327	421.1	362.6	181.1	299.3	2,437.5
<i>Kdam Chor</i>	<i>Episesarma singaporensis</i>								1.5	1.5
<i>Kdam skor krorb/Kdam sor</i>	<i>Matuta victor</i>								1	1.0
Total		307.4	292.9	246.1	327	421.1	362.6	181.1	301.8	2440

Fisher ID: 8

Local Name	Scientific Name	Catch (kg)								
		Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Total
<i>Bangkea sor</i>	(blank)	56.2	65.2	8.1			5	57	49	240.5
<i>Trey kbork</i>	<i>Moolgarda seheli</i>	0.5	5.2	26.2	40.75	24.45	28.4	3		128.5
<i>Bangkea kor gneub</i>	<i>Penaeus merguensis</i>						37.5	20.5	50.5	108.5
<i>Trey phtong phka</i>	<i>Hemiramphus far</i>			5	22	21	13.2	16	9.7	86.9
<i>Trey kantang kra-ob</i>	<i>Siganus canaliculatus</i>	0.8	1.2	17.6	41.75	16.55				77.9
<i>Trey kantang thmor</i>	<i>Siganus virgatus</i>				29.8	14.9		3.5		48.2
<i>Trey pthong prort</i>	<i>Tylosurus acus melanotus</i>			27.4						27.4
<i>Trey prorlos</i>	<i>Sillago sihama</i>			4	5.8	1.6				11.4
<i>Kdam Thmor</i>	<i>Scylla serrata</i>			0.25	0.4		0.7	3.1	5	9.5
<i>Trey karav</i>	<i>Eleutheronema tetradactylum</i>			0.6	4.8	2.9	0.1			8.4
<i>Trey mrech</i>	(blank)			1.5	5					6.5
<i>Trey damlong</i>	(blank)		1.85	1.95			0.2	0.8	0.4	5.2
<i>Trey kantang phleung</i>	<i>Siganus guttatus</i>			0.1	0.2	2		0.2		2.5
<i>Trey phtong prortkheat</i>	<i>Ablennes hians</i>			1.5	0.35		0.1		0.5	2.5
<i>Trey kantang prorpeh</i>	<i>Siganus argenteus</i>			0.2	0.95	0.9		0.3		2.4
<i>Trey deik koul</i>	(blank)		1.25	0.1						1.4
<i>Kdam Seih</i>	<i>Portunus pelagicus</i>		1.15	0.1						1.3

<i>Trey Krorb khnol</i>	<i>Lethrinus nebulosus</i>		0.9	0.1						1.0
<i>Trey andeng poy</i>	(blank)						0.7			0.7
<i>Trey prorkong</i>	(blank)						0.7			0.7
<i>Bangkea para/kloy</i>	<i>Penaeus semisulcatus</i>							0.3	0.2	0.5
<i>Trey kbork angkam</i>	<i>Valamugil buchanani</i>				0.5					0.5
<i>Trey do angkor</i>	<i>Gerres abbreviatus</i>			0.5						0.5
<i>Bangkorng pak</i>	(blank)			0.1		0.1				0.2
<i>Trey khej/proy veng</i>	<i>Pentaprion longimanus</i>				0.2					0.2
<i>X-others</i>	<i>X-others</i>	18.2	19.6	8.05	19.4	27	19.6	30.5	43.2	185.6
Total		76	96	103	172	111	106	135	159	959

Fisher ID: 9

Local Name	Scientific Name	Catch (kg)								
		Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Total
<i>Trey kamong khoun khley</i>	<i>Rastrelliger brachysoma</i>	462.0	859.0	1,988.0	7,533.0	2,121.0	2,594.0	2,460.0	2,021.0	20,038.0
<i>Trey karav</i>	<i>Eleutheronema tetradactylum</i>	12.5	55.0							67.5
<i>Trey chhlam chhkout</i>	<i>Chiloscyllium griseum</i>	12.0	15.0							27.0
<i>Bangkea para/kloy</i>	<i>Penaeus semisulcatus</i>			6.0						6.0
<i>Kdam Thmor</i>	<i>Scylla serrata</i>	2.2	1.9							4.1
<i>Bangkea sor</i>	(blank)			3.5						3.5
<i>X-others</i>	<i>X-others</i>	19.0	59.0	125.0	179.0	225.0	208.0	212.0	344.5	1,371.5
Total		508	990	2,123	7,712	2,346	2,802	2,672	2,366	21,518

Fisher ID: 10

Local Name	Scientific Name	Catch (kg)								
		Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Total
<i>Trey kbork</i>	<i>Moolgarda seheli</i>	1	7.5	47.3	93.3	47.5	38.1		7	241.7
<i>Bangkea kor gneub</i>	<i>Penaeus merguensis</i>						14	70.6	66.5	151.1

<i>Bangka sor</i>	(blank)	58.3	64.5	7		2		4		135.8
<i>Trey kantong kra-ob</i>	<i>Siganus canaliculatus</i>	0.3	2.6	45.7	5		1.5			55.1
<i>Trey prorleuk</i>	<i>Leiognathus splendens</i>					12		11.9	24.5	48.4
<i>Trey kantong thmor</i>	<i>Siganus javus</i>				29.5	4.5		0.3		34.3
<i>Bangka para/kloy</i>	<i>Penaeus semisulcatus</i>		12.5			2	16	2.3	1	33.8
<i>Trey damlong</i>	(blank)	4.2	14.8	3			7.5			29.5
<i>Trey mrech</i>	(blank)			8	16.3	1				25.3
<i>Kdam Seih</i>	<i>Portunus pelagicus</i>			12.3						12.3
<i>Kdam Thmor</i>	<i>Scylla serrata</i>	0.3	0.8	0.8	0.6	0.2	2.4	5.3	1.7	12.1
<i>Trey deik koul</i>	(blank)		8.05	1						9.1
<i>Trey prorlos</i>	<i>Sillago sihama</i>			6.8						6.8
<i>Trey pthong prort</i>	<i>Tylosurus acus melanotus</i>	0.6	3.8	0.8						5.2
<i>Trey kantong phleung</i>	<i>Siganus guttatus</i>				2		2.9			4.9
<i>Trey prorkong</i>	(blank)				2		2.5			4.5
<i>Trey Krorb khnol</i>	<i>Lethrinus nebulosus</i>			4.5						4.5
<i>Trey karav</i>	<i>Eleutheronema tetradactylum</i>			0.3	0.8				2.8	3.9
<i>Trey kantong prorpeh</i>	<i>Siganus argenteus</i>					2.5				2.5
<i>Trey kbork angkam</i>	<i>Moolgarda buchanani</i>				1			1		2.0
<i>Trey kanchanh chras</i>	(blank)				2					2.0
<i>Trey spong</i>	<i>Psammoperca waigiensis</i>					2				2.0
<i>Trey phtong phka</i>	<i>Hemiramphus far</i>				0.5			1		1.5
<i>Trey khej/proy veng</i>	<i>Pentaprion longimanus</i>				0.3				1	1.3
<i>Meuk snok</i>	<i>Sepiella inermis</i>	0.9	0.3							1.2
<i>Trey ka-ok</i>	<i>Arius maculatus</i>			1						1.0
<i>Trey andeng poy</i>	(blank)			1						1.0
<i>Trey andat chhke</i>	<i>Paraplagusia bilineata</i>						1			1.0
<i>Trey balang</i>	<i>Rastrelliger faughni</i>								1	1.0
<i>Trey do angkor</i>	<i>Gerres abbreviatus</i>		1							1.0

<i>Trey angrei</i>	<i>Sphyraena putnamae</i>					0.9				0.9
<i>Trey kantong thmor</i>	<i>Siganus virgatus</i>						0.3			0.3
<i>Meuk bompong/meit</i>	<i>Uroteuthis duvaucelii</i>		0.2							0.2
<i>X-others</i>	<i>X-others</i>	18.4	26.6	29.2	40.5	40.8	28.5	33.8	15.5	233.3
Total		84	143	169	194	115	115	130	121	1,070

Fisher ID: 11

Local Name	Scientific Name	Catch (kg)								
		Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Total
<i>Trey seik</i>	<i>Scarus spp.</i>	3.7	7	25	46	57	45	85	104	372.7
<i>Trey phneik Lan</i>	<i>Priacanthus hamrur</i>	0.25	2.5	7	11	21	33	50	67	191.8
<i>Trey teukei ouch kiev</i>	<i>Plectropomus oligacanthus</i>	1	3	12	20	24	24	52	51	187.0
<i>Trey kantong phleung</i>	<i>Siganus guttatus</i>	3.55	5.4	21	32	35	51	8	28	184.0
<i>Trey kantong kra-ob</i>	<i>Siganus canaliculatus</i>						36	56	45	137.0
<i>Trey seik thmor</i>	<i>Scarus spp.</i>	0.25					22	38	53	113.3
<i>Trey kantong prorpeh</i>	<i>Siganus argenteus</i>	2.15	2.5		11	18			68	101.7
<i>Trey teukei para</i>	<i>Epinephelus quoyanus</i>	0.5		6	8	7.5	8	11	9	50.0
<i>Trey teukei koav</i>	<i>Epinephelus coioides</i>		0.5				13	15	17	45.5
<i>Trey teukeokao</i>	(blank)				9	10				19.0
<i>Ngveav sor</i>	<i>Periglypta sp.</i>			13						13.0
<i>Trey teukei para krorthorm</i>	<i>Epinephelus fasciatus</i>			7		2				9.0
<i>Kdam neak</i>	<i>Charybdis natator</i>							3		3.0
<i>Trey teukei thmor</i>	<i>Aethaloperca rogaea</i>	0.5								0.5
<i>X-others</i>	<i>X-others</i>			1.5						1.5
Total		12	21	93	137	175	232	318	442	1,429

Fisher ID: 12

LocalName	Scientific Name	Catch (kg)								
		Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Total
<i>Meuk huysay</i>	<i>Sepioteuthis lessoniana</i>	47.1	108	72.3	93.2	85.3	65.3	127.2	119.3	717.7
<i>Kdam Thmor</i>	<i>Scylla serrata</i>		0.5			5.6				6.1
<i>Kdam Seih</i>	<i>Portunus pelagicus</i>		2.5			3.5				6.0
<i>Meuk bompong/meit</i>	<i>Uroteuthis duvaucelii</i>	0.5	4.1							4.6
<i>Kdam Sor/Dangkeab Sor</i>	<i>Charybdis anisodon</i>					2				2.0
<i>Kdam krahorm</i>	(blank)		1							1.0
Total		48	116	72	93	96	65	127	119	737

Fisher ID: 13

LocalName	Scientific Name	Catch (kg)								
		Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Total
<i>Meuk huysay</i>	<i>Sepioteuthis lessoniana</i>	20.4	47.4	28.8	28.2	20.1	15.3	32.8	33.8	226.8
<i>Meuk bompong/meit</i>	<i>Uroteuthis duvaucelii</i>	0.2	4.6	3.1	3.3			1.6	1.2	14.0
Total		21	52	32	32	20	15	34	35	241

Fisher ID: 14

LocalName	Scientific Name	Catch (kg)								
		Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Total
<i>Trey seik</i>	<i>Scarus spp.</i>	1.5	10.7	10.6	15.1	21.5	20.5	27.4	26.6	133.9
<i>Trey teukei ouch kiev</i>	<i>Plectropomus oligacanthus</i>	0.5	16.9	7.5	12.6	16.4	16.4	19	28.3	117.6
<i>Trey kantong phleung</i>	<i>Siganus guttatus</i>	0.5	4.2	10.7	11.2	19.4	19.1	21.2	22.4	108.7
<i>Trey teukei para krorhorm</i>	<i>Epinephelus fasciatus</i>	0.3	3	3.2	6.3	9	9.9	11.9	11.6	55.2
<i>Trey ka chy</i>	<i>Diagramma pictum</i>		1.2	3.6	6.9	7.1	8.4	8.1	11.8	47.1
<i>Trey teukei para</i>	<i>Epinephelus quoyanus</i>	0.2	0.7	0.6		5.9	16.8	8.8	8.1	41.1
<i>Trey kantong kra-ob</i>	<i>Siganus canaliculatus</i>				3.7					3.7

<i>Trey borbel ork</i>	<i>Aetobatus narinari</i>							2.7		2.7
<i>Trey borbel khleing</i>	<i>Mobula japanica</i>		2.3							2.3
<i>Kdam Thmor</i>	<i>Scylla serrata</i>				1.6					1.6
<i>Trey teukeokao</i>	(blank)				1					1.0
Total		3	39	36	58	79	91	99	109	515

Fisher ID: 15

LocalName	Scientific Name	Catch (kg)								
		Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Total
<i>Meuk huysay</i>	<i>Sepioteuthis lessoniana</i>	17.5	164.6	55.2	130	121.1	50.1	32.8	163.8	735.1
<i>Meuk bompong/meit</i>	<i>Uroteuthis duvaucelii</i>	0.7	28.2	2.6	12.6	8.9	1.2		6	60.2
Total		18	193	58	143	130	51	33	170	795

Fisher ID: 16

LocalName	Scientific Name	Catch (kg)								
		Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Total
<i>Meuk dong</i>	<i>Sepiola sp.</i>		9	55	38	38	200	320	280	940.0
Total			9	55	38	38	200	320	280	940

Fisher ID: 17

LocalName	Scientific Name	Catch (kg)								
		Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Total
<i>Kdam Thmor</i>	<i>Scylla serrata</i>	23.7	37.8	43.3	50.7	60.1	65.6	61.7	58.9	401.8
Total		24	38	43	51	60	66	62	59	401.8

Fisher ID: 18

LocalName	Scientific Name	Catch (kg)								
		Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Total
<i>Kdam Thmor</i>	<i>Scylla serrata</i>	19.1	39.3	62.8	63.1	58.73	57.5	50.8	56.7	408.0
Total		19	39	63	63	59	58	51	57	408

Fisher ID: 19

LocalName	Scientific Name	Catch (kg)								
		Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Total
<i>Ngveav sor</i>	<i>Periglypta sp.</i>	172	272.5	203	215	176	236	286		1,561
<i>Kdam Thmor</i>	<i>Scylla serrata</i>							1.1		1
Total		172	273	203	215	176	236	287		1,562

Fisher ID: 20

LocalName	Scientific Name	Catch (kg)								
		Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Total
<i>Kdam Chor</i>	<i>Episesarma singaporensis</i>	29.1	40.4	49.1	53.9		47.7	33.7	52.4	306.3
<i>Kdam Thmor</i>	<i>Scylla serrata</i>	1.4	6.1	3.8	14.6		5.7	2.2	1.4	35.2
<i>Trey phtourk</i>	<i>Rachycentron canadum</i>					24.8				24.8
<i>Kreing chheam</i>	<i>Anadara granosa</i>	3.5	10							13.5
<i>Ngveav sor</i>	<i>Periglypta sp.</i>	5								5.0
<i>Bangkea sor</i>	(blank)	2								2.0
<i>Khyorng Krachorkseih</i>	(blank)	0.5		1.1						1.6
<i>Khyorng dong</i>	<i>Melo melo</i>							0.6		0.6
<i>Trey kantang kra-ob</i>	<i>Siganus canaliculatus</i>	0.6								0.6
Total		42	57	54	69	25	53	37	54	390

Fisher ID: 21

LocalName	Scientific Name	Catch (kg)								
		Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Total
<i>Kdam Chor</i>	<i>Episesarma singaporensis</i>	36.1	49.5	49.9	57.1	50.3	48.7	26.9	24.1	342.6
<i>Kdam Thmor</i>	<i>Scylla serrata</i>	3.1	3.4	4.1	17.3	6.4	3.7	1.3	0.8	40.1
<i>Ngeav phourk</i>	<i>Polymesoda erosa</i>		2							2.0
Total		39	55	54	74	57	52	28	25	385

Fisher ID: 22

LocalName	Scientific Name	Catch (kg)								
		Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Total
<i>Kdam Seih</i>	<i>Portunus pelagicus</i>		306.7	268.4	121.2	246.5	87.9	30.7	163.7	1,225.1
<i>Kdam Thmor</i>	<i>Scylla serrata</i>		93.2	109.45	134.4	94.1	96.7	168.5	78.9	775.3
<i>Kdam Sor/Dangkeab Sor</i>	<i>Charybdis anisodon</i>		53.9	63.2	67.8	55.5	55.3			295.7
<i>Kdam skor krorb/Kdam sor</i>	<i>Matuta victor</i>							76.8		76.8
<i>Bangkae sor</i>	(blank)								42.5	42.5
Total		0	454	441	323	396	240	276	285	2,415

Fisher ID: 23

LocalName	Scientific Name	Catch (kg)								
		Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Total
<i>Trey beika</i>	<i>Auxis thazard</i>			0.2				551.1	984.3	1,535.6
<i>Trey kamoy</i>	<i>Anodontostoma chacunda</i>		3	0.5	126.1	67.5	192.8	202.2	127	719.1
<i>Trey kamong khoun khley</i>	<i>Rastrelliger brachysoma</i>			168.6	75	12.8	196			452.4
<i>Trey Sambor Hear</i>	(blank)					53.5	79	40.8	68.1	241.4
<i>Trey sleuk russey</i>	<i>Thryssa hamiltonii</i>	3.75	4	1.3	3		32.5	19.8		64.4

<i>Bangka para/kloy</i>	<i>Penaeus semisulcatus</i>		0.5	0.75		57.3	3.2	1.2		63.0
<i>Bangka sor</i>	(blank)	8	40.2	1.1		3.7		2.3		55.3
<i>Trey khut/trorchiek damrey</i>	<i>Ehippus orbis</i>			32.9						32.9
<i>Trey andat chhke</i>	<i>Paraplagusia bilineata</i>				5.5	19.3				24.8
<i>Trey karav</i>	<i>Eleutheronema tetradactylum</i>			0.5	1		14.5	5.6		21.6
<i>Meuk dong</i>	<i>Sepiola sp.</i>								20	20.0
<i>Trey borbel moan</i>	<i>Himantura imbricata</i>		0.8		12					12.8
<i>Bangka kor gneub</i>	<i>Penaeus merguensis</i>								12.3	12.3
<i>Trey ka-ok</i>	<i>Arius maculatus</i>					5.6	5.2			10.8
<i>Trey udouy</i>	<i>Selaroides leptolepis</i>					8.1				8.1
<i>Trey andeng poy</i>	(blank)				4					4.0
<i>Trey damlong</i>	(blank)							1.3		1.3
<i>Kdam Seih</i>	<i>Portunus pelagicus</i>	0.5								0.5
<i>Meuk snok</i>	<i>Sepiella inermis</i>	0.4								0.4
<i>Trey Kor/Romeas</i>	<i>Monacanthus chinensis</i>	0.1								0.1
<i>X-others</i>	<i>X-others</i>	2.35	15.7	6.5	12.3	22.4				59.3
Total		15	64	212	239	250	523	824	1,212	3,340

Fisher ID: 24

LocalName	Scientific Name	Catch (kg)								
		Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Total
<i>Kdam Thmor</i>	<i>Scylla serrata</i>	89.15	252.7	142	163	131	111.9	107	150.7	1,147.5
<i>Kdam Seih</i>	<i>Portunus pelagicus</i>	92.9	347	132.1	130.2	104.1	83.4	91.7	108	1,089.4
<i>Kdam Sor/Dangkeab Sor</i>	<i>Charybdis anisodon</i>	22	206	104	110	76	54			572.0
<i>Bangka sor</i>	(blank)								78.9	78.9
<i>Kdam skor krorb/Kdam sor</i>	<i>Matuta victor</i>							56.35	10	66.4
Total		204	806	378	403	311	249	255	348	2,954

Fisher ID: 25

LocalName	Scientific Name	Catch (kg)								
		Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Total
<i>Kdam Seih</i>	<i>Portunus pelagicus</i>	11	67.3	27.6	19.1	1	5.8	19.5	10.83	162.1
<i>Kdam Thmor</i>	<i>Scylla serrata</i>	4.2	12.6	27.9	16.5	19.05	16.2	18.98	13.33	128.8
<i>Trey borbel moan</i>	<i>Himantura imbricata</i>		6.8	14.6	14.4	18.8	21.6	11.5	10.5	98.2
<i>Trey beika</i>	<i>Auxis thazard</i>			0.3		2.5	10.5	19.6	7	39.9
<i>Trey balang</i>	<i>Rastrelliger faughni</i>								24.5	24.5
<i>Trey kalang</i>	<i>Scomberoides commersonianus</i>							19.5		19.5
<i>Trey andat chhke</i>	<i>Paraplagusia bilineata</i>		1.2	1.1		7.1	2.9	4.6	0.6	17.5
<i>Trey Kor/Romeas</i>	<i>Monacanthus chinensis</i>			1.1	1	4	5.8			11.9
<i>Trey kamong khoun khley</i>	<i>Rastrelliger brachysoma</i>			4.8			4.4			9.2
<i>Trey karav</i>	<i>Eleutheronema tetradactylum</i>			0.4	1.5		3	0.5	1.1	6.5
<i>Trey kantuy krabey</i>	(blank)		1.2	1	1.2	2.5				5.9
<i>Trey ka-ok</i>	<i>Arius maculatus</i>			3.2	1.2	1				5.4
<i>Trey ka-ok kbal heang/Ka-ok prak</i>	<i>Osteogeneiosus militaris</i>								5.2	5.2
<i>Trey kantong kra-ob</i>	<i>Siganus canaliculatus</i>			0.4		3.2	1.3			4.9
<i>Trey andeng</i>	(blank)					1		3.4		4.4
<i>Trey sleuk russey</i>	<i>Thryssa hamiltonii</i>				0.3		2.1			2.4
<i>Kdam kom</i>	(blank)							1		1.0
<i>Kdam skor krorb/Kdam sor</i>	<i>Matuta victor</i>					1				1.0
X-others	X-others		2		9.9	8.83	3.4	12.3	2	38.4
Total		15	91	82	65	70	77	111	75	587

Fisher ID: 26

LocalName	Scientific Name	Catch (kg)								
		Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Total
<i>Kreing chheam</i>	<i>Anadara granosa</i>	20	34.9	41	28	38	26.5	20.7	25.9	235.0
<i>Trey kantang phleung</i>	<i>Siganus guttatus</i>				2					2.0
<i>Trey seik</i>	<i>Scarus spp.</i>				1.8					1.8
<i>Trey teukei ouch kiev</i>	<i>Plectropomus oligacanthus</i>				1.7					1.7
<i>Trey ka chy</i>	<i>Diagramma pictum</i>				1.5					1.5
<i>Trey teukei para krorthorm</i>	<i>Epinephelus fasciatus</i>				0.5					0.5
Total		20	35	41	36	38	27	21	26	242.5

**Annex 6: Monthly landing value (Riel) by species and by fishers over the period from April to November 2011
(ranking according to total value)**

Fisher ID: 1

Local Name	Scientific Name	Value (Riel)								
		Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Total
<i>Trey do angkor</i>	<i>Gerres abbreviatus</i>			880,250	689,250	1,221,300	1,471,200	351,500	1,276,000	5,889,500
<i>Trey prorleuk</i>	<i>Leiognathus splendens</i>						710,400	1,521,600	451,450	2,683,450
<i>Trey Krorb khnol</i>	<i>Lethrinus nebulosus</i>			14,000	333,750	534,250	496,850	189,650	79,000	1,647,500
<i>Trey prorlos</i>	<i>Sillago sihama</i>			307,500	296,500	757,800				1,361,800
<i>Trey krorhorm srakakras</i>	<i>Scolopis taeniopterus</i>						693,200	330,000	266,450	1,289,650
<i>Trey krorhorm kbalkor</i>	(blank)				307,150	786,350		28,150	83,350	1,205,000
<i>Trey kamoy</i>	<i>Anodontostoma chacunda</i>	458,350	608,700	29,200	2,700		4,500			1,103,450
<i>Trey kamong khoun khley</i>	<i>Rastrelliger brachysoma</i>	453,000	367,700	46,750						867,450
<i>Trey chek</i>	<i>Seriolina nigrofasciata</i>	193,000	72,000	38,500	174,500	74,750	57,500	5,000	125,000	740,250
<i>Trey udouy</i>	<i>Selaroides leptolepis</i>	15,000		9,150	19,000	87,100	235,750	45,600	18,000	429,600
<i>Trey Kor/Romeas</i>	<i>Monacanthus chinensis</i>				111,000	80,400	93,900	30,750	64,450	380,500
<i>Trey angrei</i>	<i>Sphyrnaena putnamae</i>					88,100	217,300	31,000	36,800	373,200
<i>Trey krorhorm</i>	<i>Lutjanus malabaricus</i>	4,000		162,850	89,200	78,700				334,750
<i>Trey phneik Lan</i>	<i>Priacanthus hamrur</i>			220,450	36,500				7,000	263,950
<i>Trey koun</i>	<i>Hilsa keele</i>	62,500	61,500	12,000				83,900	18,650	238,550
<i>Trey kantang kra-ob</i>	<i>Siganus canaliculatus</i>			59,900	36,750	38,200	13,500	5,500	13,600	167,450
<i>Trey chhlam chhkout</i>	<i>Chiloscyllium griseum</i>			23,000	34,000	59,200	14,000		7,600	137,800
<i>Trey andeng</i>	(blank)			95,250						95,250
<i>Kdam Seih</i>	<i>Portunus pelagicus</i>	2,500	5,600	28,400	3,585	28,000	10,100	600	10,150	88,935
<i>Trey prorkong</i>	(blank)				87,750					87,750
<i>Kdam skor krorb/Kdam sor</i>	<i>Matuta victor</i>			24,000	27,800	10,340				62,140
<i>Trey andeng poy</i>	(blank)					49,000				49,000
<i>Kdam neak</i>	<i>Charybdis natator</i>	40,000	1,000	4,900						45,900

<i>Meuk huysay</i>	<i>Sepioteuthis lessoniana</i>	1,500	43,600							45,100
<i>Kdam krahorm</i>	(blank)		23,000							23,000
<i>Trey deik koul</i>	(blank)							8,000		8,000
<i>Trey trorsork</i>	<i>Terapon theraps</i>							7,000		7,000
<i>Trey sach</i>	(blank)	6,000								6,000
<i>Trey kalang</i>	<i>Scomberoides commersonianus</i>								4,800	4,800
<i>Trey balang</i>	<i>Rastrelliger faughni</i>								4,000	4,000
<i>Bangkorng kandob</i>	<i>Clorida decorata</i>			2,000						2,000
<i>Trey kbork</i>	<i>Moolgarda seheli</i>								700	700
<i>X-others</i>	<i>X-others</i>	113,700	59,200	97,000	4,500		39,300	674,090	68,300	1,056,090
Total		1,349,550	1,242,300	2,055,100	2,253,935	3,893,490	4,057,500	3,312,340	2,535,300	20,699,515

Fisher ID: 2

Local Name	Scientific Name	Value (Riel)								
		Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Total
<i>Kdam Seih</i>	<i>Portunus pelagicus</i>	1,230,000	2,453,000	1,927,700	2,732,900	2,929,200	2,896,300	2,386,800	1,599,800	18,155,700
<i>Khyorng thnot</i>	<i>Conus vexillum</i>	90,000	168,700	133,910	280,960	342,650	546,000	509,300	273,000	2,344,520
<i>Khyorng dong</i>	<i>Melo melo</i>	45,500	142,500	193,500	353,650	368,850	212,850	420,300	225,000	1,962,150
<i>Trey chhlam chhkout</i>	<i>Chiloscyllium griseum</i>	19,950	95,000	11,250	18,400	76,000	15,400	47,000	15,000	298,000
<i>Trey borbel moan</i>	<i>Himantura imbricata</i>				20,050	22,500	76,500	23,200	11,600	153,850
<i>Trey borbel ork</i>	<i>Aetobatus narinari</i>	9,000	62,500	6,000						77,500
<i>Trey kam kourch</i>	<i>Carangoides ferdau</i>		36,000							36,000
<i>Trey chenchas/pakchiv</i>	<i>Alectis ciliaris</i>								21,000	21,000
Total		1,394,450	2,957,700	2,272,360	3,405,960	3,739,200	3,747,050	3,386,600	2,145,400	23,048,720

Fisher ID: 3

Local Name	Scientific Name	Value (Riel)								
		Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Total
<i>Kdam Seih</i>	<i>Portunus pelagicus</i>	958,000	1,144,200	1,445,830	325,660	681,530	829,690	1,878,020	4,379,300	11,642,230
<i>Bangkorng kandob</i>	<i>Clorida decorata</i>			7,500	33,475	187,650	347,525	354,630	220,100	1,150,880
<i>Kdam Thmor</i>	<i>Scylla serrata</i>	103,820	99,840	79,400	13,280	12,925	2,425	10,345		322,035
<i>Bangkorng pak</i>	(blank)	3,200	5,600		32,800	75,430	162,300	17,150	3,125	299,605
<i>Kdam krahorm</i>	(blank)	28,000	98,780	94,779	16,580	4,145		7,105	1,265	250,654
<i>Trey antanou</i>	<i>Pentapodus setosus</i>								151,340	151,340
<i>Bangkea sor</i>	(blank)	61,000	15,800	14,050		22,500	28,900		4,045	146,295
<i>Kdam skor krorb/Kdam sor</i>	<i>Matuta victor</i>		18,200		19,120	18,035	19,145	10,820	32,170	117,490
<i>Kdam neak</i>	<i>Charybdis natator</i>			1,130	1,600	16,900		2,900	19,500	42,030
<i>Bangkea kor gneub</i>	<i>Penaeus merguensis</i>		11,200				25,700			36,900
<i>Kdam Sor/Dangkeab Sor</i>	<i>Charybdis anisodon</i>			14,930						14,930
<i>Meuk bompong/meit</i>	<i>Uroteuthis duvaucelii</i>							2,000	4,550	6,550
<i>Trey Kor/Romeas</i>	<i>Monacanthus chinensis</i>								6,220	6,220
<i>Khyorng thnot</i>	<i>Conus vexillum</i>					125				125
<i>Kdam pingpeang</i>	<i>Arcania undecimspinosa</i>			-	-	-				
Total		1,154,020	1,393,620	1,657,619	442,515	1,019,240	1,415,685	2,282,970	4,821,615	14,187,283

Fisher ID: 4

Local Name	Scientific Name	Value (Riel)								
		Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Total
<i>Bangkea kor gneub</i>	<i>Penaeus merguensis</i>	162,130	42,360	301,820	127,570	204,150	144,990	110,070	84,390	1,177,480
<i>Bangkea sor</i>	(blank)	87,120	7,700	1,750	9,580	57,550	20,300	15,480	56,250	255,730
<i>Trey prorkong</i>	(blank)					20,500	29,000			49,500
<i>Bangkea para/kloy</i>	<i>Penaeus semisulcatus</i>	2,000		28,160	3,200	800				34,160
<i>Trey phtong prortkheat</i>	<i>Ablennes hians</i>				2,800	7,000	20,550			30,350

<i>Trey mrech</i>	(blank)					25,970	750			26,720
<i>Kdam Thmor</i>	<i>Scylla serrata</i>	3,600	840					1,760	10,630	16,830
<i>Trey angkeuy</i>	<i>Lutjanus johnii</i>			15,000						15,000
<i>Meuk bompong/meit</i>	<i>Uroteuthis duvaucelii</i>								1,920	1,920
<i>X-others</i>	<i>X-others</i>	9,740	2,360	9,120	15,350	5,480	16,300	21,900	5,900	86,150
Total		264,590	53,260	355,850	158,500	321,450	231,890	149,210	159,090	1,693,840

Fisher ID: 5

Local Name	Scientific Name	Value (Riel)								
		Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Total
<i>Trey mrech</i>	(blank)	14,650	81,700	319,400	90,600	50,500	50,800	21,600	78,900	708,150
<i>Bangkea sor</i>	(blank)	84,000	150,000	30,700	24,900	9,200	23,000	61,200	196,000	579,000
<i>Bangkea phourk</i>	(blank)	14,000	116,000	21,000	24,700	16,300	22,000	8,000	78,400	300,400
<i>Trey prorkong</i>	(blank)		11,500	124,800	33,900	27,100	24,400	5,000	72,600	299,300
<i>Kdam Thmor</i>	<i>Scylla serrata</i>	48,000	35,200					48,700	96,500	228,400
<i>Bangkea kor gneub</i>	<i>Penaeus merguensis</i>							9,300	160,680	169,980
<i>Trey damlong</i>	(blank)			1,800	21,900	16,800	16,900	28,500	79,200	165,100
<i>Trey kanchanh chras</i>	(blank)				36,500	9,750	11,450		5,500	63,200
<i>Trey kranh</i>	<i>Lobotes surinamensis</i>		9,900	1,400	13,200	10,300	9,800	3,000	6,000	53,600
<i>Kdam Seih</i>	<i>Portunus pelagicus</i>		33,800							33,800
<i>Trey prorlos</i>	<i>Sillago sihama</i>		21,300	8,100						29,400
<i>Trey chek</i>	<i>Seriolina nigrofasciata</i>					1,500	1,200		23,750	26,450
<i>Trey andeng</i>	(blank)		5,800			1,800	4,700			12,300
<i>Trey kantang kra-ob</i>	<i>Siganus canaliculatus</i>					3,700			7,200	10,900
<i>Trey do angkor</i>	<i>Gerres abbreviatus</i>		8,000							8,000
<i>Trey andeng poy</i>	(blank)					5,300				5,300
<i>Bangkea para/kloy</i>	<i>Penaeus semisulcatus</i>							5,000		5,000
<i>Trey trorsork</i>	<i>Terapon theraps</i>								4,000	4,000

<i>Trey kbork</i>	<i>Moolgarda seheli</i>					1,600				1,600
<i>X-others</i>	<i>X-others</i>	14,600	186,100	87,700	21,080	10,300	7,900		8,000	335,680
Total		175,250	659,300	594,900	266,780	164,150	172,150	190,300	816,730	3,039,560

Fisher ID: 6

Local Name	Scientific Name	Value (Riel)								
		Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Total
<i>Trey kamong khoun khley</i>	<i>Rastrelliger brachysoma</i>	90,000	1,005,000	3,853,000	4,587,500	1,857,500	6,975,000	7,087,500	7,602,000	33,057,500
<i>Trey kamong khoun veng</i>	<i>Rastrelliger kanagurta</i>	668,000	2,008,000	400,000	3,050,000	7,900,000	2,125,000	2,640,000	4,375,000	23,166,000
<i>Trey krorthorm kbalkor</i>	(blank)		465,000	232,500				1,222,500	1,975,000	3,895,000
<i>Trey krorthorm srakakras</i>	<i>Scolopis taeniopterus</i>				682,500	975,000	462,500	266,000	402,500	2,788,500
<i>Trey krorthorm</i>	<i>Lutjanus malabaricus</i>		40,000		625,000	1,630,000				2,295,000
<i>Trey beika</i>	<i>Auxis thazard</i>	18,600	175,000	113,800	8,000	176,000	28,000	280,000	24,000	823,400
<i>Trey krahorn phneak thom</i>	<i>Priacanthus macracanthus</i>		231,000	450,000			20,000	120,000		821,000
<i>Trey Kor/Romeas</i>	<i>Monacanthus chinensis</i>						642,500			642,500
<i>Trey krorthorm phneklien</i>	<i>Priacanthus tayenus</i>		12,500		180,000	320,000				512,500
<i>Bangkae sor</i>	(blank)	28,000	37,500	236,000						301,500
<i>Trey karav</i>	<i>Eleutheronema tetradactylum</i>	46,000		13,000						59,000
<i>Bangkae para/kloy</i>	<i>Penaeus semisulcatus</i>			52,000						52,000
<i>Trey borbel ork</i>	<i>Aetobatus narinari</i>		40,000							40,000
<i>Kdam Seih</i>	<i>Portunus pelagicus</i>	12,000	5,000							17,000
<i>Trey kamoy</i>	<i>Anodontostoma chacunda</i>	2,250								2,250
<i>X-others</i>	<i>X-others</i>	55,700	70,150	117,100	113,400	187,500	196,000	131,000	128,000	998,850
Total		920,550	4,089,150	5,467,400	9,246,400	13,046,000	10,449,000	11,747,000	14,506,500	69,472,000

Fisher ID: 7

Local Name	Scientific Name	Value (Riel)								
		Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Total

<i>Kdam Seih</i>	<i>Portunus pelagicus</i>	2,962,650	2,848,350	2,324,200	3,698,750	5,139,900	4,168,800	1,601,600	2,628,150	25,372,400
<i>Kdam skor krorb/Kdam sor</i>	<i>Matuta victor</i>								12,000	12,000
<i>Kdam Chor</i>	<i>Episesarma singaporensis</i>								6,750	6,750
Total		2,962,650	2,848,350	2,324,200	3,698,750	5,139,900	4,168,800	1,601,600	2,646,900	25,391,150

Fisher ID: 8

Local Name	Scientific Name	Value (Riel)								
		Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Total
<i>Bangkae sor</i>	(blank)	286,200	254,800	32,400			25,000	338,000	252,000	1,188,400
<i>Trey kbork</i>	<i>Moolgarda seheli</i>	500	31,200	160,200	275,600	171,150	196,300	21,000		855,950
<i>Bangkae kor gneub</i>	<i>Penaeus merguensis</i>						187,500	117,500	291,500	596,500
<i>Trey kantang kra-ob</i>	<i>Siganus canaliculatus</i>	800	2,000	38,700	96,600	41,975				180,075
<i>Trey phtong phka</i>	<i>Hemiramphus far</i>			10,000	44,000	43,000	26,400	32,000	19,400	174,800
<i>Trey kantang thmor</i>	<i>Siganus virgatus</i>				69,900	42,800		10,500		123,200
<i>Kdam Thmor</i>	<i>Scylla serrata</i>			2,500	10,000		8,200	43,100	57,600	121,400
<i>Trey phtong prort</i>	<i>Tylosurus acus melanotus</i>			74,300						74,300
<i>Trey karav</i>	<i>Eleutheronema tetradactylum</i>			1,400	9,850	11,000	500			22,750
<i>Trey kantang prorpeh</i>	<i>Siganus argenteus</i>			2,000	5,400	9,000		3,000		19,400
<i>Trey kantang phleung</i>	<i>Siganus guttatus</i>			800	400	15,500		2,000		18,700
<i>Trey prorlos</i>	<i>Sillago sihama</i>			4,300	8,700	1,600				14,600
<i>Bangkae para/kloy</i>	<i>Penaeus semisulcatus</i>							3,600	6,000	9,600
<i>Trey mrech</i>	(blank)			2,250	5,000					7,250
<i>Trey phtong prortkheat</i>	<i>Ablennes hians</i>			4,300	700		300		1,000	6,300
<i>Trey damlong</i>	(blank)		1,850	1,950			200	800	400	5,200
<i>Kdam Seih</i>	<i>Portunus pelagicus</i>		4,750	400						5,150
<i>Bangkornng pak</i>	(blank)			3,000		2,000				5,000
<i>Trey andeng poy</i>	(blank)						3,000			3,000
<i>Trey kbork angkam</i>	<i>Moolgarda buchanani</i>				3,000					3,000

<i>Trey prorkong</i>	(blank)						2,800			2,800
<i>Trey Krorb khnol</i>	<i>Lethrinus nebulosus</i>		2,100	200						2,300
<i>Trey deik koul</i>	(blank)		1,250	100						1,350
<i>Trey do angkor</i>	<i>Gerres abbreviatus</i>			500						500
<i>Trey kheï/proy veng</i>	<i>Pentaprion longimanus</i>				400					400
<i>X-others</i>	<i>X-others</i>	18,200	20,350	11,250	19,400	27,000	19,600	30,500	43,200	189,500
Total		305,700	318,300	350,550	548,950	365,025	469,800	602,000	671,100	3,631,425

Fisher ID: 9

Local Name	Scientific Name	Value (Riel)								
		Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Total
<i>Trey kamong khoun khley</i>	<i>Rastrelliger brachysoma</i>	693,000	1,046,000	3,079,600	9,428,100	2,766,600	4,136,000	3,322,000	4,042,000	28,513,300
<i>Trey karav</i>	<i>Eleutheronema tetradactylum</i>	158,250	763,000							921,250
<i>Trey chhlam chhkout</i>	<i>Chiloscyllium griseum</i>	156,000	222,000							378,000
<i>Kdam Thmor</i>	<i>Scylla serrata</i>	66,000	60,500							126,500
<i>Bangka para/kloy</i>	<i>Penaeus semisulcatus</i>			72,000						72,000
<i>Bangka sor</i>	(blank)			42,000						42,000
<i>X-others</i>	<i>X-others</i>	19,000	49,400	88,200	143,200	151,400	166,400	164,000	275,600	1,057,200
Total		1,092,250	2,140,900	3,281,800	9,571,300	2,918,000	4,302,400	3,486,000	4,317,600	31,110,250

Fisher ID: 10

Local Name	Scientific Name	Value (Riel)								
		Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Total
<i>Trey kbork</i>	<i>Moolgarda seheli</i>	5,000	44,700	301,300	609,400	311,200	224,800		46,000	1,542,400
<i>Bangka sor</i>	(blank)	367,920	302,500	35,000		12,000		24,000		741,420
<i>Bangka kor gneub</i>	<i>Penaeus merguensis</i>						78,000	356,000	289,000	723,000
<i>Trey prorleuk</i>	<i>Leiognathus splendens</i>					48,000		53,550	110,250	211,800
<i>Bangka para/kloy</i>	<i>Penaeus semisulcatus</i>		50,000			12,000	80,000	40,200	24,700	206,900

<i>Trey kantang kra-ob</i>	<i>Siganus canaliculatus</i>	2,000	13,600	134,100	15,000		3,000			167,700
<i>Kdam Thmor</i>	<i>Scylla serrata</i>	840	8,800	15,000	9,000	3,000	25,500	70,500	19,950	152,590
<i>Kdam Seih</i>	<i>Portunus pelagicus</i>			123,600						123,600
<i>Trey kantang thmor</i>	<i>Siganus javus</i>				83,500	14,500		750		98,750
<i>Trey mrech</i>	(blank)			15,000	36,400	2,500				53,900
<i>Trey damlong</i>	(blank)	5,850	18,200	5,000			14,500			43,550
<i>Trey kantang phleung</i>	<i>Siganus guttatus</i>				14,000		23,200			37,200
<i>Trey phtong prort</i>	<i>Tylosurus acus melanotus</i>	3,400	24,500	1,800						29,700
<i>Trey kantang prorpeh</i>	<i>Siganus argenteus</i>					20,000				20,000
<i>Trey prorlos</i>	<i>Sillago sihama</i>			17,200						17,200
<i>Trey prorkong</i>	(blank)				6,000		10,000			16,000
<i>Trey karav</i>	<i>Eleutheronema tetradactylum</i>			1,500	3,200				8,400	13,100
<i>Trey Krorb khnol</i>	<i>Lethrinus nebulosus</i>			10,500						10,500
<i>Trey ka-ok</i>	<i>Arius maculatus</i>			10,000						10,000
<i>Trey deik koul</i>	(blank)		8,050	1,000						9,050
<i>Trey kbork angkam</i>	<i>Moolgarda buchanani</i>				4,000			3,250		7,250
<i>Trey phtong phka</i>	<i>Hemiramphus far</i>				2,500			2,000		4,500
<i>Meuk snok</i>	<i>Sepiella inermis</i>	2,800	1,200							4,000
<i>Trey andeng poy</i>	(blank)			4,000						4,000
<i>Trey angrei</i>	<i>Sphyraena putnamae</i>					3,600				3,600
<i>Trey andat chhke</i>	<i>Paraplagusia bilineata</i>						3,000			3,000
<i>Trey balang</i>	<i>Rastrelliger faughni</i>								3,000	3,000
<i>Trey khej/proy veng</i>	<i>Pentaprion longimanus</i>				1,200				1,500	2,700
<i>Trey kantang thmor</i>	<i>Siganus virgatus</i>						2,400			2,400
<i>Trey do angkor</i>	<i>Gerres abbreviatus</i>		2,000							2,000
<i>Trey kanchanh chras</i>	(blank)				2,000					2,000
<i>Trey spong</i>	<i>Psammoperca waigiensis</i>					2,000				2,000
<i>Meuk bompong/meit</i>	<i>Uroteuthis duvaucelii</i>		800							800

<i>X-others</i>	<i>X-others</i>	19,800	29,600	36,000	40,500	40,800	28,500	40,550	15,500	251,250
Total		407,610	503,950	711,000	826,700	469,600	492,900	590,800	518,300	4,520,860

Fisher ID: 11

Local Name	Scientific Name	Value (Riel)								
		Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Total
<i>Trey teukei ouch kiev</i>	<i>Plectropomus oligacanthus</i>	100,000	300,000	1,090,000	1,680,000	1,920,000	2,016,000	3,976,000	3,280,000	14,362,000
<i>Trey teukei koav</i>	<i>Epinephelus coioides</i>		0				416,000	372,000	544,000	1,332,000
<i>Trey seik</i>	<i>Scarus spp.</i>	7,400	11,500	50,000	91,000	114,000	90,000	170,000	208,000	741,900
<i>Trey teukeokao</i>	(blank)				288,000	320,000				608,000
<i>Trey teukei para</i>	<i>Epinephelus quoyanus</i>	6,000		48,000	116,000	60,000	64,000	88,000	72,000	454,000
<i>Trey kantang phleung</i>	<i>Siganus guttatus</i>	9,150	9,600	46,000	64,000	75,000	102,000	16,000	56,000	377,750
<i>Trey phneik Lan</i>	<i>Priacanthus hamrur</i>	500	3,750	12,000	19,500	42,000	66,000	100,000	134,000	377,750
<i>Trey kantang kra-ob</i>	<i>Siganus canaliculatus</i>						72,000	112,000	90,000	274,000
<i>Trey teukei para krorthorm</i>	<i>Epinephelus fasciatus</i>			164,500		64,000				228,500
<i>Trey seik thmor</i>	<i>Scarus spp.</i>	500					44,000	76,000	106,000	226,500
<i>Trey kantang prorpheh</i>	<i>Siganus argenteus</i>	4,950	3,750		22,000	36,000			136,000	202,700
<i>Kdam neak</i>	<i>Charybdis natator</i>							96,000		96,000
<i>Ngveav sor</i>	<i>Periglypta sp.</i>			7,800						7,800
<i>Trey teukei thmor</i>	<i>Aethaloperca rogae</i>	1,000								1,000
<i>X-others</i>	<i>X-others</i>			2,250						2,250
Total		129,500	328,600	1,420,550	2,280,500	2,631,000	2,870,000	5,006,000	4,626,000	19,292,150

Fisher ID: 12

Local Name	Scientific Name	Value (Riel)								
		Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Total
<i>Meuk huysay</i>	<i>Sepioteuthis lessoniana</i>	576,300	1,403,000	941,700	1,227,200	1,151,200	882,400	1,647,600	1,896,000	9,725,400
<i>Kdam Thmor</i>	<i>Scylla serrata</i>		7,000			84,000				91,000

<i>Kdam Seih</i>	<i>Portunus pelagicus</i>		45,000			21,500				66,500
<i>Meuk bompong/meit</i>	<i>Uroteuthis duvaucelii</i>	5,200	49,200							54,400
<i>Kdam krahorm</i>	(blank)		2,000							2,000
<i>Kdam Sor/Dangkeab Sor</i>	<i>Charybdis anisodon</i>					1,000				1,000
Total		581,500	1,506,200	941,700	1,227,200	1,257,700	882,400	1,647,600	1,896,000	9,940,300

Fisher ID: 13		Value (Riel)								
Local Name	Scientific Name	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Total
<i>Meuk huysay</i>	<i>Sepioteuthis lessoniana</i>	269,600	597,600	359,900	366,300	257,400	199,500	447,200	506,600	3,004,100
<i>Meuk bompong/meit</i>	<i>Uroteuthis duvaucelii</i>	2,000	55,200	37,200	39,600			19,200	14,400	167,600
Total		271,600	652,800	397,100	405,900	257,400	199,500	466,400	521,000	3,171,700

Fisher ID: 14		Value (Riel)								
Local Name	Scientific Name	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Total
<i>Trey teukei ouch kiev</i>	<i>Plectropomus oligacanthus</i>	40,000	1,352,000	600,000	1,008,000	1,312,000	1,312,000	1,520,000	2,264,000	9,408,000
<i>Trey teukei para krrohorm</i>	<i>Epinephelus fasciatus</i>	9,600	96,000	102,400	201,600	293,200	321,600	390,800	371,200	1,786,400
<i>Trey teukei para</i>	<i>Epinephelus quoyanus</i>	2,400	8,400	7,200		94,400	268,800	140,800	129,600	651,600
<i>Trey kantang phleung</i>	<i>Siganus guttatus</i>	1,000	12,100	32,100	33,600	58,200	57,300	63,600	67,200	325,100
<i>Trey seik</i>	<i>Scarus spp.</i>	3,000	21,400	21,200	30,200	43,000	41,000	54,800	53,200	267,800
<i>Trey ka chy</i>	<i>Diagramma pictum</i>		6,000	18,000	34,500	35,500	42,000	40,500	59,000	235,500
<i>Trey teukeokao</i>	(blank)				32,000					32,000
<i>Kdam Thmor</i>	<i>Scylla serrata</i>				24,000					24,000
<i>Trey borbel ork</i>	<i>Aetobatus narinari</i>							13,500		13,500
<i>Trey kantang kra-ob</i>	<i>Siganus canaliculatus</i>				11,100					11,100
<i>Trey borbel khleing</i>	<i>Mobula japanica</i>		9,900							9,900
Total		56,000	1,505,800	780,900	1,375,000	1,836,300	2,042,700	2,224,000	2,944,200	12,764,900

Fisher ID: 15

		Value (Riel)								
Local Name	Scientific Name	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Total
<i>Meuk huysay</i>	<i>Sepioteuthis lessoniana</i>	233,800	1,952,400	601,900	1,442,500	1,482,600	612,900	440,800	2,046,800	8,813,700
<i>Meuk bompong/meit</i>	<i>Uroteuthis duvaucelii</i>	7,000	282,000	26,000	126,000	89,000	12,000		60,000	602,000
Total		240,800	2,234,400	627,900	1,568,500	1,571,600	624,900	440,800	2,106,800	9,415,700

Fisher ID: 16

		Value (Riel)								
Local Name	Scientific Name	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Total
<i>Meuk dong</i>	<i>Sepioloa sp.</i>		85,500	522,500	380,000	456,000	2,400,000	3,400,000	2,800,000	10,044,000
Total			85,500	522,500	380,000	456,000	2,400,000	3,400,000	2,800,000	10,044,000

Fisher ID: 17

		Value (Riel)								
Local Name	Scientific Name	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Total
<i>Kdam Thmor</i>	<i>Scylla serrata</i>	246,600	492,800	625,100	776,300	922,500	1,176,300	1,071,500	1,049,700	6,360,800
Total		246,600	492,800	625,100	776,300	922,500	1,176,300	1,071,500	1,049,700	6,360,800

Fisher ID: 18

		Value (Riel)								
Local Name	Scientific Name	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Total
<i>Kdam Thmor</i>	<i>Scylla serrata</i>	195,000	465,000	730,600	759,300	724,100	671,000	598,400	649,800	4,793,200
Total		195,000	465,000	730,600	759,300	724,100	671,000	598,400	649,800	4,793,200

Fisher ID: 19

		Value (Riel)								

Local Name	Scientific Name	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Total
<i>Ngveav sor</i>	<i>Periglypta sp.</i>	103,200	163,500	121,800	132,300	108,200	147,600	171,996		948,596
<i>Kdam Thmor</i>	<i>Scylla serrata</i>							19,800		19,800
Total		103,200	163,500	121,800	132,300	108,200	147,600	191,796		968,396

Fisher ID: 20

		Value (Riel)								
Local Name	Scientific Name	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Total
<i>Kdam Chor</i>	<i>Episesarma singaporensis</i>	116,400	169,750	220,950	242,550		199,300	134,800	234,900	1,318,650
<i>Kdam Thmor</i>	<i>Scylla serrata</i>	12,000	50,400	47,800	159,600		73,800	26,000	18,200	387,800
<i>Trey phtourk</i>	<i>Rachycentron canadum</i>					198,400				198,400
<i>Kreing chheam</i>	<i>Anadara granosa</i>	21,000	59,000							80,000
<i>Bangkea sor</i>	(blank)	16,000								16,000
<i>Khyorng Krachorkseih</i>	(blank)	2,500		6,600						9,100
<i>Khyorng dong</i>	<i>Melo melo</i>							3,000		3,000
<i>Ngveav sor</i>	<i>Periglypta sp.</i>	3,000								3,000
<i>Trey kantang kra-ob</i>	<i>Siganus canaliculatus</i>	2,400								2,400
Total		173,300	279,150	275,350	402,150	198,400	273,100	163,800	253,100	2,018,350

Fisher ID: 21

		Value (Riel)								
Local Name	Scientific Name	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Total
<i>Kdam Chor</i>	<i>Episesarma singaporensis</i>	144,400	198,000	221,950	256,950	226,350	194,800	107,600	108,450	1,458,500
<i>Kdam Thmor</i>	<i>Scylla serrata</i>	33,800	40,550	48,700	252,100	78,000	46,800	14,400	9,400	523,750
<i>Ngveav phourk</i>	<i>Polymesoda erosa</i>		1,200							1,200
Total		178,200	239,750	270,650	509,050	304,350	241,600	122,000	117,850	1,983,450

Fisher ID: 22

Local Name	Scientific Name	Value (Riel)								
		Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Total
<i>Kdam Thmor</i>	<i>Scylla serrata</i>		1,547,100	1,941,750	2,189,900	1,598,200	1,568,980	2,931,300	1,501,000	13,278,230
<i>Kdam Seih</i>	<i>Portunus pelagicus</i>		2,278,050	1,924,700	834,300	2,045,350	669,450	232,450	1,105,600	9,089,900
<i>Kdam Sor/Dangkeab Sor</i>	<i>Charybdis anisodon</i>		12,200	28,531	33,900	27,750	27,650			130,031
<i>Kdam skor krorb/Kdam sor</i>	<i>Matuta victor</i>							38,400		38,400
<i>Bangkea sor</i>	(blank)								21,250	21,250
Total		0	3,837,350	3,894,981	3,058,100	3,671,300	2,266,080	3,202,150	2,627,850	22,557,811

Fisher ID: 23

Local Name	Scientific Name	Value (Riel)								
		Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Total
<i>Trey beika</i>	<i>Auxis thazard</i>			600				928,750	814,280	1,743,630
<i>Trey kamoy</i>	<i>Anodontostoma chacunda</i>		6,000	1,000	251,400	137,500	385,600	339,500	128,800	1,249,800
<i>Trey kamong khoun khley</i>	<i>Rastrelliger brachysoma</i>			337,200	150,000	25,600	422,000			934,800
<i>Bangkea sor</i>	(blank)	70,000	311,000	11,000		30,700		17,000		439,700
<i>Bangkea para/kloy</i>	<i>Penaeus semisulcatus</i>		3,000	4,500		343,800	19,200	8,400		378,900
<i>Trey Sambor Hear</i>	(blank)					53,500	80,800	34,000	34,050	202,350
<i>Meuk dong</i>	<i>Sepiola sp.</i>								200,000	200,000
<i>Trey karav</i>	<i>Eleutheronema tetradactylum</i>			2,500	6,000		72,500	28,000		109,000
<i>Trey sleuk russey</i>	<i>Thryssa hamiltonii</i>	5,625	5,850	1,950	4,500		54,200	29,700		101,825
<i>Bangkea kor gneub</i>	<i>Penaeus merguensis</i>								86,100	86,100
<i>Trey khut/trorchiek damrey</i>	<i>Ephippus orbis</i>			65,800						65,800
<i>Trey borbel moan</i>	<i>Himantura imbricata</i>		4,800		60,000					64,800
<i>Trey andat chhke</i>	<i>Paraplagusia bilineata</i>				8,250	28,950				37,200
<i>Trey ka-ok</i>	<i>Arius maculatus</i>					16,800	15,600			32,400
<i>Trey udouy</i>	<i>Selaroides leptolepis</i>					24,300				24,300
<i>Trey andeng poy</i>	(blank)				12,000					12,000

<i>Trey damlong</i>	(blank)							6,500		6,500
<i>Kdam Seih</i>	<i>Portunus pelagicus</i>	4,000								4,000
<i>Meuk snok</i>	<i>Sepiella inermis</i>	1,200								1,200
<i>Trey Kor/Romeas</i>	<i>Monacanthus chinensis</i>	250								250
<i>X-others</i>	<i>X-others</i>	2,350	78,350	6,500	12,550	30,400				130,150
Total		83,425	409,000	431,050	504,700	691,550	1,049,900	1,391,850	1,263,230	5,824,705

Fisher ID: 24

		Value (Riel)								
Local Name	Scientific Name	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Total
<i>Kdam Thmor</i>	<i>Scylla serrata</i>	1,810,500	4,617,000	2,251,600	2,708,700	2,126,700	1,856,000	1,691,000	2,469,100	19,530,600
<i>Kdam Seih</i>	<i>Portunus pelagicus</i>	748,400	2,530,500	899,400	806,800	680,400	520,100	579,300	681,000	7,445,900
<i>Bangkea sor</i>	(blank)								39,450	39,450
<i>Kdam skor krorb/Kdam sor</i>	<i>Matuta victor</i>							28,175	5,000	33,175
<i>Kdam Sor/Dangkeab Sor</i>	<i>Charybdis anisodon</i>	11,000	98,500	52,000	55,000	38,000	27,000			281,500
Total		2,569,900	7,246,000	3,203,000	3,570,500	2,845,100	2,403,100	2,298,475	3,194,550	27,330,625

Fisher ID: 25

		Value (Riel)								
Local Name	Scientific Name	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Total
<i>Kdam Thmor</i>	<i>Scylla serrata</i>	62,000	148,000	434,200	228,000	280,500	297,000	256,800	181,100	1,887,600
<i>Kdam Seih</i>	<i>Portunus pelagicus</i>	77,500	476,551	219,750	168,900	9,000	52,200	166,050	82,140	1,252,091
<i>Trey kalang</i>	<i>Scomberoides commersonianus</i>							479,500		479,500
<i>Trey balang</i>	<i>Rastrelliger faughni</i>								431,000	431,000
<i>Trey borbel moan</i>	<i>Himantura imbricata</i>		29,635	41,300	43,200	56,400	64,800	34,500	31,500	301,335
<i>Trey beika</i>	<i>Auxis thazard</i>			600		34,000	157,500	39,200	10,500	241,800
<i>Trey andat chhke</i>	<i>Paraplagusia bilineata</i>		1,500	3,100		21,300	8,700	13,800	1,800	50,200
<i>Trey karav</i>	<i>Eleutheronema tetradactylum</i>			1,200	7,500		13,900	2,000	3,300	27,900

<i>Trey ka-ok kbal heang/Ka-ok prak</i>	<i>Osteogeneiosus militaris</i>								26,000	26,000
<i>Trey andeng</i>	(blank)					5,000		17,700		22,700
<i>Trey Kor/Romeas</i>	<i>Monacanthus chinensis</i>			0	0	9,000	11,600			20,600
<i>Trey kantang kra-ob</i>	<i>Siganus canaliculatus</i>			800		6,400	13,000			20,200
<i>Trey ka-ok</i>	<i>Arius maculatus</i>			13,300	3,600	2,000				18,900
<i>Trey kamong khoun khley</i>	<i>Rastrelliger brachysoma</i>			9,600			8,800			18,400
<i>Trey kantuy krabey</i>	(blank)		1,440	2,000	3,600	9,100				16,140
<i>Trey sleuk russey</i>	<i>Thryssa hamiltonii</i>				450		4,200			4,650
<i>Kdam kom</i>	(blank)							2,500		2,500
<i>Kdam skor krorb/Kdam sor</i>	<i>Matuta victor</i>					2,000				2,000
X-others	X-others		3,000		18,250	17,160	6,300	44,100	2,000	90,810
Total		139,500	660,126	725,850	473,500	451,860	638,000	1,056,150	769,340	4,914,326

Fisher ID: 26

Local Name	Scientific Name	Value (Riel)								
		Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Total
<i>Kreing chheam</i>	<i>Anadara granosa</i>	80,000	144,100	164,700	126,000	174,050	119,250	141,750	119,250	1,069,100
<i>Trey teukei ouch kiev</i>	<i>Plectropomus oligacanthus</i>				136,000					136,000
<i>Trey teukei para krorthorm</i>	<i>Epinephelus fasciatus</i>				16,000					16,000
<i>Trey ka chy</i>	<i>Diagramma pictum</i>				7,500					7,500
<i>Trey kantang phleung</i>	<i>Siganus guttatus</i>				6,000					6,000
<i>Trey seik</i>	<i>Scarus spp.</i>				3,600					3,600
Total		80,000	144,100	164,700	295,100	174,050	119,250	141,750	119,250	1,238,200

Annex 7: Average price (Riel/kg) of catch by species by fisher (April-November 2011)

Fisher ID: 1									
Local Name	Scientific Name	Average price (Riel/kg)							
		Apr	May	Jun	Jul	Aug	Sep	Oct	Nov
<i>Trey do angkor</i>	<i>Gerres abbreviatus</i>			5,364.11	5,536.14	6,347.71	6,920.04	6,695.24	4,087.12
<i>Trey prorleuk</i>	<i>Leiognathus splendens</i>						12,000.00	6,200.49	6,108.93
<i>Trey Krorb khnol</i>	<i>Lethrinus nebulosus</i>			3,500.00	4,700.70	5,380.16	5,302.56	5,327.25	3,434.78
<i>Trey prorlos</i>	<i>Sillago sihama</i>			8,221.93	7,802.63	9,665.82			
<i>Trey krorhorm srakakras</i>	<i>Scolopis taeniopterus</i>						5,710.05	3,882.35	3,330.63
<i>Trey krorhorm kbalkor</i>	(blank)				5,076.86	5,415.63		2,706.73	3,671.81
<i>Trey kamoy</i>	<i>Anodontostoma chacunda</i>	3,045.51	3,308.15	4,000.00	9,000.00		1,000.00		
<i>Trey kamong khoun khley</i>	<i>Rastrelliger brachysoma</i>	5,089.89	5,036.99	5,500.00					
<i>Trey chek</i>	<i>Seriolina nigrofasciata</i>	5,078.95	4,800.00	4,812.50	5,287.88	4,950.33	5,227.27	5,000.00	5,000.00
<i>Trey udouy</i>	<i>Selaroides leptolepis</i>	3,000.00		5,382.35	7,600.00	7,444.44	8,046.08	8,000.00	6,000.00
<i>Trey Kor/Romeas</i>	<i>Monacanthus chinensis</i>				3,363.64	2,830.99	2,377.22	2,050.00	1,906.80
<i>Trey angrei</i>	<i>Sphyaena putnamae</i>					9,898.88	5,986.23	5,166.67	8,000.00
<i>Trey krorhorm</i>	<i>Lutjanus malabaricus</i>	4,000.00		3,735.09	7,495.80	7,640.78			
<i>Trey phneik Lan</i>	<i>Priacanthus hamrur</i>			4,444.56	5,214.29				2,000.00
<i>Trey koun</i>	<i>Hilsa keele</i>	2,403.85	3,416.67	4,000.00				2,550.15	1,963.16
<i>Trey kantang kra-ob</i>	<i>Siganus canaliculatus</i>			4,404.41	6,125.00	4,775.00	4,500.00	5,500.00	4,857.14
<i>Trey chhlam chhkout</i>	<i>Chiloscyllium griseum</i>			5,111.11	6,071.43	6,103.09	7,000.00	#DIV/0!	4,000.00
<i>Trey andeng</i>	(blank)			8,282.61					
<i>Kdam Seih</i>	<i>Portunus pelagicus</i>	5,000.00	8,000.00	8,875.00	1,937.84	9,655.17	5,771.43	3,000.00	7,807.69
<i>Trey prorkong</i>	(blank)				4,743.24				
<i>Kdam skor krorb/Kdam sor</i>	<i>Matuta victor</i>			2,285.71	1,077.52	1,641.27			
<i>Trey andeng poy</i>	(blank)					7,000.00			
<i>Kdam neak</i>	<i>Charybdis natator</i>	8,000.00	1,000.00	4,900.00					
<i>Meuk huysay</i>	<i>Sepioteuthis lessoniana</i>	1,500.00	13,212.12						
<i>Kdam krahorm</i>	(blank)		7,666.67						

<i>Trey deik koul</i>	(blank)							2,000.00	
<i>Trey trorsork</i>	<i>Terapon theraps</i>							1,000.00	
<i>Trey sach</i>	(blank)	6,000.00							
<i>Trey kalang</i>	<i>Scomberoides commersonianus</i>								8,000.00
<i>Trey balang</i>	<i>Rastrelliger faughni</i>								8,000.00
<i>Bangkong kandob</i>	<i>Clorida decorata</i>			10,000.00					
<i>Trey kbork</i>	<i>Moolgarda seheli</i>								1,000.00
<i>X-others</i>	<i>X-others</i>	1,439.24	1,208.16	1,267.97	1,500.00		1,880.38	1,564.74	710.72
Average		3,407.95	3,580.12	4,585.23	5,094.21	6,243.57	6,388.25	3,552.11	3,647.39

Fisher ID: 2

Local Name	Scientific Name	Average price (Riel/kg)							
		Apr	May	Jun	Jul	Aug	Sep	Oct	Nov
<i>Kdam Seih</i>	<i>Portunus pelagicus</i>	15,769.23	15,624.20	17,413.73	19,933.63	17,069.93	16,722.29	17,865.27	18,843.35
<i>Khyornng thnot</i>	<i>Conus vexillum</i>	3,103.45	3,307.84	3,266.10	3,602.05	4,055.03	5,000.00	5,458.74	5,000.00
<i>Khyornng dong</i>	<i>Melo melo</i>	7,000.00	6,785.71	7,087.91	7,755.48	8,160.40	8,723.36	9,000.00	9,000.00
<i>Trey chhlam chhkout</i>	<i>Chiloscyllium griseum</i>	6,333.33	6,785.71	7,500.00	5,575.76	7,600.00	7,000.00	10,000.00	10,000.00
<i>Trey borbel moan</i>	<i>Himantura imbricata</i>				4,010.00	4,090.91	4,781.25	4,000.00	4,000.00
<i>Trey borbel ork</i>	<i>Aetobatus narinari</i>	3,600.00	5,000.00	4,000.00					
<i>Trey kam kourch</i>	<i>Carangoides ferdau</i>		8,000.00						
<i>Trey chenchas/pakchiv</i>	<i>Alectis ciliaris</i>								9,545.45
Average		11,703.32	11,375.77	12,485.49	12,661.56	11,803.03	11,529.38	11,920.45	12,538.87

Fisher ID: 3

Local Name	Scientific Name	Average price (Riel/kg)							
		Apr	May	Jun	Jul	Aug	Sep	Oct	Nov
<i>Kdam Seih</i>	<i>Portunus pelagicus</i>	18,423.08	18,395.50	22,949.68	13,026.40	17,328.49	19,944.47	20,763.07	42,724.88
<i>Bangkong kandob</i>	<i>Clorida decorata</i>			4,687.50	5,356.00	10,973.68	18,053.25	14,185.20	15,391.61
<i>Kdam Thmor</i>	<i>Scylla serrata</i>	17,900.00	12,637.97	12,406.25	8,300.00	8,913.79	2,204.55	8,995.65	
<i>Bangkong pak</i>	(blank)	8,000.00	5,600.00		27,333.33	19,592.21	18,872.09	13,720.00	12,500.00

<i>Kdam krahorm</i>	(blank)	3,294.12	3,020.80	4,715.35	2,810.17	1,507.27		1,432.46	816.13
<i>Trey antanou</i>	<i>Pentapodus setosus</i>								7,801.03
<i>Bangkea sor</i>	(blank)	10,000.00	5,096.77	3,512.50		15,000.00	17,000.00		770.48
<i>Kdam skor krorb/Kdam sor</i>	<i>Matuta victor</i>		4,550.00		1,856.31	1,425.69	2,008.92	924.79	3,807.10
<i>Kdam neak</i>	<i>Charybdis natator</i>			2,260.00	4,000.00	4,630.14		4,142.86	7,800.00
<i>Bangkea kor gneub</i>	<i>Penaeus merguensis</i>		8,615.38				11,173.91		
<i>Kdam Sor/Dangkeab Sor</i>	<i>Charybdis anisodon</i>			1,658.89					
<i>Meuk bompong/meit</i>	<i>Uroteuthis duvaucelii</i>							4,000.00	6,500.00
<i>Trey Kor/Romeas</i>	<i>Monacanthus chinensis</i>								2,392.31
<i>Khyornng thnot</i>	<i>Conus vexillum</i>					500.00			
<i>Kdam pingpeang</i>	<i>Arcania undecimspinosa</i>								
Average		15,851.92	12,420.86	15,786.84	8,575.87	12,349.93	16,837.36	16,822.42	30,613.43

Fisher ID: 4

Local Name	Scientific Name	Average price (Riel/kg)							
		Apr	May	Jun	Jul	Aug	Sep	Oct	Nov
<i>Bangkea kor gneub</i>	<i>Penaeus merguensis</i>	7,908.78	6,139.13	11,563.98	7,087.22	7,674.81	6,712.50	8,275.94	5,444.52
<i>Bangkea sor</i>	(blank)	8,625.74	5,133.33	3,500.00	5,042.11	7,284.81	4,720.93	3,600.00	7,031.25
<i>Trey prorkong</i>	(blank)					3,796.30	9,666.67		
<i>Bangkea para/kloy</i>	<i>Penaeus semisulcatus</i>	4,000.00		7,220.51	4,571.43	4,000.00			
<i>Trey phtong prortkheat</i>	<i>Ablennes hians</i>				4,000.00	7,000.00	6,629.03		
<i>Trey mrech</i>	(blank)					3,763.77	1,500.00		
<i>Kdam Thmor</i>	<i>Scylla serrata</i>	3,600.00	4,200.00					4,400.00	5,315.00
<i>Trey angkeuy</i>	<i>Lutjanus johnii</i>			15,000.00					
<i>Meuk bompong/meit</i>	<i>Uroteuthis duvaucelii</i>								4,800.00
<i>X-others</i>	<i>X-others</i>	983.84	907.69	1,036.36	1,421.30	1,053.85	1,646.46	2,460.67	880.60
Average		6,299.76	4,755.36	8,830.02	4,937.69	6,042.29	5,469.10	5,546.84	4,880.06

Fisher ID: 5

Local Name	Scientific Name	Average price (Riel/kg)							
		Apr	May	Jun	Jul	Aug	Sep	Oct	Nov
<i>Trey mrech</i>	(blank)	4,185.71	3,281.12	2,985.05	3,000.00	3,176.10	3,235.67	4,235.29	4,407.82
<i>Bangkea sor</i>	(blank)	9,882.35	9,375.00	7,309.52	6,552.63	6,571.43	6,388.89	9,000.00	9,849.25
<i>Bangkea phourk</i>	(blank)	7,000.00	6,705.20	5,833.33	6,024.39	4,657.14	5,116.28	8,000.00	8,909.09
<i>Trey prorkong</i>	(blank)		3,833.33	2,964.37	3,000.00	3,519.48	3,873.02	5,000.00	6,000.00
<i>Kdam Thmor</i>	<i>Scylla serrata</i>	20,000.00	16,000.00					19,480.00	12,697.37
<i>Bangkea kor gneub</i>	<i>Penaeus merguensis</i>							6,200.00	21,713.51
<i>Trey damlong</i>	(blank)			3,000.00	3,000.00	3,230.77	3,380.00	4,830.51	5,462.07
<i>Trey kanchanh chras</i>	(blank)				2,172.62	2,031.25	1,877.05		1,222.22
<i>Trey kranh</i>	<i>Lobotes surinamensis</i>		2,828.57	2,000.00	3,000.00	3,218.75	2,800.00	3,000.00	3,000.00
<i>Kdam Seih</i>	<i>Portunus pelagicus</i>		9,657.14						
<i>Trey prorlos</i>	<i>Sillago sihama</i>		3,179.10	3,681.82					
<i>Trey chek</i>	<i>Seriolina nigrofasciata</i>					3,000.00	3,000.00		3,044.87
<i>Trey andeng</i>	(blank)		7,250.00			6,000.00	5,875.00		
<i>Trey kantong kra-ob</i>	<i>Siganus canaliculatus</i>					4,625.00			4,000.00
<i>Trey do angkor</i>	<i>Gerres abbreviatus</i>		4,000.00						
<i>Trey andeng poy</i>	(blank)					6,625.00			
<i>Bangkea para/kloy</i>	<i>Penaeus semisulcatus</i>							10,000.00	
<i>Trey trorsork</i>	<i>Terapon theraps</i>								2,000.00
<i>Trey kbork</i>	<i>Moolgarda seheli</i>					4,000.00			
<i>X-others</i>	<i>X-others</i>	3,476.19	2,916.93	2,108.17	2,129.29	1,661.29	1,880.95		2,666.67
Average		8,507.28	4,588.03	2,945.05	3,038.50	3,237.67	3,449.90	7,521.74	7,472.37

Fisher ID: 6

Local Name	Scientific Name	Average price (Riel/kg)							
		Apr	May	Jun	Jul	Aug	Sep	Oct	Nov
<i>Trey kamong khoun khley</i>	<i>Rastrelliger brachysoma</i>	2,571.43	2,955.88	1,653.65	1,606.83	2,972.00	3,000.00	3,500.00	3,500.00
<i>Trey kamong khoun veng</i>	<i>Rastrelliger kanagurta</i>	2,141.03	4,123.20	5,000.00	5,000.00	5,000.00	5,000.00	5,000.00	4,679.14

<i>Trey krorthorm kbalkor</i>	(blank)		1,788.46	1,500.00				2,500.00	2,500.00
<i>Trey krorthorm srakakras</i>	<i>Scolopis taeniopterus</i>				3,174.42	3,362.07	2,983.87	3,800.00	3,500.00
<i>Trey krorthorm</i>	<i>Lutjanus malabaricus</i>		2,000.00		2,717.39	2,546.88			
<i>Trey beika</i>	<i>Auxis thazard</i>		5,000.00	5,896.37	8,000.00	8,000.00	8,000.00	8,000.00	8,000.00
<i>Trey krahorm phneak thom</i>	<i>Priacanthus macracanthus</i>		3,000.00	3,000.00			4,000.00	4,000.00	
<i>Trey Kor/Romeas</i>	<i>Monacanthus chinensis</i>						2,500.00		
<i>Trey krorthorm phneklie</i>	<i>Priacanthus tayenus</i>		2,500.00		4,000.00	4,000.00			
<i>Bangkea sor</i>	(blank)	11,200.00	12,500.00	15,225.81					
<i>Trey karav</i>	<i>Eleutheronema tetradactylum</i>	2,000.00		6,500.00					
<i>Bangkea para/kloy</i>	<i>Penaeus semisulcatus</i>			10,000.00					
<i>Trey borbel ork</i>	<i>Aetobatus narinari</i>		8,000.00						
<i>Kdam Seih</i>	<i>Portunus pelagicus</i>	12,000.00	10,000.00						
<i>Trey kamoy</i>	<i>Anodontostoma chacunda</i>	1,500.00							
<i>X-others</i>	<i>X-others</i>	687.65	1,241.59	770.39	810.00	1,186.71	1,000.00	1,000.00	1,000.00
Average		2,000.33	3,172.34	1,879.48	2,257.42	3,842.71	3,103.82	3,551.09	3,501.45

Fisher ID: 7

Local Name	Scientific Name	Average price (Riel/kg)							
		Apr	May	Jun	Jul	Aug	Sep	Oct	Nov
<i>Kdam Seih</i>	<i>Portunus pelagicus</i>	9,637.77	9,724.65	9,444.13	11,311.16	12,205.89	11,496.97	8,843.73	8,780.99
<i>Kdam skor krorb/Kdam sor</i>	<i>Matuta victor</i>								12,000.00
<i>Kdam Chor</i>	<i>Episesarma singaporensis</i>								4,500.00
Total		9,637.77	9,724.65	9,444.13	11,311.16	12,205.89	11,496.97	8,843.73	8,770.38

Fisher ID: 8

Local Name	Scientific Name	Average price (Riel/kg)							
		Apr	May	Jun	Jul	Aug	Sep	Oct	Nov
<i>Bangkea sor</i>	(blank)	5,092.53	3,907.98	4,000.00			5,000.00	5,929.82	5,142.86
<i>Trey kbork</i>	<i>Moolgarda seheli</i>	1,000.00	6,000.00	6,114.50	6,763.19	7,000.00	6,911.97	7,000.00	

<i>Bangkea kor gneub</i>	<i>Penaeus merguensis</i>						5,000.00	5,731.71	5,772.28
<i>Trey kantang kra-ob</i>	<i>Siganus canaliculatus</i>	1,000.00	1,666.67	2,198.86	2,313.77	2,536.25			
<i>Trey phtong phka</i>	<i>Hemiramphus far</i>			2,000.00	2,000.00	2,047.62	2,000.00	2,000.00	2,000.00
<i>Trey kantang thmor</i>	<i>Siganus virgatus</i>				2,345.64	2,872.48		3,000.00	
<i>Kdam Thmor</i>	<i>Scylla serrata</i>			10,000.00	25,000.00		11,714.29	13,903.23	11,520.00
<i>Trey phtong prort</i>	<i>Tylosurus acus melanotus</i>			2,711.68					
<i>Trey karav</i>	<i>Eleutheronema tetradactylum</i>			2,333.33	2,052.08	3,793.10	5,000.00		
<i>Trey kantang prorpeh</i>	<i>Siganus argenteus</i>			10,000.00	5,684.21	10,000.00		10,000.00	
<i>Trey kantang phleung</i>	<i>Siganus guttatus</i>			8,000.00	2,000.00	7,750.00		10,000.00	
<i>Trey prorlos</i>	<i>Sillago sihama</i>			1,075.00	1,500.00	1,000.00			
<i>Bangkea para/kloy</i>	<i>Penaeus semisulcatus</i>							12,000.00	30,000.00
<i>Trey mrech</i>	(blank)			1,500.00	1,000.00				
<i>Trey phtong prortkheat</i>	<i>Ablennes hians</i>			2,866.67	2,000.00		3,000.00		2,000.00
<i>Trey damlong</i>	(blank)		1,000.00	1,000.00			1,000.00	1,000.00	1,000.00
<i>Kdam Seih</i>	<i>Portunus pelagicus</i>		4,130.43	4,000.00					
<i>Bangkorng pak</i>	(blank)			30,000.00		20,000.00			
<i>Trey andeng poy</i>	(blank)						4,285.71		
<i>Trey kbork angkam</i>	<i>Moolgarda buchanani</i>				6,000.00				
<i>Trey prorkong</i>	(blank)						4,000.00		
<i>Trey Krorb khnol</i>	<i>Lethrinus nebulosus</i>		2,333.33	2,000.00					
<i>Trey deik koul</i>	(blank)		1,000.00	1,000.00					
<i>Trey do angkor</i>	<i>Gerres abbreviatus</i>			1,000.00					
<i>Trey khej/proy veng</i>	<i>Pentaprion longimanus</i>				2,000.00				
<i>X-others</i>	X-others	1,000.00	1,038.27	1,397.52	1,000.00	1,000.00	1,000.00	1,000.00	1,000.00
Average		4,038.31	3,303.58	3,391.87	3,193.43	3,276.71	4,423.73	4,452.66	4,234.07

Fisher ID: 9

Local Name	Scientific Name	Average price (Riel/kg)							
		Apr	May	Jun	Jul	Aug	Sep	Oct	Nov
<i>Trey kamong khoun khley</i>	<i>Rastrelliger brachysoma</i>	1,500.00	1,217.69	1,549.09	1,251.57	1,304.38	1,594.45	1,350.41	2,000.00

<i>Trey karav</i>	<i>Eleutheronema tetradactylum</i>	12,660.00	13,872.73						
<i>Trey chhlam chhkout</i>	<i>Chiloscyllium griseum</i>	13,000.00	14,800.00						
<i>Kdam Thmor</i>	<i>Scylla serrata</i>	30,000.00	31,842.11						
<i>Bangkea para/kloy</i>	<i>Penaeus semisulcatus</i>			12,000.00					
<i>Bangkea sor</i>	(blank)			12,000.00					
<i>X-others</i>	<i>X-others</i>	1,000.00	837.29	705.60	800.00	672.89	800.00	773.58	800.00
Average		2,151.37	2,162.74	1,546.20	1,241.09	1,243.82	1,535.47	1,304.64	1,825.24

Fisher ID: 10

Local Name	Scientific Name	Average price (Riel/kg)							
		Apr	May	Jun	Jul	Aug	Sep	Oct	Nov
<i>Trey kbork</i>	<i>Moolgarda seheli</i>	5,000.00	5,960.00	6,369.98	6,531.62	6,551.58	5,900.26		6,571.43
<i>Bangkea sor</i>	(blank)	6,310.81	4,689.92	5,000.00		6,000.00		6,000.00	
<i>Bangkea kor gneub</i>	<i>Penaeus merguensis</i>						5,571.43	5,042.49	4,345.86
<i>Trey prorleuk</i>	<i>Leiognathus splendens</i>					4,000.00		4,500.00	4,500.00
<i>Bangkea para/kloy</i>	<i>Penaeus semisulcatus</i>		4,000.00			6,000.00	5,000.00	17,478.26	24,700.00
<i>Trey kantong kra-ob</i>	<i>Siganus canaliculatus</i>	6,666.67	5,230.77	2,934.35	3,000.00		2,000.00		
<i>Kdam Thmor</i>	<i>Scylla serrata</i>	2,800.00	11,000.00	18,750.00	15,000.00	15,000.00	10,625.00	13,301.89	11,735.29
<i>Kdam Seih</i>	<i>Portunus pelagicus</i>			10,048.78					
<i>Trey kantong thmor</i>	<i>Siganus javus</i>				2,830.51	3,222.22		2,500.00	
<i>Trey mrech</i>	(blank)			1,875.00	2,233.13	2,500.00			
<i>Trey damlong</i>	(blank)	1,392.86	1,229.73	1,666.67			1,933.33		
<i>Trey kantong phleung</i>	<i>Siganus guttatus</i>				7,000.00		8,000.00		
<i>Trey pthong prort</i>	<i>Tylosurus acus melanotus</i>	5,666.67	6,447.37	2,250.00					
<i>Trey kantong prorpeh</i>	<i>Siganus argenteus</i>					8,000.00			
<i>Trey prorlos</i>	<i>Sillago sihama</i>			2,529.41					
<i>Trey prorkong</i>	(blank)				3,000.00		4,000.00		
<i>Trey karav</i>	<i>Eleutheronema tetradactylum</i>			5,000.00	4,000.00				3,000.00
<i>Trey Krorb khnol</i>	<i>Lethrinus nebulosus</i>			2,333.33					
<i>Trey ka-ok</i>	<i>Arius maculatus</i>			10,000.00					

<i>Trey deik koul</i>	(blank)		1,000.00	1,000.00					
<i>Trey kbork angkam</i>	<i>Moolgarda buchanani</i>				4,000.00			3,250.00	
<i>Trey phtong phka</i>	<i>Hemiramphus far</i>				5,000.00			2,000.00	
<i>Meuk snok</i>	<i>Sepiella inermis</i>	3,111.11	4,000.00						
<i>Trey andeng poy</i>	(blank)			4,000.00					
<i>Trey angrei</i>	<i>Sphyraena putnamae</i>					4,000.00			
<i>Trey andat chhke</i>	<i>Paraplagusia bilineata</i>						3,000.00		
<i>Trey balang</i>	<i>Rastrelliger faughni</i>								3,000.00
<i>Trey khe/proy veng</i>	<i>Pentaprion longimanus</i>				4,000.00				1,500.00
<i>Trey kantong thmor</i>	<i>Siganus virgatus</i>						8,000.00		
<i>Trey do angkor</i>	<i>Gerres abbreviatus</i>		2,000.00						
<i>Trey kanchanh chras</i>	(blank)				1,000.00				
<i>Trey spong</i>	<i>Psammoperca waigiensis</i>					1,000.00			
<i>Meuk bompong/meit</i>	<i>Uroteuthis duvaucelii</i>		4,000.00						
<i>X-others</i>	<i>X-others</i>	1,076.09	1,112.78	1,232.88	1,000.00	1,000.00	1,000.00	1,199.70	1,000.00
Average		4,852.50	3,532.77	4,214.58	4,265.74	4,069.32	4,297.30	4,537.63	4,283.47

Fisher ID: 11

Local Name	Scientific Name	Average price (Riel/kg)							
		Apr	May	Jun	Jul	Aug	Sep	Oct	Nov
<i>Trey teukei ouch kiev</i>	<i>Plectropomus oligacanthus</i>	100,000.00	100,000.00	90,833.33	84,000.00	80,000.00	84,000.00	76,461.54	64,313.73
<i>Trey teukei koav</i>	<i>Epinephelus coioides</i>		0.00				32,000.00	24,800.00	32,000.00
<i>Trey seik</i>	<i>Scarus spp.</i>	2,000.00	1,642.86	2,000.00	1,978.26	2,000.00	2,000.00	2,000.00	2,000.00
<i>Trey teukeokao</i>	(blank)				32,000.00	32,000.00			
<i>Trey teukei para</i>	<i>Epinephelus quoyanus</i>	12,000.00		8,000.00	14,500.00	8,000.00	8,000.00	8,000.00	8,000.00
<i>Trey kantong phleung</i>	<i>Siganus guttatus</i>	2,577.46	1,777.78	2,190.48	2,000.00	2,142.86	2,000.00	2,000.00	2,000.00
<i>Trey phneik Lan</i>	<i>Priacanthus hamrur</i>	2,000.00	1,500.00	1,714.29	1,772.73	2,000.00	2,000.00	2,000.00	2,000.00
<i>Trey kantong kra-ob</i>	<i>Siganus canaliculatus</i>						2,000.00	2,000.00	2,000.00
<i>Trey teukei para krorthorm</i>	<i>Epinephelus fasciatus</i>			23,500.00		32,000.00			
<i>Trey seik thmor</i>	<i>Scarus spp.</i>	2,000.00					2,000.00	2,000.00	2,000.00

<i>Trey kantang prorpeh</i>	<i>Siganus argenteus</i>	2,302.33	1,500.00		2,000.00	2,000.00			2,000.00
<i>Kdam neak</i>	<i>Charybdis natator</i>							32,000.00	
<i>Ngveav sor</i>	<i>Periglypta sp.</i>			600.00					
<i>Trey teukei thmor</i>	<i>Aethaloperca rogaa</i>	2,000.00							
<i>X-others</i>	<i>X-others</i>			1,500.00					
Average		10,882.35	15,722.49	15,357.30	16,645.99	15,077.36	12,370.69	15,742.14	10,466.06

Fisher ID: 12

LocalName	Scientific Name	Average price (Riel/kg)							
		Apr	May	Jun	Jul	Aug	Sep	Oct	Nov
<i>Meuk huysay</i>	<i>Sepioteuthis lessoniana</i>	12,235.67	12,990.74	13,024.90	13,167.38	13,495.90	13,513.02	12,952.83	15,892.71
<i>Kdam Thmor</i>	<i>Scylla serrata</i>		14,000.00			15,000.00			
<i>Kdam Seih</i>	<i>Portunus pelagicus</i>		18,000.00			6,142.86			
<i>Meuk bompong/meit</i>	<i>Uroteuthis duvaucelii</i>	10,400.00	12,000.00						
<i>Kdam krahorm</i>	(blank)		2,000.00						
<i>Kdam Sor/Dangkeab Sor</i>	<i>Charybdis anisodon</i>					500.00			
Average		12,216.39	12,973.30	13,024.90	13,167.38	13,046.68	13,513.02	12,952.83	15,892.71

Fisher ID: 13

LocalName	Scientific Name	Average price (Riel/kg)							
		Apr	May	Jun	Jul	Aug	Sep	Oct	Nov
<i>Meuk huysay</i>	<i>Sepioteuthis lessoniana</i>	13,215.69	12,607.59	12,496.53	12,989.36	12,805.97	13,039.22	13,634.15	14,988.17
<i>Meuk bompong/meit</i>	<i>Uroteuthis duvaucelii</i>	10,000.00	12,000.00	12,000.00	12,000.00			12,000.00	12,000.00
Average		13,184.47	12,553.85	12,448.28	12,885.71	12,805.97	13,039.22	13,558.14	14,885.71

Fisher ID: 14

LocalName	Scientific Name	Average price (Riel/kg)							
		Apr	May	Jun	Jul	Aug	Sep	Oct	Nov
<i>Trey teukei ouch kiev</i>	<i>Plectropomus oligacanthus</i>	80,000.00	80,000.00	80,000.00	80,000.00	80,000.00	80,000.00	80,000.00	80,000.00
<i>Trey teukei para krrohorm</i>	<i>Epinephelus fasciatus</i>	32,000.00	32,000.00	32,000.00	32,000.00	32,577.78	32,484.85	32,840.34	32,000.00

<i>Trey teukei para</i>	<i>Epinephelus quoyanus</i>	12,000.00	12,000.00	12,000.00		16,000.00	16,000.00	16,000.00	16,000.00
<i>Trey kantang phleung</i>	<i>Siganus guttatus</i>	2,000.00	2,880.95	3,000.00	3,000.00	3,000.00	3,000.00	3,000.00	3,000.00
<i>Trey seik</i>	<i>Scarus spp.</i>	2,000.00	2,000.00	2,000.00	2,000.00	2,000.00	2,000.00	2,000.00	2,000.00
<i>Trey ka chy</i>	<i>Diagramma pictum</i>		5,000.00	5,000.00	5,000.00	5,000.00	5,000.00	5,000.00	5,000.00
<i>Trey teukeokao</i>	(blank)				32,000.00				
<i>Kdam Thmor</i>	<i>Scylla serrata</i>				15,000.00				
<i>Trey borbel ork</i>	<i>Aetobatus narinari</i>							5,000.00	
<i>Trey kantang kra-ob</i>	<i>Siganus canaliculatus</i>				3,000.00				
<i>Trey borbel khleing</i>	<i>Mobula japonica</i>		4,304.35						
Average		18,666.67	38,610.26	21,571.82	23,544.52	23,156.37	22,422.61	22,441.98	27,060.66

Fisher ID: 15

LocalName	Scientific Name	Average price (Riel/kg)							
		Apr	May	Jun	Jul	Aug	Sep	Oct	Nov
<i>Meuk huysay</i>	<i>Sepioteuthis lessoniana</i>	13,360.00	11,861.48	10,903.99	11,096.15	12,242.77	12,233.53	13,439.02	12,495.73
<i>Meuk bompong/meit</i>	<i>Uroteuthis duvaucelii</i>	10,000.00	10,000.00	10,000.00	10,000.00	10,000.00	10,000.00		10,000.00
Average		13,230.77	11,589.21	10,863.32	10,999.30	12,089.23	12,181.29	13,439.02	12,407.54

Fisher ID: 16

LocalName	Scientific Name	Average price (Riel/kg)							
		Apr	May	Jun	Jul	Aug	Sep	Oct	Nov
<i>Meuk dong</i>	<i>Sepiola sp.</i>		9,500.00	9,500.00	10,000.00	12,000.00	12,000.00	10,625.00	10,000.00

Fisher ID: 17

LocalName	Scientific Name	Average price (Riel/kg)							
		Apr	May	Jun	Jul	Aug	Sep	Oct	Nov
<i>Kdam Thmor</i>	<i>Scylla serrata</i>	10,405.06	13,037.04	14,436.49	15,311.64	15,349.42	17,931.40	17,366.29	17,821.73

Fisher ID: 18

LocalName	Scientific Name	Average price (Riel/kg)							
		Apr	May	Jun	Jul	Aug	Sep	Oct	Nov
<i>Kdam Thmor</i>	<i>Scylla serrata</i>	10,209.42	11,832.06	11,633.76	12,033.28	12,329.30	11,669.57	11,779.53	11,460.32

Fisher ID: 19

LocalName	Scientific Name	Average price (Riel/kg)							
		Apr	May	Jun	Jul	Aug	Sep	Oct	Nov
<i>Ngveav sor</i>	<i>Periglypta sp.</i>	600.00	600.00	600.00	615.35	614.77	625.42	601.38	
<i>Kdam Thmor</i>	<i>Scylla serrata</i>							18,000.00	
Average		600.00	600.00	600.00	615.35	614.77	625.42	668.05	

Fisher ID: 20

LocalName	Scientific Name	Average price (Riel/kg)							
		Apr	May	Jun	Jul	Aug	Sep	Oct	Nov
<i>Kdam Chor</i>	<i>Episesarma singaporensis</i>	4,000.00	4,201.73	4,500.00	4,500.00		4,178.20	4,000.00	4,482.82
<i>Kdam Thmor</i>	<i>Scylla serrata</i>	8,571.43	8,262.30	12,578.95	10,931.51		12,947.37	11,818.18	13,000.00
<i>Trey phtourk</i>	<i>Rachycentron canadum</i>					8,000.00			
<i>Kreing chheam</i>	<i>Anadara granosa</i>	6,000.00	5,900.00						
<i>Bangkea sor</i>	(blank)	8,000.00							
<i>Khyorng Krachorkseih</i>	(blank)	5,000.00		6,000.00					
<i>Khyorng dong</i>	<i>Melo melo</i>							5,000.00	
<i>Ngveav sor</i>	<i>Periglypta sp.</i>	600.00							
<i>Trey kantang kra-ob</i>	<i>Siganus canaliculatus</i>	4,000.00							
Average		4,116.39	4,940.71	5,099.07	5,870.80	8,000.00	5,114.23	4,487.67	4,704.46

Fisher ID: 21

LocalName	Scientific Name	Average price (Riel/kg)							
		Apr	May	Jun	Jul	Aug	Sep	Oct	Nov
<i>Kdam Chor</i>	<i>Episesarma singaporensis</i>	4,000.00	4,000.00	4,447.90	4,500.00	4,500.00	4,000.00	4,000.00	4,500.00
<i>Kdam Thmor</i>	<i>Scylla serrata</i>	10,903.23	11,926.47	11,878.05	14,572.25	12,187.50	12,648.65	11,076.92	11,750.00
<i>Ngeav phourk</i>	<i>Polymesoda erosa</i>		600.00						
Average		4,545.92	4,367.03	5,012.04	6,842.07	5,367.72	4,610.69	4,326.24	4,732.93

Fisher ID: 22

LocalName	Scientific Name	Average price (Riel/kg)							
		Apr	May	Jun	Jul	Aug	Sep	Oct	Nov
<i>Kdam Thmor</i>	<i>Scylla serrata</i>		16,599.79	17,740.98	16,293.90	16,984.06	16,225.23	17,396.44	19,024.08
<i>Kdam Seih</i>	<i>Portunus pelagicus</i>		7,427.62	7,171.01	6,883.66	8,297.57	7,616.04	7,571.66	6,753.82
<i>Kdam Sor/Dangkeab Sor</i>	<i>Charybdis anisodon</i>		226.35	451.43	500.00	500.00	500.00		
<i>Kdam skor krorb/Kdam sor</i>	<i>Matuta victor</i>							500.00	
<i>Bangkea sor</i>	(blank)								500.00
Average			8,456.04	8,831.15	9,456.09	9,268.62	9,445.94	11,601.99	9,217.29

Fisher ID: 23

LocalName	Scientific Name	Average price (Riel/kg)							
		Apr	May	Jun	Jul	Aug	Sep	Oct	Nov
<i>Trey beika</i>	<i>Auxis thazard</i>			3,000.00				1,685.27	827.27
<i>Trey kamoy</i>	<i>Anodontostoma chacunda</i>		2,000.00	2,000.00	1,993.66	2,037.04	2,000.00	1,679.03	1,014.17
<i>Trey kamong khoun khley</i>	<i>Rastrelliger brachysoma</i>			2,000.00	2,000.00	2,000.00	2,153.06		
<i>Bangkea sor</i>	(blank)	8,750.00	7,736.32	10,000.00		8,297.30		7,391.30	
<i>Bangkea para/kloy</i>	<i>Penaeus semisulcatus</i>		6,000.00	6,000.00		6,000.00	6,000.00	7,000.00	
<i>Trey Sambor Hear</i>	(blank)					1,000.00	1,022.78	833.33	500.00
<i>Meuk dong</i>	<i>Sepiola sp.</i>								10,000.00
<i>Trey karav</i>	<i>Eleutheronema tetradactylum</i>			5,000.00	6,000.00		5,000.00	5,000.00	
<i>Trey sleuk russey</i>	<i>Thryssa hamiltonii</i>	1,500.00	1,462.50	1,500.00	1,500.00		1,667.69	1,500.00	

<i>Bangkea kor gneub</i>	<i>Penaeus merguensis</i>								7,000.00
<i>Trey khut/trorchiek damrey</i>	<i>Ephippus orbis</i>			2,000.00					
<i>Trey borbel moan</i>	<i>Himantura imbricata</i>		6,000.00		5,000.00				
<i>Trey andat chhke</i>	<i>Paraplagusia bilineata</i>				1,500.00	1,500.00			
<i>Trey ka-ok</i>	<i>Arius maculatus</i>					3,000.00	3,000.00		
<i>Trey udouy</i>	<i>Selaroides leptolepis</i>					3,000.00			
<i>Trey andeng poy</i>	(blank)				3,000.00				
<i>Trey damlong</i>	(blank)							5,000.00	
<i>Kdam Seih</i>	<i>Portunus pelagicus</i>	8,000.00							
<i>Meuk snok</i>	<i>Sepiella inermis</i>	3,000.00							
<i>Trey Kor/Romeas</i>	<i>Monacanthus chinensis</i>	2,500.00							
<i>X-others</i>	<i>X-others</i>	1,000.00	4,990.45	1,000.00	1,020.33	1,357.14			
Average		5,524.83	6,370.72	2,029.90	2,112.60	2,763.99	2,006.69	1,688.52	1,042.53

Fisher ID: 24

LocalName	Scientific Name	Average price (Riel/kg)							
		Apr	May	Jun	Jul	Aug	Sep	Oct	Nov
<i>Kdam Thmor</i>	<i>Scylla serrata</i>	20,308.47	18,270.68	15,856.34	16,617.79	16,234.35	16,586.24	15,803.74	16,384.21
<i>Kdam Seih</i>	<i>Portunus pelagicus</i>	8,055.97	7,292.51	6,808.48	6,196.62	6,536.02	6,236.21	6,317.34	6,305.56
<i>Bangkea sor</i>	(blank)								500.00
<i>Kdam skor krorb/Kdam sor</i>	<i>Matuta victor</i>							500.00	500.00
<i>Kdam Sor/Dangkeab Sor</i>	<i>Charybdis anisodon</i>	500.00	478.16	500.00	500.00	500.00	500.00		
Average		12,594.46	8,993.42	8,471.30	8,855.41	9,145.29	9,639.39	9,011.86	9,190.30

Fisher ID: 25

LocalName	Scientific Name	Average price (Riel/kg)							
		Apr	May	Jun	Jul	Aug	Sep	Oct	Nov
<i>Kdam Thmor</i>	<i>Scylla serrata</i>	14,761.90	11,746.03	15,562.72	13,818.18	14,724.41	18,333.33	13,530.03	13,585.90
<i>Kdam Seih</i>	<i>Portunus pelagicus</i>	7,045.45	7,081.00	7,961.96	8,842.93	9,000.00	9,000.00	8,515.38	7,584.49
<i>Trey kalang</i>	<i>Scomberoides commersonianus</i>							24,589.74	

<i>Trey balang</i>	<i>Rastrelliger faughni</i>								17,591.84
<i>Trey borbel moan</i>	<i>Himantura imbricata</i>	4,358.09	2,828.77	3,000.00	3,000.00	3,000.00	3,000.00	3,000.00	3,000.00
<i>Trey beika</i>	<i>Auxis thazard</i>		2,000.00		13,600.00	15,000.00	2,000.00	1,500.00	
<i>Trey andat chhke</i>	<i>Paraplagusia bilineata</i>	1,250.00	2,818.18		3,000.00	3,000.00	3,000.00	3,000.00	
<i>Trey karav</i>	<i>Eleutheronema tetradactylum</i>		3,000.00	5,000.00		4,633.33	4,000.00	3,000.00	
<i>Trey ka-ok kbal heang/Ka-ok prak</i>	<i>Osteogeneiosus militaris</i>							5,000.00	
<i>Trey andeng</i>	(blank)				5,000.00		5,205.88		
<i>Trey Kor/Romeas</i>	<i>Monacanthus chinensis</i>		0.00	0.00	2,250.00	2,000.00			
<i>Trey kantong kra-ob</i>	<i>Siganus canaliculatus</i>		2,000.00		2,000.00	10,000.00			
<i>Trey ka-ok</i>	<i>Arius maculatus</i>		4,156.25	3,000.00	2,000.00				
<i>Trey kamong khoun khley</i>	<i>Rastrelliger brachysoma</i>		2,000.00			2,000.00			
<i>Trey kantuy krabey</i>	(blank)	1,200.00	2,000.00	3,000.00	3,640.00				
<i>Trey sleuk russey</i>	<i>Thryssa hamiltonii</i>			1,500.00		2,000.00			
<i>Kdam kom</i>	(blank)						2,500.00		
<i>Kdam skor krorb/Kdam sor</i>	<i>Matuta victor</i>				2,000.00				
<i>X-others</i>	<i>X-others</i>	1,500.00		1,843.43	1,943.37	1,852.94	3,585.37	1,000.00	
Average		9,177.63	7,246.17	8,808.86	7,273.43	6,456.99	8,285.71	9,525.16	10,249.67

Fisher ID: 26

LocalName	Scientific Name	Average price (Riel/kg)							
		Apr	May	Jun	Jul	Aug	Sep	Oct	Nov
<i>Kreing chheam</i>	<i>Anadara granosa</i>	4,000.00	4,128.94	4,017.07	4,500.00	4,580.26	4,500.00	6,847.83	4,604.25
<i>Trey teukei ouch kiev</i>	<i>Plectropomus oligacanthus</i>				80,000.00				
<i>Trey teukei para krorthorm</i>	<i>Epinephelus fasciatus</i>				32,000.00				
<i>Trey ka chy</i>	<i>Diagramma pictum</i>				5,000.00				
<i>Trey kantong phleung</i>	<i>Siganus guttatus</i>				3,000.00				
<i>Trey seik</i>	<i>Scarus spp.</i>				2,000.00				
Average		4,000.00	4,128.94	4,017.07	8,312.68	4,580.26	4,500.00	6,847.83	4,604.25

Annex 8: Average maximum length (cm) versus frequencies by fisher measured over the period from April to November 2011

Fisher ID	Local Name	Scientific Name	Maximum length		StdDev
			Frequencies	Average (cm)	
1	<i>Bangkorng kandob</i>	<i>Clorida decorata</i>	1	2.10	
	<i>Kdam krahorm</i>		2	12.00	1.41
	<i>Kdam neak</i>	<i>Charybdis natator</i>	6	10.92	1.96
	<i>Kdam Seih</i>	<i>Portunus pelagicus</i>	29	11.26	1.05
	<i>Kdam skor krorb/Kdam sor</i>	<i>Matuta victor</i>	21	4.24	2.10
	<i>Meuk huysay</i>	<i>Sepioteuthis lessoniana</i>	5	15.60	3.36
	<i>Trey andeng</i>	(blank)	2	22.00	1.41
	<i>Trey andeng poy</i>	(blank)	1	26.00	
	<i>Trey angrei</i>	<i>Sphyræna putnamae</i>	27	39.05	13.11
	<i>Trey balang</i>	<i>Rastrelliger faughni</i>	1	20.50	
	<i>Trey chek</i>	<i>Seriolina nigrofasciata</i>	57	12.74	1.18
	<i>Trey chhlam chhkout</i>	<i>Chiloscyllium griseum</i>	17	44.98	23.14
	<i>Trey deik koul</i>	(blank)	1	10.10	
	<i>Trey do angkor</i>	<i>Gerres abbreviatus</i>	102	12.88	3.88
	<i>Trey kalang</i>	<i>Scomberoides commersonianus</i>	1	50.20	
	<i>Trey kamong khoun khley</i>	<i>Rastrelliger brachysoma</i>	33	14.64	1.39
	<i>Trey kamoy</i>	<i>Anodontostoma chacunda</i>	37	11.95	2.39
	<i>Trey kantang kra-ob</i>	<i>Siganus canaliculatus</i>	17	10.91	0.93
	<i>Trey kbork</i>	<i>Moolgarda seheli</i>	1	18.90	
	<i>Trey Kor/Romeas</i>	<i>Monacanthus chinensis</i>	71	12.57	0.97
	<i>Trey koun</i>	<i>Hilsa keele</i>	33	11.94	1.96
	<i>Trey Krorb khnol</i>	<i>Lethrinus nebulosus</i>	66	12.53	1.06
	<i>Trey krorhorm</i>	<i>Lutjanus malabaricus</i>	25	17.01	3.31
	<i>Trey krorhorm kbalkor</i>	(blank)	51	16.66	2.90
	<i>Trey krorhorm srakakras</i>	<i>Scolopis taeniopterus</i>	54	15.63	2.51
	<i>Trey phneik Lan</i>	<i>Priacanthus hamrur</i>	18	16.44	1.63
	<i>Trey prorkong</i>	(blank)	4	14.73	1.13
	<i>Trey prorleuk</i>	<i>Leiognathus splendens</i>	64	16.55	4.01
	<i>Trey prorlos</i>	<i>Sillago sihama</i>	39	18.72	2.16
	<i>Trey sach</i>	(blank)	1	23.00	
	<i>Trey trorsork</i>	<i>Terapon theraps</i>	1	10.10	
<i>Trey udouy</i>	<i>Selaroides leptolepis</i>	26	17.27	2.76	
2	<i>Kdam Seih</i>	<i>Portunus pelagicus</i>	270	11.56	2.67
	<i>Khyorng dong</i>	<i>Melo melo</i>	104	19.66	3.84
	<i>Khyorng thnot</i>	<i>Conus vexillum</i>	125	18.26	1.96
	<i>Trey borbel moan</i>	<i>Himantura imbricata</i>	13	16.46	1.39
	<i>Trey borbel ork</i>	<i>Aetobatus narinari</i>	9	17.78	2.22
	<i>Trey chenchas/pakchiv</i>	<i>Alectis ciliaris</i>	2	31.00	1.41
	<i>Trey chhlam chhkout</i>	<i>Chiloscyllium griseum</i>	20	31.25	6.85
<i>Trey kam kourch</i>	<i>Carangoides ferdau</i>	1	27.00		

3	<i>Bangkea kor gneub</i>	<i>Penaeus merguensis</i>	3	12.00	1.73
	<i>Bangkea sor</i>	(blank)	36	6.34	4.89
	<i>Bangkorng kandob</i>	<i>Clorida decorata</i>	70	13.09	0.86
	<i>Bangkorng pak</i>	(blank)	37	17.41	1.98
	<i>Kdam krahorm</i>		76	2.22	1.19
	<i>Kdam neak</i>	<i>Charybdis natator</i>	16	10.00	1.55
	<i>Kdam pingpeang</i>	<i>Arcania undecimspinosa</i>			
	<i>Kdam Seih</i>	<i>Portunus pelagicus</i>	194	8.84	2.05
	<i>Kdam skor krorb/Kdam sor</i>	<i>Matuta victor</i>	84	2.51	1.98
	<i>Kdam Sor/Dangkeab Sor</i>	<i>Charybdis anisodon</i>	13	3.38	2.63
	<i>Kdam Thmor</i>	<i>Scylla serrata</i>	46	11.73	4.09
	<i>Khyorng thnot</i>	<i>Conus vexillum</i>	1	2.00	
	<i>Meuk bompong/meit</i>	<i>Uroteuthis duvaucelii</i>	3	7.33	1.15
	<i>Trey antanou</i>	<i>Pentapodus setosus</i>	10	9.80	1.81
	<i>Trey Kor/Romeas</i>	<i>Monacanthus chinensis</i>	3	8.00	0.00
4	<i>Bangkea kor gneub</i>	<i>Penaeus merguensis</i>	143	4.10	1.80
	<i>Bangkea para/kloy</i>	<i>Penaeus semisulcatus</i>	5	7.24	1.73
	<i>Bangkea sor</i>	(blank)	58	8.26	1.71
	<i>Kdam Thmor</i>	<i>Scylla serrata</i>	10	11.49	5.30
	<i>Meuk bompong/meit</i>	<i>Uroteuthis duvaucelii</i>	1	9.00	
	<i>Trey angkeuy</i>	<i>Lutjanus johnii</i>			
	<i>Trey mrech</i>	(blank)	6	4.17	0.41
	<i>Trey phtong prortkheat</i>	<i>Ablennes hians</i>	5	8.78	4.31
	<i>Trey prorkong</i>	(blank)	7	5.71	1.11
5	<i>Bangkea kor gneub</i>	<i>Penaeus merguensis</i>	13	13.46	2.96
	<i>Bangkea para/kloy</i>	<i>Penaeus semisulcatus</i>	1	19.00	
	<i>Bangkea phourk</i>	(blank)	83	8.93	2.67
	<i>Bangkea sor</i>	(blank)	109	9.51	3.67
	<i>Kdam Seih</i>	<i>Portunus pelagicus</i>	10	14.70	1.34
	<i>Kdam Thmor</i>	<i>Scylla serrata</i>	29	12.52	4.01
	<i>Trey andeng</i>	(blank)	6	15.67	3.44
	<i>Trey andeng poy</i>	(blank)	2	17.00	2.83
	<i>Trey chek</i>	<i>Seriolina nigrofasciata</i>	13	9.38	1.33
	<i>Trey damlong</i>	(blank)	69	10.59	2.92
	<i>Trey do angkor</i>	<i>Gerres abbreviatus</i>	2	16.50	4.95
	<i>Trey kanchanh chras</i>	(blank)	18	7.11	2.14
	<i>Trey kantong kra-ob</i>	<i>Siganus canaliculatus</i>	5	7.60	3.13
	<i>Trey kbork</i>	<i>Moolgarda seheli</i>	1	15.00	
	<i>Trey kranh</i>	<i>Lobotes surinamensis</i>	35	10.83	3.49
	<i>Trey mrech</i>	(blank)	142	11.97	4.06
	<i>Trey prorkong</i>	(blank)	94	10.36	2.35
	<i>Trey prorlos</i>	<i>Sillago sihama</i>	8	14.88	1.64
	<i>Trey trorsork</i>	<i>Terapon theraps</i>	2	9.50	0.71
6	<i>Bangkea para/kloy</i>	<i>Penaeus semisulcatus</i>			
	<i>Bangkea sor</i>	(blank)			
	<i>Kdam Seih</i>	<i>Portunus pelagicus</i>	2	15.00	4.24

	<i>Trey beika</i>	<i>Auxis thazard</i>	29	26.28	2.83
	<i>Trey borbel ork</i>	<i>Aetobatus narinari</i>	1	40.00	
	<i>Trey kamong khoun khley</i>	<i>Rastrelliger brachysoma</i>	139	9.35	2.99
	<i>Trey kamong khoun veng</i>	<i>Rastrelliger kanagurta</i>	80	12.58	1.46
	<i>Trey kamoy</i>	<i>Anodontostoma chacunda</i>	1	10.00	
	<i>Trey karav</i>	<i>Eleutheronema tetradactylum</i>	4	13.88	9.70
	<i>Trey Kor/Romeas</i>	<i>Monacanthus chinensis</i>	12	10.42	0.51
	<i>Trey krahorm phneak thom</i>	<i>Priacanthus macracanthus</i>	6	10.00	1.10
	<i>Trey krorkhorm</i>	<i>Lutjanus malabaricus</i>	27	11.22	1.45
	<i>Trey krorkhorm kbalkor</i>	(blank)	29	11.21	1.01
	<i>Trey krorkhorm phneklien</i>	<i>Priacanthus tayenus</i>	8	10.00	2.45
	<i>Trey krorkhorm srakakras</i>	<i>Scolopis taeniopterus</i>	30	10.17	0.38
7	<i>Kdam Chor</i>	<i>Episesarma singaporensis</i>	1	5.00	
	<i>Kdam Seih</i>	<i>Portunus pelagicus</i>	430	13.08	5.74
	<i>Kdam skor krorb/Kdam sor</i>	<i>Matuta victor</i>	1	19.00	
8	<i>Bangkea kor gneub</i>	<i>Penaeus merguensis</i>	30	8.87	1.21
	<i>Bangkea para/kloy</i>	<i>Penaeus semisulcatus</i>	4	14.00	1.15
	<i>Bangkea sor</i>	(blank)	69	7.79	1.48
	<i>Bangkornng pak</i>	(blank)	2	16.50	2.12
	<i>Kdam Seih</i>	<i>Portunus pelagicus</i>	5	8.80	4.66
	<i>Kdam Thmor</i>	<i>Scylla serrata</i>	35	7.96	2.73
	<i>Trey andeng poy</i>	(blank)	2	29.50	0.71
	<i>Trey damlong</i>	(blank)	15	4.65	2.93
	<i>Trey deik koul</i>	(blank)	6	2.90	0.72
	<i>Trey do angkor</i>	<i>Gerres abbreviatus</i>	1	5.00	
	<i>Trey kantang kra-ob</i>	<i>Siganus canaliculatus</i>	42	8.58	3.10
	<i>Trey kantang phleung</i>	<i>Siganus guttatus</i>	6	19.33	7.55
	<i>Trey kantang prorpeh</i>	<i>Siganus argenteus</i>	8	15.44	4.48
	<i>Trey kantang thmor</i>	<i>Siganus virgatus</i>	32	10.07	3.94
	<i>Trey karav</i>	<i>Eleutheronema tetradactylum</i>	17	16.76	6.08
	<i>Trey kbork</i>	<i>Moolgarda seheli</i>	64	22.08	5.53
	<i>Trey kbork angkam</i>	<i>Moolgarda bucharani</i>	1	12.00	
	<i>Trey khe/proy veng</i>	<i>Pentaprion longimanus</i>	1	25.50	
	<i>Trey Krorb khnol</i>	<i>Lethrinus nebulosus</i>	3	5.33	1.53
	<i>Trey mrech</i>	(blank)	2	8.00	2.83
	<i>Trey phtong phka</i>	<i>Hemiramphus far</i>	46	14.23	13.80
	<i>Trey phtong prortkheat</i>	<i>Ablennes hians</i>	6	27.83	5.08
	<i>Trey prorkong</i>	(blank)	2	9.50	2.12
	<i>Trey prorlos</i>	<i>Sillago sihama</i>	14	9.00	3.44
	<i>Trey pthong prort</i>	<i>Tylosurus acus melanotus</i>	12	14.16	8.11
9	<i>Bangkea para/kloy</i>	<i>Penaeus semisulcatus</i>			
	<i>Bangkea sor</i>	(blank)			
	<i>Kdam Thmor</i>	<i>Scylla serrata</i>	4	12.50	4.36
	<i>Trey chhlam chhkout</i>	<i>Chiloscyllium griseum</i>	5	70.00	7.07
	<i>Trey kamong khoun khley</i>	<i>Rastrelliger brachysoma</i>	207	6.63	1.49

		<i>Eleutheronema tetradactylum</i>	10	25.80	11.82	
10	<i>Trey karav</i>	<i>Penaeus merguensis</i>	39	7.00	0.97	
	<i>Bangkea para/kloy</i>	<i>Penaeus semisulcatus</i>	21	10.90	5.95	
	<i>Bangkea sor</i>	(blank)	42	8.77	1.09	
	<i>Kdam Seih</i>	<i>Portunus pelagicus</i>	3	13.00	7.21	
	<i>Kdam Thmor</i>	<i>Scylla serrata</i>	32	8.80	2.20	
	<i>Meuk bompong/meit</i>	<i>Uroteuthis duvaucelii</i>	1	5.30		
	<i>Meuk snok</i>	<i>Sepiella inermis</i>	5	5.20	0.84	
	<i>Trey andat chhke</i>	<i>Paraplagusia bilineata</i>				
	<i>Trey andeng poy</i>	(blank)	1	41.00		
	<i>Trey angrei</i>	<i>Sphyræna putnamae</i>	2	27.00	0.00	
	<i>Trey balang</i>	<i>Rastrelliger faughni</i>	1	7.00		
	<i>Trey damlong</i>	(blank)	37	8.55	2.41	
	<i>Trey deik koul</i>	(blank)	15	5.33	1.50	
	<i>Trey do angkor</i>	<i>Gerres abbreviatus</i>	1	4.00		
	<i>Trey kanchanh chras</i>	(blank)	1	3.00		
	<i>Trey kantong kra-ob</i>	<i>Siganus canaliculatus</i>	35	10.11	3.10	
	<i>Trey kantong phleung</i>	<i>Siganus guttatus</i>	8	15.38	2.20	
	<i>Trey kantong prorpeh</i>	<i>Siganus argenteus</i>	4	12.50	1.73	
	<i>Trey kantong thmor</i>	<i>Siganus javus</i>	26	11.73	2.15	
	<i>Trey kantong thmor</i>	<i>Siganus virgatus</i>	1	14.00		
	<i>Trey ka-ok</i>	<i>Arius maculatus</i>	1	38.00		
			<i>Eleutheronema tetradactylum</i>	6	16.17	1.72
		<i>Trey kbork</i>	<i>Moolgarda seheli</i>	85	22.22	4.15
		<i>Trey kbork angkam</i>	<i>Moolgarda buchanani</i>	4	13.50	5.80
		<i>Trey khei/proy veng</i>	<i>Pentaprion longimanus</i>	2	11.00	9.90
		<i>Trey Krorb khnol</i>	<i>Lethrinus nebulosus</i>	5	12.30	1.48
		<i>Trey mrech</i>	(blank)	17	14.41	2.94
	<i>Trey phtong phka</i>	<i>Hemiramphus far</i>	2	14.00	5.66	
	<i>Trey prorkong</i>	(blank)	4	8.00	2.71	
	<i>Trey prorleuk</i>	<i>Leiognathus splendens</i>	11	8.82	2.09	
	<i>Trey prorlos</i>	<i>Sillago sihama</i>	7	12.43	1.72	
	<i>Trey pthong prort</i>	<i>Tylosurus acus melanotus</i>	12	16.79	6.18	
	<i>Trey spong</i>	<i>Psammoperca waigiensis</i>	1	4.00		
11	<i>Kdam neak</i>	<i>Charybdis natator</i>	1	23.00		
	<i>Ngveav sor</i>	<i>Periglypta sp.</i>	1	9.00		
	<i>Trey kantong kra-ob</i>	<i>Siganus canaliculatus</i>	28	20.29	0.46	
	<i>Trey kantong phleung</i>	<i>Siganus guttatus</i>	46	18.55	5.50	
	<i>Trey kantong prorpeh</i>	<i>Siganus argenteus</i>	24	19.71	2.26	
	<i>Trey phneik Lan</i>	<i>Priacanthus hamrur</i>	52	20.41	4.09	
	<i>Trey seik</i>	<i>Scarus spp.</i>	69	20.91	6.21	
	<i>Trey seik thmor</i>	<i>Scarus spp.</i>	30	21.20	0.71	
	<i>Trey teukei koav</i>	<i>Epinephelus coioides</i>	32	21.97	5.18	
	<i>Trey teukei ouch kiev</i>	<i>Plectropomus oligacanthus</i>	65	20.88	5.06	
	<i>Trey teukei para</i>	<i>Epinephelus quoyanus</i>	42	17.39	5.73	

	<i>Trey teukei para krorthorm</i>	<i>Epinephelus fasciatus</i>	6	5.75	7.35
	<i>Trey teukei thmor</i>	<i>Aethaloperca rogaa</i>	1	10.00	
	<i>Trey teukeokao</i>	(blank)	16	22.13	1.26
12	<i>Kdam krahorm</i>		1	2.00	
	<i>Kdam Seih</i>	<i>Portunus pelagicus</i>	3	5.10	2.59
	<i>Kdam Sor/Dangkeab Sor</i>	<i>Charybdis anisodon</i>	1	2.00	
	<i>Kdam Thmor</i>	<i>Scylla serrata</i>	4	6.30	3.53
	<i>Meuk bompong/meit</i>	<i>Uroteuthis duvaucelii</i>	7	22.43	6.63
	<i>Meuk huysay</i>	<i>Sepioteuthis lessoniana</i>	334	15.54	5.75
13	<i>Meuk bompong/meit</i>	<i>Uroteuthis duvaucelii</i>	22	20.86	4.38
	<i>Meuk huysay</i>	<i>Sepioteuthis lessoniana</i>	191	15.99	4.48
14	<i>Kdam Thmor</i>	<i>Scylla serrata</i>	2	9.50	0.71
	<i>Trey borbel khleing</i>	<i>Mobula japonica</i>	3	28.67	0.58
	<i>Trey borbel ork</i>	<i>Aetobatus narinari</i>	2	29.50	0.71
	<i>Trey ka chy</i>	<i>Diagramma pictum</i>	55	27.17	3.63
	<i>Trey kantong kra-ob</i>	<i>Siganus canaliculatus</i>	2	24.50	0.71
	<i>Trey kantong phleung</i>	<i>Siganus guttatus</i>	59	21.85	3.13
	<i>Trey seik</i>	<i>Scarus spp.</i>	62	23.91	3.14
	<i>Trey teukei ouch kiev</i>	<i>Plectropomus oligacanthus</i>	62	27.68	3.65
	<i>Trey teukei para</i>	<i>Epinephelus quoyanus</i>	42	24.25	4.02
	<i>Trey teukei para krorthorm</i>	<i>Epinephelus fasciatus</i>	55	27.57	3.89
	<i>Trey teukeokao</i>	(blank)	1	26.00	
15	<i>Meuk bompong/meit</i>	<i>Uroteuthis duvaucelii</i>	37	26.57	1.24
	<i>Meuk huysay</i>	<i>Sepioteuthis lessoniana</i>	245	17.98	5.99
16	<i>Meuk dong</i>	<i>Sepiola sp.</i>	93	5.00	0.00
17	<i>Kdam Thmor</i>	<i>Scylla serrata</i>	469	10.19	1.77
18	<i>Kdam Thmor</i>	<i>Scylla serrata</i>	335	9.36	1.73
	(blank)	(blank)			
19	<i>Kdam Thmor</i>	<i>Scylla serrata</i>	1	11.00	
	<i>Ngveav sor</i>	<i>Periglypta sp.</i>	171	8.57	0.62
	(blank)	<i>Periglypta sp.</i>	1	8.00	
20	<i>Bangkea sor</i>	(blank)	1	8.00	
	<i>Kdam Chor</i>	<i>Episesarma singaporensis</i>	141	5.12	0.63
	<i>Kdam Thmor</i>	<i>Scylla serrata</i>	86	8.54	1.42
	<i>Khyorng dong</i>	<i>Melo melo</i>			
	<i>Khyorng Krachorkseih</i>	(blank)	3	1.00	0.00
	<i>Kreing chheam</i>	<i>Anadara granosa</i>	5	6.40	1.39
	<i>Ngveav sor</i>	<i>Periglypta sp.</i>	2	6.00	1.41
	<i>Trey kantong kra-ob</i>	<i>Siganus canaliculatus</i>			
	<i>Trey phtourk</i>	<i>Rachycentron canadum</i>	18	29.56	3.60
21	<i>Kdam Chor</i>	<i>Episesarma singaporensis</i>	161	4.98	0.24
	<i>Kdam Thmor</i>	<i>Scylla serrata</i>	123	9.48	1.72
	<i>Ngeav phourk</i>	<i>Polymesoda erosa</i>	1	6.00	
22	<i>Bangkea sor</i>	(blank)	1	2.00	
	<i>Kdam Seih</i>	<i>Portunus pelagicus</i>	189	3.98	0.63
	<i>Kdam skor krorthorm/Kdam sor</i>	<i>Matuta victor</i>			

	<i>Kdam Sor/Dangkeab Sor</i>	<i>Charybdis anisodon</i>	26	2.62	0.36
	<i>Kdam Thmor</i>	<i>Scylla serrata</i>	511	5.32	1.15
23	<i>Bangkea kor gneub</i>	<i>Penaeus merguensis</i>	2	8.50	0.71
	<i>Bangkea para/kloy</i>	<i>Penaeus semisulcatus</i>	20	10.55	0.83
	<i>Bangkea sor</i>	(blank)	33	10.80	1.61
	<i>Kdam Seih</i>	<i>Portunus pelagicus</i>	1	12.00	
	<i>Meuk dong</i>	<i>Sepiola sp.</i>	1	5.00	
	<i>Meuk snok</i>	<i>Sepiella inermis</i>	1	7.50	
	<i>Trey andat chhke</i>	<i>Paraplagusia bilineata</i>	8	14.63	1.06
	<i>Trey andeng poy</i>	(blank)	1	13.00	
	<i>Trey beika</i>	<i>Auxis thazard</i>	54	13.48	0.88
	<i>Trey borbrel moan</i>	<i>Himantura imbricata</i>	2	19.00	0.00
	<i>Trey damlong</i>	(blank)	1	30.00	
	<i>Trey kamong khoun khley</i>	<i>Rastrelliger brachysoma</i>	57	13.30	1.68
	<i>Trey kamoy</i>	<i>Anodontostoma chacunda</i>	90	10.61	1.02
	<i>Trey ka-ok</i>	<i>Arius maculatus</i>	4	14.25	0.50
	<i>Trey karav</i>	<i>Eleutheronema tetradactylum</i>	10	15.10	1.73
	<i>Trey khut/trorchiek damrey</i>	<i>Ephippus orbis</i>	7	10.57	0.98
	<i>Trey Kor/Romeas</i>	<i>Monacanthus chinensis</i>	1	15.00	
	<i>Trey Sambor Hear</i>	(blank)	77	8.73	1.24
	<i>Trey sleuk russey</i>	<i>Thryssa hamiltonii</i>	30	14.07	2.45
	<i>Trey udouy</i>	<i>Selaroides leptolepis</i>	3	15.33	0.58
24	<i>Bangkea kor gneub</i>	<i>Penaeus merguensis</i>			
	<i>Bangkea sor</i>	(blank)	24	1.66	0.44
	<i>Kdam Seih</i>	<i>Portunus pelagicus</i>	345	4.45	1.20
	<i>Kdam skor krorb/Kdam sor</i>	<i>Matuta victor</i>	30	1.77	0.32
	<i>Kdam Sor/Dangkeab Sor</i>	<i>Charybdis anisodon</i>	103	2.00	0.42
	<i>Kdam Thmor</i>	<i>Scylla serrata</i>	516	5.63	1.96
25	<i>Kdam kom</i>	(blank)			
	<i>Kdam Seih</i>	<i>Portunus pelagicus</i>	79	11.92	3.23
	<i>Kdam skor krorb/Kdam sor</i>	<i>Matuta victor</i>			
	<i>Kdam Thmor</i>	<i>Scylla serrata</i>	140	10.37	4.07
	<i>Trey andat chhke</i>	<i>Paraplagusia bilineata</i>			
	<i>Trey andeng</i>	(blank)			
	<i>Trey balang</i>	<i>Rastrelliger faughni</i>	4	20.25	5.85
	<i>Trey beika</i>	<i>Auxis thazard</i>	1	25.00	
	<i>Trey borbrel moan</i>	<i>Himantura imbricata</i>			
	<i>Trey kalang</i>	<i>Scomberoides commersonianus</i>			
	<i>Trey kamong khoun khley</i>	<i>Rastrelliger brachysoma</i>	1	2.00	
	<i>Trey kantong kra-ob</i>	<i>Siganus canaliculatus</i>	1	10.00	
	<i>Trey kantuy krabey</i>	(blank)	1	25.00	
	<i>Trey ka-ok</i>	<i>Arius maculatus</i>			
	<i>Trey ka-ok kbal heang/Ka-ok prak</i>	<i>Osteogeneiosus militaris</i>			
	<i>Trey karav</i>	<i>Eleutheronema</i>	1	20.00	

		<i>tetradactylum</i>			
	<i>Trey Kor/Romeas</i>	<i>Monacanthus chinensis</i>			
	<i>Trey sleuk russey</i>	<i>Thryssa hamiltonii</i>			
26	<i>Kreing chheam</i>	<i>Anadara granosa</i>	116	3.36	0.49
	<i>Trey ka chy</i>	<i>Diagramma pictum</i>	1	29.00	
	<i>Trey kantang phleung</i>	<i>Siganus guttatus</i>	1	26.00	
	<i>Trey seik</i>	<i>Scarus spp.</i>	1	23.00	
	<i>Trey teukei ouch kiev</i>	<i>Plectropomus oligacanthus</i>	1	28.00	
	<i>Trey teukei para krorthorm</i>	<i>Epinephelus fasciatus</i>	1	25.00	

RFLP

FAO Representation Office:

5, St 370, Beung Kang Kang I
Khan Chamkar Morn, Phnom Penh, Cambodia
H/P : +855 (0) 23 216 566
Fax : +855 (0) 23 216 547
Website: www.fao.org

RFLP Office in Cambodia:

Group 4, Sangkat 2
Khan Mittapheap, Preah Sihanouk Province
H/P : +855 (0) 34 934 638
Fax : +855 (0) 34 934 638
Website: www.rflp.org