

# Allergological evaluation of a dog population in a veterinary immuno-allergology consultation: What correlates in a canine model



Dias JS<sup>1</sup>, Pereira LS<sup>2</sup>, Goicoa A<sup>3</sup>, Semião-Santos SJ<sup>4</sup>, Bento OP<sup>5,6</sup>, Martins LM<sup>6,7</sup>

1. Department of Veterinary Medicine (Lusophone University of Humanities and Technology) Lisbon, Portugal; 2. Agrarian School of Elvas, Polytechnic Institute of Portalegre, Elvas, Portugal; 3. Rof Codina University Hospital, Veterinary Faculty of Lugo, University of Santiago de Compostela, Lugo, Spain; 4. Nursing and Medicine Departments, Tiradentes University (UNIT), Aracaju, Brazil; 5. Department of Animal Sciences, School of Sciences and Technology; 6. Institute of Mediterranean Agricultural and Environmental Sciences (ICAAM); 7. Department of Veterinary Medicine, School of Sciences and Technology, University of Évora, Évora, Portugal.

## INTRODUCTION

Canine allergic dermatitis (cAD) can be framed in one of the following conditions: i) atopic dermatitis; ii) food allergy dermatitis; iii) flea allergy dermatitis; iv) *Malassezia* dermatitis, or v) contact dermatitis [1]. It is the most frequent condition in veterinary dermatology [2,3] and is associated with either type I or IV hypersensitivity [4]. Flea allergy dermatitis represents a relevant number of cases and results from the response against flea saliva antigens inoculated by the bite, triggering an inflammatory process mostly located at the lumbar and rump regions [5](Fig. 1). A more or less localized inflammatory skin condition will characterize atopic, contact [5,6] or food-induced [7,8,9] dermatitis (Fig. 2 to 4).

With pruritus and its consequences as main complaints, diagnosis is based on clinical history, observed signs and exclusion of other causes of dermatitis, such as parasitic or bacterial infections and food intolerance [10]. The identification of IgE-mediated cutaneous reactivity by intradermal tests (IDT) (Fig. 5) and serological determination of allergen-specific IgE are of considerable complementary diagnostic value [3,11,12,13]. Elimination or hypoallergenic diets are the diagnosis means for food allergy [9].

## AIM

1. Characterization of clinical signs associated with cAD in the study population; 2. Identification of possible correlations between environmental factors, clinical signs, IDT and specific IgE.

## METHODS

1. Patient selection and characterization: Veterinary Hospital of the University of Évora (Portugal) + Rof Codina University Hospital (Lugo, Spain) outpatient consultations → clinical inquiry → 55 dogs with AD (30 F; 25 M) → IDT + specific IgE. Predisposed breeds = 35; Indoor = 41.

2. Procedures: Detailed anamnesis and clinical examination. Dermatology and otology, parasitological and microbiological tests were performed when necessary for further information. IDT were performed using commercial extracts for fungi, grass, herbs and tree pollens, mites, food and flea. Specific IgE were determined according to Lee *et al.* (2009)[14] in a commercial laboratory.

3. Statistical analysis: Data normality was tested with Kolmogorov-Smirnov and homogeneity of variances checked with Levene's test. Descriptive statistics was performed by the percentage of occurrence for all variables. To access the relationship between environmental factors, clinical signs, IDT and specific IgE the parametric Pearson's R test was applied. The level of significance has been set at  $p < 0.05$ . Statistical Package for Social Sciences (SPSS) version 17.0 was used.

## RESULTS and DISCUSSION

- Age at first signs <3 years old in 82% of patients, in accordance to Favrot *et al.* (2010)[15], and <1 year old in 24%.
- Signs: Itching in 70% of patients (generalized in 66%, with 78% of paw licking and chewing).
- Seasonal worsening in 64% of patients.
- Dermatitis was already present in 69.1% of patients, 50% presented external otitis (Fig. 4 and 6) and 28.9% self-induced alopecia. "Intense itching" was found in 10.5%, "medium itching" in 81.6% and "mild itching" in 5.26% of the patients, according to Rybníček *et al.* (2009)[16].
- Positive IDT: 37.3% to *L. destructor*, 29.41% to *D. farinae*, 27.5% to *D. pteronyssinus*, 25.5% to *D. glomerata* and 21.6% to *Malassezia spp.* From the 37 dogs submitted to food IDT, 40.5% revealed positive to beef, 27% to chicken or porc and 5.4% to lamb.
- Specific IgE >150 ELISA Absorbance Units (EAU) was found in 84% of dogs to indoor allergen sources and in 68% to pollens.
- Positive correlation was found between pollen sensitization, seasonality of signs ( $R=0.518$ ;  $p=0.001$ ) and positivity for *D. glomerata* ( $R=0.352$ ;  $p=0.007$ ). Chronic otitis correlated positively with alopecia (fairly:  $R=0.043$ ;  $p=0.008$ ) and reactivity to *L. destructor* ( $R=0.446$ ;  $p=0.008$ ), *P. lanceolata* ( $R=0.435$ ;  $p=0.026$ ) and *P. acerifolia* ( $R=0.402$ ;  $p=0.017$ ).
- Negative correlation was found between outdoor life and the intensity ( $R=-0.333$ ;  $p=0.033$ ) and precocity ( $R=-0.356$ ;  $p=0.026$ ) of signs.
- No correlation was found between ITD and specific IgE, as these methods evaluate different parameters – specific IgE refers to circulating IgE concentrations, while IDT detect mast cell-bound specific IgE [17,18].

## CONCLUSIONS

- Correlation between different clinical signs and positive testing for some allergenic sources may occur, as well as between sensitization to pollens and the beginning, the severity and the seasonality of clinical signs.
- Outdoor life may be a protective factor to the severity of signs.

Contact:  
lmlm@uevora.pt



## REFERENCES

- Martins L, Bento O, Inácio F. Veterinary allergy diagnosis: the past, the present and the improving perspectives. *Allergo J Int.* 2016 (in press).
- Prélaud, P. Dermite atopique canine. *EMC-Vétérinaire.* 2005; 2:14-29.
- Olivry T, DeBoer DJ, Favrot F, Jackson HA, Mueller RS, Nuttall T, Prélaud P. Treatment of canine atopic dermatitis: 2010 clinical practice guidelines from the International Task Force on Canine Atopic Dermatitis. *Vet Dermatol.* 2010; 21:233-248.
- WAO/EAACI Allergy Definitions ([http://www.eaaci.org/attachments/304\\_English.pdf](http://www.eaaci.org/attachments/304_English.pdf)). Consulted May 18, 2016.
- Beale KM. Atopic dermatitis: clinical signs and diagnosis. *Proceedings of the North American Veterinary Conference; Orlando, USA.* 2006; 20:363-4.
- Martins L, Goicoa A, López J. Allergy diagnosis – An application to dog. *Exp Path Health Sci* 2008; 2(2):51-9.
- Lloyd D. Diagnosis and management of adverse food reactions in the dog. *Proceedings of the World Congress WSAVA/ FECAVA/CSAVA.* Prague, Czech Republic, 2006; 232-5.
- Picco F, Zini E, Netti C, Naegeli C, Bigler B, Rufenacht S, Roosje P, Gutzwiller ME, Wilhelm S, Pfister J, Meng E, Favrot C. A prospective study on canine atopic dermatitis and food-induced allergic dermatitis in Switzerland. *Vet Dermatol.* 2008; 19(3):150-5.
- Mueller RS, Olivry T, Prélaud P. Critically appraised topic on adverse food reactions of companion animals (1): duration of elimination diets. *BMC Vet Res.* 2015; 11:225.
- Hensel P, Santoro D, Favrot C, Hill P, Griffin C. Canine atopic dermatitis: detailed guidelines for diagnosis and allergen identification. *BMC Vet Res.* 2015; 11:196.
- Hill P. Clinical signs and diagnosis of atopic dermatitis. *Proceedings of the European Veterinary Conference: Voorjaarsdagen.* Amsterdam, Netherlands, 2009.
- Rosje P. Canine atopic dermatitis: new concepts. *Eur J Comp Anim Pract.* 2005; 15(2):189-95.
- Kang MH, Kim HJ, Jang HJ, Park HM. Sensitization rates of causative allergens for dogs with atopic dermatitis: detection of canine allergen-specific IgE. *J Vet Sci.* 2014; 15(4):545-550.
- Lee KW, Blankenship KD, McCurry ZM. Performance characteristics of a monoclonal antibody cocktail-based ELISA for detection of allergen-specific IgE in dogs and comparison with a high affinity IgE receptor-based ELISA. *Vet Dermatol.* 2009; 20(3):157-64.
- Favrot C, Steffan J, Seewald W, Picco F. A prospective study on the clinical features of chronic canine atopic dermatitis and its diagnosis. *Vet Dermatol.* 2010; 21(1):23-31.
- Rybníček J, Lau-Gillard PJ, Harvey R, Hill PB. Further validation of a pruritus severity scale for use in dogs. *Vet Dermatol.* 2009; 20(2):115-22.
- Loewenstein C, Mueller RS. A review of allergen-specific immunotherapy in human and veterinary medicine. *Vet Dermatol.* 2009; 20(2):84-96.
- DeBoer DJ. Advances in canine atopic dermatitis. *Proceedings of the 35th World Small Animal Veterinary Congress WSAVA.* Geneva, Switzerland, 2010.

This work is funded by National Funds through FCT - Foundation For Science and Technology under the Project UID/AGR/00115/2013. In relation to this presentation, I declare that there are no conflicts of interest.