

Can birds play a role as High Nature Value indicators of *montado* system?

L. Catarino · C. Godinho · P. Pereira ·
A. Luís · J. E. Rabaça

Received: 21 April 2014 / Accepted: 29 October 2014
© Springer Science+Business Media Dordrecht 2014

Abstract *Montados* form a heterogeneous landscape of wooded matrix dominated by cork and/or holm oak with open areas characterized by fuzzy boundaries. *Montado* supports a high biological diversity associated to low intensity management and a landscape diversity provided by a continuous gradient of land cover. Among other features this permits the classification of *montados* as a High Nature Value (HNV) system. We assessed the role of birds as HNV indicators for *montado*, and tested several bird groups—farmland, edge, forest generalists and forest specialists species; and some universal

indicators such as species conservation status, Shannon's diversity index and species richness. Our study areas covered the North–South distribution of cork oak in Portugal, and we surveyed the breeding bird communities across 117 sampling sites. In addition to variables related to management and sanitary status, we considered variables that characterize the landscape heterogeneity inside the *montado*—trees and shrub density and richness of woody vegetation. Our results suggest that specific bird guilds can be used as HNV indicators of particular typologies of *montado*, and highlight the need to develop an indicator that could be transversally applied to all types of *montado*.

Electronic supplementary material The online version of this article (doi:[10.1007/s10457-014-9761-y](https://doi.org/10.1007/s10457-014-9761-y)) contains supplementary material, which is available to authorized users.

Keywords *Montados* · Birds · HNV farmland · Indicators · Landscape composition

L. Catarino · C. Godinho · P. Pereira · J. E. Rabaça
LabOr – Laboratório de Ornitologia, ICAAM,
Universidade de Évora, 7002-554 Évora, Portugal

C. Godinho (✉) · P. Pereira · J. E. Rabaça
ICAAM – Instituto de Ciências Agrárias e Ambientais
Mediterrânicas, Universidade de Évora, Núcleo da Mitra,
Ap. 94, 7002-554 Évora, Portugal
e-mail: capg@uevora.pt

A. Luís
Departamento de Biologia, Universidade de Aveiro,
Campus de Santiago, 3810-193 Aveiro, Portugal

J. E. Rabaça
Departamento de Biologia, Universidade de Évora,
7002-554 Évora, Portugal

Introduction

Portuguese *montados* and Spanish *dehesas* are silvo-pastoral systems of anthropogenic origin derived from ancient Mediterranean forests of cork oaks (*Quercus suber*) or holm oaks (*Quercus rotundifolia*). These systems can combine the use of woodland products (e.g. timber, charcoal and cork) with cereal crops and livestock grazing in the understory (Blondel and Aronson 1999). Such diversified activities allow *montados* to form a heterogeneous landscape of wooded matrix with open areas, scattered woodlands

and semi-natural patches of Mediterranean forest and scrublands, resulting in a system with high biological diversity (Rabaça 1990; Blondel and Aronson 1999; Tellería 2001; Díaz et al. 2003; Telleria et al. 2003; Harrop 2007). This mosaic of habitats has been widely recognized as an hotspot of farmland biodiversity and exponents of landscape multifunctionality (Pinto-Correia et al. 2011a).

We focused our study on cork oak *montados* due to the relevance of the area they occupy in Portugal—c. 737,000 ha (DGRF 2007)—and its influence on national economy, representing 1 % of the GNP (APCOR 2012). Currently, Portugal harbors 34 % of the world distribution of *montados*, holding half of the cork production of the world. Cork exploitation is highly sustainable, since the cork oak tree possesses the particular ability of producing a cork layer in the bark that regenerates after the debarking process. This biological feature allows the extraction of the bark from the tree in nine-year cycles without resulting in the death of the tree.

Despite its economic and environmental relevance, *montados* have been exposed to several pressures and threats, mainly intensification, overgrazing, land abandonment and the spreading of pathogenic agents (Plieninger 2007), but also extensification through lower grazing density and lower shrub control (Pinto-Correia and Almeida 2013). Therefore, management options able to conciliate the maintenance of biodiversity and the economic values of the *montado* are critical, and objective criteria must be described for the functioning of the system and for the services it provides (Andersen et al. 2003). This is particularly relevant regarding the High Nature Value (HNV) classification, which aims to integrate biodiversity and environmental concerns in the agricultural sector (Beaufoy and Cooper 2008). The HNV concept involves low intensity farming, presence of semi-natural vegetation (e.g. hedgerows, uncultivated fields/patch's, shrubs, scattered trees) and diversity of land cover, that sustains or is associated to areas with high species diversity or to the presence of priority species for conservation at European, national or regional scale (Andersen et al. 2003; Hoffmann and Greef 2003; Kleijn et al. 2009).

According to the classification proposed by the European Environment Agency, most of the silvopastoral systems in the Iberian Peninsula are considered HNV systems (Hoogeveen et al. 2004). However,

unlike forestry systems such as pine (*Pinus* spp.) plantations (Scarascia-Mugnozza et al. 2000), the diverse and most common management activities conducted in *montados* throughout the year (e.g. agriculture, pasture, grazing, livestock, game, etc.) are known to increase the structural complexity of the system, producing several types of *montado* in different natural conditions and different management contexts (Pinto-Correia 1993; Pinto-Correia and Almeida 2013). Hence, the question raised by Pinto-Correia and Almeida (2013) remains: can all *montados* typologies be classified as HNV systems?

It is recognized that *montados* exhibit the highest richness of communities of breeding birds associated with forested areas in the Iberian Peninsula (Tellería 2001). Moreover, many bird species seem to be well adapted to this system and several species even show a tolerance to debark (Godinho and Rabaça 2011; Leal et al. 2011).


When compared with other HNV systems, *montados* have a major forest component which increases its structural complexity. Since communities of breeding birds have long been recognized as good indicators of the structural complexity of forested areas (e.g. Willson 1974; Rabaça 1990; Whelan 2001, Pereira et al. 2014), we used assemblages of breeding birds to evaluate: (1) which features of *montado* contribute to higher species diversity and richness, (2) if these parameters—diversity and richness—and/or bird guilds can be used to identify *montado* areas to be classified as HNV, and (3) if a specific bird guild can better represent *montado* natural values than community parameters like species richness or diversity.

Methods

Study area

We sampled four areas covering the main distribution range of cork oak in Portugal (Fig. 1): the Site of Community Importance (SCI) of Romeu (hereafter Romeu) with several private owners (7°1'–7°6'W and 41°33'–41°28'N), Companhia das Lezírias S.A. (Lezírias) a public ownership farm (8°48'W and 38°50'N), SCI Serra de Monfurado (Monfurado) (7°40'–8°16'W and 38°27'–38°41'N) and Serra de Grândola (Grândola) (8°34'–8°38'W and 38°9'–

Fig. 1 Location of the study areas in Portugal




38°08'N) both with several private owners. These areas reflect the most common typologies of *montados*, half of the national distribution of cork oak has a tree coverage between 10 and 30 %, a quarter between 30 and 50 % and a quarter superior to 50 % (Carreiras et al. 2006). Our sampling sites represent these three categories in a similar proportion. Lezírias and Monfurado had sites belonging to all categories; sites with higher tree cover were absent and in Romeu and Grândola (Fig. 2). These areas are located in the Mediterranean part of the country (Northeast and the entire Southern half of Portugal), characterized by hot and dry summers and moderate rainy winters. Altitude ranges between 15 m (Lezírias) and 600 m (Romeu). Romeu and Monfurado showed the lowest mean annual temperature (12.3 and 12.5 °C, respectively) and the highest levels of mean annual precipitation (760 and 800 mm, respectively). Lezírias and Grândola showed a mean annual temperature of 15.7 and 15.6 °C, respectively, and lower levels of mean annual precipitation (644 and 500 mm, correspondingly). The woodland area is dominated by cork oaks, but holm oak settlements can be found in Monfurado and mixed stands with cork oak, maritime pine (*Pinus pinaster*) and stone pine (*Pinus pinea*) occur in Lezírias. Other

common land uses are: rice fields, vineyards and pine woods in Lezírias; olive groves, small orchards, dry cereal fields and fallows in Monfurado; and olive groves and vineyards in Romeu. Riparian galleries are present in all areas.

Bird census

We carried out bird censuses in Monfurado (2011) and in the other areas (2012) during the breeding seasons (between April and May). We gathered data on bird species using 10 min point counts (e.g. Bibby et al. 2000) with 100 m radius. We conducted surveys between 6:00 and 11:00 a.m., when birds are more active, and we avoided days with hard wind and rain. In each study area we surveyed 30 sites except in Romeu where we sampled 27 points. Our four areas covered a wide range of the different *montado* typologies present in Portugal (Fig. 2), and sites were randomly selected, as long as they satisfied two criteria: (1) accessibility and (2) being situated at least 500 m apart from each other to minimize the probability of double counting birds. Three experienced observers with similar skills of bird detection conducted the surveys.

Fig. 2 Distribution of sampling sites along a gradient of *montado* coverage



We excluded aerial-feeding birds (*e.g.* barn swallow *Hirundo rustica*) from data treatment, as well as species with known large home-ranges (*e.g.* carrion crow *Corvus corone* and raven *C. corax*).

We used the species richness of each of the following four ecological guilds (farmland species, edge species, forest generalists and forest specialists) defined according to their habitat preferences in the Mediterranean part of Portugal during the breeding season (Table 1). Species classification is related to the specialization degree taking into account the habitats occupied along a gradient of natural terrestrial habitats, and may not be applicable outside this geographic area or time of year. This gradient can be represented by an increased complexity of vegetation structure, from structurally simpler habitats such as grasslands to intermediate habitats as heathlands and scrublands culminating in oak forests (such as cork oak or holm oak). To this end, we used the available information regarding places and strategies used to capture food, sites used for territorial defence as well as nesting sites. Farmland species mostly occur in association with open fields or, at most, with scattered shrubs or trees; edge species are associated with transitional areas that can occur at boundaries between two or more habitats (*e.g.* forest and open areas); forest generalists occur at any woody areas, regardless of the density and height of the tree formations; forest specialists tend to occupy stable forest environments and are therefore more sensitive to human disturbance than generalist forest species. Most occur in natural or

semi-natural forests with the following features: (1) high canopy of great maturity, (2) large stratification of woody vegetation, and (3) humid microclimate.

In addition to ecological guilds we also considered Shannon's diversity index (ShInd), total species richness (SpRich) and a variable that takes into account the richness of species with conservation status (ConSt). This variable was estimated as the richness by point count of species listed in at least one of these lists: the Red Book of Vertebrates of Portugal (Cabral et al. 2005), Species of European Conservation Concern (BirdLife International 2004) and IUCN Red List for Birds (BirdLife International 2014).

HNV features and explanatory variables

Due to their nature, *montado* landscapes are distinguished by fuzzy boundaries with overlapping land cover classes (diverse combinations of forest cover, grass and shrubs) (Pinto-Correia and Vos 2004), they are often characterized by gradual changes in shrub and tree densities, resulting from the combination of different levels of land use and of variable, extensive land use practices. The fuzziness of the boundaries is inherent to the land use system and should be accepted as such. Small differences in terms of tree density and shrub cover reflect important differences in abiotic factors (Joffre et al. 1999), types of the management in the past and present (Pinto-Correia 1993) and levels of biodiversity (Ojeda et al. 1995).

Table 1 Ecological guilds based on species habitat preferences

Bird guild	Definition criteria	Species
Forest specialists	Species that occur only in certain forest types	<i>Dendrocopos minor</i> , <i>Troglodytes troglodytes</i> , <i>Erithacus rubecula</i> , <i>Phoenicurus phoenicurus</i> , <i>Sylvia atricapilla</i> , <i>Phylloscopus bonelli</i> , <i>Phylloscopus ibericus</i> , <i>Aegithalos caudatus</i> , <i>Sitta europaea</i> , <i>Coccothraustes coccothraustes</i>
Forest generalists	Forest species that occur in any woody areas, regardless of the density and height of the plant formations, as well as bioclimatic conditions of the site	<i>Columba palumbus</i> , <i>Cuculus canorus</i> , <i>Dendrocopos major</i> , <i>Luscinia megarhynchos</i> , <i>Turdus merula</i> , <i>Sylvia melanocephala</i> , <i>Parus cristatus</i> , <i>Parus caeruleus</i> , <i>Parus major</i> , <i>Certhia brachydactyla</i> , <i>Garrulus glandarius</i> , <i>Oriolus oriolus</i> , <i>Fringilla coelebs</i> , <i>Serinus serinus</i> , <i>Carduelis chloris</i>
Edge species	Species that occur at the boundary between two or more habitats	<i>Streptopelia turtur</i> , <i>Clamator glandarius</i> , <i>Jynx torquilla</i> , <i>Lullula arborea</i> , <i>Hippolais polyglotta</i> , <i>Sylvia undata</i> , <i>Sylvia cantillans</i> , <i>Lanius meridionalis</i> , <i>Lanius senator</i> , <i>Passer montanus</i> , <i>Petronia petronia</i> , <i>Carduelis carduelis</i> , <i>Carduelis cannabina</i> , <i>Emberiza cirrus</i> , <i>Emberiza cia</i>
Farmland species	Farmland species that tolerate low tree densities	<i>Alectoris rufa</i> , <i>Coturnix coturnix</i> , <i>Upupa epops</i> , <i>Galerida theklae</i> , <i>Oenanthe hispanica</i> , <i>Saxicola torquata</i> , <i>Cisticola juncidis</i> , <i>Sturnus unicolor</i> , <i>Passer domesticus</i> , <i>Emberiza calandra</i>

We recorded environmental variables for each sampling site that, according to our rationale, could represent the heterogeneity within the *montado*: (1) tree density; (2) shrub density; (3) woody richness; (4) type of edge and; (5) distance to edge (Table 2). We also recorded geographical variables (location and altitude), management variables (trunk diameter at breast height, debark), and variables associated with the sanitary status of the settlements (occurrence of pests outbreaks and fungal presence) (Table 2).

Immediately after conducting the bird censuses, we evaluated the vegetation features within a 100 m radius around the centre of the point count.

At each point of the study areas, the sanitary status of the oak woodlands was evaluated through visual assessment of five oaks separated 10 m from each other, avoiding trees with adjoined crowns. We detected the damage made by buprestids *Coroebus florentinus* and *Coroebus undatus* through the presence of typical dead branches on outer-canopy and the presence of feeding galleries of larvae on the cork layer, respectively. These measures were used as individual variables, but also incorporated together with insect trunk holes of bark beetle (Platypodidae and Scolytidae), cerambycids (Cerambycidae) and signs of fungal disease *Biscogniauxia mediterranea* to create a sanitary index (*San*). We created the sanitary

index for *montado* according to the level of harm done by insect pests on the sampling sites. For each insect pest and fungus we attributed a coefficient of impact according to the intensity of their damage on the tree: 1—high aesthetic impact with low economic or ecological relevance (*C. florentinus*), 3—high economic relevance due to loss of cork value (*C. undatus*), 5—associated with tree decline or death (bark beetles, cerambycids and *B. mediterranea*).

We created the sanitary index through the sum of the proportion of each pest or disease, which was multiplied by the respective impact coefficient.

$$San = 1 \left(\frac{OA_{Cflo}}{5 \text{ oaks}} \right) + 3 \left(\frac{OA_{Cund}}{5 \text{ oaks}} \right) + 5 \left(\frac{OA_{Barkb}}{5 \text{ oaks}} \right) + 5 \left(\frac{OA_{Bmed}}{5 \text{ oaks}} \right)$$

OA is the oaks affected by each pest or disease.

Data analysis

Prior to data analysis and in order to avoid multicollinearity among variables, we performed data reduction procedures. According to Tabachnick and Fidell (2001), we assessed all pairwise correlations through Spearman correlation coefficients and retained only one in each pair of highly correlated variables

Table 2 Environmental variables recorded at each one of the sampling sites

Environmental variables	Code
Geographical variables	
Geographical location (coordinates)	Point
Identification of the four areas surveyed	Study area
Altitude (m)	Alt
Vegetation features	
Shrubs density	Shrubs
Trees density	Trees
Woody richness including tree and shrub species (categorical; 1-one or two woody species; 2-three or more woody species)	Woody
Management practices	
Trunk diameter at breast height (cm)	DBH
Year of the last cork removal	Debark
Type of edge (categorical; 1-open area; 2-shrubs and vineyards; 3-eucalyptus, pine plantations, orchards, olive groves)	Edge
Distance to the edge (m)	EdgeD
Sanitary status	
Presence/absence of <i>Biscogniauxia mediterranea</i>	Bmed
Presence/absence of <i>Coroebus florentinus</i> damage	Cflo
Presence/absence of <i>Coroebus undatus</i> damage	Cund
Presence/absence of bark beetles (Platypodidae and Scolytidae) and cerambycids (Cerambycidae) damage	BarkB

($r > |0.7|$) for further analyses. Only a pair of variables showed strong collinearity *Altitude* and *Study Area* ($r = 0.934$). *Altitude* was removed due to the auto-correlation of this variable between sampling sites belonging to the same study area.

We used one-way ANOVA (Zar 1999) to determine if there were differences between areas in terms of ecological guilds and environmental variables.

We modelled the effects of environmental variables in function of groups of species through linear mixed-effects models (Pinheiro and Bates 2000). We treated *Study Area* as random effect and all other explanatory variables as fixed effects. To deal with model selection uncertainty we analysed our data based on the information theoretic approach (ITA) (Burnham and Anderson 2002). In the analyses of species groups (bird richness of every guild), Shannon's diversity index and total species richness, we considered as fixed effects the following six variables which are

representative of the heterogeneity inside the *montado*: *shrubs density*; *trees density*; *woody richness*; *type of edge*; *distance to edge*; and *sanitary index*.

We generated all possible models combining from none to seven explanatory variables. We used this option, classified as data dredging (Burnham and Anderson 2002), because all explanatory variables considered could potentially influence the response variables. Using all possible combinations we guaranteed that the explanatory variables were included in the model-averaging procedure in identical manner. In accordance with Burnham and Anderson (2002), we fitted the models one by one and ordered them by their values of AICc (second-order Akaike's information criterion). We used AICc as a measure of information loss for each candidate model, with the best fitting model having the lowest AICc and the highest Akaike weight (w_i), which measures the posterior probability of a given model being true, given the data and the set of competing candidate models (Burnham and Anderson 2002). Additionally, we also calculated the number of parameters (degrees of freedom), log-likelihood value and AICc difference ($\Delta AICc$), and the model-averaged coefficients of all explanatory variables for each model (Burnham and Anderson 2002; Lukacs et al. 2010; Symonds and Moussalli 2011). Finally, we estimated the relative importance of each explanatory variable, by adding the Akaike weights of all models in which the variable appeared (Burnham and Anderson 2002). The relative importance of the variables that appear in all top models tends towards 1. In variables that only appear in less likely models, their relative importance tends towards 0. We then ranked the explanatory variables according to their relative importance, and the direction and magnitude of the effect of each variable was based on the model-averaged coefficients (Burnham and Anderson 2002).

We carried out the statistical analysis using SPSS 21 for Windows (IBM Corp. 2012) and software R v. 3.0.2 (R Core Team 2013), with package MuMIn (Barton 2014) and nlme (Pinheiro et al. 2014).

Results

General results

From 74 species recorded in surveys, we used 50 in the analyses and hereafter we will refer only to those.

Almost 40 % of these species show some level of threat, according to the Species of European Conservation Concern (SPEC) (BirdLife International 2004) or have conservation status by the Portuguese Red Book of Vertebrates (Cabral et al. 2005) or the IUCN Red List (BirdLife International 2014) (Appendix I). We detected 17 of the 50 species in all areas. On the other hand, we detected five bird species exclusive from Romeu, three exclusive from Monfurado and Lezírias and one exclusive from Grândola (Appendix I). Average species richness and standard deviation, per point count, was 6.8 ± 1.3 in Grândola, 7.3 ± 1.2 in Romeu, 8.5 ± 2.1 in Lezírias and 11.2 ± 2.4 in Monfurado.

Environmental variables

In order to evaluate how our environmental variables range along the four areas we run one-way ANOVAs for each variable, with a Bonferroni correction post hoc test (Table 3). The average age of cork removal (Debark) and the distance to the edge (EdgeD) were similar between all areas. On the contrary, *Altitude* was significantly different among all areas.

Although shrub coverage was higher in Grândola, being almost the double than in Lezírias and Monfurado, tree coverage and woody diversity were the lowest observed. Monfurado was the area with higher tree coverage and significantly different from all the others. Diversity of woody vegetation was higher in Lezírias and Monfurado and statistically different from Grândola and Romeu. The oldest settlements were observed in Romeu and Lezírias (based on trunk diameter at breast high—DBH), being statistically different from Monfurado.

We observed a similar pattern for both *Coroebus* species, with lower values of affectation in Romeu and higher values in Grândola and Monfurado. In this sense, these areas were significantly different for both pests. Romeu emerges as the area with less evidence of affectation in regard to the sanitary status index.

Bird guilds

In order to evaluate how our species ranged along the four areas we run a one-way ANOVA among areas for each species guild, with a Bonferroni correction post hoc test. Significant differences were observed between areas for all guilds analyzed, and also for

total bird richness, species diversity and conservation status (Table 4). Monfurado was the area with highest species richness for all bird guilds, with the exception of farmland species. Lezírias had the highest values for this guild whereas it appears to be residual in Romeu. Romeu had the lowest value for species with conservation status, and Monfurado presented the highest species richness of this guild. As expected, based on guilds results, Monfurado stood out from the other areas regarding species diversity and total species richness. The other areas did not present significant statistical differences, with the exception of higher species richness in Lezírias than in Grândola.

Modelling bird guilds

We ranked the candidate models for each one of the response variables based on the AICc ($\Delta AICc < 2.00$) and also estimated the relative importance of each variable (Table 5) (Burnham and Anderson 2002).

Farmland species: three variables—tree density, edge type, and woody species—were included in the set of best models (Table VI—supplementary material). Tree density assumed the greatest importance (0.92), being negatively associated with the richness of farmland species. Edge type also had a high relative importance (0.69), with these species being negatively influenced by the presence of non-agricultural edges. Woody species still had some relative importance. Patches with more woody species showed a greater richness of farmland birds.

Edge species: none of the variables showed an association with this group of species, the top ranked model being the null model (Table VI).

Forest generalists: although the null model was ranked in the best set of models (in fourth position—Table VI), we found it useful to discuss the variables that enter in the other models. These included three variables—sanitary index, woody species and tree density (Table 5); all positively associated with this guild.

Forest specialists: the most relevant variables in the candidate models were tree density, edge type, shrubs density and woody species. Sites with higher tree cover were most important to this species (1.00) being present in all models.

Conservation status: two variables—edge type and tree density—were included in the models, both influencing negatively the richness of species with

Table 3 One-way ANOVA for environmental variables (Shrubs—density of shrubs, Trees—density of trees, Woody—woody richness including trees and shrubs, DebarK—year of the last cork removal, DBH—trunk diameter at breast high, Cund—trunk diameter at breast high, Cflo—damage by *Coroebus indatus*, Cflo—damage by *Coroebus florentinus*, San—sanitary index, Alt—altitude and EdgeD—distance to edge) according to each area (G—Serra de Grândola, L—Companhia das Lezírias, M—Serra de Monfurado and R—Romeu)

Areas	Shrubs	Trees	Woody	DebarK	DBH	Cund	Cflo	San	Alt	EdgeD
G (n = 30)	60 ± 30.73	28 ± 10.64	1.13 ± 0.35	5.13 ± 3.27	35.28 ± 5.76	2.13 ± 0.73	1.70 ± 0.88	7.63 ± 2.35	205.83 ± 43.56	115.27 ± 85.03
L (n = 30)	32.90 ± 27.96	40 ± 15.97	1.87 ± 0.35	6.13 ± 2.54	39.51 ± 8.22	1.73 ± 0.74	1.73 ± 0.91	6.95 ± 1.08	32.27 ± 8.24	86.53 ± 53.26
M (n = 30)	38.50 ± 32.56	55 ± 18.75	1.87 ± 0.35	5.23 ± 1.94	33.16 ± 5.74	2.23 ± 0.86	1.80 ± 0.55	7.31 ± 1.11	294.33 ± 52.89	77.30 ± 76.83
R (n = 27)	52.96 ± 26.86	30.19 ± 11.31	1.15 ± 0.36	6.89 ± 2.59	39.04 ± 7.37	1.48 ± 0.51	1.15 ± 0.36	5.15 ± 1.65	454 ± 91.08	106.41 ± 101.06
F	5.53	20.87	42.08	2.81	5.84	6.67	4.87	12.86	282.58	1.41
P	<0.001	<0.001	<0.001	<0.05	<0.001	<0.001	<0.01	<0.001	<0.001	0.244
Bonferroni correction										
GxR	-	-	-	-	-	<0.01	<0.05	<0.001	<0.001	-
GxL	<0.01	<0.05	<0.001	-	-	-	-	-	<0.001	-
GxM	<0.05	<0.001	<0.001	-	-	-	-	-	<0.001	-
RxL	-	-	<0.001	-	-	-	<0.05	<0.001	<0.001	-
RxM	-	<0.001	<0.001	-	<0.01	<0.01	<0.01	<0.001	<0.001	-
LxM	-	<0.05	-	-	<0.01	-	-	-	<0.001	-

Values represent average richness and respectively standard deviation, and results with significant differences after applying the Bonferroni correction

conservation status. The types of edge that negatively influenced species with conservation status were forest edges (e.g. eucalyptus, pine, orchards) and shrub edges.

Shannon’s diversity index: two variables assumed higher importance, and positively influence species diversity: sanitary index (0.95) and woody species (0.88).

Species richness: as in the case of the Shannon’s diversity index, the variables included in the models were woody species and sanitary index.

Discussion

In a first approach we characterized areas and associated bird guilds in order to have a better perception of their similarities and differences, allowing us to distinguish eventual patterns arising from regional influence. The analysis showed the amplitude of *montados* evaluated throughout the country (Table 3), and even inside of each area, this is particularly relevant when we try to assess associations between bird guilds and HNV features.

The highest richness for most guilds (except for farmland species) was associated with Monfurado (Table 4), this area revealing high amplitude in variables associated with landscape heterogeneity (trees, woody and shrubs; Table 3). These results are in line with what has been documented by other authors (e.g. Blondel and Aronson 1999; Díaz et al. 2003; Harrop 2007). Actually, the heterogeneous pattern of wooded matrix with open areas, scattered woodlands and undisturbed patches of Mediterranean forest and scrublands, creates a patchwork of habitats which is a trait of *montado* landscape, and induces the highest richness in breeding bird communities in the Iberian Peninsula (e.g. Tellería 2001). So, areas that encompass higher densities of shrubs and trees and that are more diverse in woody species have greater potential of being considered good habitats for a large number of species, including species of conservation concern (e.g. redstart *Phoenicurus phoenicurus* and crested tit *Parus cristatus*).

Species belonging to the farmland guild, including thekla lark (*Galerida theklae*), stonechat (*Saxicola torquata*) and corn bunting (*Emberiza calandra*), are tolerant to the presence of trees in their breeding sites. Therefore, they can occur in *montados*, contrarily to

Table 4 One-way ANOVA for species richness by ecological guild (Fa—farmland species, ES—edge species, FoG—forest generalists, FoS—forest specialists, ConSt—conservation status), Shannon's diversity index (ShInd) and total species

richness (SpRich) according to each area (G—Serra de Grândola, L—Companhia das Lezírias, M—Serra de Monfurado and R—Romeu)

Areas	Fa	ES	FoG	FoS	ConSt	ShInd	SpRich
G (n = 30)	0.97 ± 1.10	0.50 ± 0.63	4.03 ± 1.16	1.30 ± 0.70	1.20 ± 0.85	0.79 ± 0.10	6.80 ± 1.32
L (n = 30)	1.90 ± 1.21	1.20 ± 0.93	4.37 ± 1.38	1.07 ± 0.94	1.57 ± 0.90	0.86 ± 0.12	8.53 ± 2.10
M (n = 30)	1.77 ± 1.43	1.80 ± 1.00	5.77 ± 2.37	1.90 ± 1.37	2.23 ± 1.41	0.99 ± 0.15	11.23 ± 2.43
R (n = 27)	0.15 ± 0.36	0.85 ± 0.95	5.19 ± 1.39	1.15 ± 0.86	0.63 ± 0.74	0.83 ± 0.08	7.33 ± 1.24
<i>F</i>	14.96	11.69	6.77	4.17	12.62	16.56	33.61
<i>P</i>	<0.001	<0.001	<0.001	<0.01	<0.001	<0.001	<0.001
Bonferroni correction							
GxR	<0.05	–	–	–	–	–	–
GxL	<0.01	<0.05	–	–	–	–	<0.01
GxM	<0.05	<0.001	<0.01	–	<0.01	<0.001	<0.001
RxL	<0.001	–	–	–	<0.01	–	–
RxM	<0.001	<0.01	–	<0.05	<0.001	<0.001	<0.001
LxM	–	–	<0.01	<0.05	–	<0.001	<0.001

Values represent average richness and respectively standard deviation, and results with significant differences after applying the Bonferroni correction

what happens to strict farmland species (Moreira et al. 2005). This guild is associated with scattered *montados* and to agricultural edges, decreasing its densities with the proximity of forested edges, like eucalyptus or pine plantations. Such tendency was also recorded by Reino et al. (2009) with the same group of species but in a farmland context. According to such relations between birds and habitat, the farmland species occurring in *montados* can be defined as generalist farmland species. Therefore, the diversity of woody vegetation may provide them with more ecological niches; however, shrub coverage must be low since most of these species nests in open ground.

The fuzziness of the *montado* may be the reason for the absence of strong relations between the variables considered and edge species. In *montado* these species are associated with the decreasing density of vegetation (e.g. trees and shrubs) more than with the abrupt transition between farmland and forestry patches or early-successional habitats as in the case of Central and Northern Europe (e.g. Imbeau et al. 2003; Storch et al. 2005). Additionally, inside the guild there are species associated with different kinds of forest interfaces, for example: turtle dove (*Streptopelia turtur*), melodious warbler (*Hippolais polyglotta*) and cirl bunting (*Emberiza cirlus*) are mainly associated with forest—farmland edges; wood lark (*Lullula*

arborea), Iberian grey shrike (*Lanius meridionalis*) and rock sparrow (*Petronia petronia*) are mainly associated with scattered arboreal vegetation such as sparse *montados*, which can be considered transitional habitats between forested and open areas. The diversity of species-habitat associations in this guild is reflected in the uncertainty of the variables considered in the models (Table VII; supplementary material).

Into a more comprehensive view, the forest is the most important component inside the cork oak *montados*, and the occurrence of generalist forest species along a wide range of *montado* typologies is to be expected. Our data reveals a preference for diversity of woody vegetation and tree density, although the association with sites with higher pest affectation (reflecting more degraded areas) may indicate the avoidance of sites with higher tree coverage.

The occurrence of forest specialist species was associated with higher densities of trees and shrubs, and with forested edges. These features characterize old and complex *montados*. In a Mediterranean context, species like wren (*Troglodytes troglodytes*), blackcap (*Sylvia atricapilla*) and European robin (*Erithacus rubecula*) use the remaining patches of ancient oak forest with several vegetation strata (e.g. Pérez-Tris et al. 2004) which is a rare feature across

Table 5 Results from the multi-model inference procedure for the parameters describing species guilds associations to environmental variables: relative variable importance (RVI) and trend of the relationship

	Farmland species		Edge species		Forest generalists		Forest specialists		Conservation status		Shannon's diversity index		Species richness	
	RVI	Trend	RVI	Trend	RVI	Trend	RVI	Trend	RVI	Trend	RVI	Trend	RVI	Trend
Edge	0.69*	-	0.23	-	0.10	+	0.37*	+	0.89*	-	<0.01	+	0.23	-
Distance to edge	<0.01	+	<0.01	-	<0.01	+	<0.01	+	<0.01	+	<0.01	+	<0.01	-
Sanitary index	0.04	+	0.05	+	0.48*	+	0.05	+	0.06	+	0.99*	+	0.60*	+
Shrubs density	0.11	-	0.01	-	0.01	+	0.33*	+	0.01	-	<0.01	+	0.01	+
Trees density	0.92*	-	0.08	-	0.27*	+	1*	+	0.86*	-	<0.01	+	0.02	+
Woody	0.29*	+	0.17	+	0.42*	+	0.30*	+	0.21	+	0.94*	+	0.96*	+

The asterisk indicates the variables included in the best models ($\Delta AICc < 2.00$)

montado. Besides, these patches allow an increase of niche availability for species and can be used as an important evaluation element of HNV in Europe (Morelli 2013).

The occurrence of priority species for conservation is one of the key points for a site to be recognized as HNV. In our study these species are associated with more sparse *montados* and with farmland edge, in other words they tend to avoid more mature and complex sites. This trend is directly influenced by the existing imbalance between the species of this group: 16 out of 19 species (84 %) with some criteria of conservation are farmland or edge species associated with the interface forest-farmland. We suggest that this result should be considered with some caution because most of the threatened forest species that nesting in *montado* were not targeted by our study since they have large territories (e.g. raptors, black stork) and this could be another factor of bias. The inclusion of other conservation criteria must be considered, such as those proposed by Tavares (2009), based on species of which Portugal hosts significant populations at European level, thus having a responsibility for their conservation (e.g. Iberian chiffchaff *Phylloscopus ibericus* or serin *Serinus serinus*).

Species richness and species diversity showed the same trend, being positively associated with areas that have higher diversity of woody vegetation and higher values of sanitary index. The association between these two indices and the sanitary condition may reflect (1) the current state of conservation of *montados* and (2) the influence of the forest generalist species in the species taken as a whole. Most of these species were recorded along several sampling sites (e.g. blue tit *Parus caeruleus*, short-toed treecreeper *Certhia brachyactyla*, blackbird *Turdus merula*) which suggests that the global indexes were primarily influenced by the presence of forest generalist species. Cork oak decline has been reported in Southwestern Portugal since the 1890s (Cabral et al. 1992) and therefore, in the light of our results, it seems plausible to state that most of the surveyed areas are under some kind of threat.

Conclusion

The continuous gradient of land cover and fuzzy boundaries characteristic of *montado* (Pinto-Correia

et al. 2011) are well expressed in the variables of density and diversity of vegetation and their association with the bird guilds evaluated. With the exception of the edge species, these variables were important to all the other bird groups (Table 5). The guilds under consideration mainly characterized a gradient of forest complexity, from the farmland species to the forest specialists, and perhaps they can be individually used as HNV indicators of a particular typology of *montado*, based on tree coverage: farmland species for scattered areas; and, on the opposite side of the range, forest specialists for more mature settlements or small well-preserved forest patches. We cannot say if all *montado* typologies can be classified as HNV, but we can define characteristic bird guilds for several typologies, and through the ratio between the species observed and the expected pool of species it should be possible to evaluate if a site may be classified as HNV. At a broader scale the universal measures of species diversity and species richness could also be used as HNV indicators.

The next steps on research should be focused on: (1) the creation and testing of a compound index with farmland, forest generalists and forest specialists species in order to create a reliable indicator of HNV for *montados* applicable to several scales, (2) the assessment of other HNV parameters for *montado* such as stone piles, ponds or fences, and (3) the integration of other species conservation criteria in addition to the traditional red lists.

Acknowledgments We would like to thank two anonymous reviewers whose comments helped to improve the manuscript. We also thank to Rui Lourenço, Pedro Salgueiro and Luís Gomes for invaluable comments and suggestions. This study was partially supported by Companhia das Lezírias S.A. and by the Portuguese Science Foundation through the Doctoral grants SFRH/BD/81602/2011 (Carlos Godinho) and SFRH/BD/87340/2012 (Pedro Pereira).

References

- Andersen E, Baldock D, Bennett H, Beaufoy G, Bignal E, Brouwer F, Elbersen B, Eiden G, Godeschalk F, Jones G, McCracken D, Nieuwenhuizen W, van Eupen M, Henneskens S, Zervas G (2003) Developing a High Nature Value Farming area indicator. Internal report for the European Environment Agency.
- APCOR (2012) Anuário da APCOR. Plenimagem
- Barton K (2014) Multi-model inference. Version 1.10.0. <http://cran.r-project.org/web/packages/MuMIn/index.html>. Accessed 24 Sept 2014
- Beaufoy G, Cooper T (2008) Guidance document to the member states on the application of the High Nature Value impact indicator. Institute for European Environmental Policy, Brussels
- Bibby CJ, Burgess ND, Hill DA, Mustoe S (2000) Bird census techniques, 2nd edn. Academic Press, London
- BirdLife International (2004) Birds in the European Union: a status assessment. BirdLife International, Wageningen
- BirdLife International (2014) IUCN Red List for birds. <http://www.birdlife.org>. Accessed 4 Aug 2014
- Blondel J, Aronson J (1999) Biology and wildlife of the Mediterranean region. Oxford University Press, New York
- Burnham KP, Anderson DR (2002) Model selection and multimodel inference: a practical information-theoretic approach. Springer, New York
- Cabral MT, Ferreira MC, Moreira T, Carvalho EC, Diniz AC (1992) Diagnóstico das causas da anormal mortalidade dos sobreiros a Sul do Tejo. *Scientia Gerundensis* 18:205–214
- Cabral M, Almeida J, Almeida P, Dellinger T, Almeida N, Oliveira M, Palmeirim J, Queiroz A, Rogado L, Reis M (2005) Livro Vermelho dos Vertebrados de Portugal. Instituto da Conservação da Natureza, Lisboa
- Carreiras JMB, Pereira JMC, Pereira JS (2006) Estimation of tree canopy cover in evergreen oak woodlands using remote sensing. *For Ecol Manag* 223:45–53
- DGRF (2007) Inventário Florestal Nacional 2005/06. Direcção Geral dos Recursos Florestais, Lisboa
- Díaz M, Pulido FJ, Marañón T (2003) Diversidad biológica y sostenibilidad ecológica y económica de los sistemas adehesados. *Ecosistemas* XII (3)
- Godinho C, Rabaça JE (2011) Birds like it Corky: the influence of habitat features and management of ‘montados’ in breeding bird communities. *Agrofor Syst* 82(2):138–195
- Harrop SR (2007) Traditional agricultural landscapes as protected areas in international law and policy. *Agric Ecosyst Environ* 121(3):296–307
- Hoffmann J, Greef JM (2003) Mosaic indicators—theoretical approach for the development of indicators for species diversity in agricultural landscapes. *Agric Ecosyst Environ* 98(1–3):387–394
- Hoogeveen Y, Petersen J-E, Balazs K, Higuero I (2004) High Nature Value farmland: characteristics, trends and policy challenges. European Environment Agency, Luxembourg
- IBM Corp. (2012) IBM SPSS Statistics for Windows, Version 21.0. IBM Corp., Armonk, NY
- Imbeau L, Drapeau P, Mönkkönen M (2003) Are forest birds categorised as “edge species” strictly associated with edges? *Ecography* 26(4):514–520
- Joffre R, Rambal S, Ratte JP (1999) The dehesa system of southern Spain and Portugal as a natural ecosystem mimic. *Agrofor Syst* 45(1–3):57–79
- Kleijn D, Kohler F, Báldi A, Batáry P, Concepción ED, Clough Y, Díaz M, Gabriel D, Holzschuh A, Knop E, Kovács A, Marshall EJP, Tschamtké T, Verhulst J (2009) On the relationship between farmland biodiversity and land-use intensity in Europe. *Proc R Soc B* 276(1658):903–909
- Leal AI, Correia RA, Granadeiro JP, Palmeirim JM (2011) Impact of cork extraction on birds: relevance for

- conservation of Mediterranean biodiversity. *Biol Conserv* 144(5):1655–1662
- Lukacs PM, Burnham KP, Anderson DR (2010) Model selection bias and Freedman's paradox. *Ann Inst Stat Math* 62: 117–125
- Moreira F, Beja P, Morgado R, Reino L, Gordinho L, Delgado A, Borralho R (2005) Effects of field management and landscape context on grassland wintering birds in Southern Portugal. *Agric Ecosyst Environ* 109(1–2):59–74
- Morelli F (2013) Relative importance of marginal vegetation (shrubs, hedgerows, isolated trees) surrogate of HNV farmland for bird species distribution in Central Italy. *Ecol Eng* 57:261–266
- Ojeda F, Arroyo J, Marañón T (1995) Biodiversity components and conservation of mediterranean healthlands in Southern Spain. *Biol Conserv* 72(1):61–72
- Pereira P, Godinho C, Roque I, Marques A, Branco M, Rabaça JE (2014) Time to rethink the management intensity in a Mediterranean oak woodland: the response of insectivorous birds and leaf-chewing defoliators as key groups in the forest ecosystem. *Ann For Sci* 71(1):25–32
- Pérez-Tris J, Bensch S, Carbonell R, Helbig A, Tellería J (2004) Historical diversification of migration patterns in a passerine bird. *Evolution* 58(8):1819–1832
- Pinheiro JC, Bates DM (2000) Mixed-effects models in S and S-PLUS. Springer, New York
- Pinheiro J, Bates D, DebRoy S, Sarkar D and R Core Team (2014) nlme: Linear and nonlinear mixed effects models. R package version 3.1-117, <http://CRAN.R-project.org/package=nlme>. Accessed 24 Sept 2014
- Pinto-Correia T (1993) Threatened landscape in Alentejo, Portugal: the 'montado' and other 'agro-silvo-pastoral' systems. *Landsc Urban Plan* 24(1–4):43–48
- Pinto-Correia T, Almeida M (2013) Tentative identification procedure for HNV Montados. In: ICAAM International Conference 2013: Acknowledging the MONTADOS and DEHESAS as High Nature Value Farming Systems, Évora, Portugal, 6–8 February.
- Pinto-Correia T, Vos W (2004) Multifunctionality in Mediterranean landscapes—past and future. *New Dimens Eur Lands* 4:135–164
- Pinto-Correia T, Ribeiro N, Sá-Sousa P (2011) Introducing the *montado*, the cork and holm oak agroforestry system of Southern Portugal. *Agrofor Syst* 82(2):99–104
- Plieninger T (2007) Compatibility of livestock grazing with stand regeneration in Mediterranean holm oak parklands. *J Nat Conserv* 15(1):1–9
- Rabaça JE (1990) The influence of shrubby understory in breeding bird communities of cork oak (*Quercus suber*) woodlands in Portugal. *Portugaliae Zoologica* 1(1):6
- RCoreTeam (2013) A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria
- Reino L, Beja P, Osborne PE, Morgado R, Fabião A, Rotenberry JT (2009) Distance to edges, edge contrast and landscape fragmentation: interactions affecting farmland birds around forest plantations. *Biol Conserv* 142(4):824–838
- Scarascia-Mugnozza G, Oswald H, Piusi P, Radoglou K (2000) Forests of the Mediterranean region: gaps in knowledge and research needs. *For Ecol Manag* 132(1):97–109
- Storch I, Woitke E, Krieger S (2005) Landscape-scale edge effect in predation risk in forest-farmland mosaics of Central Europe. *Landsch Ecol* 20(8):927–940. doi:10.1007/s10980-005-7005-2
- Symonds MR, Moussalli A (2011) A brief guide to model selection, multimodel inference and model averaging in behavioural ecology using Akaike's information criterion. *Behav Ecol Sociobiol* 65(1):13–21
- Tabachnick BG, Fidell LS (2001) Using multivariate statistics. Allyn and Bacon, Boston
- Tavares JP (2009) Uma perspectiva internacional sobre as prioridades de conservação da avifauna Portuguesa. Actas do VI Congresso de Ornitologia da Sociedade Portuguesa para o Estudo das Aves. Elvas
- Tellería JL (2001) Passerine bird communities of Iberian dehesas: a review. *Anim Biodivers Conserv* 24(2):67
- Tellería JL, Baquero R, Santos T (2003) Effects of forest fragmentation on European birds: implications of regional differences in species richness. *J Biogeogr* 30:621–628
- Whelan CJ (2001) Foliage structure influences foraging of insectivorous forest birds: an experimental study. *Ecology* 82:219–231
- Willson MF (1974) Avian community organization and habitat structure. *Ecology* 55:1017–1029
- Zar JH (1999) Biostatistical analysis, 4th edn. Prentice Hall, New Jersey