

Plant/Leaf traits and adaptive strategies of *Cistus* species to Mediterranean drought and insolation in southern Portugal

FONSECA, Elsa; SIMÕES, Maria Paula¹; GAZARINI, Luiz¹; COCHARD, Hervé²; VAZ, Margarida^{1,3}

¹Instituto de Ciências Agrárias e Ambientais Mediterrânicas (ICAAM)/Dep. Biologia, Universidade de Évora, Apartado 94, 7002-554 Évora, Portugal

²INRA, UMR 547 PIAF, F-63100 Clermont-Ferrand, France

³ Corresponding author (mvaz@uevora.pt)



INTRODUCTION

The effects of climate change can result in dramatic consequences in specific ecosystems such as *montados* that are seriously threatened by the absence of cork and holm oak (*Quercus suber* and *Q. rotundifolia*) natural regeneration. Shrubs of the genus *Cistus*, which are among the most important elements of encroached *montados*, seem to promote soil rehabilitation and enhance oak regeneration (Simões et al. 2009). In this context, we compared the life strategies and evaluated the potential ability of *Cistus* species to adapt to the increasing drought expected for the Mediterranean region, and thus their role on the sustainability of cork oak *montados*.



Fig. 1 – Location of the study area.

STUDY AREA

The study was carried out in a *montado* located in southern Portugal, close to Évora (Fig. 1). The area has the typical winter-wet, summer dry pattern of the Mediterranean-type climate. Mean annual rainfall is 609.4 mm and mean annual temperature is 15.9°C. The vegetation of the study site is an open cork oak woodland with an understorey of shrubs, dominated by *Cistus ladanifer* (hereafter CLAD), *C. monspeliensis* (hereafter CMON), *C. populifolius* (hereafter CPOP), and *C. psilosepalus* (hereafter CPSI) (Fig. 2), which accounted for >70% of the community cover.

METHODS

- Stomatal conductance (g_s) was measured with a portable steady-state photosynthetic system (Li-6400; Li-Cor, Lincoln, NE) in spring, late-summer and autumn 2012.
- Xylem vulnerability curves were obtained by the 'Cavitron' technique (Cochard 2002; Cochard et al. 2005), with a Sorvall RC5 high speed centrifuge and the Cavisoft software, by measuring the percentage of loss of conductance (PLC) after air infiltration, in mature branches of the year (2012).
- Predawn and mid-day leaf water potential (Ψ_{IPD} , Ψ_{IMD}) was measured on a monthly basis, from April to November 2012, with a Scholander pressure chamber (PMS 1000, PMS Instruments, Corvallis, USA).

RESULTS

- CPSI and CLAD had the highest stomatal conductance (g_s) in both spring and autumn, while CMON with the lowest spring value showed the greatest recovery after the autumn rains (Figs. 3 and 4). CPOP presented the smallest g_s seasonal variation.




Fig. 3 – Mean air temperature (T) and rainfall (R) during the study period, by a meteorological station located in the study area.




Fig. 4 - Stomatal conductance (g_s) in the study *Cistus* species. Values are mean \pm SE (n=24). Significant differences (P<0.05) between species are indicated by different letters.



Fig. 2 – *C. ladanifer* - CLAD (A), *C. monspeliensis* - CMON (B), *C. populifolius* - CPOP (C), and *C. psilosepalus* - CPSI (D).

- CMON was the most resistant to increasing xylem pressure (P50 was -10.2 ± 0.16 MPa), followed by CLAD (-8.95 ± 0.03 MPa) and CPOP (-8.14 ± 0.05 MPa). CPSI (-6.51 ± 0.09 MPa) was the most sensitive species (Fig. 5).




Fig. 5 – Xylem vulnerability curves of the study *Cistus* species. PLC - percentage of loss of conductance.

- Seasonal variation of predawn and mid-day leaf water potential (Ψ_{IPD} , Ψ_{IMD}) followed the same pattern (Fig. 6). However, the significant (P<0.05) decline observed in summer with rain scarcity was less marked for CPSI than for the other species. After autumn rains, Ψ recovered to spring values in all species.




Fig. 6 - (A) Predawn leaf water potential (Ψ_{IPD}) and (B) mid-day leaf water potential (Ψ_{IMD}) of the study *Cistus* species. Values are mean \pm SE (n=12). Significant differences (P<0.05) between species are indicated by different letters.

CONCLUSIONS

- All studied *Cistus* showed a water status and stomatal conductance with maximum values in spring followed by a progressive decline or interruption during the summer drought and a partial recovery in response to autumn rains.
- C. monspeliensis* and *C. ladanifer*, exhibited a drought-tolerance mechanism, *C. psilosepalus* an avoidance mechanism and *C. populifolius* is in-between avoidance and tolerance mechanisms.

REFERENCES

- Cochard H (2002) A technique for measuring xylem hydraulic conductance under high negative pressures. *Plant Cell Environ* 25: 815–819
- Cochard H, Damour G, Bodet C, Tharwat I, Poirier M, Améglio T (2005) Evaluation of a new centrifuge technique for rapid generation of xylem vulnerability curves. *Physiologia Plantarum* 124: 410–418
- Simões MP, Madeira M, Gazarini L (2009) Ability of *Cistus* L. shrubs to promote soil rehabilitation in extensive oak woodlands of Mediterranean areas. *Plant and Soil* 323(1):249–265
- Acknowledgements: The authors are grateful to "Centro de Geofísica da Universidade de Évora" for meteorological data. Thanks are due also to ICAAM for financial support.