

ORIGINAL ARTICLE

Sodium and potassium urinary excretion and dietary intake:
a cross-sectional analysis in adolescentsCarla Gonçalves^{1*}, Sandra Abreu², Patrícia Padrão^{1,3}, Olívia Pinho^{1,4}, Pedro Graça^{1,5},
João Breda⁶, Rute Santos^{2,7,8} and Pedro Moreira^{1,2,3}

¹Faculty of Nutrition and Food Sciences, University of Porto, Porto, Portugal; ²Research Centre in Physical Activity, Health and Leisure, Faculty of Sport, University of Porto, Porto, Portugal; ³Institute of Public Health, University of Porto (ISPUP), EPIUnit-Epidemiology Research Unit, Porto, Portugal; ⁴LAQV/REQUIMTE, University of Porto, Porto, Portugal; ⁵Directorate General of Health, Lisbon, Portugal; ⁶Division of Noncommunicable Diseases and Health through the Life-Course, World Health Organization Regional Office for Europe, UN City, Denmark; ⁷Maia University Institute, Maia, Portugal; ⁸Early Start Research Institute, School of Education, Faculty of Social Sciences, University of Wollongong, Australia

Abstract

Background: Hypertension is the leading cause for heart disease and stroke, for mortality and morbidity worldwide, and a high sodium-to-potassium intake ratio is considered a stronger risk factor for hypertension than sodium alone.

Objective: This study aims to evaluate sodium and potassium urinary excretion, and assess the food sources of these nutrients in a sample of Portuguese adolescents.

Design: A cross-sectional study with a sample of 250 Portuguese adolescents. Sodium and potassium excretion were measured by one 24-h urinary collection, and the coefficient of creatinine was used to validate completeness of urine collections. Dietary sources of sodium and potassium were assessed using a 24-h dietary recall.

Results: Valid urine collections were provided by 200 adolescents (118 girls) with a median age of 14.0 in both sexes ($p = 0.295$). Regarding sodium, the mean urinary excretion was 3,725 mg/day in boys and 3,062 mg/day in girls ($p < 0.01$), and 9.8% of boys and 22% of girls met the World Health Organization (WHO) recommendations for sodium intake. Concerning potassium, the mean urinary excretion was 2,237 mg/day in boys and 1,904 mg/day in girls ($p < 0.01$), and 6.1% of boys and 1.7% of girls met the WHO recommendations for potassium intake. Major dietary sources for sodium intake were cereal and cereal products (41%), meat products (16%), and milk and milk products (11%); and for potassium intake, main sources were milk and milk products (21%), meat products (17%), and vegetables (15%).

Conclusions: Adolescents had a high-sodium and low-potassium diet, well above the WHO recommendations. Health promotion interventions are needed in order to decrease sodium and increase potassium intake.

Keywords: *urinary sodium; urinary potassium; intake; adolescents; salt*

Responsible Editor: Per Ole Iversen, Oslo University, Norway.

Received: 15 August 2015; Revised: 23 February 2016; Accepted: 4 March 2016; Published: 11 April 2016

Evidence demonstrates that high sodium intake increases blood pressure and has an impact in endothelial dysfunction, cardiovascular function and structure, kidney disease, and cardiovascular morbidity and mortality (1, 2). Hypertension is recognized as a primary risk factor of heart disease and stroke, both leading causes of death worldwide (3). Recent data regarding sodium intake show that populations around the world are consuming far more sodium than is physiologically necessary (4). In response, the World Health Organization (WHO) considers the reduction of sodium intake

to be a priority concern (5), and WHO member states have agreed to work toward a global reduction of 30% until 2025 (6).

At the same time, interest in potassium intake has grown, mainly because potassium attenuates sodium's negative effects, by reducing stroke rates and cardiovascular risk (7) and increasing urinary sodium excretion (8). Furthermore, low dietary intakes of potassium potentiate the sodium sensitivity of blood pressure (9) and the risk of hypertension, while the relationship between sodium and blood pressure strengthens if the urinary

sodium-to-potassium ratio (Na^+/K^+) is considered instead of only sodium excretion rate (10). High Na^+/K^+ ratio intake is considered to be a stronger risk factor of hypertension and cardiovascular disease than each of these nutrients alone (11, 12), and the benefits of a higher intake of potassium are particularly important when sodium intake is high (13).

Data from around the world suggest that the average potassium consumption in the populations of many countries is less than the 2,730–3,120 mg/day, the reference value recommended by the 2002 Joint WHO/Food and Agriculture Organization expert consultation (14).

The WHO recommends a maximum sodium intake of 2,000 mg/day (15) for children and a potassium intake of at least 3,510 mg/day, which should be adjusted downward for children based on their energy requirements (16).

In Portugal, cerebrovascular and cardiovascular diseases are the major causes of death (17), and nearly 42% of the adult population has hypertension (18). In the adult population, the latest data indicate an average consumption of 4,200 mg sodium/day and 2,900 mg potassium/day (18), whereas data regarding the youngest population remain scarce. To the best of our knowledge, no study has yet characterized sodium and potassium intake in Portuguese adolescents using the method of 24-h urine collection (19). This method is considered the ‘gold standard’ for assessing the distribution and average intake of sodium in a representative population (20, 21).

Accurate estimates of sodium and potassium intake are essential for monitoring the effectiveness of current actions to reduce sodium intake and to improve efforts to increase potassium consumption.

In this study, we thus aimed to 1) describe dietary intakes of sodium and potassium and Na^+/K^+ ratio; 2) assess their compliance with sodium and potassium intake guidelines; and 3) investigate the main food sources of total dietary sodium and potassium intake among Portuguese adolescents.

Methods

Study design

Data for the present cross-sectional study came from a 3-year follow up study with Portuguese children and adolescents, the Longitudinal Analysis of Biomarkers and Environmental Determinants of Physical activity (LabMed Physical Activity Study). The LabMed Physical Activity Study aimed to evaluate independent and combined associations of dietary intake, physical activity, and sedentary behavior on fitness levels and other factors in 12- to 18-year-old children and adolescents.

From the participating schools all students enrolled in the seventh and tenth grade in 2011 scholar year were invited to participate in the study ($n = 1,678$). The sample for this analysis consisted of a convenience subgroup of

LabMed Physical Activity Study aged 13 to 18 years ($n = 398$) that was invited to perform 24-h urine collection and 250 subjects (63%) voluntarily agreed to perform 24-h urine collections. Urine collections were performed between 2012 and 2014. After a validation control of 24-h urine collection (described below), 50 urine collections were rejected (20%). Thus, the final sample consisted of 200 adolescents (82 boys) with both valid urine collection and corresponding dietary recall.

This study was conducted according to the guidelines laid down in the Declaration of Helsinki and all procedures were approved by the Ethical Commission of University of Porto. Written informed consent was obtained from all subjects and caregivers.

24-h Urine collection

Participants were asked to complete a 24-h urine collection. Participants and caregivers received oral and written instructions on how to collect complete 24-h urine samples. They were instructed to discard the first morning void and to collect all urine over the following 24-h, including the first void on the next morning, and the time of the start and finish collection was recorded in a questionnaire. Considering the participant’s comfort and for feasibility purposes, urine collections were held on Sundays and, during the collection period, subjects were asked to store the collected urine in a cool place. Urine samples were sent to a certified laboratory to be analyzed for urinary creatinine (mg/day) (Jaffé reaction, Siemens Advia 1650), urinary sodium (mEq/day), and urinary potassium (mEq/day) (indirect ion-selective electrodes methodology, Siemens Advia 1800). The maximum coefficient of variation of laboratorial methods was 0.71% for sodium, 1.2% for potassium, and 4.55% for creatinine. Sodium and potassium excretion was reported in mEq/day; however, for comparative purposes, it was converted to mg/day by using their molecular weight. Na^+/K^+ ratio was also calculated as sodium (mg/day) divided by potassium (mg/day).

For 24-h urine collection validation, quality control was used calculating 24-h urinary creatinine excretion in relation to body weight according to age group (22), and valid collections were considered when creatinine was between 0.1–0.245 and 0.117–0.294 mmol/kg/day for boys at 9–13 years and 14–18 years, respectively; and between 0.117–0.244 and 0.129–0.238 mmol/kg/day for girls at 9–13 years and 14–18 years, respectively. If the urine collection was incomplete ($n = 50$) or if the subjects took medication on the day of collection ($n = 2$), the urine specimen was rejected and not considered for analysis. The subjects who reported taking medication on the day of collection agreed to collect 24-h urine again.

Dietary record

A 24-h dietary recall refers to the day urine was collected by trained interviewers using a photographic book and

household measures to quantify portion sizes (23). Energy and nutritional intake were estimated using an adapted Portuguese version of the nutritional analysis software Food Processor Plus (ESHA Research Inc., Salem, OR, USA). The nutrient content of local food was taken from standard nutrient tables (24), whereas the content of commercial food (e.g. pizza and ready-to-eat-food) was derived from labeled ingredients and nutrients.

The food codes used were categorized into 13 major food groups (25): 1) cereal and cereal products (bread; breakfast cereals; biscuits, cakes, puddings, scones, and doughnuts; pasta, rice, and other cereal-based products); 2) meat products (chicken and turkey dishes; sausages; bacon and ham; red meats); 3) milk and milk products (milks, yogurts, and cheeses); 4) vegetable and potato products (potato products); 5) soups and sauces (vegetable soups and sauces); 6) fruits; 7) fish and seafood dishes (cod fish); 8) oils and fats; 9) egg and egg dishes; 10) sugars, preserves, and confectionery; 11) fast food (pizza; sandwiches, burgers, filled wraps; salted snacks and fried snacks; fried potatoes); 12) beverages (soft drinks; water; hot beverages); 13) other foods (beans, pulses, canned fruit, and pickles). The contribution of these 13 food groups to sodium and potassium intake was calculated. An additional analysis was performed to evaluate the contribution of each food group to total sodium and potassium intake only in participants who ingested foods included in that food group.

For validation of records and to check for under-reporting, the ratio of reported energy intake (EI) and estimated basal metabolic rate (BMR) according to Schofield (26) was used taking into account age, sex, body weight, and height. Using the formulas proposed by Goldberg et al. (27), we calculated individual specific cutoffs with the following modifications: using the new estimates of the physical activity level assuming light physical activity given for the different age/sex groups of adolescents (28), using coefficients of variation for EI (23%) (29), and the use one recorded day.

Therefore, records with EI:BMR ratios below the cutoff values, depending on the subject's age and sex, were not considered a plausible measurement of the actual one day EI in our further analysis and were excluded ($n = 22$, 11%).

Anthropometric measures

For weight and height measurements, we used a digital scale (Tanita Inner Scan BC 532, Tokyo, Japan) and a portable stadiometer (Seca 213, Hamburg, Germany), respectively. All measurements were performed with participants in light clothing, without shoes, and according to standard procedures (30).

According to body mass index (BMI)-for-age z-scores from the WHO, participants were classified as thin (< -2 SD), normal weight, overweight ($> +1$ SD), or obese ($> +2$ SD) (31). The classification was performed using WHO AnthroPlus Software (32).

Socioeconomic status

As an indicator of the socioeconomic status of the household, the Family Affluence Scale (FAS) was used (33). The FAS is a four-item questionnaire that helps students report their family income objectively: it evaluates the sum of scores regarding whether the family owns a car, whether the student has his or her own bedroom, the number of family vacations during the past 12 months, and the number of computers the family owns. The final score ranges from 0 to 9 points, with higher scores indicating higher socioeconomic status.

Statistical analysis

All statistical analyses were performed using SPSS 21.0 Inc. A p -value < 0.05 was considered to indicate statistical significance. The Kolmogorov–Smirnov test was used to assess the assumption of normality. Independent sample t -test or Mann–Whitney U test were performed to compare continuous variables, and the Chi-Square test was used for categorical variables. Dietary sources of sodium and potassium were reported through the 13 food and beverages categories. A proportion to total food sodium or potassium was used to report results as mean percentage contribution of each category. Nutrients and food intake were also energy-adjusted according to residual method (34), and adjusted values were used to evaluate the differences between sexes. Spearman's correlation coefficient was used to identify and test the strength of a relationship between urinary sodium with dietary sodium and urinary potassium with dietary potassium.

Table 1. Sample characteristics of the study sample (13–18 years) ($n = 200$)

	Boys	Girls	p
Age (years) ^{a,b}	14.7 ± 1.3	14.9 ± 1.4	0.30
Weight (kg) ^{a,b}	60.2 ± 12.2	55.1 ± 9.3	0.01
Height (m) ^{a,b}	1.7 ± 0.1	1.6 ± 0.1	<0.01
Weight status (%) ^c			
Thin	1.2	0.8	
Normal weight	73.2	74.6	
Overweight	17.1	17.8	
Obese	8.5	6.8	<0.01
FAS ^{a,b}	6.5 ± 1.8	6.7 ± 1.6	0.70

FAS, Family Affluence Scale.

^a Values are mean ± standard deviation.

^b Between-sex analysis by Mann–Whitney U test.

^c Analysis by χ^2 for categorical variables.

Table 2. Urinary data on sodium and potassium excretion by sex (13–18 years)^a

	Boys <i>n</i> = 82	Girls <i>n</i> = 118	<i>p</i>
Sodium (mg/day) ^{a,b}	3,725 ± 1,445	3,062 ± 1,379	< 0.01
% of sodium excretion below 2,000 mg/day ^c	9.8	22	
Potassium (mg/day) ^{a,b}	2,237 ± 704	1,904 ± 593	< 0.01
% of potassium excretion above 3,510 mg/day ^d	6.1	1.7	
Na/K ratio ^{a,b}	1.7 ± 0.6	1.6 ± 0.6	0.11

Na/K ratio, sodium-to-potassium ratio.

^aValues are mean ± standard deviation.

^bAnalysis by Mann–Whitney U test.

^cAccording to the WHO guidelines for sodium intake (15).

^dAccording to the WHO guidelines for potassium intake (16).

Results

Descriptive characteristics of the sample are presented in Table 1. Adolescents were on average 14 years old and most exhibited normal BMI.

Results from urine collection are shown in Table 2. Boys showed a greater median sodium and potassium excretion than girls ($p < 0.01$). Of all participants, 83% exhibited sodium intake above upper-limit recommendations (2,000 mg/day), while 96.1% showed potassium intake below recommendations (3,510 mg/day). The Na⁺/K⁺ ratio were similar in both sexes (1.7 in boys and 1.6 in girls, $p = 0.11$). The coefficient of variation was 43.2% for sodium excretion, 32.3% for potassium excretion, and 26.8% for creatinine excretion.

In this study, sodium and potassium intakes determined by dietary records correlated (weak correlations) with 24-h sodium and potassium excretion values ($r = 0.152$, $p = 0.03$ for sodium and $r = 0.229$, $p < 0.01$ for potassium).

As shown in Table 3, boys reported higher EI than girls ($p < 0.01$). From dietary records, mean sodium intake was 2,649 mg/day for boys and 2,106 mg/day for girls ($p < 0.01$), which corresponds to a mean estimated salt intake of 7 g/day for boys and 5 g/day for girls. Mean potassium intake was 2,998 mg/day for boys and 2,471 mg/day for girls ($p = 0.01$).

The major food sources of sodium in participants' diet (Fig. 1) were cereal and cereal products (mean 41%: including bread, 16%; pasta, rice, and other cereal-based products, 12%; and breakfast cereals, 5%); meat products (mean 16%: including bacon and ham, 7%; sausages, 4%; red meats, 3%; and chicken and turkey dishes, 2%); and milk and milk products (mean 11%: including milk, 4%, and cheese, 4%). Fast food contributed a mean of 9% to total sodium ingestion, including the contribution of pizza (mean 3%), salted and fried snacks (mean 1%), fried

potatoes (mean 3%) and sandwiches, and burgers and filled wraps (mean 2%).

By contrast, milk and milk products (mean 21%: including milk, 14%, and cheese, 1%), meat products (mean 17%: including red meats, 9%; chicken and turkey dishes, 5%; bacon and ham, 2%; and sausages, 1%) and vegetables (mean 15%) were major sources of total potassium intake (Fig. 2). Fast food contributed 10% to total potassium ingestion, including the large contribution of fried potatoes (mean 8%).

The contribution of food groups to total sodium and potassium intake by sex, considering only participants who ingested foods included in the groups is shown in Table 4.

Discussion

To the best of our knowledge, this study has marked the first in Portugal to estimate sodium and potassium intake in a large group of adolescents aged 13–18 years using 24-h urine excretion. Our results have shown that 83% of participants exceed the recommended sodium intake and that 96.1% did not meet recommendations for potassium intake.

Portuguese data related to sodium excretion are scarce. A study of Portuguese children aged 10–12 years showed that sodium excretion was 3,072 ± 985 mg/day (35), while a more recent study with children aged 8–10 years reported that median sodium excretion was 2,737 mg/day in boys and 2,104 mg/day in girls (36). Comparing our results to those representing other European adolescents, Portuguese adolescent boys seem to have a mean sodium excretion greater than that of Italian (2,967 mg/day) (37), Spanish (3,270.6 mg/day) (38), English (i.e. London) (3,401.47 mg/day) (25), and German (3,013 mg/day) adolescents (39). Similar results were found between Portuguese and other European adolescent girls, except the Spanish (2,888.8 mg/day) (38). Across the board, sodium excretion was greater in boys than in girls (25, 36–39), probably due to their higher food intake.

Conversely, other studies have evaluated sodium intake in children and adolescents via dietary recall. In Australian adolescents (aged 14–16 years), mean dietary sodium intake was 3,190 mg/day, a value that increased with age (40). In French adolescents, median sodium intake was 2,245 mg/day (41) and in Korean adolescents mean sodium ingestion was 4,100 mg/day (42). Results from our study related to sodium intake via dietary recall are more similar to French adolescents, which is lower than Korean and Australian adolescents.

The amount of potassium excretion in our study was well below the minimum value recommended by the WHO, but it is comparable with those reported by Geleijnse et al. (43) regarding 233 Dutch children aged 5–17 years, among whom mean potassium excretion was

Table 3. Dietary intake of the sample by sex (13–18 years) crude and adjusted to total energy intake ($n = 178$)

	Boys	Girls	<i>p</i>	<i>p</i> *
Nutritional intake				
Total energy (kcal/day) ^{a,b}	2,449 ± 929	1,981 ± 665	<0.01	<0.01
Carbohydrates (g/day) ^{a,b}	285 ± 113	242 ± 83	0.01	0.31
Carbohydrates (%TEI) ^{a,b}	47 ± 7	49 ± 8	0.08	0.20
Sugars (g/day) ^{a,b}	131 ± 64	115 ± 57	0.02	0.41
Protein (g/day) ^{a,b}	103 ± 53	79 ± 32	<0.01	0.61
Protein (%TEI) ^{a,b}	17 ± 4	16 ± 4	0.53	0.38
Fat (g/day) ^{a,b}	93 ± 42	73 ± 34	<0.01	0.91
Fat (% TEI) ^{a,c}	34 ± 7	33 ± 7	0.21	0.81
Dietary fiber (g/1,000 kcal) ^{a,b}	6 ± 3	7 ± 3	0.07	0.09
Total food (g/day) ^{a,b}	2,540 ± 916	2,219 ± 735	0.02	0.89
Total water (g/day) ^{a,b,d}	1,781 ± 713	1,632 ± 609	0.22	0.46
Potassium (mg/day) ^{a,b}	2,998 ± 1,368	2,471 ± 1,068	0.01	0.52
Sodium (mg/day) ^{a,b,e}	2,649 ± 1,586	2,106 ± 1,204	<0.01	0.29
Salt (g/day) ^{a,b,e}	7 ± 4	5 ± 3	<0.01	0.29
Na/K ratio ^{a,b}	1.7 ± 0.6	1.6 ± 0.6	0.19	0.35
Dietary intake				
Cereal and cereal products (g/day) ^{b,f}	286 (227, 357)	225 (162, 309)	0.01	0.22
Breads (g/day) ^{b,f}	93 (45, 143)	80 (45, 120)	0.26	0.67
Breakfast cereals (g/day) ^{b,f}	30 (30, 40)	30 (30, 40)	0.12	0.52
Biscuits, cakes, puddings, scones, doughnuts (g/day) ^{b,f}	60 (29, 124)	75 (40, 117)	0.52	0.05
Pasta, rice, and other cereal-based products (g/day) ^{b,f}	151 (97, 224)	97 (43, 180)	<0.01	0.01
Meat products (g/day) ^{b,f}	160 (120, 252)	132 (80, 212)	0.01	0.80
Chicken and turkey dishes (g/day) ^{b,f}	118 (93, 220)	105 (99, 173)	0.44	0.87
Sausages (g/day) ^{b,f}	40 (20, 60)	25 (14, 45)	0.12	0.17
Bacon and ham (g/day) ^{b,f}	30 (23, 60)	30 (23, 45)	0.43	0.83
Red meats (g/day) ^{b,f}	119 (90, 195)	112 (56, 160)	0.10	0.33
Milk and milk products (g/day) ^{b,f}	545 (292, 841)	471 (304, 659)	0.07	0.17
Milk (g/day) ^{b,f}	516 (283, 841)	400 (280, 545)	0.01	0.01
Cheeses (g/day) ^{b,f}	30 (28, 60)	30 (28, 56)	0.86	0.71
Vegetable and potato products (g/day) ^{b,f}	123 (73, 175)	121 (63, 224)	0.84	0.18
Potato products (g/day) ^{b,f}	155 (93, 350)	155 (116, 231)	0.77	0.86
Soups and sauces (g/day) ^{b,f}	363 (117, 541)	363 (174, 368)	0.50	0.71
Vegetable soups (g/day) ^{b,f}	363 (363, 541)	363.0 (261.5, 458.8)	0.10	0.18
Sauces (g/day) ^{b,f}	15 (11, 68)	10 (8, 20)	0.18	0.32
Fruits (g/day) ^{b,f}	301 (174, 487)	174 (132, 328)	0.10	0.19
Fish and seafood dishes (g/day) ^{b,f}	184 (108, 272)	120 (75, 227)	0.14	0.24
Cod fish (g/day) ^{b,f}	236 (71.0, -)	114 (68, 233)	0.50	0.55
Oils and fats (g/day) ^{b,f}	10 (4, 27)	10 (5, 21)	0.53	0.18
Egg and egg dishes (g/day) ^{b,f}	100 (58, 100)	46 (22, 50)	0.03	0.03
Sugars, preserves, and confectionery (g/day) ^{b,f}	16 (10, 37)	12 (8, 25)	0.07	0.92
Fast food (g/day) ^{b,f}	150 (100, 333)	150 (82, 202)	0.85	0.37
Pizza (g/day) ^{b,f}	220 (191, 445)	301 (200, 403)	0.94	0.29
Sandwiches, burgers, filled wraps (g/day) ^{b,f}	219 (219, 219)	219 (80, 219)	0.40	0.75
Salted snacks and fried snacks (g/day) ^{b,f}	104 (52, 229)	144 (88, 147)	0.77	0.46
Fried potatoes (g/day) ^{b,f}	100 (100, 150)	100 (50, 123)	0.21	0.26
Beverages (g/day) ^{b,f}	720 (476, 1,140)	825 (540, 1,047)	0.58	0.04
Soft drinks (g/day) ^{b,f}	435 (200, 615)	495 (220, 707)	0.37	0.11

Table 3. (Continued)

	Boys	Girls	p	p*
Water (g/day) ^{b,f}	410 (90, 720)	440 (140, 846)	0.54	0.73
Hot beverages (g/day) ^{b,f}	30 (4, 122)	102 (10, 295)	0.20	0.34
Other foods (g/day) ^{b,f}	69 (22, 185)	81 (21, 127)	0.83	0.93

TEI, Total Energy Intake.

^aValues are mean ± standard deviation.

^bAnalysis by Mann–Whitney U test.

^cAnalysis by Student’s t-test for continuous variables.

^dWater from foods, beverages, and metabolic water.

^eEstimated from dietary records without considering household salt, 393 mg sodium = 1 g salt.

^fValues are median (P25, P75).

*Adjusted for energy intake.

1,704.3 mg/day and was higher in boys. Among Portuguese children (8–10 years old), mean potassium excretion was 1,701 ± 594 mg/day in boys and 1,682 ± 541 mg/day in girls (36).

Na⁺/K⁺ ratio intake is a critical risk factor for hypertension (44), and in our study, mean urinary Na⁺/K⁺ ratio was 1.7 in boys and 1.6 in girls; these values are higher than Na⁺/K⁺ < 0.59 mg/mg the WHO recommends (45). Higher Na⁺/K⁺ ratio is associated with Western diets heavily reliant on processed foods, high in added sodium, and low in potassium (38).

Na⁺/K⁺ ratio excretion among our study’s participants was smaller than the median ratio found in Portuguese children (2.73 in boys and 2.33 in girls) (36) and in Dutch adolescents (ratio was 3.3) (43), as well as in Italian children and adolescents (ratio was 3.79 ± 1.68) (46). However, the ratio in our participants is still three times greater than 0.59 mg/mg, which raises concerns about poor dietary behavior early in childhood.

Participants in our study did not meet the recommended intake for sodium or potassium, and major dietary

contributors of sodium and potassium were identified in order to improve food-based intervention programs.

Correlations between dietary and urinary data found in our study were comparable with those reported by other authors (41, 47). The possible explanation of this result could be the difficulty in maintaining food records that accurately estimate food quantities and food ingredients, especially if the subjects eat out of home.

Taking into account the overall intake reported for our sample of adolescents, the food groups that contribute most to total sodium intake are cereals and cereal products (mean 41%) and meat products (mean 16%). Meaning that, in addition to bread (mean 16%), sodium added during the cooking process is the chief source of sodium intake as in pasta, rice, and other cereal-based products (mean 12%), meat products (mean 16%), fish and seafood products (mean 5%), and vegetable soup (mean 10%). Our data show that in individuals who consumed vegetable soup (approximately 15% of participants), soup contributed roughly a third to their total sodium intake (29.6% for boys and 31.4% for girls in relation to total dietary sodium




Fig. 1. Contribution (%) of dietary sources for total sodium intake to total sample of subjects (n = 178). The contribution of each food category is expressed in mean percentage.




Fig. 2. Contribution (%) of dietary sources for total potassium intake to total sample of subjects (n = 178). The contribution of each food category is expressed in mean percentage.

Table 4. Contribution (%) of dietary sources for total sodium and potassium intake by sex considering subjects that consume those food groups ($n = 178$)

	% contribution to sodium intake						% contribution to potassium intake					
	<i>n</i>	Boys	<i>n</i>	Girls	<i>p</i>	<i>p</i> *	<i>n</i>	Boys	<i>n</i>	Girls	<i>p</i>	<i>p</i> *
Cereal and cereal products	73	39.5 ± 16.2	105	42.5 ± 18.0	0.37	0.54	73	10.9 ± 7.7	105	11.3 ± 8.8	0.92	0.59
Breads	62	18.1 ± 10.7	85	19.4 ± 11.6	0.63	0.76	62	4.6 ± 3.1	85	5.1 ± 3.9	0.68	0.63
Breakfast cereals	36	9.0 ± 5.9	47	10.6 ± 7.6	0.26	0.87	36	3.4 ± 3.6	47	4.1 ± 3.0	0.03	0.67
Biscuits, cakes, puddings, scones, doughnuts	45	10.0 ± 7.6	75	16.4 ± 10.4	<0.01	<0.01	45	4.6 ± 4.6	75	4.6 ± 4.6	0.73	0.68
Pasta, rice, and other cereal-based products	60	16.5 ± 11.7	83	14.1 ± 10.8	0.13	0.11	60	3.3 ± 3.7	83	3.4 ± 5.5	0.07	0.23
Meat products	71	18.3 ± 14.1	95	15.7 ± 13.5	0.16	0.69	71	19.3 ± 11.8	95	18.2 ± 12.8	0.25	0.58
Chicken and turkey dishes	25	4.4 ± 2.7	42	5.3 ± 4.0	0.36	0.65	25	12.3 ± 9.1	42	12.6 ± 8.7	0.63	0.17
Sausages	22	17.9 ± 12.7	14	16.3 ± 12.4	0.63	0.82	22	3.6 ± 2.6	14	3.0 ± 1.9	0.77	0.73
Bacon and ham	31	15.1 ± 9.9	42	18.2 ± 8.2	0.06	0.12	31	5.2 ± 4.0	42	5.4 ± 4.6	0.75	0.79
Red meats	48	6.9 ± 8.4	54	4.8 ± 6.9	0.09	0.30	48	16.9 ± 11.3	54	15.9 ± 11.5	0.63	0.28
Milk and milk products	71	10.4 ± 7.8	97	12.0 ± 8.9	0.31	0.38	71	21.6 ± 13.0	97	22.7 ± 13.4	0.85	0.63
Milk	61	5.2 ± 5.0	85	5.2 ± 4.2	0.37	0.73	61	17.6 ± 10.7	85	16.9 ± 11.7	0.47	0.15
Cheeses	26	10.5 ± 6.5	35	12.4 ± 7.4	0.23	0.35	26	3.1 ± 2.3	35	3.2 ± 2.8	0.92	0.78
Vegetable and potato products	45	5.3 ± 11.6	77	1.7 ± 3.9	0.93	0.63	45	16.1 ± 16.8	77	23.3 ± 19.2	0.16	0.13
Potato products	24	8.8 ± 15.2	47	2.0 ± 4.6	0.11	0.33	24	27.3 ± 16.8	47	31.5 ± 17.8	0.36	0.98
Soups and sauces	34	23.0 ± 15.2	42	28.6 ± 15.1	0.15	0.96	34	16.3 ± 11.9	42	15.2 ± 10.8	0.89	0.52
Vegetable soups	24	29.6 ± 12.0	32	31.4 ± 12.9	0.70	0.29	24	21.3 ± 9.6	32	19.2 ± 9.1	0.39	0.10
Sauces	9	4.8 ± 5.3	7	2.3 ± 1.1	0.96	0.87	9	2.9 ± 3.6	7	1.3 ± 2.2	0.79	0.63
Fruits	36	0.2 ± 0.4	61	0.1 ± 0.1	0.31	0.02	36	17.9 ± 11.7	61	17.7 ± 15.3	0.64	0.12
Fish and seafood dishes	18	17.0 ± 17.6	36	17.2 ± 15.3	0.74	0.75	18	17.1 ± 14.5	36	12.9 ± 11.8	0.36	0.12
Cod fish	3	22.9 ± 13.9	13	24.3 ± 19.1	0.84	0.84	3	1.6 ± 1.1	13	2.4 ± 2.5	0.64	0.74
Oils and fats	34	2.2 ± 2.9	61	2.2 ± 3.8	0.91	0.61	34	0.1 ± 0.1	61	0.1 ± 0.2	0.48	0.83
Egg and egg dishes	5	6.8 ± 5.4	17	4.0 ± 3.3	0.29	0.78	5	3.3 ± 1.4	17	3.1 ± 2.3	0.72	0.29
Sugars, preserves, and confectionery	42	1.3 ± 2.1	59	0.8 ± 1.0	0.57	0.43	42	2.6 ± 6.2	59	1.4 ± 2.0	0.73	0.94
Fast food	38	18.5 ± 17.3	35	24.0 ± 20.1	0.28	0.11	38	25.9 ± 15.3	35	26.7 ± 14.9	0.83	0.92
Pizza	5	36.6 ± 19.8	8	41.3 ± 13.0	0.56	0.72	5	23.7 ± 19.3	8	19.9 ± 8.9	0.59	0.81
Sandwiches, burgers, filled wraps	7	27.0 ± 16.6	7	22.2 ± 19.7	0.75	0.41	7	15.9 ± 13.0	7	13.6 ± 6.0	0.66	0.57
Salted snacks and fried snacks	4	21.2 ± 11.2	4	27.1 ± 19.4	0.56	0.99	4	10.5 ± 3.0	4	7.7 ± 6.5	0.51	0.99
Fried potatoes	32	7.7 ± 4.7	26	9.4 ± 8.1	0.73	0.78	32	23.0 ± 10.1	26	24.8 ± 13.2	0.90	0.55
Beverages	71	1.2 ± 1.1	105	1.4 ± 1.2	0.09	0.08	71	2.6 ± 7.4	105	2.4 ± 5.3	0.02	<0.01
Soft drinks	45	0.9 ± 1.2	53	1.2 ± 1.5	0.20	0.20	45	0.3 ± 0.3	53	1.4 ± 3.9	0.84	0.49
Water	55	0.7 ± 0.6	85	0.9 ± 0.7	0.19	0.41	55	0.6 ± 1.6	85	0.5 ± 1.0	0.36	0.05
Hot beverages	9	0.6 ± 0.8	26	0.2 ± 0.2	0.76	0.49	9	14.9 ± 15.7	26	5.2 ± 7.4	0.23	0.38
Other foods	16	6.7 ± 8.2	28	7.2 ± 9.3	0.63	0.53	16	8.3 ± 11.0	28	7.0 ± 10.1	0.48	0.76

The contribution of each food category is expressed in mean percentage ± standard deviation.

*Adjusted for energy intake.

intake; Table 4). Conversely, in individuals who consumed processed foods such as fast food (about 41% of participants), such foods became an important contributor to their total sodium intake.

Since the largest proportion of sodium is added during food manufacture or preparation, including for vegetable soup and bread, changes in recipes to reduce sodium content could be possible without disturbing other nutrient values (48) and consumer preferences. In a study of children and the elderly, 30% salt reduction in vegetable soup was achieved without compromising perceived salti-

ness (49). This result is especially important in Portugal where the amount of sodium in a 300 g portion of vegetable soup cooked outside home may be as high as 1,316 mg (50).

To diminish salt added during cooking processes, it is necessary to devise nutrition education strategies that increase public awareness of salt reduction and develop technology to control the amounts of added salt.

At the same time, industry and policy makers need to engage efforts to make processed foods with low sodium content available on the market. Portugal's salt reduction

plan (51) includes a strategic objective to increase the availability of products with low salt content, via the participation of the industry. Therefore, since 2009 in Portugal, sodium content in bread was legally limited to 550 mg/100 g (52), although this limit was exceptionally permissive compared with that of other countries' legislation (53, 54). Bread is probably the most important staple food of modern Portuguese diet (55) as well as of the traditional Mediterranean diet of the early 1960s (56). Although salt (i.e. sodium chloride) has a technical and functional role in the manufacture of bread (57), the production of low-salt bread is feasible (57), and incremental reductions of 30 to 50% have proven to be acceptable to consumers (58–60).

Surprisingly, the top sources of potassium among all sample of participants were milk and milk products (21%) and meat products (17%), as explained by the low intake of fruit and vegetables (2 and 9%, respectively; data not shown). These data may reflect the low proportion of adolescents who follow the recommendation to eat at least 400 g of fruit and vegetables daily (61). The intake of fruits and vegetables as a natural source of potassium is inversely related to the risk of stroke (62) and is associated with more varied and higher quality diet (63).

Food processing reduces the natural amount of potassium in many food products (64). Adolescent diets in our study were high in processed foods and low in fresh fruits and vegetables with high sodium intake. It suggests that Portuguese adolescents have moved away from a typical Mediterranean diet toward the dietary patterns common in industrialized countries (41). Developing strategies to improve fruit and vegetable consumption would thus increase potassium intake, as well as offer other beneficial health effects (65).

In the context of our study, vegetable soup deserves particular attention, for it is a staple food of the Mediterranean diet and of Portuguese dietary habits, consumed daily by 67% of population (66) and promoted by healthy eating education programs (67). Portuguese-style soups are rich in vegetables, low in energy density, and high in dietary fiber, all of which are associated with a lower risk of obesity (68). In our study, vegetable soup was an important contributor to total potassium intake; however, its contribution to total sodium was exceptionally high and therefore detrimental. This typical food seems to show the potential for improving potassium intake if its consumption could be stimulated in adolescents and added salt were diminished.

A strategy that emphasizes reducing sodium added to foods, promotes potassium-rich foods, and promotes reformulation of processed and packaged foods could achieve greater public health benefits than restricting sodium alone (69).

Our study is strengthened by the 24-h urine collection, the clinical gold standard method, and by our sample

size (considering that a representative sample of 120 to 240 individuals from each population subgroup is needed to estimate population-level sodium intake according to WHO (21)). The 24-h urine collection objectively measured dietary sodium intake from food and from salt added during cooking and at the table. Bias stemming from under collection or overcollection was decreased by 24-h urine creatinine excretion quality control and self-reported total daily collection. The fact that we performed only one urine collection per participant could be a limitation; however, urine collections were measured in a large quantity of participants and showed a response rate of 63% similar to that of other studies (19).

Another potential limitation of our study is the possible underestimation of sodium and potassium intake in 24-h dietary recall, as has been previously reported (47, 70). Nevertheless, 24-h recalls were administered by trained researchers using a photographic book and household measures to quantify portion sizes, which minimized this potential limitation. It should also be noted that the food composition data in software used for calculating nutrient intakes might also introduce bias into dietary data. To avoid this, we carefully checked the sodium and potassium composition in the foods consumed by participants. We should additionally note that the collection of urine and the respective diet recall refers to Sunday. Although variation in day-to-day dietary intake exists, at their age, our participants were more likely concerned about what their peers think about them, and the thought of completing urine collection during weekdays was roundly rejected by adolescents and their parents at the outset of the study. For this reason, we performed urine data collection on Sunday only.

Conclusions

We found that Portuguese adolescents have high sodium intake and low potassium intake compared with WHO recommendations. The low potassium intake raises concerns about the potential detachment of Portuguese adolescents' dietary habits from the traditional Mediterranean diet with increased intake of processed foods and few vegetables. Sodium reduction in widely consumed staple foods such as cereal products/bread and vegetable soup could be effective for decreasing sodium intake, while the promotion of milk, fruit, and vegetable intake could similarly increase potassium intake.

Disclaimer

João Breda is a staff member of the WHO. The authors are responsible for the content and writing of this paper, which does not necessarily represent the decisions, stated policy, or views of the WHO.

Acknowledgements

The authors are grateful for the participation of all adolescents and their parents, teachers, and schools in the LabMed Study. They also thank the volunteers for their cooperation and the Research Centre in Physical Activity, Health and Leisure (University of Porto), for sponsoring the LabMed Study.

Conflict of interest and funding

None of the authors declare conflict of interest. This study was supported by WHO and FCT grants: PTDC/DES/116586/2010 and BPD/102381/2014. The Research Centre on Physical Activity Health and Leisure (CIA-FEL) is supported by UID/DTP/00617/2013. The author Rute Santos has a Discovery Early Career Research Award from the Australian Research Council (DE150101921).

References

- He FJ, MacGregor GA. A comprehensive review on salt and health and current experience of worldwide salt reduction programmes. *J Hum Hypertens* 2009; 23(6): 363–84.
- Poggio R, Gutierrez L, Matta MG, Elorriaga N, Irazola V, Rubinstein A. Daily sodium consumption and CVD mortality in the general population: systematic review and meta-analysis of prospective studies. *Public Health Nutr* 2015; 18(4): 695–704.
- Chockalingam A, Campbell NR, Fodor JG. Worldwide epidemic of hypertension. *Can J Cardiol* 2006; 22(7): 553–5.
- Brown IJ, Tzoulaki I, Candeias V, Elliott P. Salt intakes around the world: implications for public health. *Int J Epidemiol* 2009; 38(3): 791–813.
- World Health Organization (2013). Vienna declaration on nutrition and noncommunicable diseases in the context of health 2020. WHO European Ministerial Conference on Nutrition and Noncommunicable Diseases in the Context of Health 2020. Vienna: World Health Organization.
- Webster J, Trieu K, Dunford E, Hawkes C. Target salt 2025: a global overview of national programs to encourage the food industry to reduce salt in foods. *Nutrients* 2014; 6(8): 3274–87.
- Aaron KJ, Sanders PW. Role of dietary salt and potassium intake in cardiovascular health and disease: a review of the evidence. *Mayo Clin Proc* 2013; 88(9): 987–95.
- Morris RC, Jr, Schmidlin O, Frassetto LA, Sebastian A. Relationship and interaction between sodium and potassium. *J Am Coll Nutr* 2006; 25(3 Suppl): 262s–70.
- Kotchen TA, Kotchen JM. Nutrition, diet, and hypertension. In: Shils ME, Shike M, Ross AC, Caballero B, Cousins RJ (eds.). *Modern nutrition in health and disease*. 10th ed. Philadelphia, PA: Lippincott Williams & Wilkins; 2006, pp. 1095–107.
- Intersalt Cooperative Research Group. Intersalt: an international study of electrolyte excretion and blood pressure. Results for 24 hour urinary sodium and potassium excretion. Intersalt Cooperative Research Group. *BMJ (Clinical research ed)* 1988; 297(6644): 319–28.
- Yang Q, Zhang Z, Kuklina EV, Fang J, Ayala C, Hong Y, et al. Sodium intake and blood pressure among US children and adolescents. *Pediatrics* 2012; 130(4): 611–19.
- Umesawa M, Iso H, Date C, Yamamoto A, Toyoshima H, Watanabe Y, et al. Relations between dietary sodium and potassium intakes and mortality from cardiovascular disease: the Japan Collaborative Cohort Study for Evaluation of Cancer Risks. *Am J Clin Nutr* 2008; 88(1): 195–202.
- Thornton SN. Salt in health and disease – a delicate balance. *N Engl J Med* 2013; 368(26): 2531.
- WHO (2003). Diet, nutrition and the prevention of chronic diseases. World Health Organization technical report series. 2003/05/29 ed. Geneva: World Health Organization (WHO). pp. i–viii, 1–149, backcover.
- World Health Organization (2012). Guideline: sodium intake for adults and children. Available from: http://www.who.int/nutrition/publications/guidelines/sodium_intake/en/ [cited 5 September 2014].
- World Health Organization (2012). Guideline: potassium intake for adults and children. Available from: http://www.who.int/nutrition/publications/guidelines/potassium_intake/en/ [cited 5 September 2014].
- George F. Causes of deaths in Portugal and challenges in prevention. *Acta Med Port* 2012; 25(2): 61–3.
- Polonia J, Martins L, Pinto F, Nazare J. Prevalence, awareness, treatment and control of hypertension and salt intake in Portugal: changes over a decade. The PHYSA study. *J Hypertens* 2014; 32(6): 1211–21.
- McLean RM. Measuring population sodium intake: a review of methods. *Nutrients* 2014; 6(11): 4651–62.
- Elliott P, Brown I. Sodium intakes around the world. Geneva, Switzerland: World Health Organization; 2007.
- World Health Organization/Pan American Health Organization Regional Expert Group for Cardiovascular Disease Prevention through Population-wide Dietary Salt Reduction (2010). Protocol for population level sodium determination in 24-hour urine samples. Available from: <http://new.paho.org/hq/dmdocuments/2010/pahosaltprotocol.pdf> [cited 5 September 2014].
- Remer T, Neubert A, Maser-Gluth C. Anthropometry-based reference values for 24-h urinary creatinine excretion during growth and their use in endocrine and nutritional research. *Am J Clin Nutr* 2002; 75(3): 561–9.
- Marques M, Pinho O, Almeida MVD. Manual de quantificação de alimentos. Porto: Rocha/Artes Gráficas, Lda; 1996.
- Martins I. Tabela da Composição de Alimentos. Lisboa: Editorial do Ministério da Educação; 2007.
- Marrero NM, He FJ, Whincup P, Macgregor GA. Salt intake of children and adolescents in South London: consumption levels and dietary sources. *Hypertension* 2014; 63(5): 1026–32.
- Schofield WN. Predicting basal metabolic rate, new standards and review of previous work. *Hum Nutr Clin Nutr* 1985; 39 (Suppl 1): 5–41.
- Goldberg GR, Black AE, Jebb SA, Cole TJ, Murgatroyd PR, Coward WA, et al. Critical evaluation of energy intake data using fundamental principles of energy physiology: 1. Derivation of cut-off limits to identify under-recording. *Eur J Clin Nutr* 1991; 45(12): 569–81.
- Torun B, Davies PS, Livingstone MB, Paolisso M, Sackett R, Spurr GB. Energy requirements and dietary energy recommendations for children and adolescents 1 to 18 years old. *Eur J Clin Nutr* 1996; 50(Suppl 1): S37–80; discussion S-1.
- Nelson M, Black AE, Morris JA, Cole TJ. Between- and within-subject variation in nutrient intake from infancy to old age: estimating the number of days required to rank dietary intakes with desired precision. *Am J Clin Nutr* 1989; 50(1): 155–67.
- Lohman T, Roache A, Martorell R. Anthropometric standardization reference manual. *Med Sci Sports Exerc* 1992; 24(8): 952.
- de Onis M, Onyango AW, Borghi E, Siyam A, Nishida C, Siekmann J. Development of a WHO growth reference for school-aged children and adolescents. *Bull World Health Organ* 2007; 85(9): 660–7.

32. WHO (2009). *AnthroPlus for personal computers manual: software for assessing growth of the world's children and adolescents*. Geneva: WHO.
33. Currie C, Molcho M, Boyce W, Holstein B, Torsheim T, Richter M. Researching health inequalities in adolescents: the development of the Health Behaviour in School-Aged Children (HBSC) family affluence scale. *Soc Sci Med* 2008; 66(6): 1429–36.
34. Willett W. *Nutritional epidemiology*. New York: Oxford University Press; 2012.
35. Cotter J, Cotter MJ, Oliveira P, Cunha P, Polonia J. Salt intake in children 10–12 years old and its modification by active working practices in a school garden. *J Hypertens* 2013; 31(10): 1966–71.
36. Oliveira AC, Padrao P, Moreira A, Pinto M, Neto M, Santos T, et al. Potassium urinary excretion and dietary intake: a cross-sectional analysis in 8-10 year-old children. *BMC Pediatr* 2015; 15(1): 60.
37. Campanozzi A, Avallone S, Barbato A, Iacone R, Russo O, De Filippo G, et al. High sodium and low potassium intake among Italian children: relationship with age, body mass and blood pressure. *PLoS One* 2015; 10(4): e0121183.
38. Maldonado-Martin A, Garcia-Matarin L, Gil-Extremera B, Avivar-Oyonarte C, Garcia-Granados ME, Gil-Garcia F, et al. Blood pressure and urinary excretion of electrolytes in Spanish schoolchildren. *J Hum Hypertens* 2002; 16(7): 473–8.
39. Shi L, Krupp D, Remer T. Salt, fruit and vegetable consumption and blood pressure development: a longitudinal investigation in healthy children. *Br J Nutr* 2014; 111(4): 662–71.
40. Grimes CA, Riddell LJ, Campbell KJ. Dietary salt intake, sugar-sweetened beverage consumption, and obesity risk. *Pediatrics* 2013; 131: 14–21.
41. Meneton P, Lafay L, Tard A, Dufour A, Ireland J, Menard J, et al. Dietary sources and correlates of sodium and potassium intakes in the French general population. *Eur J Clin Nutr* 2009; 63(10): 1169–75.
42. Lee HS, Duffey KJ, Popkin BM. Sodium and potassium intake patterns and trends in South Korea. *J Hum Hypertens* 2013; 27(5): 298–303.
43. Geleijnse JM, Grobbee DE, Hofman A. Sodium and potassium intake and blood pressure change in childhood. *BMJ (Clinical research ed)* 1990; 300(6729): 899–902.
44. Adrogue HJ, Madias NE. The impact of sodium and potassium on hypertension risk. *Semin Nephrol* 2014; 34(3): 257–72.
45. Yi SS, Curtis CJ, Angell SY, Anderson CA, Jung M, Kansagra SM. Highlighting the ratio of sodium to potassium in population-level dietary assessments: cross-sectional data from New York City, USA. *Public Health Nutr* 2014; 17(11): 2484–8.
46. De Santo NG, Dilorio B, Capasso G, Russo F, Stamler J, Stamler R, et al. The urinary sodium/potassium ratio in children from southern Italy living in Cimitile: a case for concern. *Int J Pediatr Nephrol* 1987; 8(3): 153–8.
47. Schachter J, Harper PH, Radin ME, Caggiula AW, McDonald RH, Diven WF. Comparison of sodium and potassium intake with excretion. *Hypertension* 1980; 2(5): 695–9.
48. Witschi JC, Capper AL, Hosmer DW, Jr, Ellison RC. Sources of sodium, potassium, and energy in the diets of adolescents. *J Am Diet Assoc* 1987; 87(12): 1651–5.
49. Gonçalves C, Monteiro S, Padrao P, Rocha A, Abreu S, Pinho O, et al. Salt reduction in vegetable soup does not affect saltiness intensity and liking in the elderly and children. *Food Nutr Res* 2014; 58: 24825, doi: <http://dx.doi.org/10.3402/fnr.v58.24825>
50. Gonçalves C, Silva G, Pinho O, Camelo S, Amaro L, Teixeira V, et al. Sodium content in vegetable soups prepared outside the home: identifying the problem. In: Arezes P, Baptista J, Barroso M, Carneiro P, Cordeiro P, Costa N, et al., eds. *Occupational Safety and Hygiene – SHO 2012 – Book of Abstracts*. Vila Nova de Gaia: Portuguese Society of Occupational Safety and Hygiene (SPOSHO); 2012, 278–281.
51. Graça P. *Relatório Estratégia para a redução do consumo de sal na alimentação em Portugal*. Direção Geral de Saúde; 2013.
52. Assembleia da Republica. *Lei n 75/2009 Estabelece normas com vista à redução do teor de sal no pão bem como informação na rotulagem de alimentos embalados destinados ao consumo humano*. Diário da República 2009.
53. Dunford EK, Eyles H, Mhurchu CN, Webster JL, Neal BC. Changes in the sodium content of bread in Australia and New Zealand between 2007 and 2010: implications for policy. *Med J Aust* 2011; 195(6): 346–9.
54. Wyness LA, Buttriss JL, Stanner SA. Reducing the population's sodium intake: the UK Food Standards Agency's salt reduction programme. *Public Health Nutr* 2012; 15(2): 254–61.
55. Burton PM, Monro JA, Alvarez L, Gallagher E. Glycemic impact and health: new horizons in white bread formulations. *Crit Rev Food Sci Nutr* 2011; 51(10): 965–82.
56. D'Alessandro A, De Pergola G. Mediterranean diet pyramid: a proposal for Italian people. *Nutrients* 2014; 6(10): 4302–16.
57. Belz MC, Ryan LA, Arendt EK. The impact of salt reduction in bread: a review. *Crit Rev Food Sci Nutr* 2012; 52(6): 514–24.
58. Bolhuis DP, Temme EH, Koeman FT, Noort MW, Kremer S, Janssen AM. A salt reduction of 50% in bread does not decrease bread consumption or increase sodium intake by the choice of sandwich fillings. *J Nutr* 2011; 141(12): 2249–55.
59. La Croix KW, Fiala SC, Colonna AE, Durham CA, Morrissey MT, Drum DK, et al. Consumer detection and acceptability of reduced-sodium bread. *Public Health Nutr* 2015; 18(8): 1412–18.
60. Willems AA, van Hout DH, Zijlstra N, Zandstra EH. Effects of salt labelling and repeated in-home consumption on long-term liking of reduced-salt soups. *Public Health Nutr* 2014; 17(5): 1130–7.
61. World Health Organization. Diet, nutrition and the prevention of chronic diseases. *World Health Organ Tech Rep Ser* 2003; 916(i–viii): 1–149.
62. Hu D, Huang J, Wang Y, Zhang D, Qu Y. Fruits and vegetables consumption and risk of stroke: a meta-analysis of prospective cohort studies. *Stroke* 2014; 45(6): 1613–19.
63. Rodriguez-Rodriguez E, Ortega RM, Andres Carvajales P, Gonzalez-Rodriguez LG. Relationship between 24 h urinary potassium and diet quality in the adult Spanish population. *Public Health Nutr* 2015; 18(5): 850–9.
64. Webster JL, Dunford EK, Neal BC. A systematic survey of the sodium contents of processed foods. *Am J Clin Nutr* 2010; 91(2): 413–20.
65. He FJ, MacGregor GA. Beneficial effects of potassium on human health. *Physiol Plant* 2008; 133(4): 725–35.
66. Instituto Nacional de Estatística I.P., Instituto Nacional de Saúde Doutor Ricardo Jorge I.P. (2009). *Inquérito Nacional de Saúde 2005/2006*. Portugal, Lisboa: Instituto Nacional de Estatística IP.
67. Rodrigues SS, Franchini B, Graça P, de Almeida MD. A new food guide for the Portuguese population: development and technical considerations. *J Nutr Educ Behav* 2006; 38(3): 189–95.
68. Moreira P, Padrao P. Educational, economic and dietary determinants of obesity in Portuguese adults: a cross-sectional study. *Eat Behav* 2006; 7(3): 220–8.
69. Yang Q, Liu T, Kuklina EV, Flanders WD, Hong Y, Gillespie C, et al. Sodium and potassium intake and mortality among us adults: prospective data from the third national health and

- nutrition examination survey. *Arch Intern Med* 2011; 171(13): 1183–91.
70. Espeland MA, Kumanyika S, Wilson AC, Reboussin DM, Easter L, Self M, et al. Statistical issues in analyzing 24-hour dietary recall and 24-hour urine collection data for sodium and potassium intakes. *Am J Epidemiol* 2001; 153(10): 996–1006.

***Carla Gonçalves**

Faculty of Nutrition and Food Sciences
University of Porto
Rua Roberto Frias, 4200-465 Porto
Portugal
Email: carlagoncalves.pt@gmail.com