

2016

The disability rate of 5-year post-stroke and its correlation factors: A national survey in China

Y Yang

Y-Z Shi

N Zhang

S Wang

G Ungvari

University of Notre Dame Australia, Gabor.Ungvari@nd.edu.au

See next page for additional authors

Follow this and additional works at: http://researchonline.nd.edu.au/med_article



Part of the [Medicine and Health Sciences Commons](#)

This article was originally published as:

Yang, Y., Shi, Y., Zhang, N., Wang, S., Ungvari, G., Ng, C., Wang, Y., Zhao, X., Wang, Y., Wang, C., & Xiang, Y. (2016). The disability rate of 5-year post-stroke and its correlation factors: A national survey in China. *PLoS One*, 11 (11).

Original article available here:

<http://journals.plos.org/plosone/article?id=10.1371/journal.pone.0165341>

This article is posted on ResearchOnline@ND at
http://researchonline.nd.edu.au/med_article/770. For more information,
please contact researchonline@nd.edu.au.



Authors

Y Yang, Y-Z Shi, N Zhang, S Wang, G Ungvari, C Ng, Y-L Wang, X-Q Zhao, Y-J Wang, C-X Wang, and Y-T Xiang

This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source.

See: <https://creativecommons.org/licenses/by/4.0/>

No changes have been made to this article was originally published in *PLoS One* available at: <http://journals.plos.org/plosone/article?id=10.1371/journal.pone.0165341>

Yang, Y., Shi, Y-Z, Zhang, N., Wang, S., Ungvari, G., Ng, C., Wang, Y-L, Zhao, X-Q, Wang, Y-J, Wang, C-X and Xiang, Y-T (2016) The disability rate of 5-year post-stroke and its correlation factors: A national survey in China. *PLoS One*, 11(11). doi: 10.1371/journal.pone.0165341

RESEARCH ARTICLE

The Disability Rate of 5-Year Post-Stroke and Its Correlation Factors: A National Survey in China

Yang Yang^{1,2,3,4,5}, Yu-Zhi Shi^{1,3,4,5}, Ning Zhang^{1,2,3,4,5}, Shuo Wang^{1,3,4,5}, Gabor S. Ungvari^{6,7}, Chee H. Ng⁸, Yi-Long Wang^{1,3,4,5}, Xing-Quan Zhao^{1,3,4,5}, Yong-Jun Wang^{1,3,4,5}, Chun-Xue Wang^{1,2,3,4,5*}, Yu-Tao Xiang^{9*}

1 Department of Neurology, Beijing Tian Tan Hospital, Capital Medical University, Beijing, China, **2** Department of Neuropsychiatry & Behavioral Neurology and Clinical Psychology, Beijing Tian Tan Hospital, Capital Medical University, Beijing, China, **3** China National Clinical Research Center for Neurological Diseases, Beijing, China, **4** Center of Stroke, Beijing Institute for Brain Disorders, Beijing, China, **5** Beijing Key Laboratory of Translational Medicine for Cerebrovascular Disease, Beijing, China, **6** University of Notre Dame Australia / Marian Centre, Perth, Australia, **7** School of Psychiatry & Clinical Neurosciences, University of Western Australia, Perth, Australia, **8** Department of Psychiatry, University of Melbourne, Melbourne, Victoria, Australia, **9** Unit of Psychiatry, Faculty of Health Sciences, University of Macau, Macao SAR, China

* snowsen@126.com (CXW); xyutly@gmail.com (YTX)



OPEN ACCESS

Citation: Yang Y, Shi Y-Z, Zhang N, Wang S, Ungvari GS, Ng CH, et al. (2016) The Disability Rate of 5-Year Post-Stroke and Its Correlation Factors: A National Survey in China. PLoS ONE 11 (11): e0165341. doi:10.1371/journal.pone.0165341

Editor: Xiaoying Wang, Massachusetts General Hospital, UNITED STATES

Received: August 1, 2016

Accepted: October 10, 2016

Published: November 8, 2016

Copyright: © 2016 Yang et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: The Clinical Research Ethics Committee of Beijing Tiantan Hospital that approved the study prohibits the authors from making the research data set publicly available. Readers and all interested researchers may contact Dr. Chunxue Wang (Email address: snowsen@126.com) for details. Dr. Wang could apply to the Clinical Research Ethics Committee of Beijing Tiantan Hospital for the release of the data.

Funding: This study was funded by the Ministry of Science and Technology and the Ministry of Health of the People's Republic of China. Individual grants

Abstract

Few studies on long-term functional outcome have been conducted in post-stroke patients in China. The objective of this study was to conduct a nationwide survey in China to investigate the 5-year prevalence of post-stroke disability and its correlation factors. A total of 893 patients with ischemic stroke were included. Demographic, clinical and neuro-imaging information were collected with standardized instruments that assessed stroke severity, depression, cognitive impairment, stroke recurrence and physical disability. Disability was assessed with the modified Ranking Score (mRS), of which a cutoff score ≥ 2 indicates disability. Statistical analysis included chi-square tests, two independent samples t-tests, Mann-Whitney U test and multiple logistic regression analysis. The frequency of disability in this study population was 45%. Multivariate analyses revealed that older age, lower education level, previous history of stroke, stroke severity at admission, depression, cognitive impairment at 3 months, and stroke recurrence within 5 years follow up were all significantly associated with post-stroke disability. The disability rate in 5-year post-stroke was high in Chinese patients. Treatment of depression, secondary prevention of stroke and rehabilitation may benefit disabled patients with stroke in China.

Introduction

In 2010, approximately 16.9 million persons worldwide developed first-ever stroke. There was an estimate of 33 million stroke survivors [1], of whom a substantial proportion suffered from disability. Disability after stroke is a significant outcome that leads to personal suffering and

include the National 11th & 12th Five-year S & T Major Project (2006BAI01A11, 2011BAI08B01, 2011BAI08B02), the National Key Technology Research and Development Program of the Ministry of Science and Technology of China (2013BAI09B03), the Ministry of Science and Technology of the People's Republic of China (2012ZX09303-005-001), Beijing Institute for Brain Disorders (BIBD-PXM2013_014226_07_000084), Beijing Biobank of Cerebral Vascular Disease (D131100005313003), Beijing Brain Research (Z16110000216131) and Beijing Municipal Science & Technology Commission (Z151100004015127 and Z151100003915117).

Competing Interests: The authors have declared that no competing interests exist.

burden on their families [2]. Most studies have focused on short-term data within 3 months post-stroke [3]. However, long-term functional status, as measured by the activities of daily living (ADL) in one particular study, does not remain stable between 3 months and 1 year post-stroke [4]. Another 5-year follow-up study found a delayed but significant functional decline in stroke survivors [5]. Growing evidence supports the importance of early rehabilitation intervention after stroke, but strategies to reduce the risk of long-term post-stroke disability beyond a year remain unclear. Long-term post-stroke outcomes are usually determined by different factors compared to short-term outcomes [6, 7]. In addition, post-stroke disability significantly increases the risk for adverse outcomes, such as recurrent stroke and increased mortality [8, 9]. Thus, the identification of post-stroke disability, associated risk factors, and appropriate rehabilitation strategies are important for long-term prognosis of stroke survivors.

The objective of this study was to conduct a nationwide survey in China to investigate the 5-year prevalence of post-stroke disability and its associated risk factors.

Methods

Study Settings and Patients

This 5-year follow-up study of stroke patients is a national project entitled the “Prospective Cohort Study on the Incidence and Outcome of Patients with Post-stroke Depression in China (PRIOD)” [10]. At baseline, patients in 56 hospitals nationwide were consecutively recruited and enrolled if they were aged 18 years or older, had an acute stroke according to WHO criteria [11] confirmed by CT or MRI within 14 days, could complete a clinical interview and provide informed consent. Patients with severe cognitive impairment and history of drug dependence were excluded. The study protocol was approved by the medical ethics committees of Beijing Tiantan Hospital and the participating institutions, respectively. All participants provided written informed consent.

Measurement Tools and Evaluation

In this study we used data at baseline, 3 months and 5 years after the index stroke. Basic socio-demographic, clinical and neuro-imaging features at baseline were recorded using a standard form, and supplemented by medical records review and a clinical interview by qualified neurologists.

Stroke severity was assessed at baseline by the National Institutes of Health Stroke Scale (NIHSS) [12]. Stroke lesions was classified as left, right, frontal, temporal, occipital-parietal lobes, basal ganglia, thalamus, brainstem and cerebellum based on MRI or CT scans reported by radiologists. Current smoking was defined as at least 1 cigarette in the past month. Moderate/heavy drinking was defined as ≥ 2 standard drinks/day.

Post-stroke depression at 3 months follow up was based on the checklist of the DSM-IV criteria [13]. The Chinese version of the Hamilton Rating Scale for Depression (HRSD) [14, 15] was used to assess severity of depression. Cognition function was measured by the Mini Mental State Examination (MMSE) [16] at 3 months after stroke. Cognitive impairment was defined as the MMSE total score < 24 [17, 18].

The 5-year assessments including stroke recurrence and physical disability were conducted via a telephone interview. Recurrent stroke was defined as any type of stroke diagnosed at re-admission during the past 5 years [19]. Post-stroke functional status was assessed using the modified Ranking Score (mRS), with a score ≥ 2 as meeting the criteria for disability [20].

Statistical analyses

Statistical analyses were performed with SPSS Statistics 21.0 for Windows. The comparisons between the disability and non-disability groups with regard to demographic, clinical and neuro-imaging variables were conducted with chi-square tests, two independent samples t-tests and Mann-Whitney U test, as appropriate. Multiple logistic regression analysis used the “enter” method (i.e., all specified independent variables were entered at one time) to determine the independent relationships between disability at 5 years follow up and variables that significantly differed in the above univariate analysis. The level of significance was set at 0.05 (two-tailed).

Results

Out of 2,324 ischemic stroke patients who fulfilled study entry criteria at baseline, 248 died, 696 could not be located via the information recorded in medical records, and 474 refused or did not complete the telephone interview at the 5-year assessment. Finally, data on 893 patients were included in the statistical analyses (Fig 1). Compared to patients who completed the




Fig 1. Flow diagram of the recruitment.

doi:10.1371/journal.pone.0165341.g001

5-year assessment, those who were excluded for any reasons were older and more likely to be single marital status (S1 Table). In addition, there were missing data in several variables (cardiac disease, current smoking and family history of stroke) with the percentages of less than 0.5%, except for hyperlipidemia (1.5%).

Altogether, 402 post-stroke patients (45.0%) met the criteria of disability. Table 1 shows the basic demographic and clinical characteristics of the whole sample, together with comparative data between the groups with and without disability. Compared to the group without disability, the group with disability at 5 years follow up were more likely to be older, female gender, have diabetes, cardiac disease, previous history of stroke, cognitive impairment and depression at 3 months, and stroke recurrence within the past 5 years, and have higher NIHSS score at baseline, higher HRSD score at 3 months, lower MMSE score at 3 months, but lower education level and less current smoking and moderate/heavy drinking.

Table 2 shows the independent correlates of disability in stroke patients. Older age, low education level, previous history of stroke, more severe stroke at baseline, greater depression and cognitive impairment at 3 months, and stroke recurrence within the past 5 years were significantly associated with disability ($R^2 = 0.47$; $p < 0.001$).

Discussion

To the best of our knowledge, this is the first national large-scale longitudinal study in China to examine the disability rate of 5-year post-stroke and its associated factors.

In this study, 45% of post-stroke patients had disability, which is consistent with the range (36%-71%) reported in other Western surveys [21–23]. Variation between studies may partly be attributed to differences in definitions of disability (e.g. Barthel Index vs. mRS), length of follow up, post-stroke illness stage and sample size. In this study, those who dropped out were older than those who participated. Given the impact of advanced age on post-stroke disability, the prevalence rate in our study could be an under-estimation.

In this study cognitive impairment at 3-month post-stroke significantly contributed to the disability at 5-year follow up, which is consistent with earlier findings [18, 24, 25]. One study found that cognitive impairment, as defined by the MMSE score ≤ 24 at 3 months after the index stroke, independently predicted functional outcomes as measured by the Frenchay Activities Index up to 3 years in stroke survivors [18]. The mechanism for the association between cognitive impairment and poor functional outcome is unclear [26, 27]. Cognitive impairment was observed in multiple cognitive domains, especially in executive function, attention and processing speed, both of which are closely associated with physical function [24, 28]. Of note, both comprehensive neuropsychological instruments (e.g., the digit span test and frontal assessment battery) and brief screening tests (e.g., the MMSE or Montreal Cognitive Assessment) on cognition could detect vascular cognitive impairment [29].

The association between post-stroke depression and clinical outcomes in previous studies have been mixed [30]. Similar to some studies [31, 32], but not others [18, 33], we found that depression at 3 months post-stroke was highly predictive of disability at 5-year follow up. A recent study involving 1,101 stroke survivors in south London registry also found that depression at 3 months after stroke was predictive for disability up to 5 years follow up [32]. Depression causes both behavioral (e.g. decreased motivation or poor medication adherence) and biological changes (e.g. dysregulation in autonomic system activation, hypothalamic-pituitary-adrenocortical axis and inflammation states), which could increase the risk of disability [2, 32]. Post-stroke depression is likely to have a recurrent and chronic pattern which may partly explain the impact of depression after stroke on long-term disability [34].

Table 1. Comparison between stroke patients with and without disability at 5-year follow up with respect to demographic and clinical characteristics.

	Total sample (n = 893)		No-disability (n = 491)		Disability (n = 402)		Statistics		
	N	%	N	%	N	%	χ^2	df	p
Male	614	68.8	352	71.7	262	65.2	4.3	1	0.03
Married/ cohabited	842	94.3	469	95.5	373	92.8	3.0	1	0.08
High school or above	367	41.1	219	44.6	148	36.8	5.5	1	0.01
Diabetes	217	24.3	106	21.6	111	27.6	4.3	1	0.03
Hyperlipidemia	249	28.3	130	27.0	119	29.9	0.9	1	0.33
Hypertension	620	69.4	328	66.8	292	72.6	3.5	1	0.06
Cardiac disease	189	21.3	89	18.2	100	25.0	6.0	1	0.01
Current smoking	333	37.4	205	41.9	128	31.9	9.4	1	0.002
Moderate/Heavy-drinking	138	15.5	94	19.1	44	10.9	11.3	1	0.001
Stroke History	197	22.1	70	14.3	127	31.6	38.6	1	0.001
Stroke Family History	179	20.1	104	21.2	75	18.8	0.8	1	0.36
Toast Classification									
Large-Artery etiology	570	63.8	308	62.7	262	65.2	0.5	1	0.44
Cardio-embolism etiology	24	2.7	9	1.8	15	3.7	3.0	1	0.08
Small-vessels etiology	216	24.2	118	24.0	98	24.4	0.01	1	0.90
Other determined or undetermined etiology	41	4.6	27	5.5	14	3.5	2.0	1	0.15
Neuro-imaging characteristics									
Left lesions	328	36.7	180	36.7	148	36.8	0.002	1	0.96
Right lesions	335	37.5	192	39.1	143	35.6	1.1	1	0.27
Frontal lobe	114	12.8	63	12.8	51	12.7	0.004	1	0.94
Temporal lobe	77	8.6	42	8.6	35	8.7	0.007	1	0.93
Parietal lobe	87	9.7	51	10.4	36	9.0	0.5	1	0.47
Occipital lobe	63	7.1	34	6.9	29	7.2	0.02	1	0.86
Basal ganglia	433	48.5	240	48.9	193	48.0	0.06	1	0.79
Thalamus	84	9.4	49	10.0	35	8.7	0.4	1	0.51
Brainstem	186	20.8	102	20.8	84	20.9	0.002	1	0.96
Cerebellum	68	7.6	36	7.3	32	8.0	0.1	1	0.72
Cerebral Lesions	237	26.5	133	27.1	104	25.9	0.1	1	0.68
Sub-cortical Lesions	495	55.4	274	55.8	221	55.0	0.06	1	0.80
Infra-tentorial Lesions	232	26.0	128	26.1	104	25.9	0.005	1	0.94
Cognitive Impairment at 3 months	129	14.4	41	8.4	88	21.9	32.7	1	<0.001
Depression at 3 months	198	22.2	90	18.3	108	26.9	9.3	1	0.002
Stroke recurrence within 5-years	211	23.6	65	13.2	146	36.3	65.2	1	<0.001
	Mean	SD	Mean	SD	Mean	SD	T / Z	df	p
Age (yrs)	60.6	10.7	57.6	10.2	64.2	10.2	-9.5	891	<0.001
NIHSS score at admission	4.4	3.6	3.9	3.2	5.0	3.9	-4.0	— ^a	<0.001
HRSD total at 3 months	4.8	4.8	4.3	4.5	5.3	5.0	-3.2	— ^a	0.001
MMSE total at 3 months	27.2	3.6	28.1	2.4	26.2	4.4	-7.3	— ^a	<0.001

Bold values are p<0.05; a = Mann-Whitney U test; HRSD = Hamilton Rating Scale for Depression-17; MMSE = Mini Mental State Examination; NIHSS = National Institutes of Health Stroke Scale; Toast = Trial of Org 10172 in Acute Stroke Treatment

doi:10.1371/journal.pone.0165341.t001

Consistent with previous findings [35–37], older age, more severe stroke, previous history of stroke and stroke recurrence are found to be significant risk factors for long-term disability. Previous history of stroke and stroke recurrence may cause a culmination of deficits related to

Table 2. Independent correlates of disability at 5 years in stroke patients (multiple logistic regression analysis with the group without disability as reference group).

Variables	Stroke patients*		
	p	OR	95% CI
Age	<0.001	1.06	1.04, 1.08
Male	0.20	0.7	0.4, 1.1
High school or above	0.02	0.6	0.4, 0.9
Diabetes	0.75	1.06	0.7, 1.6
Cardiac Disease	0.69	1.09	0.7, 1.6
Current smoking	0.09	1.4	0.9, 2.2
Moderate/Heavy Drinking	0.65	1.1	0.6, 1.9
Stroke History	<0.001	2.6	1.7, 4.1
NIHSS score at admission	<0.001	1.1	1.05, 1.1
Depression at 3 months	0.009	1.8	1.1, 2.9
Cognitive impairment at 3 months	<0.001	2.7	1.6, 4.7
Stroke Recurrence within 5 years	<0.001	4.1	2.7, 6.3

Bold values are p<0.05

* adjusting for study site; NIHSS = National Institutes of Health Stroke Scale

doi:10.1371/journal.pone.0165341.t002

gait and impaired motor function arising from other brain lesions [2, 38]. Higher education was correlated with less disability, which is consistent with a study reported in India [39]. Well-educated patients may have less stigma, be more conscious about outcomes of stroke, and more likely to adhere to medications, and actively participate in physical rehabilitation [30], which could reduce the risk for long-term disability.

The major merits of this study include the large, multi-site, nationwide sample, the long-term 5-year outcome evaluation and the use of standardized assessments. However, there are several limitations in this study. First, only 38.4% of the baseline eligible patients completed the 5-year assessment. Due to the rapid urbanization and huge population migration in China, the high dropout rate in large scale cohort studies was unavoidable. Second, the mRS may not be sensitive enough to detect subtle changes in physical functioning. However, its contents are clinically distinct and responsive to patient-centered outcomes in stroke patients [40]. Third, it should be noted that the MMSE is not a stroke-specific cognitive measure. Fourth, several important variables, such as poststroke rehabilitation therapy, treatment adherence and the size of infarcts, white matter lesions and brain atrophy, were not evaluated at the 5-year assessment due to unavailable data. In addition, due to significant missing data of medication used, this was not included for analyses. Finally, patients who could not complete a clinical interview were excluded at baseline, which could lead to potential selection bias. Despite these methodological shortcomings, the findings are likely to approximate the general long-term profile of stroke patients across China.

In conclusion, long-term disability is highly prevalent in Chinese post-stroke patients. Neuropsychiatric (depression and cognitive impairment) and sociodemographic (older age and lower educational level) factors, poor stroke management (previous stroke history and recurrence stroke) and previous stroke severity are negatively associated with functional outcome. Adequate treatment of depression, secondary prevention of strokes and appropriate rehabilitation strategy need to be considered to reduce disability in post-stroke patients in China.

Supporting Information

S1 Table. Comparison of baseline characteristics between study completers and non-completers at 5-year.
(DOC)

Acknowledgments

We would like to thank all clinicians, patients and their families involved in this project.

Author Contributions

Conceptualization: YJW CXW YLW XQZ.

Data curation: YY YZS NZ SW.

Formal analysis: YY YTX.

Funding acquisition: YJW CXW.

Investigation: YY YZS NZ SW.

Methodology: YJW CXW YTX.

Project administration: YJW CXW YLW XQZ.

Resources: YJW CXW YLW.

Supervision: YJW CXW.

Writing – original draft: YY CXW YTX.

Writing – review & editing: GSU CHN.

References

1. Feigin VL, Forouzanfar MH, Krishnamurthi R, Mensah GA, Connor M, Bennett DA, et al. Global and regional burden of stroke during 1990–2010: findings from the Global Burden of Disease Study 2010. *Lancet*. 2014; 383(9913):245–54. PMID: [24449944](#)
2. Dhamoon MS, McClure LA, White CL, Lakshminarayan K, Benavente OR, Elkind MS. Long-term disability after lacunar stroke: secondary prevention of small subcortical strokes. *Neurology*. 2015; 84(10):1002–8. doi: [10.1212/WNL.0000000000001331](#) PMID: [25663222](#)
3. Quinn TJ, Dawson J, Walters MR, Lees KR. Functional outcome measures in contemporary stroke trials. *Int J Stroke*. 2009; 4(3):200–5. doi: [10.1111/j.1747-4949.2009.00271.x](#) PMID: [19659822](#)
4. Ullberg T, Zia E, Petersson J, Norrving B. Changes in functional outcome over the first year after stroke: an observational study from the Swedish stroke register. *Stroke*. 2015; 46(2):389–94. doi: [10.1161/strokeaha.114.006538](#) PMID: [25538204](#)
5. Dhamoon MS, Moon YP, Paik MC, Boden-Albala B, Rundek T, Sacco RL, et al. Long-term functional recovery after first ischemic stroke: the Northern Manhattan Study. *Stroke*. 2009; 40(8):2805–11. doi: [10.1161/strokeaha.109.549576](#) PMID: [19556535](#)
6. Feigin VL, Barker-Collo S, Parag V, Senior H, Lawes CM, Ratnasabapathy Y, et al. Auckland Stroke Outcomes Study. Part 1: Gender, stroke types, ethnicity, and functional outcomes 5 years poststroke. *Neurology*. 2010; 75(18):1597–607. doi: [10.1212/WNL.0b013e3181fb44b3](#) PMID: [21041783](#)
7. Woo J, Kay R, Yuen YK, Nicholls MG. Factors influencing long-term survival and disability among three-month stroke survivors. *Neuroepidemiology*. 1992; 11(3):143–50. PMID: [1407250](#)
8. Huybrechts KF, Caro JJ, Xenakis JJ, Vemmos KN. The prognostic value of the modified Rankin Scale score for long-term survival after first-ever stroke. Results from the Athens Stroke Registry. *Cerebrovasc Dis*. 2008; 26(4):381–7. doi: [10.1159/000151678](#) PMID: [18753743](#)
9. Park JH, Ovbiagele B. Relationship of functional disability after a recent stroke with recurrent stroke risk. *Eur J Neurol*. 2016; 23(2):361–7. doi: [10.1111/ene.12837](#) PMID: [26493027](#)

10. Zhang N, Wang CX, Wang AX, Bai Y, Zhou Y, Wang YL, et al. Time course of depression and one-year prognosis of patients with stroke in mainland China. *CNS Neurosci Ther*. 2012; 18(6):475–81. doi: [10.1111/j.1755-5949.2012.00312.x](https://doi.org/10.1111/j.1755-5949.2012.00312.x) PMID: [22672300](https://pubmed.ncbi.nlm.nih.gov/22672300/)
11. Kunitz SC, Gross CR, Heyman A, Kase CS, Mohr JP, Price TR, et al. The pilot Stroke Data Bank: definition, design, and data. *Stroke*. 1984; 15(4):740–6. PMID: [6464070](https://pubmed.ncbi.nlm.nih.gov/6464070/)
12. Lyden P, Brott T, Tilley B, Welch KM, Mascha EJ, Levine S, et al. Improved reliability of the NIH Stroke Scale using video training. NINDS TPA Stroke Study Group. *Stroke*. 1994; 25(11):2220–6. PMID: [7974549](https://pubmed.ncbi.nlm.nih.gov/7974549/)
13. First MB. Diagnostic and statistical manual of mental disorders. DSM IV-4th edition APA. 1994:1994.
14. Hamilton M. A rating scale for depression. *J Neurol Neurosurg Psychiatry*. 1960; 23(1):56.
15. Zheng YP, Zhao JP, Phillips M, Liu JB, Cai MF, Sun SQ, et al. Validity and reliability of the Chinese Hamilton Depression Rating Scale. *Br J Psychiatry*. 1988; 152:660–4. PMID: [3167442](https://pubmed.ncbi.nlm.nih.gov/3167442/)
16. Folstein MF, Maiberger R, McHugh PR. Mood disorder as a specific complication of stroke. *J Neurol Neurosurg Psychiatry*. 1977; 40(10):1018–20. PMID: [591971](https://pubmed.ncbi.nlm.nih.gov/591971/)
17. Liman TG, Heuschmann PU, Endres M, Floel A, Schwab S, Kolominsky-Rabas PL. Changes in cognitive function over 3 years after first-ever stroke and predictors of cognitive impairment and long-term cognitive stability: the Erlangen Stroke Project. *Dement Geriatr Cogn Disord*. 2011; 31(4):291–9. doi: [10.1159/000327358](https://doi.org/10.1159/000327358) PMID: [21502760](https://pubmed.ncbi.nlm.nih.gov/21502760/)
18. Liman TG, Heuschmann PU, Endres M, Floel A, Schwab S, Kolominsky-Rabas PL. Impact of low minimal status on health outcome up to 5 years after stroke: the Erlangen Stroke Project. *J Neurol*. 2012; 259(6):1125–30. doi: [10.1007/s00415-011-6312-6](https://doi.org/10.1007/s00415-011-6312-6) PMID: [22109634](https://pubmed.ncbi.nlm.nih.gov/22109634/)
19. Jia Q, Zhao X, Wang C, Wang Y, Yan Y, Li H, et al. Diabetes and poor outcomes within 6 months after acute ischemic stroke: the China National Stroke Registry. *Stroke*. 2011; 42(10):2758–62 doi: [10.1161/strokeaha.111.621649](https://doi.org/10.1161/strokeaha.111.621649) PMID: [21852614](https://pubmed.ncbi.nlm.nih.gov/21852614/)
20. Uyttenboogaart M, Stewart RE, Vroomen PC, De Keyser J, Luijckx GJ. Optimizing cutoff scores for the Barthel index and the modified Rankin scale for defining outcome in acute stroke trials. *Stroke*. 2005; 36(9):1984–7. doi: [10.1161/01.str.0000177872.87960.61](https://doi.org/10.1161/01.str.0000177872.87960.61) PMID: [16081854](https://pubmed.ncbi.nlm.nih.gov/16081854/)
21. Jonsson AC, Delavaran H, Iwarsson S, Stahl A, Norrving B, Lindgren A. Functional status and patient-reported outcome 10 years after stroke: the Lund Stroke Register. *Stroke*. 2014; 45(6):1784–90. doi: [10.1161/strokeaha.114.005164](https://doi.org/10.1161/strokeaha.114.005164) PMID: [24803595](https://pubmed.ncbi.nlm.nih.gov/24803595/)
22. Hardie K, Hankey GJ, Jamrozik K, Broadhurst RJ, Anderson C. Ten-year risk of first recurrent stroke and disability after first-ever stroke in the Perth Community Stroke Study. *Stroke*. 2004; 35(3):731–5. doi: [10.1161/01.str.0000116183.50167.d9](https://doi.org/10.1161/01.str.0000116183.50167.d9) PMID: [14764929](https://pubmed.ncbi.nlm.nih.gov/14764929/)
23. Hankey GJ, Jamrozik K, Broadhurst RJ, Forbes S, Anderson CS. Long-term disability after first-ever stroke and related prognostic factors in the Perth Community Stroke Study, 1989–1990. *Stroke*. 2002; 33(4):1034–40. PMID: [11935057](https://pubmed.ncbi.nlm.nih.gov/11935057/)
24. Barker-Collo S, Feigin VL, Parag V, Lawes CM, Senior H. Auckland Stroke Outcomes Study. Part 2: Cognition and functional outcomes 5 years poststroke. *Neurology*. 2010; 75(18):1608–16. doi: [10.1212/WNL.0b013e3181fb44c8](https://doi.org/10.1212/WNL.0b013e3181fb44c8) PMID: [21041784](https://pubmed.ncbi.nlm.nih.gov/21041784/)
25. Narasimhalu K, Ang S, De Silva DA, Wong MC, Chang HM, Chia KS, et al. The prognostic effects of poststroke cognitive impairment no dementia and domain-specific cognitive impairments in nondisabled ischemic stroke patients. *Stroke*. 2011; 42(4):883–8. doi: [10.1161/strokeaha.110.594671](https://doi.org/10.1161/strokeaha.110.594671) PMID: [21330625](https://pubmed.ncbi.nlm.nih.gov/21330625/)
26. Jokinen H, Melkas S, Ylikoski R, Pohjasvaara T, Kaste M, Erkinjuntti T, et al. Post-stroke cognitive impairment is common even after successful clinical recovery. *Eur J Neurol*. 2015; 22(9):1288–94. doi: [10.1111/ene.12743](https://doi.org/10.1111/ene.12743) PMID: [26040251](https://pubmed.ncbi.nlm.nih.gov/26040251/)
27. Nys GM, van Zandvoort MJ, de Kort PL, van der Worp HB, Jansen BP, Algra A, et al. The prognostic value of domain-specific cognitive abilities in acute first-ever stroke. *Neurology*. 2005; 64(5):821–7. doi: [10.1212/01.wnl.0000152984.28420.5a](https://doi.org/10.1212/01.wnl.0000152984.28420.5a) PMID: [15753416](https://pubmed.ncbi.nlm.nih.gov/15753416/)
28. Middleton LE, Lam B, Fahmi H, Black SE, McIlroy WE, Stuss DT, et al. Frequency of domain-specific cognitive impairment in sub-acute and chronic stroke. *NeuroRehabilitation*. 2014; 34(2):305–12. doi: [10.3233/nre-131030](https://doi.org/10.3233/nre-131030) PMID: [24401826](https://pubmed.ncbi.nlm.nih.gov/24401826/)
29. Dong Y, Venketasubramanian N, Chan BP, Sharma VK, Slavin MJ, Collinson SL, et al. Brief screening tests during acute admission in patients with mild stroke are predictive of vascular cognitive impairment 3–6 months after stroke. *J Neurol Neurosurg Psychiatry*. 2012; 83(6):580–5. doi: [10.1136/jnnp-2011-302070](https://doi.org/10.1136/jnnp-2011-302070) PMID: [22496580](https://pubmed.ncbi.nlm.nih.gov/22496580/)
30. Kutlubaev MA, Hackett ML. Part II: predictors of depression after stroke and impact of depression on stroke outcome: an updated systematic review of observational studies. *Int J Stroke*. 2014; 9(8):1026–36. doi: [10.1111/ijvs.12356](https://doi.org/10.1111/ijvs.12356) PMID: [25156411](https://pubmed.ncbi.nlm.nih.gov/25156411/)

31. Ayerbe L, Ayis S, Wolfe CD, Rudd AG. Natural history, predictors and outcomes of depression after stroke: systematic review and meta-analysis. *Br J Psychiatry*. 2013; 202(1):14–21. doi: [10.1192/bjp.bp.111.107664](https://doi.org/10.1192/bjp.bp.111.107664) PMID: [23284148](https://pubmed.ncbi.nlm.nih.gov/23284148/)
32. Ayerbe L, Ayis S, Crichton S, Wolfe CD, Rudd AG. The long-term outcomes of depression up to 10 years after stroke; the South London Stroke Register. *J Neurol Neurosurg Psychiatry*. 2014; 85(5):514–21. doi: [10.1136/jnnp-2013-306448](https://doi.org/10.1136/jnnp-2013-306448) PMID: [24163430](https://pubmed.ncbi.nlm.nih.gov/24163430/)
33. Donnellan C, Hickey A, Hevey D, O'Neill D. Effect of mood symptoms on recovery one year after stroke. *Int J Geriatr Psychiatry*. 2010; 25(12):1288–95. doi: [10.1002/gps.2482](https://doi.org/10.1002/gps.2482) PMID: [21086539](https://pubmed.ncbi.nlm.nih.gov/21086539/)
34. Ayerbe L, Ayis S, Crichton S, Wolfe CD, Rudd AG. The natural history of depression up to 15 years after stroke: the South London Stroke Register. *Stroke*. 2013; 44(4):1105–10. doi: [10.1161/strokeaha.111.679340](https://doi.org/10.1161/strokeaha.111.679340) PMID: [23404719](https://pubmed.ncbi.nlm.nih.gov/23404719/)
35. Kelly-Hayes M, Beiser A, Kase CS, Scaramucci A, D'Agostino RB, Wolf PA. The influence of gender and age on disability following ischemic stroke: the Framingham study. *J Stroke Cerebrovasc Dis*. 2003; 12(3):119–26. doi: [10.1016/s1052-3057\(03\)00042-9](https://doi.org/10.1016/s1052-3057(03)00042-9) PMID: [17903915](https://pubmed.ncbi.nlm.nih.gov/17903915/)
36. Knoflach M, Matosevic B, Rucker M, Furtner M, Mair A, Wille G, et al. Functional recovery after ischemic stroke—a matter of age: data from the Austrian Stroke Unit Registry. *Neurology*. 2012; 78(4):279–85. doi: [10.1212/WNL.0b013e31824367ab](https://doi.org/10.1212/WNL.0b013e31824367ab) PMID: [22238419](https://pubmed.ncbi.nlm.nih.gov/22238419/)
37. Wu L, Wang A, Wang X, Zhao X, Wang C, Liu L, et al. Factors for short-term outcomes in patients with a minor stroke: results from China National Stroke Registry. *BMC neurology*. 2015; 15:253. doi: [10.1186/s12883-015-0505-z](https://doi.org/10.1186/s12883-015-0505-z) PMID: [26645312](https://pubmed.ncbi.nlm.nih.gov/26645312/)
38. Moon HI, Pyun SB, Tae WS, Kwon HK. Neural substrates of lower extremity motor, balance, and gait function after supratentorial stroke using voxel-based lesion symptom mapping. *Neuroradiology*. 2016; 58(7):723–31. doi: [10.1007/s00234-016-1672-3](https://doi.org/10.1007/s00234-016-1672-3) PMID: [26961307](https://pubmed.ncbi.nlm.nih.gov/26961307/)
39. Ghosal MK, Burman P, Singh V, Das S, Paul N, Ray BK, et al. Correlates of functional outcome among stroke survivors in a developing country—a prospective community-based study from India. *J Stroke Cerebrovasc Dis*. 2014; 23(10):2614–21. doi: [10.1016/j.jstrokecerebrovasdis.2014.06.001](https://doi.org/10.1016/j.jstrokecerebrovasdis.2014.06.001) PMID: [25238928](https://pubmed.ncbi.nlm.nih.gov/25238928/)
40. Wallace D, Duncan PW, Lai SM. Comparison of the responsiveness of the Barthel Index and the motor component of the Functional Independence Measure in stroke: the impact of using different methods for measuring responsiveness. *J Clin Epidemiol*. 2002; 55(9):922–8. PMID: [12393081](https://pubmed.ncbi.nlm.nih.gov/12393081/)