

1993

Environmental Analysis of the Caddo River and its Tributaries: Comparison of Water Quality During 1992 with 1974-75

Kelly L. House


Jimmy D. Bragg

Clark Kuyper

T. Kent Thomas

C. Renn Tumlison

Follow this and additional works at: <http://scholarworks.uark.edu/jaas>

 Part of the [Fresh Water Studies Commons](#), and the [Water Resource Management Commons](#)

Recommended Citation

House, Kelly L.; Bragg, Jimmy D.; Kuyper, Clark; Thomas, T. Kent; and Tumlison, C. Renn (1993) "Environmental Analysis of the Caddo River and its Tributaries: Comparison of Water Quality During 1992 with 1974-75," *Journal of the Arkansas Academy of Science*: Vol. 47, Article 16.

Available at: <http://scholarworks.uark.edu/jaas/vol47/iss1/16>

This article is available for use under the Creative Commons license: Attribution-NoDerivatives 4.0 International (CC BY-ND 4.0). Users are able to read, download, copy, print, distribute, search, link to the full texts of these articles, or use them for any other lawful purpose, without asking prior permission from the publisher or the author.

This Article is brought to you for free and open access by ScholarWorks@UARK. It has been accepted for inclusion in Journal of the Arkansas Academy of Science by an authorized editor of ScholarWorks@UARK. For more information, please contact scholar@uark.edu.

Environmental Analysis of the Caddo River and its Tributaries: Comparison of Water Quality During 1992 with 1974-75

Kelly L. House, J. D. Bragg, Clark Kuyper, T. Kent Thomas and Renn Tumblison
Garland County Community College, Hot Springs, AR, Henderson State University,
Arkadelphia, AR and Ouachita Baptist University, Arkadelphia, AR.

Abstract

Environmental data related to water quality of the Caddo River and its tributaries were collected from March - October, 1992, and compared with data from August, 1974 - May, 1975. Bacterial, chemical and physical parameters were investigated at six river locations and thirteen tributary sites. Ammonia, nitrates, soluble phosphorus, turbidity and fecal coliform were significantly lower, and sodium and potassium were significantly higher in 1992 than during the previous study. Bacterial loading exceeded EPA criteria at some locations during both studies.

Introduction

Environmental data related to water quality of the Caddo River and its tributaries were collected from March - October, 1992, and compared with a previous study from August, 1974 - May, 1975 (Nix et al., 1975). The Caddo River above DeGray Reservoir drains a portion of the southeastern flank of the Ouachita Mountains in southwest Arkansas. There are two wastewater facilities, several chicken houses, a barite mine and septic drainages located on the watershed, as well as non-point sources such as grazing livestock.

In recent years considerable concern has been shown for the quality of Arkansas waterways. In 1989 it was reported that almost one-fourth of the miles of streams in the state have impaired quality due to pollution (Ridlehoover, 1992). This study was directed toward determination of current water quality of the Caddo River system, and elucidation of any changes which have occurred in quality since the mid-1970s.

Methods and Materials

Study sites are listed in Table 1. Bacterial parameters included enumeration of total coliforms (TC), fecal coliforms (FC) and fecal streptococci (FS). Chemical and physical parameters included soluble phosphorous, ammonia, nitrates, sulfates, chloride, manganese, sodium, potassium, iron, calcium, magnesium, alkalinity, pH, conductivity and turbidity.

Samples were collected and analyzed according to standard methods (American Public Health Association, 1989). Bacterial analyses were by membrane filtration on mEndo (TC), mFC (FC) and mEnterococcus (FS) media (Difco). A 500 ml raw water sample was collected for testing pH, turbidity, alkalinity and conductivity.

Approximately 2 ml of 1:1 hydrochloric acid was added to a 250 ml sample of water for analyses of soluble phosphorus and ammonia. A 175 ml filtered sample was used for analyses of chloride, sulfates and nitrates. A 20 ml sample was acidified with 2 ml of concentrated sulfuric acid for determination of iron, calcium, sodium, potassium, magnesium and manganese.

Table 1. Location of sampling stations for the Caddo River, Arkansas

| River Stations | River Tributaries |
|----------------|------------------------|
| Black Springs | Beech Creek |
| Norman | Polk Creek |
| Caddo Gap | Lick Creek |
| Glenwood | Huddleston Creek |
| Amity | Collier Creek |
| Highway 84 | Smith Creek |
| | Gap Creek |
| | Mill Creek |
| | South Fork Caddo River |
| | Mudlick Creek |
| | Sweetwater Creek |
| | Rock Creek |
| | Caney Creek |

The data were analyzed by use of the Statistical Analysis System (SAS). Analysis of variance (ANOVA) was used to evaluate differences in physio-chemical and bacteriological parameters. When significant, Tukey-Kramer tests (Sokal and Rohlf, 1981) were used to determine which locations were different. Two-way ANOVA was used to evaluate variations due to year of sample and location.

of sample.

Results and Discussion

Data from the river sites collected during 1992 were averaged and compared to that of 1974-75 (Table 2). Ammonia, nitrates, soluble phosphorus, turbidity and fecal coliform were significantly lower, and sodium and potassium were significantly higher in 1992. Alkalinity and conductivity were significantly higher in the upper river. Previous studies have noted the presence of limestone in the upper watershed, and its absence in the lower reaches of the river (Nix et al., 1975). Therefore, dilution by the tributaries tend to occur in the lower regions of the river.

The means of the physio-chemical and bacteriological parameters measured for the entire river were compared to Environmental Protection Agency (EPA) quality criteria (EPA, 1986). The values for these parameters were within EPA criteria except for 9% of the samples of fecal coliform.

Table 2. Physio-chemical and bacteriological data from Caddo River stations. *Significant differences between years (0.05 level)

| Variable | 1992 | | 1974-75 | |
|--------------------|-------------|----------------|-------------|----------------|
| | No. samples | Mean/Std. Dev. | No. samples | Mean/Std. Dev. |
| chloride (mg/L) | 42 | 1.77/0.34 | na | na |
| pH | 42 | 7.44/0.30 | 42 | 7.38/0.19 |
| ammonia (mg/L)* | 42 | 0.06/0.03 | 42 | 0.11/0.06 |
| manganese (ug/L) | 42 | 0.07/0.02 | na | na |
| alkalinity (mg/L) | 42 | 49.0/10.3 | 42 | 45.4/10.6 |
| nitrates (mg/L)* | 41 | 0.10/0.07 | 42 | 0.17/0.18 |
| conductivity | 42 | 107.0/19.2 | 42 | 106.2/29.3 |
| phosphorus (ug/L)* | 36 | 0.02/0.008 | 42 | 0.03/0.06 |
| sodium (mg/L)* | 42 | 2.22/0.87 | 42 | 1.19/0.34 |
| sulfates (mg/L) | 42 | 5.16/0.67 | na | na |
| potassium (mg/L)* | 42 | 0.92/0.17 | 42 | 0.59/0.35 |
| iron (mg/L) | 41 | 0.59/0.19 | na | na |
| calcium (mg/L) | 42 | 16.3/4.45 | 36 | 15.19/5.14 |
| magnesium (mg/L) | 42 | 1.81/0.35 | 30 | 1.82/0.40 |
| turbidity* | 42 | 2.25/1.20 | 30 | 4.30/3.76 |
| FC (cfu/100ml)* | 47 | 58/882 | 48 | 861/1202 |
| FS (cfu/100ml) | 48 | 173/380 | na | na |
| TC (cfu/100ml) | 35 | 17312/8390 | na | na |

na = not available

The data from each specific sampling stie were grouped and compared with EPA recommendation. All chemical and physical parameters were within EPA criteria. However, fecal coliform bacteria surpassed EPA crite-

ria at Black Springs (25% of samples) and at Glenwood (13% of samples) (Table 3).

Table 3. Bacterial data from river stations. EPA criterion for Fecal Coliforms=200/100ml. *Some samples exceeded EPA criterion.

| Site | Variable | No. Samples | Mean/Standard Deviation |
|---------------|----------|-------------|-------------------------|
| Black Springs | FC | 8 | 109.125* |
| | FS | 8 | 165/108 |
| | TC | 6 | 10750/2840 |
| Norman | FC | 8 | 75/48 |
| | FS | 8 | 106/70 |
| | TC | 5 | 18650/6147 |
| Caddo Gap | FC | 7 | 43/45 |
| | FS | 8 | 209/437 |
| | TC | 6 | 24558/12902 |
| Glenwood | FC | 8 | 83/151* |
| | FS | 8 | 387/817 |
| | TC | 6 | 10200/8453 |
| Amity | FC | 8 | 19.28 |
| | FS | 8 | 52/41 |
| | TC | 6 | 14583/7150 |
| Highway 84 | FC | 8 | 21/18 |
| | FS | 8 | 115/96 |
| | TC | 6 | 15133/2988 |

Thirteen tributaries were investigated for bacterial loading (Table 4). Fecal coliform bacteria did not exceed the EPA criterion in any samples from Beech and Huddleston Creeks, but 13% of samples from Caney Creek and South Fork Caddo River and 14% of samples from Gap, Smith and Polk Creeks exceeded EPA criteria. Excessive bacteria were also present in 25% of samples from Lick and Collier Creeks, 38% of samples from Mill and Sweetwater Creeks, 63% of samples from Mudlick Creek and 86% of samples from Rock Creek. Sweetwater and Mudlick tributaries are located above the Glenwood site in the river proper and would relate to the high counts at the river station.

Chemical and physical parameters of the Caddo River and its tributaries are generally within acceptable limits. However, there appear to be excessive bacterial loading in some tributaries and around the middle reaches of the river proper. Statistically significant changes seem to have occurred since the 1970s study. However, one must exercise care in accepting such data at face value, particularly because of the impact of heavy rainfall runoff on such a small river system.

Table 4. Bacterial data from thirteen tributaries during 1992. EPA criterion for FC=200/100ml. *Some samples exceeded EPA criterion.

| Site | Variable | No. Samples | Mean/Std Dev. |
|---------------------------|----------|-------------|---------------|
| Beech | FC | 8 | 37/41 |
| | FS | 8 | 157/135 |
| | TC | 6 | 15126/6994 |
| Polk | FC | 7 | 190/348* |
| | FS | 7 | 1210/2428 |
| | TC | 4 | 29050/28049 |
| Lick | FC | 8 | 190/194* |
| | FS | 8 | 205/125 |
| | TC | 6 | 17400/9014 |
| Huddleston | FC | 8 | 36/32 |
| | FS | 8 | 336/473 |
| | TC | 6 | 13790/9833 |
| Collier | FC | 8 | 162/241* |
| | FS | 8 | 118/71 |
| | TC | 6 | 16875/5337 |
| Mill | FC | 8 | 504/723* |
| | FS | 8 | 335/292 |
| | TC | 6 | 24256/14283 |
| Smith | FC | 8 | 158/157* |
| | FS | 8 | 291/324 |
| | TC | 6 | 23006/15047 |
| Gap | FC | 7 | 102/185* |
| | FS | 8 | 199/177 |
| | TC | 6 | 7615/3689 |
| South Fork Caddo River | FC | 8 | 59/78* |
| | FS | 8 | 88/73 |
| | TC | 6 | 15325/3648 |
| Mudlick | FC | 8 | 1927/4214* |
| | FS | 8 | 143/130 |
| | TC | 6 | 26566/27306 |
| Sweetwater | FC | 8 | 320/336* |
| | FS | 8 | 458/358 |
| | TC | 6 | 19767/11344 |
| Rock | FC | 7 | 402/389* |
| | FS | 7 | 265/142 |
| | TC | 5 | 18650/5871 |
| Caney | FC | 8 | 45/76* |
| | FS | 6 | 377/424 |
| | TC | 6 | 1651/3998 |

Literature Cited

American Public Health Association. 1989. Standard methods of the examination of water and wastewater, 17th Edition. APHA, Inc. New York.

EPA. 1986. Quality Criteria for Water. Environmental Protection Agency PB-263943. Washington D.C.

Nix, J.F., R.L. Meyer, E.H. Schmitz, J.D. Bragg and R. Brown 1975. Collection of environmental data on DeGray Reservoir and the watershed of the Caddo River, Arkansas. U.S. Army Engineer Contract No. DACW39-75-C-0025.

Ridlehoover, Bobbi. 1992. State gets failing marks on many clean tests. Arkansas Democrat-Gazette. Feb. 9, p 22-23A.

Sokal, Robert R. and R. James Rohlf. 1981. Biometry. 2nd ed. W.H. Freeman and Company. 859 pp.

Acknowledgements

We would like to thank Dr. Joe Nix, Professor of Chemistry, Ouachita Baptist University for his assistance and use of his laboratory. This study was supported in part by a Henderson State University faculty research grant. Thanks are extended to Thomas House who spent many hours helping in the field.