

A PRELIMINARY LIST OF THE MYXOMYCETES OF CEDAR POINT.¹

E. L. FULLMER.

Eight species were represented by specimens in the herbarium at Cedar Point at the beginning of the present season, as follows:

Arcyria nutans (Bull.) Grev.
Dictydium cancellatum (Batsch) Macbr.
Diderma crustaceum Peck.
Lachnobolus globosus (Schw.) Rost.
Lindbladia effusa (Ehr.) Rost.
Mucilago spongiosa (Leyss.) Morg.
Physarella oblonga (B. & C.) Morg.
Trichia inconspicua Rost.

Specimens representing the following fourteen species were added to the herbarium during the summer of 1911.

Arcyria cinerea (Bull.) Pers.
Arcyria denudata (L.) Sheld.
Badhamia orbiculata Rex.
Didymium squamulosum (A. & S.) Fr.
Didymium crustaceum Fr.
Hemitrichia intorta List.
Lycogala epidendrum (Buxb) Fr.
Lycogala flavo-fuscum (Ehr.) Rost.
Ophiotheca wrightii B. & C.
Stemonitis fenestrata Rex.
Stemonitis maxima Schw.
Stemonitis smithii Macbr.
Tilmadoche alba (Bull.) Macbr.
Tubifera ferruginosa (Batsch) Macbr.

1. Presented at the meeting of the Ohio Acad. of Sci., Dec. 1, 1911.