

Oleic acid is elevated in cell membranes during rapid cold-hardening and pupal diapause in the flesh fly *Sarcophaga crassipalpis*.

Sarcophaga crassipalpis.

M. Robert Michaud and D.L. Denlinger

Department of Entomology, Ohio State University, 318 W. 12th Avenue, Columbus, OH 43210-1242, USA. michaud.11@osu.edu

Abstract


The integrity of cellular membranes is critical to the survival of insects at low temperatures, thus there is tremendous advantage conferred to insects that can adjust their composition of membrane fatty acids (FAs). Such changes, known as homeoviscous adaptation, allow cellular membranes to maintain a liquid-crystalline state at temperatures that are normally low enough to cause the membrane to enter the gel state and lose the ability to maintain homeostasis.

Flesh flies (*Sarcophaga crassipalpis*) were subjected to two experimental conditions that elicit low temperature tolerance: rapid cold-hardening and diapause. FAs were isolated and analyzed using gas chromatography-mass spectrometry. FAs changed in response to both rapid cold-hardening and diapause. In response to rapid cold-hardening, the proportion of oleic acid (18:1n-9) in pharate adults increased from 30% to 47% of the total fatty acid pool. The proportion of almost every other FA was reduced. Diapausing pupae experienced an even greater increase in oleic acid proportion to 58% of the total FA pool. Oleic acid not only increases membrane fluidity at low temperature, but also allows the cell membrane to maintain a liquid crystalline state should the temperature increase. This is the first demonstration of homeoviscous adaptation in a cold-hardy insect with a pupal diapause.


Introduction

Preventing cellular damage due to low temperatures (cold shock) is a major challenge for insects. The flesh fly, *Sarcophaga crassipalpis*, possesses two known mechanisms by which cold shock can be prevented or attenuated: by entering into a cold-hardy pupal diapause and by rapid cold-hardening (RCH). Diapause for *S. crassipalpis* is a photoperiod-induced developmental arrest that features physiological changes that contribute to low temperature survival such as upregulation of heat shock proteins and glycerol (Lee et al. 1987, Hayward et al. 2005). RCH for the flesh fly is induced by short exposures (hrs) to temperatures between 0°C and 10°C and is characterized by a three-fold increase in whole body glycerol content. Glycerol alone probably does not fully explain the protection imparted from RCH because glycerol concentrations never reach a level shown to protect proteins and membranes (Macartney et al. 1994), and other insects that possess RCH ability do not have any detectable upregulation of polyols (Kelty and Lee 2001).

Homeoviscous adaptation, the ability to alter the composition of membrane phospholipids to maintain fluidity during temperature changes (Sinensky 1974), has been linked to cold adaptation in a number of organisms, but has not been investigated extensively in insects. In the present study we investigate the hypothesis that homeoviscous adaptation in *S. crassipalpis* is involved in the RCH and diapause responses of this flesh fly by chromatographic analysis of fatty acid constituents of phospholipids as well as by thin-layer chromatographic analysis of phospholipid head groups.



Methods



Results

- Oleic acid (18:1n-9) levels increased by 17% in response to rapid cold-hardening (Fig 1A). All other fatty acids were decreased or remained the same.
- Diapause also increased oleic acid levels by 14% of the fatty acid pool (Fig. 2). All other fatty acids were reduced or remained the same. Unlike rapid cold-hardening, the change in oleic acid increased the unsaturation index of the fatty acid pool.
- Both diapause and rapid cold-hardening cause phospholipid head group changes with an increase in the proportion of phosphatidylethanolamine in relation to phosphatidylcholine (Fig. 3).




Figure 1. Change in individual fatty acids (FA) as a result of chilling consistent with flesh fly rapid cold-hardening induction. *Sarcophaga crassipalpis* pupae were subjected to 4°C for a period of 0-8 hrs and their fatty acids were analyzed with GC-MS. Each data point represents the mean ± standard error (n=5). Each point that bears an asterisk is considered statistically significant with respect to the zero-hr treatment group (one-way ANOVA with Tukey's post-test). It is clear from the above data that oleic acid (18:1n-9) is dramatically increased (17.0% total increase) between 4 and 8 hrs at the expense of many other fatty acids.




Figure 2. Change in major (A) and minor (B) fatty acid (FA) proportions due to diapause and chilling in the flesh fly. *Sarcophaga crassipalpis*. Fatty acid methyl esters from non-diapausing pupae (ND) were compared with 30 day diapausing pupae (D) and 30 day diapausing pupae that had been chilled at 4°C for 8 hrs (D+Ch). Bars represent means ± standard error (n=5). Significance for each fatty acid peak was determined through a one-way ANOVA, and significant peaks were assigned statistical groupings (a,b,c) by Tukey's post-ANOVA t-tests.





Figure 3. Change in phospholipid class due to rapid cold-hardening and diapause in the flesh fly, *Sarcophaga crassipalpis*. Phosphatidylethanolamines (PhoEtN) are upregulated at the expense of phosphatidylcholines (PhoCho) due to rapid cold-hardening (p<0.05) and diapause (p<0.01), but chilling a fly in diapause does not enhance this response beyond that induced by diapause alone. Letters above the bars (mean ± standard error, n=5) represent statistical groupings (Tukey's). D is diapause and ND is nondiapause. Chilling was carried out by exposing the fly to 4°C for 8 hrs.

Conclusions

- Our experiments provide strong evidence for adaptation in flesh fly cell membranes during diapause. This is evidenced by an increase in degree of unsaturation of fatty acids as well as phospholipid head groups. Both promote membrane fluidity at low temperatures.
- Fatty acid change due to RCH and diapause 14-17% increase in oleic acid, a monoene without double bond located in the middle of the acyl chain. A particular fatty acid possesses the unique ability to provide a wide window of fluidity rather than the shift of a narrow window to a lower temperature (Mcllhenny 1993). In this manner, the flesh fly can maintain membrane fluidity at low temperatures as well as warmer temperatures.
- Increase in the proportion of phosphatidylethanolamine at the expense of phosphatidylcholine aids homeoviscous adaptation by reducing torsional strain on cell membranes at low temperatures drop.
- The “kinks” in the acyl chains of fatty acids without double bonds promote membrane fluidity (and homeoviscous adaptation) by increasing overall molecular motion. Lowering temperatures increase molecular motion, which promotes membrane fluidity (and homeoviscous adaptation) by increasing overall molecular motion.



References

Macartney, A. B., Maresca, A.R., Cossins. 1994. Acyl-CoA desaturation: regulation of membrane lipid composition. *Temperature adaptation membranes*, ed. by [A.R. Cossins], Portland Press, London, pps. 12-22.

Kelty, J.D. and R.E. Lee, Jr. 2001. Rapid cold-hardening of *Drosophila* (Diptera: Drosophilidae) during ecologically based thermoperiodic diapause. *Experimental Biology* 204:1659-1666.

McIlhenny, R.N., 1974. The effect of alterations in the physical state of lipids on the ability of *Acholeplasma laidlawii* to grow at various temperatures. *Molecular Biology* 84:145-157.

Sinensky, M. 1974. Homeoviscous adaptation – A homeostatic process by which viscosity of membrane lipids in *Escherichia coli*. *Proceedings of the National Academy of Sciences, U.S.A.* 71:522-525.

Lee, R.E., C.P. Chen, M.H. Meacham, and D.L. Denlinger. 1987. Cold-hardiness and glycerol production in *Sarcophaga crassipalpis*. *Physiology* 33:587-592.

Hayward, S. A. L., S. C. Pavlides, S.P. Tammarillo, J.P. Rinehart. 2005. Temporal expression patterns of diapause-associated genes during the onset of diapause through post-diapause quiescence. *Journal of Insect Physiology* 51:631-640.