

## ORIGINAL ARTICLE

# The phylogenetic composition and structure of soil microbial communities shifts in response to elevated carbon dioxide

Zhili He<sup>1</sup>, Yvette Piceno<sup>2</sup>, Ye Deng<sup>1</sup>, Meiyong Xu<sup>1,3</sup>, Zhenmei Lu<sup>1,4</sup>, Todd DeSantis<sup>2</sup>, Gary Andersen<sup>2</sup>, Sarah E Hobbie<sup>5</sup>, Peter B Reich<sup>6</sup> and Jizhong Zhou<sup>1,2</sup>

<sup>1</sup>Institute for Environmental Genomics, Department of Botany and Microbiology, University of Oklahoma, Norman, OK, USA; <sup>2</sup>Ecology Department, Earth Sciences Division, Lawrence Berkeley National Laboratory, Berkeley, CA, USA; <sup>3</sup>Guangdong Provincial Key Laboratory of Microbial Culture Collection and Application, Guangdong Institute of Microbiology, Guangzhou, China; <sup>4</sup>College of Life Sciences, Zhejiang University, Hangzhou, China; <sup>5</sup>Department of Ecology, Evolution, and Behavior, St Paul, MN, USA and <sup>6</sup>Department of Forest Resources, University of Minnesota, St Paul, MN, USA

One of the major factors associated with global change is the ever-increasing concentration of atmospheric CO<sub>2</sub>. Although the stimulating effects of elevated CO<sub>2</sub> (eCO<sub>2</sub>) on plant growth and primary productivity have been established, its impacts on the diversity and function of soil microbial communities are poorly understood. In this study, phylogenetic microarrays (PhyloChip) were used to comprehensively survey the richness, composition and structure of soil microbial communities in a grassland experiment subjected to two CO<sub>2</sub> conditions (ambient, 368 p.p.m., versus elevated, 560 p.p.m.) for 10 years. The richness based on the detected number of operational taxonomic units (OTUs) significantly decreased under eCO<sub>2</sub>. PhyloChip detected 2269 OTUs derived from 45 phyla (including two from Archaea), 55 classes, 99 orders, 164 families and 190 subfamilies. Also, the signal intensity of five phyla (Crenarchaeota, Chloroflexi, OP10, OP9/JS1, Verrucomicrobia) significantly decreased at eCO<sub>2</sub>, and such significant effects of eCO<sub>2</sub> on microbial composition were also observed at the class or lower taxonomic levels for most abundant phyla, such as Proteobacteria, Firmicutes, Actinobacteria, Bacteroidetes and Acidobacteria, suggesting a shift in microbial community composition at eCO<sub>2</sub>. Additionally, statistical analyses showed that the overall taxonomic structure of soil microbial communities was altered at eCO<sub>2</sub>. Mantel tests indicated that such changes in species richness, composition and structure of soil microbial communities were closely correlated with soil and plant properties. This study provides insights into our understanding of shifts in the richness, composition and structure of soil microbial communities under eCO<sub>2</sub> and environmental factors shaping the microbial community structure.

*The ISME Journal* (2012) 6, 259–272; doi:10.1038/ismej.2011.99; published online 28 July 2011

**Subject Category:** microbial population and community ecology

**Keywords:** soil microbial community; PhyloChip; elevated CO<sub>2</sub>; microbial composition; microbial community structure

## Introduction

The concentration of atmospheric CO<sub>2</sub> has risen by approximately 36% since the mid-19th century, largely because of human activities, such as fossil fuel combustion and land use. With the current rate of increase of 1.9 p.p.m./year, it is projected to reach 700 p.p.m. by the end of this century, which may have major consequences on carbon cycling and the functioning of terrestrial ecosystems (IPCC, 2007). The stimulating effects of elevated CO<sub>2</sub> (eCO<sub>2</sub>) on

plant growth and primary productivity are well-established (Reich *et al.*, 2001; Ainsworth and Long, 2005; Luo *et al.*, 2006). For example, eCO<sub>2</sub> has been found to increase plant growth (Curtis and Wang, 1998), enhance fine root production (Hungate *et al.*, 1997) and augment soil carbon allocation (Zak *et al.*, 1993; Hu *et al.*, 2001). However, the influence of eCO<sub>2</sub> on soil microbial communities remains poorly understood and controversial (Walther *et al.*, 2002; Parmesan and Yohe, 2003; Heath *et al.*, 2005; Carney *et al.*, 2007; Drigo *et al.*, 2007, 2009, 2010; Gruber and Galloway, 2008; Heimann and Reichstein, 2008; Lesaulnier *et al.*, 2008; Austin *et al.*, 2009; Ge *et al.*, 2010; He *et al.*, 2010b). Also, the plant growth stimulation observed under eCO<sub>2</sub> may be transient (Drake *et al.*, 1997; DeLucia *et al.*, 1999) possibly because of the depletion of available nitrogen (N)

Correspondence: J Zhou, Institute for Environmental Genomics (IEG), Department of Botany and Microbiology, University of Oklahoma, Norman, OK 73019, USA.

E-mail: [jzhou@ou.edu](mailto:jzhou@ou.edu)

Received 4 January 2011; revised 18 June 2011; accepted 20 June 2011; published online 28 July 2011

(Luo *et al.*, 2004; Reich *et al.*, 2006). In addition, multiple global change factors, such as eCO<sub>2</sub>, elevated O<sub>3</sub>, warming and/or precipitation, may interact to alter soil microbial community diversity, composition, structure and function (Chung *et al.*, 2006; Castro *et al.*, 2010). Therefore, understanding the diversity, composition and structure of soil microbial communities is necessary for us to assess how eCO<sub>2</sub> modifies ecosystem properties and functional processes.

Soil may be the most complex of all microbial communities with extremely high diversity. For example, 1 g of soil contains thousands to millions of different bacterial, archaeal and eukaryotic species (Torsvik *et al.*, 2002; Gans *et al.*, 2005) interwoven in extremely complex food webs. Furthermore, most (>99%) of those microbes are as-yet uncultured (Whitman *et al.*, 1998). Thus, characterizing the phylogenetic diversity of soil microbial communities and their responses to global change (for example, eCO<sub>2</sub>) will make a significant contribution to understanding soil ecosystems.

Conventional molecular biology approaches have demonstrated that soil microbial diversity generally increased (Mitchell *et al.*, 2003; Janus *et al.*, 2005; Sonnemann and Wolters, 2005; Jossi *et al.*, 2006; Lesaulnier *et al.*, 2008), decreased (Horz *et al.*, 2004) or remained unchanged (Barnard *et al.*, 2004; Ebersberger *et al.*, 2004; Loy *et al.*, 2004; Chung *et al.*, 2006; Gruter *et al.*, 2006; Lipson *et al.*, 2006; Drigo *et al.*, 2007, 2009; Austin *et al.*, 2009; Ge *et al.*, 2010) in response to eCO<sub>2</sub>. The apparent discrepancy of microbial responses to eCO<sub>2</sub> could be partially due to real differences among various ecosystems, but could also be due to differences in the methodologies used, such as terminal restriction-fragment length polymorphism, denaturing gradient gel electrophoresis, 16S rRNA-based sequencing, enzyme activities and phospholipid fatty acids. For example, it is possible that some methods may not be sensitive enough to resolve the differences caused by eCO<sub>2</sub> at the community level.

Recently, 16S rRNA gene-based microarray technologies have been used to obtain more comprehensive information on microbial community diversity, composition, structure and dynamics. PhyloChip (G2) consists of 506 944 probe features, and of these features, 297 851 are oligonucleotide perfect match (PM) or mismatch match (MM) probes for 16S rRNA genes (Brodie *et al.*, 2006, 2007). PhyloChip has been used to detect microorganisms in a variety of environments, such as contaminated sites (Brodie *et al.*, 2006; Rastogi *et al.*, 2010), air (Brodie *et al.*, 2007), water (Hery *et al.*, 2010), soil (Cruz-Martinez *et al.*, 2009; DeAngelis *et al.*, 2009; Yergeau *et al.*, 2009; Teixeira *et al.*, 2010), microbial fuel cell (Wrighton *et al.*, 2008) and Huanglongbing pathogen-infected citrus (Sagaram *et al.*, 2009). In addition, several studies demonstrated that PhyloChip could detect many more bacterial taxa as compared with the 16S rRNA gene-based clone library approach (DeSantis *et al.*, 2007; La Duc *et al.*,

2009; Rastogi *et al.*, 2010), suggesting that PhyloChip provides more comprehensive surveys of microbial diversity, composition and structure.

The objectives of this study were to: (i) survey the richness and composition of soil microbial communities; (ii) examine the effects of eCO<sub>2</sub> on the richness, composition and structure of soil microbial communities and (iii) link soil geochemistry and plant properties with the microbial community composition and structure using PhyloChip (Brodie *et al.*, 2006, 2007). For these purposes, this study was conducted in a constructed grassland ecosystem subjected to CO<sub>2</sub> manipulation for 10 years by using the free-air CO<sub>2</sub> enrichment (FACE) technology. The results showed that eCO<sub>2</sub> significantly altered the richness, composition and structure of soil microbial communities, especially for particular microbial populations, at the operational taxonomic unit (OTU) level. Such microbial population changes were closely correlated with soil and plant properties.

## Materials and methods

The following is a summary of the methods used in this study. More detailed information is provided in Supplementary Data-A.

### Site and sampling

This study was conducted within the BioCON (Biodiversity, CO<sub>2</sub> and Nitrogen) experiment site (<http://www.biocon.umn.edu/>) located at the Cedar Creek Ecosystem Science Reserve in Minnesota, USA. The main BioCON field experiment has a total of 296 plots with three treatments: CO<sub>2</sub> (ambient (aCO<sub>2</sub>), 368 p.p.m. versus elevated, 560 p.p.m.), N (ambient versus 4 g N per m<sup>2</sup> per year) and plant diversity (1, 4, 9 or 16 species) (Reich *et al.*, 2001). In this study, soil samples from 24 plots (12 replicates from aCO<sub>2</sub>, 12 replicates from eCO<sub>2</sub> and all with 16 species and ambient N supply) were collected in July 2007 when they had been exposed to aCO<sub>2</sub> or eCO<sub>2</sub> for 10 years, and each sample was composited from five soil cores at a depth of 0–15 cm for analysis of soil properties or DNA extraction. Additional information about the BioCON experimental site, and plant groups and species, is provided in Supplementary Data-A.

### Plant and soil analyses

The aboveground and belowground biomass, plant C and N concentrations, soil pH, volumetric soil moisture, total soil C and N concentrations, and *in situ* net N mineralization and net nitrification were measured as described previously (Reich *et al.*, 2001, 2006; He *et al.*, 2010b).

### DNA extraction, purification and quantitation

Soil DNA was extracted by freeze-grinding mechanical lysis as described previously (Zhou *et al.*, 1996).

DNA quality was assessed by the ratios of 260/280 and 260/230 nm and final DNA concentrations were quantified by the PicoGreen method (Ahn *et al.*, 1996).

#### PhyloChip analysis

The second generation of the PhyloChip (PhyloTech, San Francisco, CA, USA), which has 8741 OTUs and 842 subfamilies with 297 851 probes, was used for this study (Brodie *et al.*, 2006). PhyloChip analysis included three major steps: (i) Amplification of 24 soil genomic DNAs using universal 16S rRNA primers (27F/1492R for bacteria and 4Fa/1492R for archaea); (ii) 500 ng of bacterial and 30 ng of archaeal PCR products were hybridized to each PhyloChip (Brodie *et al.*, 2006, 2007) and (iii) hybridization data were preprocessed prior to statistical analysis as detailed in Supplementary Data-A. For eight of 12 eCO<sub>2</sub> samples with less than 30 ng of archaeal PCR products, 10 µL of concentrated archaeal amplicons were used. A mixture of amplicons at known concentrations was added to each sample prior to fragmentation, which allows for standardization/normalization of PhyloChip data. Data obtained from the CEL files (produced from GeneChip Microarray Analysis Suite, version 5.1) were scaled by setting the mixture of internal standards (spike mix) mean intensity to 2500 to compensate for slight differences in probe responses on different chips. OTU reports were generated as described in Supplementary Data-A. Because setting a positive fraction (pf) cut-off can vary the number of passing OTUs (and hence affect the reported number of OTUs for each sample), several pf cut-off values (0.86, 0.88, 0.90, 0.92, 0.94) were evaluated by using the PhyloChip data analysis pipeline PhyloTrac (<http://www.phylotracer.org/Home.html>) and statistical methods as described under Materials and methods and Supplementary Data-A. A pf cut-off of 0.9 was determined to be a reasonable choice and used to generate the final OTU report used in this study.

#### Statistical analysis

Pre-processed PhyloChip data were further analyzed by different statistical methods: (i) Response ratio (Luo *et al.*, 2006); (ii) detrended correspondence analysis of the microbial community structure; (iii) analysis of similarities (Clarke, 1993), non-parametric multivariate analysis of variance (ADONIS) (Anderson, 2001) and multi-response permutation procedure (Mielke and Berry, 2001; McCune and Grace, 2002) were used to analyze differences of microbial communities by using the Binomial index (Anderson and Millar, 2004); (iv) Mantel test and canonical correspondence analysis for linking the functional structure of microbial communities to plant or soil variables; and (v) partial Mantel test and partial canonical correspondence analysis for co-variation analysis of soil and plant variables (Zhou *et al.*, 2008; He *et al.*, 2010b).

## Results

### *Effects of eCO<sub>2</sub> on plant and soil properties*

The plant productivity measured by biomass of aboveground, roots and fine roots was significantly ( $P < 0.05$ ) stimulated by eCO<sub>2</sub> (Supplementary Table S1), which is consistent with previous studies in this site (Reich *et al.*, 2001; Adair *et al.*, 2009). Also, the whole-plot total N (g m<sup>-2</sup>) and legume biomass significantly ( $P < 0.05$ ) increased at eCO<sub>2</sub>, but the percentages of nitrogen (N) in the whole-plot plant biomass, aboveground biomass and belowground biomass significantly ( $P < 0.05$ ) decreased (Supplementary Table S1). In addition, the aboveground carbon/nitrogen (C/N) ratio significantly ( $P < 0.05$ ) increased (Supplementary Table S1), probably because of an increase in plant biomass and a decrease in the aboveground N concentration. Similarly, eCO<sub>2</sub> significantly ( $P < 0.05$ ) increased soil pH and soil moisture (at depths of 0–17, 42–59 and 83–100 cm). However, no significant ( $P > 0.05$ ) changes in soil carbon, nitrogen, C/N ratio, or rates of ammonification, nitrification or net N mineralization, were observed (Supplementary Table S2). The significant differences in plant characteristics and soil properties suggest that the diversity, composition and structure of soil bacterial communities may be shifted in response to eCO<sub>2</sub>.

### *Richness of soil microbial communities in response to eCO<sub>2</sub>*

The richness of soil microbial communities was examined by PhyloChip. A total of 2269 OTUs were detected at least in three samples, accounting for 26% OTUs on the PhyloChip. An average of 1916 OTUs were detected at aCO<sub>2</sub>, which was significantly ( $P = 0.0281$ ) higher than an average of 1864 OTUs detected at eCO<sub>2</sub> (Table 1). All detected OTUs were taxonomically derived from two archaeal phyla and 43 bacterial phyla, 55 classes, 99 orders, 164 families and 190 subfamilies; most phylotypes were detected at both aCO<sub>2</sub> and eCO<sub>2</sub>, with few detected only at aCO<sub>2</sub> or eCO<sub>2</sub> (Table 2). At the phylum level, among a total of 2269 OTUs detected, 1002 OTUs were derived from Proteobacteria, a phylum with the highest number of detectable OTUs, followed by Firmicutes with 384, Actinobacteria with 289, Bacteroidetes with 162 and Acidobacteria with 76 OTUs (Table 1). Also, based on the number of OTUs detected in each phylum, two phyla had significantly ( $P < 0.05$ ) lower numbers of OTUs detected at eCO<sub>2</sub> than at aCO<sub>2</sub>, including Chloroflexi ( $P = 0.003$ ) and OP10 ( $P = 0.007$ ) (Table 1). The results indicate that the richness of soil microbial communities was decreased at eCO<sub>2</sub>.

### *Overall taxonomic composition and structure of soil microbial communities in response to eCO<sub>2</sub>*

To examine if eCO<sub>2</sub> affects the taxonomic composition and structure of soil microbial communities, detrended correspondence analysis was performed

**Table 1** Numbers of OTUs detected by PhyloChip in major phyla under aCO<sub>2</sub> and eCO<sub>2</sub> conditions

| Phylum            | OTUs on PhyloChip | No. of OTUs detected by PhyloChip |                  |                  |              |
|-------------------|-------------------|-----------------------------------|------------------|------------------|--------------|
|                   |                   | Total (%)                         | aCO <sub>2</sub> | eCO <sub>2</sub> | P (t-test)   |
| Crenarchaeota     | 79                | 12 (15.1)                         | 12.00 ± 0.00     | 12.00 ± 0.00     | 1.000        |
| Euryarchaeota     | 224               | 2 (0.9)                           | 1.00 ± 0.00      | 1.00 ± 0.00      | 1.000        |
| Acidobacteria     | 98                | 76 (77.6)                         | 67.67 ± 6.56     | 64.25 ± 5.96     | 0.195        |
| Actinobacteria    | 810               | 289 (35.6)                        | 233.17 ± 20.32   | 225.92 ± 17.24   | 0.356        |
| Bacteroidetes     | 880               | 162 (18.4)                        | 123.42 ± 14.39   | 119.67 ± 20.00   | 0.603        |
| Chlorobi          | 21                | 11 (52.4)                         | 9.42 ± 1.78      | 8.83 ± 1.59      | 0.406        |
| Chloroflexi       | 117               | 44 (37.6)                         | 37.58 ± 4.58     | 31.33 ± 4.75     | <b>0.003</b> |
| Cyanobacteria     | 202               | 51 (25.2)                         | 45.42 ± 3.37     | 43.00 ± 3.19     | 0.085        |
| Firmicutes        | 2012              | 384 (19.1)                        | 312.25 ± 30.54   | 300.75 ± 25.39   | 0.327        |
| Gemmatimonadetes  | 15                | 9 (60.0)                          | 8.67 ± 0.49      | 8.58 ± 0.51      | 0.689        |
| Natronoanaerobium | 7                 | 5 (71.4)                          | 4.00 ± 0.74      | 3.42 ± 0.79      | 0.076        |
| Nitrospira        | 29                | 8 (27.6)                          | 7.3 ± 0.98       | 6.5 ± 1.73       | 0.161        |
| OP10              | 12                | 7 (58.3)                          | 5.67 ± 1.07      | 4.58 ± 0.67      | <b>0.007</b> |
| OP9/JS1           | 12                | 5 (41.6)                          | 4.25 ± 1.14      | 3.5 ± 1.17       | 0.125        |
| Planctomycetes    | 182               | 26 (14.3)                         | 20.00 ± 4.09     | 17.25 ± 2.22     | 0.053        |
| Proteobacteria    | 3170              | 1002 (31.6)                       | 849.75 ± 70.62   | 837.58 ± 85.72   | 0.708        |
| Spirochaetes      | 150               | 36 (24.0)                         | 33.75 ± 2.93     | 32.67 ± 2.99     | 0.380        |
| Synergistes       | 19                | 5 (26.3)                          | 5.00 ± 0.00      | 5.00 ± 0.00      | 1.000        |
| TM7               | 45                | 9 (20.0)                          | 8.83 ± 0.58      | 8.25 ± 1.29      | 0.166        |
| Verrucomicrobia   | 78                | 36 (46.1)                         | 28.33 ± 3.92     | 25.75 ± 3.05     | 0.085        |
| Others (<5 OTUs)  | 250               | 53 (21.2)                         | 46.35 ± 9.37     | 37.67 ± 6.35     | 0.173        |
| Unclassified      | 329               | 37 (11.2)                         | 32.92 ± 1.83     | 31.25 ± 3.14     | 0.126        |
| Total             | 8741              | 2269 (26.0)                       | 1916.6 ± 52.03   | 1864.1 ± 57.25   | <b>0.028</b> |

Abbreviations: aCO<sub>2</sub>, ambient CO<sub>2</sub>; eCO<sub>2</sub>, elevated CO<sub>2</sub>; OTU, operational taxonomic unit. Boldface indicates significantly changed phylotypes or all detected OTUs.


**Table 2** Phylotypes detected by PhyloChip at different taxonomic levels

|   | Domain | Phylum | Class | Order | Family | Subfamily |
|---|--------|--------|-------|-------|--------|-----------|
| Total no. detected phylotypes                   | 2      | 45     | 55    | 99    | 164    | 190       |
| Shared at aCO <sub>2</sub> and eCO <sub>2</sub> | 2      | 44     | 52    | 97    | 163    | 188       |
| Only detected at aCO <sub>2</sub>               | 0      | 1      | 2     | 1     | 1      | 2         |
| Only detected at eCO <sub>2</sub>               | 0      | 0      | 1     | 1     | 1      | 0         |

for PhyloChip signal intensity data. Overall, the majority of the samples from eCO<sub>2</sub> and aCO<sub>2</sub> were distributed in different parts of the data space, although there was some overlap. Eight of 12 aCO<sub>2</sub> samples were separated well from the eCO<sub>2</sub> samples, but four other aCO<sub>2</sub> samples seemed to be clustered closer to the eCO<sub>2</sub> than the aCO<sub>2</sub> samples (Figure 1). Based on the Binomial index (Anderson and Millar, 2004), three non-parametric, multivariate statistical tests, analysis of similarities, ADONIS and multi-response permutation procedure, showed significant ( $P=0.007$ , 0.046 and 0.018, respectively) differences between microbial communities at aCO<sub>2</sub> and eCO<sub>2</sub>. The results indicated that the overall taxonomic composition and structure of soil microbial communities was altered at eCO<sub>2</sub>.

#### Relationships between microbial communities and soil and plant properties

To link the taxonomic structure of microbial communities with soil and plant properties, Mantel tests



**Figure 1** Detrended correspondence analysis of PhyloChip data for both aCO<sub>2</sub> and eCO<sub>2</sub> samples. Only OTUs (a total of 2269) detected in three or more samples out of 12 at aCO<sub>2</sub> or eCO<sub>2</sub> were analyzed.

**Table 3** Relationships of microbial phylotypes (OTUs) detected at the class level by PhyloChip to soil and plant variables revealed by partial Mantel test

| In association with: Controlling |                       |         | Soil <sup>a</sup>  |              | Plant <sup>b</sup> |              |
|----------------------------------|-----------------------|---------|--------------------|--------------|--------------------|--------------|
| Phylum                           | Class                 | OTU no. | Plant <sup>b</sup> |              | Soil <sup>a</sup>  |              |
|                                  |                       |         | r                  | P            | r                  | P            |
| All detected                     |                       | 2269    | 0.166              | 0.091        | 0.082              | 0.227        |
| Acidobacteria                    | Acidobacteria-4       | 10      | 0.075              | 0.272        | 0.283              | <b>0.046</b> |
| Bacteroidetes                    | KSA1                  | 1       | 0.083              | 0.262        | 0.230              | 0.071        |
|                                  | Unclassified          | 8       | 0.039              | 0.306        | 0.144              | 0.053        |
| Caldithrix                       | Unclassified          | 2       | 0.122              | 0.187        | 0.271              | <b>0.040</b> |
| Chlamydiae                       | Chlamydiae            | 2       | 0.069              | 0.058        | -0.003             | 0.478        |
| Chlorobi                         | Chlorobia             | 11      | 0.143              | 0.088        | 0.096              | 0.195        |
| Chloroflexi                      | Chloroflexi-3         | 2       | 0.168              | 0.106        | 0.192              | 0.069        |
|                                  | Chloroflexi-4         | 2       | 0.138              | 0.127        | 0.213              | 0.063        |
|                                  | Dehalococcoidetes     | 7       | 0.062              | 0.289        | 0.243              | 0.065        |
|                                  | Unclassified          | 6       | -0.022             | 0.495        | 0.309              | 0.059        |
| Coprothermobacteria              | Unclassified          | 1       | 0.112              | 0.196        | 0.194              | 0.07         |
| Crenarchaeota                    | C1                    | 12      | 0.077              | 0.258        | 0.455              | <b>0.012</b> |
|                                  | Thermoprotei          | 2       | 0.085              | 0.261        | 0.455              | <b>0.009</b> |
| Cyanobacteria                    | Cyanobacteria         | 48      | 0.147              | 0.145        | 0.315              | <b>0.027</b> |
| Deferribacteres                  | Deferribacter         | 1       | 0.053              | 0.098        | -0.057             | 0.919        |
| Dictyoglomi                      | Dictyoglomi           | 1       | 0.158              | 0.174        | 0.423              | 0.029        |
| DSS1                             | Unclassified          | 1       | 0.174              | 0.095        | 0.213              | 0.067        |
| Firmicutes                       | Catabacter            | 7       | 0.131              | 0.162        | 0.249              | 0.060        |
|                                  | Symbiobacteria        | 2       | 0.154              | 0.144        | 0.354              | <b>0.023</b> |
|                                  | Unclassified          | 17      | 0.124              | 0.220        | 0.245              | 0.063        |
| Lentisphaerae                    | Unclassified          | 3       | 0.178              | 0.084        | 0.212              | 0.065        |
| Marinegroup-A                    | mgA-1                 | 2       | 0.299              | 0.054        | 0.555              | <b>0.014</b> |
| OD1                              | OP11-5                | 1       | 0.172              | <b>0.038</b> | 0.111              | 0.152        |
| OP10                             | Unclassified          | 4       | 0.316              | <b>0.015</b> | 0.049              | 0.298        |
| OP3                              | Unclassified          | 3       | 0.132              | 0.076        | 0.040              | 0.282        |
| OP8                              | Unclassified          | 1       | -0.177             | 0.920        | 0.329              | <b>0.039</b> |
| OP9JS1                           | OP9                   | 2       | 0.121              | 0.138        | 0.195              | 0.064        |
| Proteobacteria                   | Gammaproteobacteria   | 330     | 0.326              | <b>0.002</b> | 0.031              | 0.357        |
|                                  | Unclassified          | 9       | 0.027              | 0.392        | 0.181              | 0.060        |
| Spirochaetes                     | Spirochaetes          | 36      | 0.092              | 0.212        | 0.257              | <b>0.045</b> |
| SR1                              | Unclassified          | 1       | 0.072              | 0.265        | 0.452              | <b>0.020</b> |
| Synergistes                      | Unclassified          | 5       | 0.085              | 0.218        | 0.260              | <b>0.030</b> |
| Thermodesulfobacteria            | Thermodesulfobacteria | 1       | 0.049              | 0.349        | 0.249              | 0.056        |
| Thermotogae                      | Thermotogae           | 1       | 0.182              | 0.078        | 0.103              | 0.184        |
| TM6                              | Unclassified          | 1       | 0.090              | 0.264        | 0.416              | <b>0.035</b> |
| TM7                              | Unclassified          | 4       | 0.161              | 0.101        | 0.227              | 0.068        |
| Unclassified                     | Unclassified          | 37      | 0.160              | 0.150        | 0.218              | 0.068        |
| WS3                              | Unclassified          | 2       | 0.055              | 0.303        | 0.166              | 0.089        |

Abbreviation: OTU, operational taxonomic unit.

<sup>a</sup>Selected soil variables include soil %N at a depth of 10–20 cm (N10–20), soil %C and N at a depth of 10–20 cm (SCN10–20), soil pH, nitrification (mg kg<sup>-1</sup> day<sup>-1</sup>) and net N mineralization (mg kg<sup>-1</sup> day<sup>-1</sup>).

<sup>b</sup>Selected plant variables include total root biomass (TRB), species count from % cover (SCFPC), aboveground percentage carbon (APC), fine roots at a depth of 0–20 cm (FR0–20) and total biomass (TB).

Only classes with *P*-values <0.10 to soil or plant variables have been listed.

Boldface indicates significantly changed phylotypes or all detected OTUs.


and canonical correspondence analysis were performed. By using the BioENV procedure (Clarke and Ainsworth, 1993), five plant variables, including total root biomass (TRB), species count from % cover (SCFPC), aboveground percentage carbon (APC), fine roots at a depth of 0–20 cm (FR0–20) and total biomass (TB), were selected from 24 plant parameters (Supplementary Table S1). Similarly, five soil variables, including soil %N at a depth of 10–20 cm (N10–20), soil C/N ratio at a depth of 10–20 cm (SCN10–20), soil pH, nitrification rate (mg kg<sup>-1</sup> day<sup>-1</sup>) and net N mineralization rate

(mg kg<sup>-1</sup> day<sup>-1</sup>) were selected from 20 soil parameters (Supplementary Table S2).

Based on the above selected sets of plant and soil variables, partial Mantel tests were initially performed to correlate the microbial community measured by the signal intensity of all detected 2269 OTUs with those environmental factors, and such an analysis showed that the microbial community on the whole was not correlated significantly with the soil variables (*P*=0.091) or plant variables (*P*=0.227) (Table 3). Then, we examined the correlations of the plant or soil properties with

specific microbial populations at different taxonomic levels (phylum, class, order, family and subfamily). At the phylum level, 14 phylotypes significantly ( $P < 0.05$ ) correlated with the soil or/and plant properties. For example, there was a significant ( $P = 0.011$ ) correlation between Crenarchaeota and the selected plant variables, and a significant ( $P = 0.037$ ) correlation between Proteobacteria and the selected soil variables (Supplementary Table S3). At the class level, 16 classes were significantly ( $P < 0.05$ ) correlated with the soil or/and plant characteristics. For example,  $\gamma$ -Proteobacteria and OP10 were significantly ( $P = 0.002$  and  $0.015$ , respectively) correlated with the selected soil variables, whereas Cyanobacteria and Spirochaetes were significantly ( $P = 0.027$  and  $0.045$ , respectively) correlated with the selected plant variables (Table 3). Also, there were significant ( $P = 0.012$  and  $0.009$ ) correlations between the microbial community and the selected plant properties for both archaeal classes, C1 and Thermoprotei, respectively (Table 3). Similarly, 48 families were detected to be correlated with the plant or soil properties (Supplementary Table S4). For example, the signal intensities of Anaplasmataceae from  $\alpha$ -Proteobacteria and Spirochaetaceae from Spirochaetes had significant ( $P = 0.019$  and  $0.046$ , respectively) correlations with the selected plant variables, and those of Enterobacteriaceae and Vibrionaceae from  $\gamma$ -Proteobacteria had significant ( $P = 0.001$  and  $0.004$ , respectively) correlations with the selected soil variables, whereas Erysipelotrichaceae from Mollicutes was significantly correlated with both soil ( $P = 0.038$ ) and plant ( $P = 0.025$ ) properties (Supplementary Table S4). In addition, five unclassified classes and 29 unclassified families were significantly ( $P < 0.05$ ) correlated with the selected soil or plant variables, respectively, suggesting that soil and plant factors may also largely shape taxonomically uncharacterized microorganisms (Table 3 and Supplementary Table S4).

Variation partition analysis (Ramette and Tiedje, 2007) was then used to assess the contribution of CO<sub>2</sub>, soil and plant properties to the taxonomic structure of microbial communities with the same selected variables (Figure 2). When the plant and soil variables were held constant, there was a significant ( $P = 0.037$ ) correlation between community structure and CO<sub>2</sub>; when plant variables and CO<sub>2</sub> were held constant, there was a significant ( $P = 0.048$ ) correlation between community structure and soil variables; and when soil variables and CO<sub>2</sub> were held constant, the plant variables did not show a significant ( $P = 0.082$ ) correlation with microbial community. The single variable CO<sub>2</sub> was able to independently explain 5.8% of the variation observed, which was the second largest contributor based on all 11 individual variables. Five soil variables could independently explain 22.1% of the variation, and five plant variables could explain 21.5% of the variation (Figure 2). Also, the interac-



**Figure 2** Variation partition analysis of the effects of CO<sub>2</sub>, soil and plant variables on the phylogenetic structure of soil microbial communities. The BioENV procedure was used to identify common sets of soil and plant variables important to the microbial community. The same sets of soil or plant variables were used for variation partition analysis and partial Mantel tests (Table 3; Supplementary Table S3 and Supplementary Table S4). The concentrations of CO<sub>2</sub> are 368 p.p.m. for ambient and 560 p.p.m. for elevated environments; soil variables included soil %N at a depth of 10–20 cm (N10–20), soil C and N ratio at a depth of 10–20 cm (SCN10–20), soil pH, nitrification rate (mg kg<sup>-1</sup> day<sup>-1</sup>) and net N mineralization rate (mg kg<sup>-1</sup> day<sup>-1</sup>); plant variables included total root biomass (TRB), species count from % cover (SCFPC), aboveground percentage carbon (APC), fine roots at a depth of 0–20 cm (FR0–20) and total biomass (TB).

tions between CO<sub>2</sub> and soil variables, CO<sub>2</sub> and plant variables, and soil and plant variables, and among CO<sub>2</sub>, soil and plant variables, were 0.1%, 2.4%, 0.0% and 2.8%, respectively (Figure 2). In addition, 45.3% of the variation remained unexplained (Figure 2). The above statistical analyses suggest that CO<sub>2</sub> has a direct effect on the microbial community, and that both soil and plant properties are almost equally important for shaping microbial communities through indirect CO<sub>2</sub> effects in this grassland ecosystem.

#### Significantly changed and unique OTUs

To examine effects of eCO<sub>2</sub> on microbial community composition, both significantly changed and unique OTUs were identified. Among 2269 OTUs detected, 2075 were shared by aCO<sub>2</sub> and eCO<sub>2</sub> samples, and 194 unique OTUs were only detected at aCO<sub>2</sub> (123) or eCO<sub>2</sub> (71), respectively. For those shared OTUs, a response ratio was calculated for each OTU based on its signal intensity. A total of 194 OTUs were significantly ( $P < 0.05$ ) decreased and only 13 were significantly ( $P < 0.05$ ) increased at eCO<sub>2</sub> (Table 4). Most phyla, including relatively abundant ones (for example, Crenarchaeota, Acidobacteria, Chloroflexi, Planctomycetes, Verrucomicrobia), did not have OTUs detected with increased signal intensities at eCO<sub>2</sub>, and only a few phyla (for example, Actinobacteria, Bacteroidetes, Firmicutes, Proteobacteria) had OTUs with both increased and decreased signal intensities at eCO<sub>2</sub> (Table 4), which is consistent with the general trend that the richness decreased at

**Table 4** Numbers of shared OTUs detected at both aCO<sub>2</sub> and eCO<sub>2</sub> based on the hybridization signal intensity and unique OTUs detected only at aCO<sub>2</sub> or eCO<sub>2</sub>

| Phylum            | Shared OTUs               |                           | Unique OTUs      |                  |
|-------------------|---------------------------|---------------------------|------------------|------------------|
|                   | Decrease-eCO <sub>2</sub> | Increase-eCO <sub>2</sub> | aCO <sub>2</sub> | eCO <sub>2</sub> |
| Crenarchaeota     | 12                        | 0                         | 0                | 0                |
| Euryarchaeota     | 1                         | 1                         | 1                | 1                |
| Acidobacteria     | 15                        | 0                         | 3                | 2                |
| Actinobacteria    | 10                        | 1                         | 26               | 17               |
| Bacteroidetes     | 4                         | 2                         | 8                | 9                |
| Chlorobi          | 1                         | 0                         | 0                | 0                |
| Chloroflexi       | 11                        | 0                         | 5                | 0                |
| Cyanobacteria     | 4                         | 1                         | 4                | 0                |
| Firmicutes        | 29                        | 2                         | 25               | 11               |
| Gemmatimonadetes  | 2                         | 0                         | 0                | 0                |
| Natronoanaerobium | 0                         | 0                         | 1                | 0                |
| Nitrospira        | 4                         | 0                         | 0                | 0                |
| OP10              | 2                         | 0                         | 2                | 0                |
| OP9/JS1           | 3                         | 0                         | 0                | 0                |
| Planctomycetes    | 9                         | 0                         | 6                | 1                |
| Proteobacteria    | 55                        | 6                         | 33               | 29               |
| Spirochaetes      | 2                         | 0                         | 0                | 0                |
| Synergistes       | 1                         | 0                         | 0                | 0                |
| TM7               | 0                         | 0                         | 0                | 0                |
| Verrucomicrobia   | 16                        | 0                         | 4                | 1                |
| Others (<5 OTUs)  | 6                         | 0                         | 3                | 0                |
| Unclassified      | 7                         | 0                         | 2                | 0                |
| Total             | 194                       | 13                        | 123              | 71               |

Abbreviations: aCO<sub>2</sub>, ambient CO<sub>2</sub>; eCO<sub>2</sub>, elevated CO<sub>2</sub>; OTU, operational taxonomic unit.

Increase-CO<sub>2</sub> or Decrease-eCO<sub>2</sub> indicates the signal intensity of an OTU was significantly higher or lower at eCO<sub>2</sub>, respectively.


eCO<sub>2</sub>. The signal intensities of 6 and 56 OTUs were significantly ( $P < 0.05$ ) increased and decreased, respectively, in Proteobacteria. Specifically, two OTUs were increased and 12 decreased in  $\alpha$ -Proteobacteria, 2 and 2 in  $\beta$ -Proteobacteria, 1 and 2 in  $\epsilon$ -Proteobacteria, and 1 and 7 in  $\gamma$ -Proteobacteria, respectively, but all 31 OTUs derived from  $\delta$ -Proteobacteria were decreased at eCO<sub>2</sub> (Figure 3). Similarly, 29 and 2 (OTU3497 and OTU3254) OTUs were significantly ( $P < 0.05$ ) decreased and increased, respectively, in Firmicutes, which were mostly derived from two classes, Clostridia and Bacilli (Supplementary Figure S1). Among 194 unique OTUs, 123 and 71 were from aCO<sub>2</sub> and eCO<sub>2</sub>, respectively, and those OTUs were largely derived from the most abundant phyla, such as Proteobacteria, Firmicutes and Actinobacteria (Table 4 and Supplementary Table S5). The analysis of significantly changed and unique OTUs further confirms that the phylogenetic composition of soil microbial communities changed in response to eCO<sub>2</sub>.

#### Significantly changed microbial populations at eCO<sub>2</sub>

To understand what specific microbial populations may be affected by eCO<sub>2</sub>, we mapped OTUs detected to microbial populations at the phylum or lower levels, and significantly changed populations were identified by response ratio based on PhyloChip hybridization signal intensity. At the phylum level,

five phyla, including one archaeal phylum (Crenarchaeota) and four bacterial phyla (Chloroflexi, OP10, OP9/JS1, Verrucomicrobia), showed significantly ( $P < 0.05$ ) decreased signal intensities, but most abundant phyla (for example, Proteobacteria, Firmicutes, Actinobacteria, Bacteroidetes, Acidobacteria) remained unchanged at eCO<sub>2</sub> (Figure 4). A further examination of those significantly changed phyla showed that those changes occurred in some specific microbial groups at the class or lower levels. In the phylum of Chloroflexi, a significant decrease of signal intensities at eCO<sub>2</sub> was observed in three orders (Chloroflexi-1a, 1b, 1f) of the class Anaerolineae, and the class of Dehalococcoidetes, whereas the signal intensities of other classes (for example, Chloroflexi-3, Chloroflexi-4, Thermomicrobia) did not change significantly (Supplementary Figure S2A). In the phylum of Verrucomicrobia, all significant changes appeared to occur in the order of Verrucomicrobiales from the class of Verrucomicrobiae, in which three families (Verrucomicrobiaceae, Verrucomicrobia subdivision-3, Verrucomicrobia subdivision-7) and an unclassified phylotype had significantly ( $P < 0.01$ ) decreased signal intensities, although the other two families (Verrucomicrobia subdivision-5 and Xiphinematobacteraceae) detected did not show significant changes in signal intensity at eCO<sub>2</sub> (Supplementary Figure S2B). Crenarchaeota is an archaeal phylum showing significantly ( $P < 0.01$ ) decreased signal intensities at eCO<sub>2</sub>, and such decreases were observed in three orders (C1a, C1b, Cenarchaeales) from two classes (C1 and Thermoprotei) (Supplementary Figure S3A). In addition, significant decreases of signal intensity were seen in two less characterized phyla (OP10 and OP9/JS1), with one from an unclassified class in OP10 (Supplementary Figure S3B) and the other from an unclassified order of JS1 class in OP9/JS1 (Supplementary Figure S3C).

Although significant changes were not observed at eCO<sub>2</sub> for the most abundant phyla at the phylum level based on summed intensities, such significances were detected at the class or lower taxonomic levels for some phyla. In the phylum Proteobacteria, the signal intensity of the AMD clone order of  $\delta$ -Proteobacteria was significantly ( $P < 0.05$ ) decreased although no significant changes were detected at the class level (Supplementary Figure S4). In the phylum Firmicutes, the signal intensities of the family Syntrophomonadaceae in the order of Clostridiales and an unclassified order in the class of Clostridia significantly ( $P < 0.05$ ) decreased at eCO<sub>2</sub>, as did an unclassified phylotype ( $P < 0.01$ ), although no significant changes were observed in other phylotypes (for example, Bacilli, Mollicutes) (Supplementary Figure S5). Also, three groups of Actinobacteria showed significantly decreased signal intensities at eCO<sub>2</sub>, which included the order Acidimicrobiaceae and an unclassified phylotype in the class Acidimicrobiales; the order Bifidobacteriaceae in the class Bifidobacteriales and




**Figure 3** Significantly changed OTUs in the phylum of Proteobacteria at eCO<sub>2</sub> by using the response ratio method (Luo *et al.*, 2006) at 95% confidence interval.

an unclassified phylotype at the phylum level, and interestingly, no significant changes in signal intensity were observed in the largest class Actinomycetales (Supplementary Figure S6). In addition, no significant changes were observed in the phylum of Bacteroidetes (Supplementary Figure S7) or Acidobacteria even at the class, order or family level (Supplementary Figure S8). Those results indicate that eCO<sub>2</sub> significantly affected some specific microbial populations at different taxonomic levels, such as phylum, class, order and family, and those phylotypes generally appeared to have decreased signal intensities at eCO<sub>2</sub>.

## Discussion

The long-term sustainability of ecosystem productivity requires detailed knowledge of its biodiversity coupled to profound understanding of its functioning. To better understand the implications of eCO<sub>2</sub> on microbial communities, we used PhyloChip to comprehensively survey the richness, composition and structure of soil microbial communities in the BioCON grassland. Our results showed that eCO<sub>2</sub> significantly altered the microbial community diversity, composition and structure, especially for particular microbial populations at the OTU level.





**Figure 4** Average PhyloChip hybridization signal intensities for aCO<sub>2</sub> and eCO<sub>2</sub> samples at the phylum level. Significance was tested by response ratios (Luo *et al.*, 2006). \*\*\* $P < 0.01$ ; \*\* $P < 0.05$ .

Such microbial population changes were significantly correlated with soil and plant properties.

This study provides a comprehensive survey of the microbial richness and composition of grassland soil microbial communities. Previous studies with 16S rRNA-based analyses using clone libraries (Janssen, 2006; Lesaulnier *et al.*, 2008), microarrays (for example, PhyloChip) (Cruz-Martinez *et al.*, 2009; DeAngelis *et al.*, 2009; Yergeau *et al.*, 2009), pyrosequencing (Roesch *et al.*, 2007; Fulthorpe *et al.*, 2008; Campbell *et al.*, 2010; Eilers *et al.*, 2010; Uroz *et al.*, 2010) and other approaches (Drigo *et al.*, 2007, 2008, 2009, 2010; Feng *et al.*, 2009) show that soil microbial communities are highly diverse and complex. In this study, 2269 OTUs affiliated 45 phyla, 55 classes, 99 orders, 164 families and 190 subfamilies were detected. Proteobacteria was the most well-represented phylum, with  $\alpha$ -,  $\beta$ -,  $\delta$ -,  $\gamma$ - and  $\epsilon$ -Proteobacteria, as well as unclassified classes, detected. This group of bacteria has considerable morphological, physiological and metabolic diversity, which are of great importance to global carbon, nitrogen and sulfur cycling (Kersters *et al.*, 2006). Firmicutes were detected as the second most prevalent phylum in terms of the number of OTUs, and Bacilli, Clostridia and Mollicutes were found to be major classes in this grassland ecosystem. Some previous cloning analyses may underestimate this group as cells or spores are known to be difficult to lyse during DNA extraction (Janssen, 2006). The members of the Actinobacteria phylum are a group of Gram-positive bacteria that have an important role in organic matter turnover and carbon cycling, such as decomposition of cellulose and chitin, and Actinomycetales and Acidimicrobiales were found to be major classes in the BioCON site. A previous study showed that long-term organic and inorganic amendments significantly

altered the Actinobacterial community structure but not its diversity (Piao *et al.*, 2008). Bacteroidetes are the fourth most prevalent group of bacteria detected in this study, with three major classes (Bacteroidetes, Flavobacteria, Sphingobacteria). Acidobacteria are among the most dominant phyla in soil-borne microbial communities, and generally are classified into eight classes (Handelsman, 2004). The diversity of Acidobacteria in soil was recently examined using different approaches, and a higher proportion of Acidobacteria was observed in bulk soil than in rhizosphere soil (Kielak *et al.*, 2008). Therefore, this study provides a comprehensive survey of the richness and composition of soil microbial communities at this grassland ecosystem.

Elevated atmospheric CO<sub>2</sub> may affect soil microbial communities in both direct and indirect ways. In a previous study, soil CO<sub>2</sub> flux increased 0.57 mmol m<sup>-2</sup> s<sup>-1</sup> or 16% on average at eCO<sub>2</sub> conditions as compared with aCO<sub>2</sub> conditions in the BioCON site (Craine *et al.*, 2001). Also, a recent study in the same site showed that the abundance of key genes involved in microbial C and N fixation, and labile C degradation, was significantly increased at eCO<sub>2</sub> (He *et al.*, 2010b). Those results suggest that eCO<sub>2</sub> may directly impact soil microbial community structure and function. However, as CO<sub>2</sub> concentrations in the pore space of soil generally are between 2000 and 38 000 p.p.m., much higher than those in the atmosphere even under aCO<sub>2</sub> condition, the direct effects of eCO<sub>2</sub> on soil microbial communities may be negligible compared with potential indirect effects, such as increased plant carbon inputs to soil and changes in soil properties (Drigo *et al.*, 2008). The data presented here reflect this idea: CO<sub>2</sub> alone explained 5.8% of the total variation of microbial community structure, compared with soil variables at 22.1% and plant variables at 21.5%.

Also, eCO<sub>2</sub> significantly increased plant productivity, whole-plot total N, soil pH and soil moisture, and decreased whole-plot plant N, aboveground and belowground N concentrations. The results suggest eCO<sub>2</sub> may directly and indirectly affect soil microbial communities, and the indirect effects appear to make more of a contribution to shaping the soil microbial communities.

Effects of eCO<sub>2</sub> on plant and soil properties are expected to modify taxonomic microbial community composition and structure, and regulate ecosystem functioning. First, it is indicated by differential responses of soil microbial populations to eCO<sub>2</sub>. Increases in soil carbon, coupled to an increase in cellulolytic and chitinolytic activities, were noted to alter the availability of soil substrates for microbial metabolism (Larson *et al.*, 2002; Phillips *et al.*, 2002). Previous studies showed increases in the abundance of Actinobacteria and Bacteroidetes at eCO<sub>2</sub> (Sait *et al.*, 2006; Lesaulnier *et al.*, 2008). However, we did not see significant changes in the total signal intensities for Actinobacteria, Bacteroidetes or other most abundant phyla at the phylum level in response to eCO<sub>2</sub>. In  $\alpha$ -Proteobacteria, a significant increase in the abundance of OTUs related to Rhodobium and a significant decrease in the abundance of OTUs related to Bradyrhizobium were detected in the trembling aspen FACE study (Lesaulnier *et al.*, 2008), and a recent study showed a stimulation of purple phototrophic  $\alpha$ - and  $\beta$ -Proteobacteria in a flooded paddy soil by eCO<sub>2</sub> (Feng *et al.*, 2009), which are generally consistent with this study. Also, the preference of Acidobacteria in bulk soil has been suggested to be a result of the oligotrophic lifestyle for many members of this phylum (Fierer *et al.*, 2007). A higher input of organic matter into soil at eCO<sub>2</sub> may be favorable for carbon polymer-degrading or fast-growing microorganisms, which presumably outcompete Acidobacteria. If true, the signal intensity of Acidobacteria may remain unchanged or decrease at eCO<sub>2</sub>, which was observed in this study. In addition, a decrease in the abundance of Crenarchaea and Verrucomicrobia at eCO<sub>2</sub> was observed previously (Lesaulnier *et al.*, 2008), which is consistent with our observation in this study. Members of Verrucomicrobia are reported to be negatively impacted by soil moisture (Buckley and Schmidt, 2001), which has been shown to increase at eCO<sub>2</sub> in the BioCON site (Reich *et al.*, 2001; He *et al.*, 2010b) and other sites (Zavaleta *et al.*, 2003), which is largely due to reduced stomatal conductance of plants (Kandeler *et al.*, 2008). Second, significant correlations are observed between environmental factors and microbial communities. A negative impact of increased concentrations of organic matter on the growth of Acidobacteria was observed previously (Stevenson *et al.*, 2004), and consistently, Acidobacteria-4 populations were found to be correlated significantly with plant variables in this study. Also, more carbon input into soil may affect autotrophic

populations, such as Chloroflexis and Cyanobacteria. Indeed, a decrease in the signal intensity of those phyla was observed in this study. pH has been considered an important factor affecting the diversity and structure of soil microbial communities (Fierer and Jackson, 2006), and our Mantel analysis showed a significant correlation between soil properties, including pH and  $\gamma$ -Proteobacteria or OP10. In this study, soil pH was  $\sim 6.2$  in aCO<sub>2</sub> plots and it significantly increased to  $\sim 6.5$  in eCO<sub>2</sub> plots, suggesting a possible shift of microbial community composition and structure. Therefore, our results indicate that both soil and plant properties, such as soil pH, moisture and plant biomass, significantly affect the microbial richness, composition and structure, which may determine or modify ecosystem functioning.

The central hypothesis of this study was that, at eCO<sub>2</sub>, an increase in plant biomass (Reich *et al.*, 2006) and soil carbon inputs (Adair *et al.*, 2009), and associated microenvironmental changes (Reich, 2009; He *et al.*, 2010b), would stimulate microbial, especially bacterial growth, which would lead to significant changes in the richness, composition, structure and function of soil microbial communities. Previous studies of effects of eCO<sub>2</sub> on soil microbial communities showed variable responses. For example, a study conducted at a trembling aspen FACE experiment site in Wisconsin, USA, showed an increase in heterotrophic decomposers and a decrease in nitrate reducers of the domain bacteria and archaea, although the total bacterial abundance did not change (Lesaulnier *et al.*, 2008). By contrast, no detectable effects on microbial community structure, microbial activity, potential soil N mineralization or nitrification rates were observed at a sweetgum FACE experiment in Tennessee, USA (Austin *et al.*, 2009). For testing our core hypothesis, this study had several strengths: (i) It was conducted at a well-designed BioCON experimental site, with 12 replicates for each CO<sub>2</sub> condition, so that the effects of eCO<sub>2</sub> on soil microbial communities could be robustly examined; (ii) PhyloChip is considered a powerful tool for a comprehensive survey of microbial richness and composition (DeAngelis *et al.*, 2009; Rastogi *et al.*, 2010), which may overcome the limitations of cloning-based approaches and (iii) this study was conducted in a grassland ecosystem with defined plant species, which minimizes the effects of plant diversity and composition on soil microbial communities. Indeed, consistently with some previous studies in grasslands (Schortemeyer *et al.*, 1996; Drissner *et al.*, 2007), our results indicated that eCO<sub>2</sub> had significant effects on the richness, composition and structure of soil microbial communities. First, such changes are reflected in a decrease in richness, which generally agrees with previous studies (Begon *et al.*, 1996; Hughes *et al.*, 2001). Similarly, a significant decrease in richness was observed in some phyla (for example, Chloroflexi, OP10). Second, more OTUs were found

to have decreased signal intensities and fewer OTUs with increased signals at eCO<sub>2</sub> among shared OTUs, and more unique OTUs were detected at aCO<sub>2</sub> than eCO<sub>2</sub>. Third, based on PhyloChip signal intensities, although some specific microbial populations, especially from the most abundant phyla, remained unchanged at the phylum level, significant changes were apparent at the class or lower levels, suggesting the microbial composition was altered at eCO<sub>2</sub> at finer taxonomic scales. Finally, the taxonomic structure is different between aCO<sub>2</sub> and eCO<sub>2</sub> samples as shown by statistical analyses (for example, detrended correspondence analysis, analysis of similarities, ADONIS, multi-response permutation procedure). Recently, a study using a comprehensive functional gene array, GeoChip 3.0 (He *et al.*, 2010a), also demonstrated that the functional composition and structure of soil microbial communities were significantly altered at eCO<sub>2</sub> (He *et al.*, 2010b), which may be due to eCO<sub>2</sub>-induced shifts in microbial populations. The results suggest that the richness, composition and structure of soil microbial communities shift in response to eCO<sub>2</sub>.

PhyloChip has been considered a powerful tool to comprehensively and rapidly analyze microbial communities. Specifically, as such a microarray-based technology has a defined probe set and targets known populations, it minimizes or eliminates sampling artifacts, including under-sampling, unequal sampling and random sampling (Zhou *et al.*, 2008), making it a preferable approach for community-scale comparison of microbial communities, as has been demonstrated in this study. Like other high-throughput technologies, however, PhyloChip has its limitations. For example, PhyloChip only detects known sequences already present in a database at the time of probe design, so the G2 PhyloChip used in this study may not fully cover the species richness of soil microbial communities, and a follow-up study using the G3 PhyloChip could prove beneficial. To discover unknown 16S rRNA genes, future investigations may use high-quality, full-length sequencing as a complementary approach to further understand the taxonomic and phylogenetic diversity, composition, structure and function of the soil microbial communities in this grassland ecosystem.

## Acknowledgements

This work was supported by the United States Department of Agriculture (Project 2007-35319-18305) through the NSF-USDA Microbial Observatories Program; the National Science Foundation under DEB-0716587, DEB-0620652, DEB-0322057, DEB-0080382, DEB-0218039, DEB-0219104, DEB-0217631, DEB-0716587, LTREB DEB-0716587 and LTER DEB-9411972 projects; the DOE Program for Ecosystem Research; the Minnesota Environment and Natural Resources Trust Fund (DE-FG96ER2291) and the DOE under contract DE-AC02-05CH11231 through the

University of California and Lawrence Berkeley National Laboratory; and by NIH Grant U01-HG004866.

## References

- Adair EC, Reich P, Hobbie S, Knops J. (2009). Interactive effects of time, CO<sub>2</sub>, N, and diversity on total below-ground carbon allocation and ecosystem carbon storage in a grassland community. *Ecosystems* **12**: 1037–1052.
- Ahn S, Costa J, Emanuel J. (1996). PicoGreen quantitation of DNA: effective evaluation of samples pre- or post-PCR. *Nucleic Acids Res* **24**: 2623–2625.
- Ainsworth EA, Long SP. (2005). What have we learned from 15 years of free-air CO enrichment (FACE)? A meta-analytic review of the responses of photosynthesis, canopy properties and plant production to rising CO<sub>2</sub>. *New Phytol* **165**: 351–372.
- Anderson MJ. (2001). A new method for non-parametric multivariate analysis of variance. *Aust Ecol* **26**: 32–46.
- Anderson MJ, Millar RB. (2004). Spatial variation and effects of habitat on temperate reef fish assemblages in northeastern New Zealand. *J Exp Marine Biol Ecol* **305**: 191–221.
- Austin EE, Castro HF, Sides KE, Schadt CW, Classen AT. (2009). Assessment of 10 years of CO<sub>2</sub> fumigation on soil microbial communities and function in a sweet-gum plantation. *Soil Biol Biochem* **41**: 514–520.
- Barnard R, Barthes L, Le Roux X, Leadley PW. (2004). Dynamics of nitrifying activities, denitrifying activities and nitrogen in grassland mesocosms as altered by elevated CO<sub>2</sub>. *New Phytol* **162**: 365–376.
- Begon M, Harper JL, Townsend CR. (1996). *Ecology: Individuals Populations and Communities*. Blackwell Publishing Ltd: Cambridge, MA, USA.
- Brodie EL, DeSantis TZ, Joyner DC, Baek SM, Larsen JT, Andersen GL *et al.* (2006). Application of a high-density oligonucleotide microarray approach to study bacterial population dynamics during uranium reduction and reoxidation. *Appl Environ Microbiol* **72**: 6288–6298.
- Brodie EL, DeSantis TZ, Parker JP, Zubietta IX, Piceno YM, Andersen GL. (2007). Urban aerosols harbor diverse and dynamic bacterial populations. *Proc Natl Acad Sci USA* **104**: 299–304.
- Buckley DH, Schmidt TM. (2001). Environmental factors influencing the distribution of rRNA from Verrucomicrobia in soil. *FEMS Microbiol Ecol* **35**: 105–112.
- Campbell BJ, Polson SW, Hanson TE, Mack MC, Schuur EAG. (2010). The effect of nutrient deposition on bacterial communities in Arctic tundra soil. *Environ Microbiol* **12**: 1842–1854.
- Carney MC, Hungate BA, Drake BG, Megonigal JP. (2007). Altered soil microbial community at elevated CO<sub>2</sub> leads to loss of soil carbon. *Proc Natl Acad Sci USA* **104**: 4990–4995.
- Castro HF, Classen AT, Austin EE, Norby RJ, Schadt CW. (2010). Soil microbial community responses to multiple experimental climate change drivers. *Appl Environ Microbiol* **76**: 999–1007.
- Chung H, Zak D, Lilleskov E. (2006). Fungal community composition and metabolism under elevated CO<sub>2</sub> and O<sub>3</sub>. *Oecologia* **147**: 143–154.
- Clarke KR. (1993). Nonparametric multivariate analyses of changes in community structure. *Aust J Ecol* **18**: 117–143.

- Clarke KR, Ainsworth M. (1993). A method of linking multivariate community structure to environmental variables. *Mar Ecol Prog Ser* **92**: 205–219.
- Craine JM, Wedin DA, Reich PB. (2001). Grassland species effects on soil CO<sub>2</sub> flux track the effects of elevated CO<sub>2</sub> and nitrogen. *New Phytol* **150**: 425–434.
- Cruz-Martinez K, Suttle KB, Brodie EL, Power ME, Andersen GL, Banfield JF. (2009). Despite strong seasonal responses, soil microbial consortia are more resilient to long-term changes in rainfall than overlying grassland. *ISME J* **3**: 738–744.
- Curtis PS, Wang X. (1998). A meta-analysis of elevated CO<sub>2</sub> effects on woody plant mass, form, and physiology. *Oecologia* **113**: 299–313.
- DeAngelis KM, Brodie EL, DeSantis TZ, Andersen GL, Lindow SE, Firestone MK. (2009). Selective progressive response of soil microbial community to wild oat roots. *ISME J* **3**: 168–178.
- DeLucia EH, Hamilton JG, Naidu SL, Thomas RB, Andrews JA, Finzi A et al. (1999). Net primary production of a forest ecosystem with experimental CO<sub>2</sub> enrichment. *Science* **284**: 1177–1179.
- DeSantis T, Brodie E, Moberg J, Zubieta I, Piceno Y, Andersen G. (2007). High-density universal 16S rRNA microarray analysis reveals broader diversity than typical clone library when sampling the environment. *Microb Ecol* **53**: 371–383.
- Drake BG, González-Meler MA, Long SP. (1997). MORE EFFICIENT PLANTS: a consequence of rising atmospheric CO<sub>2</sub>? *Annu Rev Plant Physiol Plant Mol Biol* **48**: 609–639.
- Drigo B, Kowalchuk G, van Veen J. (2008). Climate change goes underground: effects of elevated atmospheric CO<sub>2</sub> on microbial community structure and activities in the rhizosphere. *Biol Fertil Soils* **44**: 667–679.
- Drigo B, Kowalchuk GA, Yergeau E, Bezemer TM, Boschker HTS, Van Veen JA. (2007). Impact of elevated carbon dioxide on the rhizosphere communities of *Carex arenaria* and *Festuca rubra*. *Glob Change Biol* **13**: 2396–2410.
- Drigo B, Pijl AS, Duyts H, Kielak AM, Gamper HA, Houtekamer MJ et al. (2010). Shifting carbon flow from roots into associated microbial communities in response to elevated atmospheric CO<sub>2</sub>. *Proc Natl Acad Sci USA* **107**: 10938–10947.
- Drigo B, van Veen JA, Kowalchuk GA. (2009). Specific rhizosphere bacterial and fungal groups respond differently to elevated atmospheric CO<sub>2</sub>. *ISME J* **3**: 1204–1217.
- Drissner D, Blum H, Tscherko D, Kandeler E. (2007). Nine years of enriched CO<sub>2</sub> changes the function and structural diversity of soil microorganisms in a grassland. *Eur J Soil Sci* **58**: 260–269.
- Ebersberger D, Werrnbtter N, Niklaus PA, Kandeler E. (2004). Effects of long term CO<sub>2</sub> enrichment on microbial community structure in calcareous grassland. *Plant Soil* **264**: 313–323.
- Eilers KG, Lauber CL, Knight R, Fierer N. (2010). Shifts in bacterial community structure associated with inputs of low molecular weight carbon compounds to soil. *Soil Biol Biochem* **42**: 896–903.
- Feng Y, Lin X, Wang Y, Zhang J, Mao T, Yin R et al. (2009). Free-air CO<sub>2</sub> enrichment (FACE) enhances the biodiversity of purple phototrophic bacteria in flooded paddy soil. *Plant Soil* **324**: 317–328.
- Fierer N, Bradford MA, Jackson RB. (2007). Toward an ecological classification of soil bacteria. *Ecology* **88**: 1354–1364.
- Fierer N, Jackson RB. (2006). The diversity and biogeography of soil bacterial communities. *Proc Natl Acad Sci USA* **103**: 626–631.
- Fulthorpe RR, Roesch LFW, Riva A, Triplett EW. (2008). Distantly sampled soils carry few species in common. *ISME J* **2**: 901–910.
- Gans J, Wolinsky M, Dunbar J. (2005). Computational improvements reveal great bacterial diversity and high metal toxicity in soil. *Science* **309**: 1387–1390.
- Ge Y, Chen C, Xu Z, Oren R, He J-Z. (2010). The spatial factor, rather than elevated CO<sub>2</sub>, controls the soil bacterial community in a temperate forest ecosystem. *Appl Environ Microbiol* **76**: 7429–7436.
- Gruber N, Galloway JN. (2008). An Earth-system perspective of the global nitrogen cycle. *Nature* **451**: 293–296.
- Gruter D, Schmid B, Brandl H. (2006). Influence of plant diversity and elevated atmospheric carbon dioxide levels on belowground bacterial diversity. *BMC Microbiol* **6**: 68.
- Handelsman J. (2004). Metagenomics: application of genomics to uncultured microorganisms. *Microbiol Mol Biol Rev* **68**: 669–685.
- He Z, Deng Y, Van Nostrand JD, Tu Q, Xu M, Hemme CL et al. (2010a). GeoChip 3.0 as a high-throughput tool for analyzing microbial community composition, structure and functional activity. *ISME J* **4**: 1167–1179.
- He Z, Xu M, Deng Y, Kang S, Kellogg L, Wu L et al. (2010b). Metagenomic analysis reveals a marked divergence in the structure of belowground microbial communities at elevated CO<sub>2</sub>. *Ecol Lett* **13**: 564–575.
- Heath J, Ayres E, Possell M, Bardgett RD, Black HJ, Grant H et al. (2005). Rising atmospheric CO<sub>2</sub> reduces sequestration of root-derived soil carbon. *Science* **309**: 1711–1713.
- Heimann M, Reichstein M. (2008). Terrestrial ecosystem carbon dynamics and climate feedbacks. *Nature* **451**: 289–292.
- Hery M, Sanguin H, Perez Fabiel S, Lefebvre X, Vogel TM, Paul E et al. (2010). Monitoring of bacterial communities during low temperature thermal treatment of activated sludge combining DNA phylochip and respirometry techniques. *Water Res* **44**: 6133–6143.
- Horz H-P, Barbrook A, Field CB, Bohannan BJM. (2004). Ammonia-oxidizing bacteria respond to multifactorial global change. *Proc Natl Acad Sci USA* **101**: 15136–15141.
- Hu S, Chapin FS, Firestone MK, Field CB, Chiariello NR. (2001). Nitrogen limitation of microbial decomposition in a grassland under elevated CO<sub>2</sub>. *Nature* **409**: 188–191.
- Hughes JB, Hellmann JJ, Ricketts TH, Bohannan BJM. (2001). Counting the uncountable: statistical approaches to estimating microbial diversity. *Appl Environ Microbiol* **67**: 4399–4406.
- Hungate BA, Holland EA, Jackson RB, Chapin FS, Mooney HA, Field CB. (1997). The fate of carbon in grasslands under carbon dioxide enrichment. *Nature* **388**: 576–579.
- IPCC (2007). *Global Climate Projections*. Cambridge University Press: Cambridge, United Kingdom, p 806.
- Janssen PH. (2006). Identifying the dominant soil bacterial taxa in libraries of 16S rRNA and 16S rRNA genes. *Appl Environ Microbiol* **72**: 1719–1728.
- Janus LR, Angeloni NL, McCormack J, Rier ST, Tuchman NC, Kelly JJ. (2005). Elevated atmospheric CO<sub>2</sub> alters soil microbial communities associated with trembling aspen (*Populus tremuloides*) roots. *Microb Ecol* **50**: 102–109.

- Jossi M, Fromin N, Tarnawski S, Kohler F, Gillet F, Aragno M *et al.* (2006). How elevated pCO<sub>2</sub> modifies total and metabolically active bacterial communities in the rhizosphere of two perennial grasses grown under field conditions. *FEMS Microbiol Ecol* **55**: 339–350.
- Kandeler E, Mosier AR, Morgan JA, Milchunas DG, King JY, Rudolph S *et al.* (2008). Transient elevation of carbon dioxide modifies the microbial community composition in a semi-arid grassland. *Soil Biol Biochem* **40**: 162–171.
- Kerstens K, De Vos P, Gillis M, Swings J, Vandamme P, Stackebrandt E. (2006). Introduction to the Proteobacteria. In: Dworkin M, Falkow S, Rosenberg E, Schleifer K-H, Stackebrandt E (eds) *The Prokaryotes*, 3rd edn Springer: New York, 5: 3–37.
- Kielak A, Pijl AS, van Veen JA, Kowalchuk GA. (2008). Phylogenetic diversity of Acidobacteria in a former agricultural soil. *ISME J* **3**: 378–382.
- La Duc MT, Osman S, Vaishampayan P, Piceno Y, Andersen G, Spry JA *et al.* (2009). Comprehensive census of bacteria in clean rooms by using DNA microarray and cloning methods. *Appl Environ Microbiol* **75**: 6559–6567.
- Larson JL, Zak DR, Sinsabaugh RL. (2002). Extracellular enzyme activity beneath temperate trees growing under elevated carbon dioxide and ozone. *Soil Sci Soc Am J* **66**: 1848–1856.
- Lesaulnier C, Papamichail D, McCorkle S, Ollivier B, Skiena S, Taghavi S *et al.* (2008). Elevated atmospheric CO<sub>2</sub> affects soil microbial diversity associated with trembling aspen. *Environ Microbiol* **10**: 926–941.
- Lipson DA, Blair M, Barron-Gafford G, Grieve K, Murthy R. (2006). Relationships between microbial community structure and soil processes under elevated atmospheric carbon dioxide. *Microb Ecol* **51**: 302–314.
- Loy A, Kusel K, Lehner A, Drake HL, Wagner M. (2004). Microarray and functional gene analyses of sulfate-reducing prokaryotes in low-sulfate, acidic fens reveal cooccurrence of recognized genera and novel lineages. *Appl Environ Microbiol* **70**: 6998–7009.
- Luo Y, Hui D, Zhang D. (2006). Elevated CO<sub>2</sub> stimulates net accumulations of carbon and nitrogen in land ecosystems: a meta-analysis. *Ecology* **87**: 53–63.
- Luo Y, Su B, Currie WS, Dukes JS, Finzi A, Hartwig U *et al.* (2004). Progressive nitrogen limitation of ecosystem responses to rising atmospheric carbon dioxide. *BioScience* **54**: 731–739.
- McCune B, Grace JB. (2002). *Analysis of Ecological Communities*. MJM Software Design: Gleneden Beach, OR.
- Mielke PW, Berry KJ. (2001). *Permutation Methods: A Distance Function Approach*, 2nd edn. Springer-Verlag: New York, NY.
- Mitchell E, Gilbert D, Buttler A, Amblard C, Grosvernier P, Gobat J. (2003). *Structure of Microbial Communities in Sphagnum Peatlands and Effect of Atmospheric Carbon dioxide Enrichment*, vol. 46. Springer-Verlag: New York, LLC, pp 187–199.
- Parmesan C, Yohe G. (2003). A globally coherent fingerprint of climate change impacts across natural systems. *Nature* **421**: 37–42.
- Phillips RL, Zak DR, Holmes WE, White DC. (2002). Microbial community composition and function beneath temperate trees exposed to elevated atmospheric carbon dioxide and ozone. *Oecologia* **131**: 236–244.
- Piao Z, Yang L, Zhao L, Yin S. (2008). Actinobacterial community structure in soils receiving long-term organic and inorganic amendments. *Appl Environ Microbiol* **74**: 526–530.
- Ramette A, Tiedje JM. (2007). Multiscale responses of microbial life to spatial distance and environmental heterogeneity in a patchy ecosystem. *Proc Natl Acad Sci USA* **104**: 2761–2766.
- Rastogi G, Osman S, Vaishampayan P, Andersen G, Stetler L, Sani R. (2010). Microbial diversity in uranium mining-impacted soils as revealed by high-density 16S microarray and clone library. *Microb Ecol* **59**: 94–108.
- Reich PB. (2009). Elevated CO<sub>2</sub> reduces losses of plant diversity caused by nitrogen deposition. *Science* **326**: 1399–1402.
- Reich PB, Hobbie SE, Lee T, Ellsworth DS, West JB, Tilman D *et al.* (2006). Nitrogen limitation constrains sustainability of ecosystem response to CO<sub>2</sub>. *Nature* **440**: 922–925.
- Reich PB, Knops J, Tilman D, Craine J, Ellsworth D, Tjoelker M *et al.* (2001). Plant diversity enhances ecosystem responses to elevated CO<sub>2</sub> and nitrogen deposition. *Nature* **410**: 809–812.
- Roesch LF, Fulthorpe RR, Riva A, Casella G, Hadwin AK, Kent AD *et al.* (2007). Pyrosequencing enumerates and contrasts soil microbial diversity. *ISME J* **1**: 283–290.
- Sagaram US, DeAngelis KM, Trivedi P, Andersen GL, Lu S-E, Wang N. (2009). Bacterial diversity analysis of Huanglongbing pathogen-infected citrus using PhyloChip arrays and 16S rRNA gene clone library sequencing. *Appl Environ Microbiol* **75**: 1566–1574.
- Sait M, Davis KER, Janssen PH. (2006). Effect of pH on isolation and distribution of members of subdivision 1 of the phylum Acidobacteria occurring in soil. *Appl Environ Microbiol* **72**: 1852–1857.
- Schortemeyer M, Hartwig UA, Hendrey GR, Sadowsky MJ. (1996). Microbial community changes in the rhizospheres of white clover and perennial ryegrass exposed to Free Air Carbon dioxide Enrichment (FACE). *Soil Biol Biochem* **28**: 1717–1724.
- Sonnemann I, Wolters V. (2005). The microfood web of grassland soils responds to a moderate increase in atmospheric CO<sub>2</sub>. *Glob Change Biol* **11**: 1148–1155.
- Stevenson BS, Eichorst SA, Wertz JT, Schmidt TM, Breznak JA. (2004). New strategies for cultivation and detection of previously uncultured microbes. *Appl Environ Microbiol* **70**: 4748–4755.
- Teixeira LCRS, Peixoto RS, Cury JC, Sul WJ, Pellizari VH, Tiedje J *et al.* (2010). Bacterial diversity in rhizosphere soil from Antarctic vascular plants of Admiralty Bay, maritime Antarctica. *ISME J* **4**: 989–1001.
- Torsvik V, Ovreaas L, Thingstad TF. (2002). Prokaryotic diversity—magnitude, dynamics, and controlling factors. *Science* **296**: 1064–1066.
- Uroz S, Buée M, Murat C, Frey-Klett P, Martin F. (2010). Pyrosequencing reveals a contrasted bacterial diversity between oak rhizosphere and surrounding soil. *Environ Microbiol Rep* **2**: 281–288.
- Walther G-R, Post E, Convey P, Menzel A, Parmesan C, Beebee TJC *et al.* (2002). Ecological responses to recent climate change. *Nature* **416**: 389–395.
- Whitman WB, Coleman DC, Wiebe WJ. (1998). Prokaryotes: the unseen majority. *Proc Natl Acad Sci USA* **95**: 6578–6583.
- Wrighton KC, Agbo P, Warnecke F, Weber KA, Brodie EL, DeSantis TZ *et al.* (2008). A novel ecological role of the Firmicutes identified in thermophilic microbial fuel cells. *ISME J* **2**: 1146–1156.

- Yergeau E, Schoondermark-Stolk SA, Brodie EL, Dejean S, DeSantis TZ, Goncalves O *et al.* (2009). Environmental microarray analyses of Antarctic soil microbial communities. *ISME J* **3**: 340–351.
- Zak D, Pregitzer K, Curtis P, Teeri J, Fogel R, Randlett D. (1993). Elevated atmospheric CO<sub>2</sub> and feedback between carbon and nitrogen cycles. *Plant Soil* **151**: 105–117.
- Zavaleta ES, Thomas BD, Chiariello NR, Asner GP, Shaw MR, Field CB. (2003). Plants reverse warming effect on ecosystem water balance. *Proc Natl Acad Sci USA* **100**: 9892–9893.
- Zhou J, Kang S, Schadt CW, Garten CT, Jr. (2008). Spatial scaling of functional gene diversity across various microbial taxa. *Proc Natl Acad Sci USA* **105**: 7768–7773.
- Zhou J, Bruns MA, Tiedje JM. (1996). DNA recovery from soils of diverse composition. *Appl Environ Microbiol* **62**: 316–322.

Supplementary Information accompanies the paper on The ISME Journal website (<http://www.nature.com/ismej>)