



University of Pennsylvania
ScholarlyCommons

Center for Human Modeling and Simulation

Department of Computer & Information Science

1-2008

Empirical developments in retraction

B K. Redman
Wayne State University

H N. Yarandi
Wayne State University

Jon F. Merz
University of Pennsylvania, merz@mail.med.upenn.edu

Follow this and additional works at: <http://repository.upenn.edu/hms>

Recommended Citation

Redman, B. K., Yarandi, H. N., & Merz, J. F. (2008). Empirical developments in retraction. Retrieved from <http://repository.upenn.edu/hms/113>

Reprinted from:

Empirical developments in retraction. B K Redman, H N Yarandi, and J F Merz. *J Med Ethics* 2008; 34: 807-809.

DOI:10.1136/jme.2007.023069

URL: <http://jme.bmj.com/cgi/content/abstract/34/11/807>

This paper is posted at ScholarlyCommons. <http://repository.upenn.edu/hms/113>

For more information, please contact libraryrepository@pobox.upenn.edu.

Empirical developments in retraction

Abstract

This study provides current data on key questions about retraction of scientific articles. Findings confirm that the rate of retractions remains low but is increasing. The most commonly cited reason for retraction was research error or inability to reproduce results; the rate from research misconduct is an underestimate, since some retractions necessitated by research misconduct were reported as being due to inability to reproduce. Retraction by parties other than authors is increasing, especially for research misconduct. Although retractions are on average occurring sooner after publication than in the past, citation analysis shows that they are not being recognised by subsequent users of the work. Findings suggest that editors and institutional officials are taking more responsibility for correcting the scientific record but that reasons published in the retraction notice are not always reliable. More aggressive means of notification to the scientific community appear to be necessary.

Comments

Reprinted from:

Empirical developments in retraction. B K Redman, H N Yarandi, and J F Merz. *J Med Ethics* 2008; 34: 807-809.

DOI:10.1136/jme.2007.023069

URL: <http://jme.bmj.com/cgi/content/abstract/34/11/807>



Empirical developments in retraction

B K Redman, H N Yarandi and J F Merz

J. Med. Ethics 2008;34:807-809
doi:10.1136/jme.2007.023069

Updated information and services can be found at:
<http://jme.bmj.com/cgi/content/full/34/11/807>

These include:

References

This article cites 7 articles, 5 of which can be accessed free at:
<http://jme.bmj.com/cgi/content/full/34/11/807#BIBL>

Rapid responses

You can respond to this article at:
<http://jme.bmj.com/cgi/eletter-submit/34/11/807>

Email alerting service

Receive free email alerts when new articles cite this article - sign up in the box at the top right corner of the article

Notes

To order reprints of this article go to:
<http://journals.bmj.com/cgi/reprintform>

To subscribe to *Journal of Medical Ethics* go to:
<http://journals.bmj.com/subscriptions/>

Empirical developments in retraction

B K Redman,^{1,2} H N Yarandi,² J F Merz¹

¹ Department of Medical Ethics, University of Pennsylvania, Philadelphia, Pennsylvania, USA; ² College of Nursing, Wayne State University, Detroit, Michigan, USA

Correspondence to: Barbara K Redman, College of Nursing, Wayne State University, 5557 Cass Avenue, Detroit, Michigan 48202, USA; B.Redman@wayne.edu

Received 12 September 2007
Revised 10 December 2007
Accepted 9 January 2008

ABSTRACT

This study provides current data on key questions about retraction of scientific articles. Findings confirm that the rate of retractions remains low but is increasing. The most commonly cited reason for retraction was research error or inability to reproduce results; the rate from research misconduct is an underestimate, since some retractions necessitated by research misconduct were reported as being due to inability to reproduce. Retraction by parties other than authors is increasing, especially for research misconduct. Although retractions are on average occurring sooner after publication than in the past, citation analysis shows that they are not being recognised by subsequent users of the work. Findings suggest that editors and institutional officials are taking more responsibility for correcting the scientific record but that reasons published in the retraction notice are not always reliable. More aggressive means of notification to the scientific community appear to be necessary.

Previous studies of retraction of scientific articles have shown that the dominant reason for retraction is research error or inability to reproduce results (generally referred to as “inability to reproduce”).^{1,2} Citation continues long after retraction, raising questions about the adequacy of current methods of notification.³ Here we update these findings. We report changes in rate and agent of retraction and in length of time between publication and retraction and examine the role of publicity on post-retraction citation rate.

METHODS

The National Library of Medicine’s PubMed database identifies retracted articles. Retractions for articles published in the decade 1995–2004 were retrieved on 9 June 2005 using publication type “retraction in”. Each case was coded to indicate who retracted the publication, the portion (part or whole) retracted, the reason for retraction and the length of time in months between publication and retraction. Ten per cent of the records were coded independently, yielding a 3% disagreement, which was negotiated to agreement.

The impact factor for the journal in which the article was published and retracted was obtained from *Journal Citation Reports* (2003).⁴ A journal’s impact factor is the number of citations in the current year to papers published in the previous 2 years, divided by the total number published in the same 2 years.⁵ Those with the highest impact factor are considered the most prestigious and their papers are likely to be widely cited, making it important that retractions be noted. PubMed citations for the years 1995–2004 to the search terms “retraction”, “research misconduct”, “research fraud” and “plagiarism” reported in the

news sections of *Science* and *Nature* and matched to cases in our data set were retrieved as proxy measures of publicity to the scientific community.

Citation activity was obtained from *Science Citation Index* (SCI) and *Social Science Citation Index* (SSCI), through the Web of Science (<http://portal.isiknowledge.com/portal.cgi?Init=Yes&SID=E2ni9cFg43Mik3akg6g>). In addition to citation activity before and after each retraction, a subset of all cases in the biomedical sciences in which the full article was retracted for reasons of confirmed data fabrication/falsification, research error or inability to reproduce was examined. Those articles cited at least 10 times 4 years or more after retraction (that is, with high late post-retraction citation rates) were analysed. In these cases, each citation was checked for whether it acknowledged the retraction. Descriptive statistics, analysis of frequency, regression and correlation analysis were used to analyse the data.

RESULTS

The number of records in PubMed for the study period was 5 041 587; of these, 328 (0.0065%) were retracted. We analysed the 315 cases (96%) that were in English. As shown in table 1, the largest proportion of retractions were by all authors, followed by some authors and by editors/publishers. The primary reasons for retraction were research error, inability to reproduce, research misconduct and plagiarism. Usually (in 90% of cases) the whole article was retracted. In the span of 10 years, the 315 retracted articles cumulatively were cited 3942 times before retraction and 4501 times after retraction.

Table 2 presents months between publication and retraction, citations before and after retraction and journal impact factor. Thirty-two cases of retraction (10%) resulted in publicity in the news sections of *Science* or *Nature*, most frequently reporting on cases of research misconduct.

Compared with an earlier study of retractions for the years 1966–1994,¹ the present study found (1) a significantly higher rate of retraction (0.0021% versus 0.0065%, $p < 0.0001$ by test of proportions), (2) a decrease in the mean time from publication to retraction (from 28 months to 21 months), and (3) a decrease in the proportion of retractions initiated by authors (from 81% to 67%) (see table 1), suggesting that others have become involved in oversight.

Continued citation of invalid, retracted work, already documented in 1990³ and in 1998,¹ remains a problem, even though retractions are now clearly identified in the Medline and PubMed databases.⁶ Our findings show that the studies highly cited before retraction remained highly cited post-retraction ($r = 0.60$, $p < 0.0001$), with those in

Research ethics

Table 1 Agents of retraction of scientific articles, and reasons for retraction

Variable	Budd's study ¹	Present study
	(n = 235)	(n = 315)
Agent of retraction (per cent)		
Missing	–	4
All authors	81	48
Some authors		19
Editor/publisher	19	17
Other*	–	12
Primary reason for retraction (per cent)		
Research error	39	22
Inability to reproduce	16	20
Plagiarism	–	17
Research misconduct	37†	17
No reason given	–	12
Other	9%	11

– Category not used.

*Attorney, editor, sponsor, institutional official often with author, or not clear.

†Includes presumed misconduct and may include plagiarism.

higher-impact journals more highly cited after retraction ($r = 0.61$, $p < 0.0001$). Differences in the average number of citations before ($F = 5.39$, $p < 0.0001$) and after retraction ($F = 3.02$, $p = 0.0113$) among the six reasons for retraction (table 3) were statistically significant, with significant differences between research misconduct and plagiarism ($t = 2.90$, $p = 0.0040$) and between inability to reproduce and plagiarism ($t = 2.50$, $p = 0.0131$) after retraction. The average number of citations of manuscripts whose retractions had publicity (29.38) was significantly larger than of those with no publicity (14.27) ($t = 2.98$, $p = 0.0031$), reflecting the high impact factor of the journals (*Science*, *Nature*) in which publication was tracked.

There were 10 papers with high late post-retraction citation rates and these were cited a total of 225 times 4 or more years after retraction (range 10–96). In nine of these cases, the rate of acknowledgement of the retraction in these late citations was less than 3%. The paper with 96 late post-retraction citations dealt with a supposedly settled controversy with significant environmental implications⁷ and was immediately questioned by other scientists; the retraction of this paper was acknowledged in 29% of the post-retraction citations.

DISCUSSION AND CONCLUSIONS

An early study by Parrish noted that retraction notices often obfuscated the reasons for retraction, particularly in cases of research misconduct⁸. Things haven't changed. We matched retractions required by findings of confirmed data fabrication and/or falsification by the US Office of Research Integrity during the time frame of this study (1995–2004). Twenty-six such retractions were found (26/315), involving 17 authors. Twelve retraction notices straightforwardly cited data fabrication or falsification as the reason for retraction. The rest (14)

Table 3 Citations before and after retraction, and journal impact factor (JIF), according to reasons for retraction

Reason for retraction	Mean number of citations		Mean JIF
	Before retraction	After retraction	
Research error	13.7	15.8	11.68
Inability to reproduce	10.7	20.60	11.11
Plagiarism	2.9	5.6	6.77
Research misconduct	38.3	23.6	11.82
No reason given	3.6	7.7	2.11
Other	3.3	10.8	8.13

gave as reasons “unable to reproduce”, “data are invalid”, “for complicated reasons”, “not supportable by reproducible evidence” or “appears to have been falsified” (without identifying which author was at fault), with the remainder offering no reason for the retraction. Since other agencies also sanction authors for research misconduct, a more thorough study of this issue would extend beyond the Office of Research Integrity.

In summary, rates of retraction, although very low, have increased between the periods 1966–1997 and 1995–2004, and retractions are occurring more quickly and are more frequently initiated by parties other than authors. The post-retraction citation rate remains high, especially associated with retraction for reasons of research misconduct and inability to reproduce. Some⁹ suggest that journals should require authors to attest that they have checked their manuscripts' reference list against the National Library of Medicine (NLM) master list of retracted articles, perhaps by way of a web-based program.

These findings raise implications for the scientific community, institutions, professional associations and journals. International standards are clear that the text of the retraction should explain why the article is being retracted;¹⁰ these standards are in some instances not followed. A commitment of journals to publish retraction notices for a period of time (instead of just once or not at all) would be likely to better inform the scientific community that the findings from those papers should not be relied upon. Professional associations should strongly support these efforts through their codes of conduct and through policies of the journals they sponsor. While some error is inevitable, the public's trust that error is being minimised is important to justify their continued investment in science.

Further study of the impact of continued use of retracted data on end users such as patients (who may be harmed) and on the use of research resources (which could be wasted) as well as on downstream research that depends on upstream research will provide a perspective on costs of the current system relied upon for correction of the scientific record. Asking those who cite retracted articles why they continue to cite and testing new mechanisms for effective notification of the scientific community will provide options for improving on current practices. Better understanding of the myriad kinds of research error and

Table 2 Summary measures of the retraction variables

Variable	Mean	SD	Median	Quartile		Range
				Lower	Upper	
Months between publication and retraction	20.75	17.73	15.5	8.0	28.0	0–102.0
Number of citations						
Before retraction	13.83	41.49	1.0	0.0	7.0	0–343.0
After retraction	15.81	26.23	7.0	2.0	18.0	0–270.0
Journal impact factor	10.36	11.09	4.61	2.0	15.7	0.4–34.8

inability to reproduce will be likely to help further define best scientific practices.

Competing interests: None declared.

REFERENCES

1. **Budd JM**, Sievert M, Schultz TR. Phenomena of retraction: reasons for retraction and citations to the publications. *JAMA* 1998;**280**:296–7.
2. **Nath SB**, Marcus SC, Druss BG. Retractions in the research literature: misconduct or mistakes? *Med J Aust* 2006;**185**:152–4.
3. **Pfeifer MP**, Snodgrass GL. The continued use of retracted, invalid scientific literature. *JAMA* 1990;**263**:1420–3.
4. *Journal Citation Reports*. Philadelphia: Thomson ISI, 2003.
5. **Garfield E**. The history and meaning of the journal impact factor. *JAMA* 2006;**295**:90–3.
6. **Anon**. *What's the difference between MEDLINE and PubMed?* [fact sheet]. Bethesda, Maryland: United States National Library of Medicine, 2006. http://www.nlm.nih.gov/pubs/factsheets/dif_med_pub.html (accessed 23 July 2008).
7. **Arnold SF**, Klotz DM, Collins BM, *et al*. Synergistic activation of estrogen receptor with combination of environmental chemicals. *Science* 1996;**272**:1489–92.
8. **Parrish DM**. Scientific misconduct and correcting the scientific literature. *Acad Med* 1999;**74**:221–30.
9. **Sox HC**, Rennie D. Research misconduct, retraction and cleansing the medical literature: lessons from the Poehlman case. *Ann Intern Med* 2006;**144**:609–13.
10. **International Committee of Medical Journal Editors**. Uniform requirements for manuscripts submitted to biomedical journals: writing and editing for biomedical publication. Philadelphia: ICMJE, 2007. <http://www.icmje.org/index.html#authorsur> (accessed 23 July 2008).