



4-11-1977

Critical Behavior of Random Resistor Networks

A. Brooks Harris

University of Pennsylvania, harris@sas.upenn.edu

Ronald Fisch

University of Pennsylvania

Follow this and additional works at: http://repository.upenn.edu/physics_papers

 Part of the [Physics Commons](#)

Recommended Citation

Harris, A., & Fisch, R. (1977). Critical Behavior of Random Resistor Networks. *Physical Review Letters*, 38 (15), 796-799.
<http://dx.doi.org/10.1103/PhysRevLett.38.796>

This paper is posted at ScholarlyCommons. http://repository.upenn.edu/physics_papers/314
For more information, please contact repository@pobox.upenn.edu.

Critical Behavior of Random Resistor Networks

Abstract

We present numerical data and scaling theories for the critical behavior of random resistor networks near the percolation threshold. We determine the critical exponents of a suitably defined resistance correlation function by a Padé analysis of low-concentration expansions as a function of dimensionality. We verify that $d=6$ is the critical dimensionality for the onset of mean-field behavior. We use the coherent-potential approximation to construct a mean-field scaling function for the critical region.

Disciplines

Physics

model in the scaling limit [see T. T. Wu, B. M. McCoy, C. A. Tracy, and E. Barouch, Phys. Rev. B **13**, 316 (1976), B. M. McCoy, C. A. Tracy, and T. T. Wu, J. Math. Phys. (to be published), and in *Statistical Mechanics and Statistical Methods in Theory and Application*, edited by U. Landman (Plenum, New York, 1977, to be published); E. Barouch, B. M. McCoy, and T. T. Wu, Phys. Rev. Lett. **31**, 1409 (1973); C. A. Tracy and B. M. McCoy, Phys. Rev. Lett. **31**, 1500 (1973); R. Z. Bariev, Phys. Lett. **55A**, 456 (1976); E. Barouch, B. M. McCoy, C. A. Tracy, and T. T. Wu, Phys. Lett. **57A**, 111 (1976)]. Some of these papers are A. Luther and V. J. Emery, Phys. Rev. Lett. **33**, 589 (1974); S. Coleman, Phys. Rev. D **11**, 2088 (1975); S. Mandelstam, Phys. Rev. D **11**, 3026 (1975); R. F. Dashen, B. Hasslacher, and A. Neveu, Phys. Rev. D **11**, 3424 (1975); A. Luther, Phys. Rev. B **14**, 2153 (1976); J. Fröhlich and E. Seiler, Helv. Phys. Acta **49**, 889 (1976); H. Lehmann and J. Stehr, to be published.

³J. Schwinger, Proc. Nat. Acad. Sci. U. S. A. **44**, 956 (1958); E. Nelson, J. Funct. Anal. **12**, 97 (1973); K. Osterwalder and R. Schrader, Commun. Math. Phys. **31**, 83 (1973), and **42**, 281 (1975).

⁴The free energy was computed by L. Onsager, Phys. Rev. **65**, 117 (1944). The spontaneous magnetization was computed by C. N. Yang, Phys. Rev. **85**, 808 (1952). For a presentation of all results through 1973 see B. M. McCoy and T. T. Wu, *The Two-Dimensional Ising Model* (Harvard Univ. Press, Cambridge, Mass., 1973).

⁵E. W. Montroll, R. B. Potts, and J. C. Ward, J. Math. Phys. (N.Y.) **4**, 308 (1963).

⁶T. T. Wu, Phys. Rev. **149**, 380 (1966); H. Cheng and T. T. Wu, Phys. Rev. **164**, 719 (1967).

⁷Wu *et al.*, Ref. 2; McCoy, Tracy, and Wu, Ref. 2; Barouch, McCoy, and Wu, Ref. 2; Tracy and McCoy, Ref. 2; Bariev, Ref. 2; Barouch *et al.*, Ref. 2.

⁸The leading nontrivial term for $n=3$, $T < T_c$, was first given by R. Z. Bariev, Physica (Utrecht) **83A**, 388 (1976), provided that suitable interpretation is given to his singular integrals.

⁹D. A. Pink, Can. J. Phys. **46**, 2399 (1968), and Phys. Rev. **188**, 1032 (1969); L. P. Kadanoff, Phys. Rev. **188**, 859 (1969); L. P. Kadanoff and H. Ceva, Phys. Rev. B **3**, 3918 (1971); Helen Au-Yang, Phys. Rev. B **15**, 2704 (1971).

Critical Behavior of Random Resistor Networks*

A. B. Harris and R. Fisch

Department of Physics, University of Pennsylvania, Philadelphia, Pennsylvania 19104

(Received 31 January 1977)

We present numerical data and scaling theories for the critical behavior of random resistor networks near the percolation threshold. We determine the critical exponents of a suitably defined resistance correlation function by a Padé analysis of low-concentration expansions as a function of dimensionality. We verify that $d=6$ is the critical dimensionality for the onset of mean-field behavior. We use the coherent-potential approximation to construct a mean-field scaling function for the critical region.

In this Letter we report some new ideas concerning the properties of random resistor networks near the percolation threshold.¹ The model we treat is that of an electrical network on a d -dimensional hypercubic lattice of L^d sites with conductances σ_{ij} connecting nearest neighboring pairs of lattice sites i and j . Each σ_{ij} is an independent random variable assuming the values $\sigma_<$ or $\sigma_>$ with respective probabilities $1-p$ and p . The macroscopic conductivity, Σ , is then defined to be the configurational average of σL^{2-d} , where $\sigma \equiv I/V$, where I is the current when the potential difference V is applied between two opposite ($d-1$)-dimensional faces of the hypercube. We may define clusters as being groups of sites which are connected with respect to the conductances $\sigma_>$. The statistics of cluster size and the associated pair connectedness correlation length, $\xi(p)$, were shown² to be related to the thermodynamics of the

s -state Potts model in the limit $s \rightarrow 1$, if the identification $p = 1 - e^{-J/kT}$ is made, where J is the coupling constant for nearest-neighbor interaction in the Potts model. This relation indicates that the usual exponent description for phase transitions can be applied to the percolation threshold and that the various scaling relations and universality predictions can be expected to hold as well. It was later shown^{3,4} that for $d > d_c = 6$, mean-field theory gives correct values for cluster statistics near the percolation threshold: $\alpha = -1$, $\beta = 1$, $\gamma = 1$, and $\nu = \frac{1}{2}$. In view of scaling arguments which relate the resistor network and percolation problems, de Gennes⁵ has suggested that $d_c = 6$. Here we present numerical evidence which confirms that this suggestion is correct. We also discuss several new scaling relations.

A way to determine d_c without using the renormalization group (RG) is to analyze the high-tem-

perature expansion for the susceptibility, χ , as a function of dimensionality, d .⁶ At $d=4$ an analysis for the Ising model⁷ based on a single power-law divergence yields $\gamma=1.09$. However, the RG predicts⁸ corrections to scaling of the form

$$\chi = A[1 + (1 - B/\epsilon\Delta_1)(t^{\epsilon\Delta_1} - 1)]^\theta t^{-\gamma}, \quad (1)$$

where $\epsilon = d_c - d$, $t = (T - T_c)/T_c$, A and B are non-universal constants, and Δ_1 , θ , and γ are universal exponents. Fitting to a form of this type, Van Dyke and Camp⁶ found $\gamma = 1.00 \pm 0.02$ for the Ising model at $d=4$, thus confirming $d_c=4$ for this model. We have carried out the same program for the random resistor network by analyzing a low-concentration expansion for an analogous susceptibility. In order to do this, one must identify the order parameter. For arbitrary networks, Kasteleyn and Fortuin² show that the correlation function for the $s \rightarrow 0$ state Potts model yields the resistance, R_{ij} , between lattice points i and j . This correlations function can also be obtained by averaging $x_i x_j$ over the Gaussian density matrix, $\rho = \exp[-\frac{1}{2} \sum_{mn} \sigma_{mn} (x_m - x_n)^2]$. These observations suggest that R_{ij} plays the role of the correlation function and therefore we define the resistive susceptibility as $\chi^{(r)} = \sum_j \chi_{ij}$, where

$$\chi_{ij} = [R_{ij} \nu_{ij}]_{av}, \quad (2)$$

where $\nu_{ij} = 1$ if i and j are in the same cluster and $\nu_{ij} = 0$ otherwise, and $[]_{av}$ denotes a configurational average.

We first consider the case $\sigma_c = 0$ and relate $\gamma^{(r)}$, the exponent for $\chi^{(r)}$, to the conductivity exponent μ defined by $\Sigma \sim (p - p_c)^\mu$, $p - p_c^+$. Consider the Green's function $G(\vec{r}, \vec{r}')$ which gives the voltage at \vec{r}' in response to a unit current source at \vec{r} . In the limit $q \rightarrow 0$ the Fourier transform of this Green's function is of the form

$$G(q) = P^2(p) \{ \Sigma(p) q^2 F[\xi(p)q] \}^{-1}, \quad (3)$$

where $P(p)$ is the fraction of sites in the infinite cluster and $\xi(p)$ is the correlation length. This Green's function is identical to that for transverse excitations in a Heisenberg ferromagnet at zero temperature for which Eq. (3) holds.⁹ In Eq. (3) $F(z)$ is analytic at $z=0$, so that if one writes

$$G(q) = \sum_{n=-2}^{\infty} G^{(n)} q^n,$$

then one has

$$G^{(2)} \sim P^2(p) \xi(p)^4 / \Sigma(p). \quad (4)$$

Now consider the response of finite clusters for

$p < p_c$, to a current source $I_0 e^{i\vec{q} \cdot \vec{r}}$. In order to be able to solve Kirchoff's equations we must eliminate the $q=0$ component of current. Thus we take the current source to be $I_0 [\exp(i\vec{q} \cdot \vec{r}) - \exp(i\vec{q} \cdot \vec{r}')]]$, where \vec{r}' is approximately the center of gravity of the cluster. In the limit $q \rightarrow 0$ the induced voltage $V(\vec{r}')$ is then

$$V(\vec{r}') = I_0 \sum_{\vec{r}} G(\vec{r}, \vec{r}') i\vec{q} \cdot (\vec{r} - \vec{r}'). \quad (5)$$

Here and below, the lattice sums are restricted to a single cluster. To get the response at wave vector \vec{q} we need to evaluate

$$V(\vec{q}) = \sum_{\vec{r}} V(\vec{r}') [\exp(-i\vec{q} \cdot \vec{r}') - \exp(-i\vec{q} \cdot \vec{r})]. \quad (6)$$

Here again we subtract the term $\exp(-i\vec{q} \cdot \vec{r})$, because we can arbitrarily fix the average voltage of each cluster to be zero. Thus

$$V(\vec{q})/I_0 = \sum_{\vec{r}, \vec{r}'} G(\vec{r}, \vec{r}') (\vec{q} \cdot \delta\vec{r}) (\vec{q} \cdot \delta\vec{r}'), \quad (7)$$

where $\delta\vec{r} = \vec{r} - \vec{r}'$. In a cluster the distances are of order $\xi(p)$ and $G(\vec{r}, \vec{r}')$ is related to $R_{\vec{r}, \vec{r}'}$, so that

$$G^{(2)} \sim \xi^2(p) \sum_{\vec{r}, \vec{r}'} R_{\vec{r}, \vec{r}'} = \xi^2(p) \chi^{(r)}. \quad (8)$$

If we assume $G^{(2)}$ to be governed by the same exponent for $p < p_c$ as for $p > p_c$, we obtain from Eqs. (4) and (8) that¹⁰

$$\gamma^{(r)} = \mu - 2\beta + 2\nu = \mu - (d-2)\nu + \gamma, \quad (9)$$

where $P(p) \sim (p - p_c)^\beta$ and $\xi(p) \sim |p - p_c|^{-\nu}$. The second equality in Eq. (9) follows by using the scaling relation $2\beta = d\nu - \gamma$ for the percolation exponents.

We have developed a low- p expansion for $\chi^{(r)}$ as $\chi^{(r)} = \sum a_n(d) p^n$ and have obtained the polynomials⁷ in d for $a_n(d)$ for $n \leq 10$. This was done as follows. For any cluster, Γ , we define $\chi^0(\Gamma)$ to be the value of $\chi^{(r)}$ for that cluster. The cumulant, $\chi^c(\Gamma)$, is defined recursively as

$$\chi^c(\Gamma) = \chi^0(\Gamma) - \sum_{\gamma \in \Gamma} \chi^c(\gamma),$$

where the sum is over all connected clusters contained within Γ . Then the result is $a_n(d) = \sum_{\Gamma(n)} b(\Gamma) \chi^c(\Gamma)$, where the sum is over all clusters with n bonds and $b(\Gamma)$ is the number of times the cluster Γ can be formed on a hypercubic lattice in d dimensions. The mean-field-theory value of $\gamma^{(r)}$, denoted $\gamma_{MF}^{(r)}$, can be obtained by taking d large and keeping only self-avoiding walk diagrams. For n -step walks $b \sim (2d)(2d-1)^{n-1}$ and χ^c is equal to the resistance between the ends of the walk, viz. n/σ_s . Thus

$$\chi^{(r)} \sim \sum_n p^n (n/\sigma_s) (2d-1)^n, \quad (10)$$

TABLE I. Padé approximant results for p_c and $\gamma^{(r)}$.

d	p_c	$\gamma^{(r)}$
2	1/2	3.7 ^a
3	0.2465	2.78
4	0.1600	2.46
5	0.1181	2.19 (2.22) ^b
5.5	0.1047	2.13 (2.10) ^b
6	0.0943	2.09 (2.00) ^b
6.5	0.0858	2.06
7	0.0787	2.04

^aThis value is rather uncertain (± 0.3) (see Ref. 13). The errors for the other entries (p_c and $\gamma^{(r)}$) are of order one in the last decimal place quoted.

^bValues obtained using corrections to scaling, Eq. (1).

which gives $\gamma_{MF}^{(r)} = 2$. For finite d one uses the a_n 's to construct the various Padé approximants to $\partial \ln \chi^{(r)} / \partial p \sim \gamma^{(r)} / (p_c - p)$ to determine $\gamma^{(r)}$. The results of this and similar procedures are summarized in Table I. [The coefficients $a_n(d)$ will be published elsewhere.]

The effect for d near d_c of the logarithmic corrections of Eq. (1) [with $t + (p_c - p)/p_c$] is scaled by θ . For isotropic n -vector φ^4 and φ^3 models¹¹ one has $\theta = 2(\partial \gamma^{(r)} / \partial \epsilon)|_{\epsilon=0}$ and $\Delta_1 = \frac{1}{2}$. Without including logarithmic corrections we find $\gamma^{(r)} - \gamma_{MF}^{(r)} = 0.09$ for $d = 6$, and $\gamma^{(r)} - \gamma_{MF}^{(r)} \sim \epsilon/5$ for $\epsilon \leq 1$, which gives $\theta \approx 0.4$. These values are quite similar to those for the Ising model for $d = 4$. Including logarithmic corrections as in Eq. (1) we find an acceptable fit to the series at $d = 6$ for $\gamma^{(r)} = \gamma_{MF}^{(r)}$ with $B = 0.4$. This result is again similar to that found for the Ising model, and confirms that $d_c = 6$ here.

The scaling relation, Eq. (9), is satisfied by the mean-field-theory values, viz. $\gamma^{(r)} = 2$, $\beta = 1$, $\nu = \frac{1}{2}$, and (see below) $\mu = 3$. For $d = 3$ one has¹² $\mu = 1.6$, $\beta = 0.4$, and $\nu = 0.95$, which yields $\gamma^{(r)} = 2.7$ from Eq. (9), in good agreement with our value $\gamma^{(r)} = 2.78$. For $d = 2$, the values¹² $\mu = 1.1$, $\beta = 0.15$, and $\nu = 1.35$ yield $\gamma^{(r)} = 3.5$, which is also consistent with our result, $\gamma^{(r)} = 3.7 \pm 0.3$.¹³

Finally, we propose scaling relations for the case $\sigma_c \neq 0$. We will use the coherent-potential approximation¹⁴ (CPA) to generate mean-field values of the critical exponents. Within the CPA Kirkpatrick¹⁵ finds that the network is described by an effective conductance σ_{eff} which is determined by

$$p = (\sigma_{eff} - \sigma_c) \left[\frac{1-p}{\sigma_c - \sigma_{eff}} + \frac{2}{z\sigma_{eff}} \right]. \quad (11)$$

In this approximation $p_c = p^* = 2/z$. For $|t| \ll 1$, with $t = (p - p^*)/p^*$, and $\sigma_c/\sigma_s \ll 1$, the dominant behavior of σ_{eff} is given by the scaling form

$$\sigma_{eff} = \frac{1}{2}\sigma_s \{t + [t^2 + 2(\sigma_c/\sigma_s)/(z-2)]^{1/2}\}. \quad (12)$$

If we define the exponents¹⁶ $\bar{\mu}$ by $\Sigma(p) \sim (p_c - p)^{-\bar{\mu}}$ for $p \rightarrow p_c^-$ with $\sigma_s = \infty$, and τ by $\Sigma(p_c) \sim \sigma_c^\tau \sigma_s^{1-\tau}$, then Eq. (12) implies $\bar{\mu} = 1$ and $\tau = \frac{1}{2}$. To determine μ from Eq. (12) we must be able to relate σ_{eff} to Σ , when $\sigma_c = 0$. Kirkpatrick¹⁵ assumed σ_{eff} and Σ to be equivalent. However, it is more plausible to assume that the CPA provides an approximation to the Green's function of Eq. (3), in which case one would write $G^{-1}(q) \sim \sigma_{eff} q^2$, for $q\xi(p) \ll 1$. Equation (3) would then imply that $\Sigma(p) \sim \sigma_{eff} P^2(p)$, which would give $\mu = 3$ as the mean-field result. These mean-field values of μ and $\bar{\mu}$ agree with those given, respectively, by de Gennes⁵ and Straley.¹⁶ In particular, our treatment and numerical values of $\gamma^{(r)}$ confirm de Gennes's argument⁵ that Stinchcombe's result¹⁷ ($\mu = 2$) for the conductivity of the Bethe lattice is not the appropriate value for μ_{MF} . The CPA result for τ does not agree with Straley's value,¹⁶ which, however, is based in part on the Bethe lattice value for μ .

Now we consider the regime $d < 6$, where mean-field theory is not valid. For $p < p_c$ and $\sigma_s \rightarrow \infty$, we assume that $\Sigma(p)$ obeys a scaling form

$$\Sigma(p) = \sigma_c^\tau \sigma_s^{1-\tau} F((\sigma_s/\sigma_c)(p - p_c)^\varphi), \quad (13)$$

where $F(0) = 1$. [The CPA, Eq. (12), provides the mean-field-theory result, $\varphi = 2$.] For $\sigma_s = \infty$, Eq. (13) implies that

$$\bar{\mu} = (1 - \tau)\varphi. \quad (14)$$

To develop another scaling relation we consider the regime $p < p_c$, $\sigma_c \rightarrow 0$ and study the behavior of $\chi^{(r)}$ as defined in Eq. (2). In the limit of large separation R_{ij} tends to a value proportional to $\Sigma(p)^{-1}$, since we take $\sigma_c \neq 0$. In this limit, then, one has $[R_{ij}\nu_{ij}]_{av} = [\nu_{ij}]_{av}/\Sigma(p)$, so that $\chi^{(r)} \sim \chi^{(p)}/\Sigma(p)$, where $\chi^{(p)}$ is the percolation susceptibility, i.e., the mean-square cluster size. We now assume a scaling form for $\chi^{(r)}$:

$$\chi^{(r)} = \sigma_s^{-1} (p - p_c)^{-\gamma^{(r)}} G((\sigma_c/\sigma_s)(p - p_c)^{-\varphi}) \quad (15)$$

with $G(0) = 1$ and $G(x) \sim x^{-s}$ for large x . Then for $\sigma_c/\sigma_s \gg (p - p_c)^\varphi$ we have

$$\chi^{(p)}/\Sigma(p) = (p - p_c)^{-\gamma^{(r)} + s\varphi} (\sigma_c^{-s} \sigma_s^{s-1}). \quad (16)$$

For this to agree with the definition of τ , we

must set $s = \tau$. Then we get

$$\chi^{(p)} \sim (p - p_c)^{-\gamma^{(r)} + \tau \varphi},$$

so that

$$\gamma^{(r)} - \tau \varphi = \gamma. \quad (17)$$

Combining relations (9), (14), and (17) we get¹⁸

$$\tau = [\mu - (d-2)\nu] / [\mu + \bar{\mu} - (d-2)\nu] \quad (18)$$

as contrasted to Straley's result,¹⁶ $\tau = \mu / (\mu + \bar{\mu})$. For $d=2$, our result agrees with Straley's and with the exact results,¹⁹ $\tau = \frac{1}{2}$, $\bar{\mu} = \mu$. For $d=6$, our mean-field-theory values of the exponents satisfy Eq. (18), whereas Straley's do not. The numerical evidence for $d=3$ is not yet conclusive, although Eq. (18) does not work very well for the current values, $\mu^{(r)} = 1.6$, $\bar{\mu}^{(r)} = 0.6$, and $\tau = 0.77$.¹⁹ However, since Straley's values¹⁹ of the exponents do not satisfy universality very well, they may not be sufficiently accurate for this comparison.

Our conclusions are summarized in the Abstract.

The authors have greatly profited from the continual encouragement and advice of Professor T. C. Lubensky. We would also like to thank Professor E. Siggia for advice on the calculation of the logarithmic corrections and Professor M. E. Fisher for advice on the analysis of the power series. We also thank Professor J. P. Straley for communicating his results prior to publication.

*Supported in part by National Science Foundation Grant No. DMR 76-21703, Office of Naval Research Grant No. N00014-76-0-0106, and National Science Foundation Materials Research Laboratory Program Grant No. DMR 76-00678.

¹For a review see V. K. S. Shante and S. Kirkpatrick, *Adv. Phys.* **20**, 325 (1971).

²P. W. Kasteleyn and C. M. Fortuin, *J. Phys. Soc. Jpn. Suppl.* **26**, 11 (1969).

³G. Toulouse, *Nuovo Cimento* **23**, 234 (1974).

⁴A. B. Harris, T. C. Lubensky, W. K. Holcomb, and C. Dasgupta, *Phys. Rev. Lett.* **35**, 327 (1975).

⁵P. G. de Gennes, *J. Phys. Lett.* **37**, L1 (1976).

⁶J. P. Van Dyke and W. J. Camp, in *Magnetism and Magnetic Materials—1975*, AIP Conference Proceedings No. 29, edited by J. J. Becker, G. H. Lander, and J. J. Rhyne (American Institute of Physics, New York, 1976), p. 502.

⁷M. E. Fisher and D. S. Gaunt, *Phys. Rev. B* **3**, A224 (1964).

⁸J. Rudnick and D. R. Nelson, *Phys. Rev. B* **13**, 2208 (1976).

⁹B. I. Halperin and P. C. Hohenberg, *Phys. Rev.* **188**, 898 (1969).

¹⁰There is an alternative way to derive Eq. (9). We note that if $[R_{ij} \nu_{ij}]_{av}$ is the correlation function, then the order parameter $\sigma^{(r)}$ may be defined as

$$(\sigma^{(r)})^2 = \lim_{r_{ij} \rightarrow \infty} [R_{ij} \nu_{ij}]_{av},$$

for $p > p_c$. In this limit we have that $[\nu_{ij}]_{av} = P^2(p)$ and $[R_{ij}] \sim L$, the path length between nodes, introduced by Shklovskii {B. I. Shklovskii, *Fiz. Tekh. Poluprovodn.* **8**, 1586 (1974) [*Sov. Phys. Semicond.* **8**, 1029 (1975)]} and de Gennes (Ref. 5), who find that $\Sigma(p) \sim L^{-1} \xi^{2-d}$. Thus, $(\sigma^{(r)})^2 \sim P^2(p) \xi^{2-d} / \Sigma(p)$, i.e., $2\beta^{(r)} = 2\beta + (d-2)\nu - \mu$, from which Eq. (9) follows. Even though there are doubts as to the validity of defining L , this argument may yield the correct answer, since L does not appear in the final result.

¹¹A. D. Bruce and D. J. Wallace, *J. Phys. A* **9**, 1117 (1976).

¹²A tabulation with references to recent work is given in A. B. Harris and S. Kirkpatrick, *Phys. Rev. B* (to be published).

¹³The series for $\chi^{(r)}$ is very irregular at $d=2$. This is probably a reflection of the logarithmic divergence, $R_{\vec{r}, \vec{r}'} \sim \ln |\vec{r} - \vec{r}'|$ for large $|\vec{r} - \vec{r}'|$, which occurs in the pure system for $d=2$.

¹⁴P. Soven, *Phys. Rev.* **156**, 809 (1967).

¹⁵S. Kirkpatrick, *Phys. Rev. Lett.* **27**, 1722 (1971).

¹⁶J. P. Straley, *J. Phys. C* **9**, 783 (1976).

¹⁷R. B. Stinchcombe, *J. Phys. C* **7**, 179 (1974).

¹⁸Our results suggest that in fact σ_{eff} should be identified with L^{-1} , where L is as defined in Ref. 10.

¹⁹J. P. Straley, to be published.