



The University of Bradford Institutional Repository

<http://bradscholars.brad.ac.uk>

This work is made available online in accordance with publisher policies. Please refer to the repository record for this item and our Policy Document available from the repository home page for further information.

To see the final version of this work please visit the publisher's website. Access to the published online version may require a subscription.

Link to publisher's version: <http://dx.doi.org/10.1080/13607863.2015.1021753>

Citation: Jones D, Drew P, Elsey C, Blackburn D, Wakefield S, Harkness K and Reuber M (2016) Conversational assessment in memory clinic encounters: interactional profiling for differentiating dementia from functional memory disorders. *Aging & Mental Health*. 20(5): 500-509.

Copyright statement: © 2016 Taylor & Francis. This is an Author's Original Manuscript of an article published by Taylor & Francis in *Aging & Mental Health* in 2016 available online at <http://www.tandfonline.com/doi/full/10.1080/13607863.2015.1021753>

Conversational assessment in memory clinic encounters: interactional profiling for the differential diagnosis of dementia and functional memory disorder

Danielle Jones*, Paul Drew, Chris Elsey, Daniel Blackburn, Markus Reuber

Bradford Dementia Group, University of Bradford, Bradford, UK; Department of Social Sciences, Loughborough University, Loughborough, UK; Department of Social Sciences, Loughborough University, Loughborough, UK; Academic Neurology Unit, University of Sheffield, Sheffield, UK; Sheffield Institute for Translational Neuroscience (SITraN), University of Sheffield, Sheffield, UK

Corresponding Author: Danielle Jones, Bradford Dementia Group, Faculty of Health Studies, University of Bradford, Bradford, BD7 1DP. **Telephone:** 01274 236469 **Email:** d.k.jones1@bradford.ac.uk

Paul Drew, Department of Social Sciences, Loughborough University, Epinal Way, Leicestershire, LE11 3TU. **Telephone:** 01509 223365 **Email:** p.drew@lboro.ac.uk

Chris Elsey, Department of Social Sciences, Loughborough University, Epinal Way, Leicestershire, LE11 3TU. **Email:** C.Elsey@lboro.ac.uk

Daniel Blackburn, Sheffield Institute for Translational Neuroscience (SITraN), University of Sheffield, 385A Glossop Road, Sheffield, S10 2HQ. **Telephone:** 0114 222 2267 **Email:** d.blackburn@sheffield.ac.uk

Markus Reuber, Academic Neurology Unit, University of Sheffield, Royal Hallamshire Hospital, Glossop Road, Sheffield, S10 2JF. **Telephone:** 0114 226 8688 **Email:** markus.reuber@sth.nhs.uk

Acknowledgements: This article presents independent research funded by the National Institute for Health Research (NIHR) under its Research for Patient Benefit (RfPB) Programme (Grant Reference Number PB-PG-0211-24079). The views expressed are those of the author(s) and not necessarily those of the NHS, the NIHR or the Department of Health.

Word count: 5,748 (of which 1,405 is data)

Conversational assessment in memory clinic encounters: interactional profiling for the differential diagnosis of dementia and functional memory disorder

Abstract

Objectives: In the UK dementia is under-diagnosed and there is national variation in memory clinic service provision. At present the clinical differentiation between dementia and functional (nonprogressive) memory complaints is complex and involves extensive neuropsychological testing. Government initiatives on ‘timely diagnosis’ aim to improve the rate and quality of diagnosis for those with dementia. This study seeks to improve methods of diagnostic screening by analysing communication between clinicians and patients during diagnostic assessment and establishing conversational profiles from which clinicians can establish differential diagnoses.

Method: The data corpus consists of video- and audio recording of 105 initial consultations between neurologists and patients referred to a UK memory clinic. Conversation analysis was used to explore recurrent communicative practices within these data.

Results: Two discrete conversational profiles began to emerge to help differentiate between patients with dementia and functional memory complaints based on, 1) whether the patient is able to answer questions about personal information; 2) whether they can display working memory in interaction; 3) whether they are able to respond to compound questions; 4) the time taken to respond to questions; and 5) the level of detail they offer when providing an account of their memory failure experiences.

Conclusion: Conversational profiles can differentiate patients with dementia from those with functional memory complaints. Conversational profiling has potential clinical application; using conversation as a method of diagnostic screening and assessment could hold differential diagnostic value.

Keywords: Dementia, functional memory disorder, conversation analysis, differential diagnosis, interaction,

Introduction

There has been a sharp increase in the number of people attending primary care, and being referred to specialist care services with concerns about their memory. The number of people assessed by memory clinics has risen four-fold since 2010 (Royal College of Psychiatrists, 2013). These referral rates are expected to rise further as the UK government introduces incentives for practitioners to screen for dementia (DOH, CQUIN 2012; NHS, DES 2013). The increase in referrals risks overwhelming memory clinics and other specialist services (such as neurology, gerontology and old age psychiatry). Specialist memory services are already under pressure in many areas of the United Kingdom and there is great national variation in memory clinic service provision and the time it takes for people to be diagnosed. The rate of people with dementia carrying a formal diagnosis is currently only 48% in England and varies from 32% in the worst performing areas to 75% in the best (Alzheimer’s Society, 2014). The government initiatives on ‘timely diagnosis’ aim to increase this rate by two thirds by 2015.

At present the clinical differentiation between a form of dementia and functional (nonprogressive) memory complaints is complex, and further research is needed to establish reliable biological markers to improve the differential clinical definitions and diagnostic accuracy between progressive neurodegenerative disorders and 'functional' memory concerns (Knopman et al, 2001). The diagnosis is based on a clinician's interpretation of the history given by a patient and their companion, complemented by brain scanning (Magnetic Resonance Imaging, MRI or Computerised tomography, CT) and extensive neuropsychological testing. In uncertain cases, the diagnostic process will involve a re-examination and investigation of the patient after an interval of six to twelve months. This diagnostic process requires considerable expertise, is costly and time-consuming and cannot be offered to all people complaining of memory problems. A 'suspended' diagnosis is likely to cause significant anxiety and unlikely to improve the condition of patients with functional memory complaints; and patients, as well as their family carers, often find the lengthy process of extensive testing distressing (Lai et al, 2008; Gibson & Anderson, 2011). Simpler and shorter neuropsychological screening instruments lack specificity and are only of modest diagnostic value (Boustani et al, 2005). Increasingly, medical practitioners are searching for new approaches for diagnostic assessment, which will work to reduce these problems by placing less emphasis on the extensive and expensive formal testing, and more on the conversation they have with the patient during history taking.

This paper reports the findings of a qualitative, in depth analysis of the conversations between neurologists and patients referred to a memory clinic because of concerns about their memory. The aim of this research is to assist in the diagnosis of memory complaints, particularly to assist in the diagnostic differentiation between dementia from those associated with functional memory disorder (FMD). Through careful, detailed analysis of patients' communication in their first encounters with neurologists, our research strategy is to identify features of patients' talk and interaction patterns that have the potential to help clinicians differentiate between people presenting with a progressive dementia and those presenting with other non-progressive memory complaints. A further goal is thereby to reduce the extent to which patients undergo what may be unnecessary and distressing further neuropsychological testing. Anecdotal evidence from conversations with neurologists suggests that specialist practitioners will often form a working diagnosis within the first five minutes of the opening of the consultation. This research seeks to provide the conversational evidence underpinning such clinical assessments, and explores whether interactional or linguistic features can be identified and described which could help with the diagnostic process when patients present with memory complaints.

Methods

Study design & subjects

Video- and audio recording of 105 initial consultations between neurologists and patients referred to the Department of Neurology at the Royal Hallamshire Hospital (Sheffield, UK) were collected between October 2012 and August 2014. Patients had been referred to the specialist neurology-led clinic for younger adults (minimum age 18) by primary care general practitioners (GP), neurologists and other secondary services such as psychiatry because of concerns about memory. Patients were seen by one of four neurologists with a special interest in memory disorders. Patients were routinely encouraged to bring along a family member, friend or carer to

the appointment. After receiving a 'gold standard' clinical diagnosis (confirmed on the basis of an initial assessment by a Consultant Neurologist with a special interest in memory disorders including the Addenbrooke's Cognitive Examination (ACE-R), a detailed neuropsychological test battery and magnetic resonance imaging (MRI) of the brain) 25 patients were selected for the purpose of interactional profiling- 9 with a dementia diagnosis (average ACE-R score 56/100, range from 28-80) and 16 with functional memory complaints (average ACE-R score 93/100 range from 85-99) (for a definition of FMD please see Schmidtke et al, 2008; Schmidtke & Metternich, 2009). This study represents an initial analysis of the data from the memory clinic. At the time of writing, recruitment to the study and further analysis were ongoing.

Data analysis

Recordings were transcribed in considerable detail, using the transcription conventions widely adopted in CA (Jefferson, 1983, 2004; for transcription conventions see Appendix 1). In transcribing the data, all names of people, places and other potentially identifying information have been pseudonymised.

The qualitative method of CA was used for analysing the data (for a summary see Drew, 2001 & 2005; Sidnell, 2010). This micro-analytic approach has been applied successfully in primary care services to examine the organization of medical communication (for a review Heritage & Stivers, 1999; Stivers, 2002; Heritage & Maynard, 2006; Heritage et al, 2007). More recently, research has been conducted in secondary care services revealing medical communication to be a powerful diagnostic tool for practitioners (Schwabe et al, 2007; Robson et al, 2012). Most notably, this study follows the design of research conducted in the neurological seizure clinic at the Royal Hallamshire Hospital in Sheffield, UK, by a team which has used CA methodology in the differential diagnosis of seizure disorders. This research identified diagnostically relevant linguistic, topical and interactional features that aided clinicians to differentiate between a diagnosis of epileptic and non-epileptic seizures. Furthermore, a prospective multi-rater study has confirmed the diagnostic potential of these conversational profiles in the seizure clinic setting. By using a diagnostic scoring aid (DSA) to convert qualitative assessments into a numeric score, analysts blinded to diagnosis predicted 85% of diagnoses correctly. The video-EEG recording of typical attacks ultimately proved all diagnoses (see Reuber et al, 2009). These interdisciplinary collaborations between clinicians and linguists support an applied approach whereby research findings inform medical practice (Reuber et al, 2009; Ekberg et al, 2014).

This paper will present only a small number of extracts, comprising only short communicative exchanges, but they exemplify larger patterns in the data (not statistically presented here). This is a practice routinely used in CA research to provide evidence for the findings.

Ethics

The patients recruited received written information about the study prior to their appointment date and had the opportunity to ask questions of a member of the research team (not the doctor they had come to see) prior to their initial appointment in the memory clinic. All participants gave written informed consent to participate and were informed that they could withdraw from the study at any time. Patients lacking capacity to consent were excluded from the study. The study was approved by NRES Committee Yorkshire & The Humber - South Yorkshire and was guided by the principles outlined in the Mental Capacity Act (2005).

Results

The initial history-taking phase of the encounters, which formed the basis of this analysis, lasted between 7 minutes and 28 seconds and 32 minutes and 29 seconds. 9 of the patients were male, 16 were female. The patient's age ranged from 47 to 77. Of the 25 clinical encounters selected for analysis, 11 were dyadic in type (those generated from the interaction between the neurologist and the patient) (no one with dementia attended the clinic alone whereas 11 of the 16 patients with FMD came alone) and 14 were triadic (including the talk of the patients companion) (all 9 patients with dementia attended accompanied whereas only 5 of the 16 patients with FMD came accompanied). This paper focuses on some of the dyadic features of interaction. This initial analysis of these dyadic interactions in the memory clinic data identified five features that could contribute to a differential diagnostic conversational profile of patients presenting with dementia or FMD. These features are: 1) whether the patient is able to answer questions about personal information (for example 'how old are you?' or 'where do you live?'); 2) whether they can display working memory in interaction; 3) whether they are able to respond to compound questions; 4) time taken to respond to questions; and 5) the level of detail they offer when providing an account of their memory failure experiences.

Responding to questions about personal information

Routinely, at the beginning of the consultations, neurologists ask patients a series of questions. These questions are designed to seek personal information from the patient, for example how old they are or where they live. The first two extracts demonstrate contrasting cases in which the neurologist is asking the patient's age. The first patient (Extract 1) subsequently received a medical diagnosis of FMD, the second one of dementia (Extract 2).

Extract 1

```
004_FMD
01 _ DOC .hh So first of all how old are you now:.
02 PAT Seventy-eight.
```

Extract 2

```
017_DEM
01 _ DOC So how old are you now Mr Marshall,
02 PAT Okay.=How old,
03 DOC How old. Yes.
04 PAT Er Twenty one:: e[r:
05 COM [No how old are you.=
06 =You're actual age ( ).
07 PAT Oh two thousa:[nd:
08 COM [No you're age darli[ng.
09 PAT [Ages.=
10 COM No. You're age.
11 (1.8)
12 COM How old you are.
13 (2.8)
14 PAT hhh
15 COM Not you're date of birth. How old are you,
16 (1.0)
17 PAT I'm er (2.9)
18 COM huhmm
19 (1.0)
20 PAT Well I wa:s mu hu hu
```

21 (3.0)
 22 PAT No. It's gone.
 23 (0.8)
 24 DOC Oka:y. S[o
 25 COM [Sixty nine.

The first extract (1) is typical of patients with FMD in that they are able to produce the information required by the neurologist quickly and unproblematically. The second extract (2) illustrates how patients with dementia regularly have difficulty recalling personal information (in this case the patient's wife intercedes in lines 12 and 15 to try to prompt the patient to answer and eventually in line 25 answers on his behalf). A person's ability to respond contiguously and accurately to questions seeking personal information such as their age differentiates those with FMD, from those with dementia (who often struggle to produce the correct information).

Working memory in interaction

The second feature also relates to a patient's memory functionality. Working memory is one area of cognition examined in neuropsychological tests (such as the ACE-R). This aspect of cognitive functionality can also be displayed and assessed in interaction during the history taking conversation. The patient in Extract 3 (diagnosed with FMD) is able to display memory in the interaction itself.

Extract 3

011_FMD
 01 DOC ERm: (.) So I'm Doctor Blackburn.= I'm
 02 the (0.8) registrar in Neurology, .hh
 03 d'yu'wanna tell me:: (1.0) um why you've
 04 come today and what expectations you have
 05 about the clinic.
 06 PAT .hh Well one of the reasons was because I
 07 have a partner (0.7) a:nd he was sort of
 08 reminiscing about:t (0.2) times past >like
 09 holidays and things we've had and I thought
 10 .h "well I can't remember tha:t an' I
 11 can't remember that happening." .hh An' the's
 12 there's other things where, 'cause I work in a
 13 public hou:se: (0.8) I'd be down stairs working
 14 (0.3) .h an' then somebody u'll say Oh a- a
 15 pri:me example was er on Frida:y (.) .hh ((clears
 16 throat)) when I needed to go upstairs for
 17 something (1.0) tkh an' I just set off going
 18 and we've got a telephone of the staircase going
 19 upstairs. .h An' just as I went upstairs the phone
 20 rang.(.) .hh I: had to do- somebody said "oh
 21 would you mind doing a quick survey."= So I did
 22 this quick survey .hh and I went to the top of
 23 the stairs (0.2) and thought (1.3) "what have I
 24 come up here for."
 25 ... (1 minutes 17 seconds of patient talk omitted)
 26 PAT I think that's about it really.
 27 (.)
 28 PAT .hh Expectations, (0.9) I don't know,
 29 (0.6)
 30 PAT I don't know what to: to expect >'cause
 31 I've never been in this situation before.=
 32 DOC =No. S[o ho-
 33 PAT [I don't think you'll be able to
 34 give me a magic tablet that'll make
 35 everything perfect but (1.0) there you

The neurologist asks the patient a question made up of two parts requiring two separate answers. The first, ‘d’yu’wanna tell me why you’ve come today’ (lines 3-4) pragmatically requires the patient to inform the neurologist about the reason for their attendance at the clinic; and the second, ‘(d’yu’wanna tell me) what expectations you have about the clinic’ (lines 4-5). The patient proceeds with an extended telling, in great detail, about his memory failure experiences. Prefacing the telling with ‘Well one of the reasons was...’ attends to the first part of the neurologist’s question. Having fully answered that questions after 2.5 minutes (1:17mins omitted from the transcript) the patient is perfectly able to retrieve the second part of the question and answer it, ‘Expectations. I don’t know’ (line 28). This demonstrates that the patient is aware of the requirements of both the questions, can process them and respond accordingly, and can display through interaction his cognitive functioning with regards to working memory.

Patients with FMD not only display working memory in relation to what the *other* has said in the consultation (as seen in Extract 3 – recalling the neurologist’s prior talk), but also when recalling and repeating information they have previously voiced. People with FMD do additional interactional work, marking their self-repetitive talk with ‘like I said’ or ‘as I say’, which orients to their awareness of their talk as being repetitive and displays their working memory functionality in the interaction (see Extract 4).

Extract 4

028_FMD

01 PAT .hh I seem to get- I- I do tend to get
 02 mi:graines which is: li:ght induced.
 03 (0.4)
 04 PAT If I get a flashing light or:,
 05 COM °You’ve always had that [thoug]h.°
 06 PAT [Yeah.]
 07 DOC Can you take me through a typ- typical
 08 mi:graine:. (.) for you,
 09 PAT .h For me it’s: u- >as I say< it’s usually
 10 li:ght induced a:nd it always starts with
 11 flashing li:ghts: in my right eye.

Here the patient informs the neurologist that he suffers from light induced migraines (lines 1-2), adding that he experiences a ‘flashing light’ (line 4). The neurologist asks for further information about the patients migraines (lines 6-8), in response to which the patient repeats what he had previously told the neurologist about his migraine being ‘light induced’ and their being associated with ‘flashing lights’ (lines 9-11). However he prefaces this repetition with ‘as I say’ (line 9) to mark his self-repetition and to display that the information which will proceed has been voiced by him previously. This interactional resource, which displays working memory, appears as a recurrent feature in the consultations with patients with FMD and contributes to their conversational profile.

Unlike patients with FMD, those with dementia are often unable to display memory in this way in their consultations. They are often unable to retain information about what has been said even a few seconds earlier in the interaction, either by themselves or by the neurologist. When repeating themselves, they do not indicate that they are aware of their repetitions – that is, they do not preface their repetitions with such markers of awareness as ‘As is said...’. This absence of marked self-

repetitions is therefore a part of the conversational profile of those with dementia. Repeated information is often delivered as if for the first time (Jones, 2013). These 'second first-time tellings' are not marked using prefaces such as 'as I said'. Neurologists often indicate that they are aware that the patients are repeating themselves. Extract 5 is an example of an unmarked repetition or a 'second first-time telling' where a person with dementia is repeating information previously told to the neurologist as if for the first time. This provides interactional evidence that their working memory is compromised.

Extract 5

048_DEM
01 DOC .hh And what was your first job after
02 leaving school.
03 (6.8)
04 DOC ↑ Can you remember what your first job was?
05 (3.0)
06 PAT Not off hand.
07 DOC Okay. And what's: what's your job (0.4) your
08 main career been during your working life.
09 (4.2)
10 PAT It was a: (7.8)
11 DOC So what- what sort of work were you doing
12 just before you retired.
13 (6.3)
14 DOC Do you remember the job title or what kind
15 of things you would do on a day-to-day basis.
16 PAT Uhm Laboratory supervisor.
17 DOC Right.
18 (0.4)
19 DOC .hhh [And what] so-, what sort of lab was that in.
20 PAT [Sorry]
21 (0.9)
22 PAT Chemical lab.
23 (4.4)
24 DOC So di- had you done a degree or a diploma
25 >as par-< obviously to get to that level
26 you must have done a number of profess-
27 professional qualifications.=And did you do
28 that in a block or as a day release or,
29 PAT I did it as a day release.
30 DOC Uhm hmm.
31 (6.5) ((Doctor writing))
32 DOC And did you have to do a Masters for that,
33 or:, is it a, at what level.
34 (0.8)
35 PAT Can't remember now.
36 (1.0)
37 DOC .h And how many people were you- you in
38 charge of,
39 (0.3)
40 DOC Before you'd finished work.
41 (0.6)
42 PAT Er quite a few,
43 DOC °Uhm°
44 (3.5)
45 PAT I was a laboratory supervisor you see.=So
46 I was in charge of eve[rything].
47 DOC [Yes:. Yes.

The patient told the neurologist twice within a few minutes that she was a laboratory supervisor (lines 16 and 45), once in response to a series of previously unanswered questions about her career and once in response to a question about how many staff she was in charge of. The second telling was produced with no preface orienting it as a repetition and is receipted by the neurologist with a multiple saying, ‘Yes. Yes.’ (Stivers, 2004) marking the prior as problematic due to its perseveration and revealing a disruption of cognitive functioning. The ‘you see’ (line 45) which forms part of the patient’s second time telling attends to the lack of specificity she gave in her previous answer of being in charge of ‘quite a few’ people (line 42). Being in a higher position as a ‘supervisor’, and indeed being ‘in charge of everything’ could explain why ‘quite a few’ is a legitimate answer in that she may have overseen a larger number of people. The point here is that patient’s in these consultations who have dementia are often repetitive but do not mark their repetition. None of the patients with dementia used resources such as ‘as I said’ or ‘like I say’. These features and displays of working memory (or lack thereof) in the interaction appear to be diagnostically significant and can contribute to the differentiation of diagnoses.

Compound questions

Neurologists questions during the consultations range from mono-topical, e.g. ‘How old are you?’ (extracts 1 & 2) or ‘What was your first job after leaving school?’ (Extract 5), or they may consist of multiple components (Extract 3) requiring the patient to produce several different responses to each of the multiple components. A patient’s ability to answer all parts of these compound questions offers differential diagnostic potential. It has already been established (in Extract 3) that patients with FMD are able to respond to compound questions, often over an extended period of time and following detailed talk. Here is another example of an exchange involving the neurologist asking a compound question (this time constructed with three parts) and the patient with FMD responding relevantly to all three parts of the question.

Extract 6

010_FMD
01 _ DOC .hh Can you tell me a >little bit<
02 about your background.=Wher- Where you’re
03 from originally and um: (0.6) where did
04 you go to un- college or,
05 PAT I’m: from (city name).=
06 DOC =Hm hmm.
07 PAT .h er::m: an’ I came into: erm: .hh when
08 I first started wo:rk I wor- I worked
09 in an office and did (.) varying things
10 like that..h As my chil- as I had my
11 children erm: tch I’ve done retail.= So
12 I worked for Marks & Spencer and Boo:ts.
13 .hh And then it were only at thirty-fi:ve
14 that I came as a mature student to be a
15 nurse.
16 (0.2)
17 PAT .hh So: I’ve done my: training if you
18 like, er: .h I went from:: little
19 part time work to going to full
20 ti:me,.h w[ith shi]fts and studying.
21 DOC [Mm hmm.]
22 PAT .hh Erm: so I’ve s- I’ve been t- it’s
23 (City name) erm University that I’ve
24 been to.

25 (0.2)
26 PAT For me nur:sin'.

The patient recalls and is able to process and respond accurately to all parts of the neurologist's compound question – telling her where she was from originally (line 5), giving a bit of background about herself (lines 7-20) and where she went to university (line 23) – the three things the neurologist asked for. Her answer demonstrates that she is able to respond in detail to personal information-seeking questions, is able to display working memory in interaction and is able to process and respond to compound questions. These features are characteristic of individuals with FMD.

People with dementia frequently have difficulty comprehending questions, in addition to those they experience in recalling information; together these difficulties mean that conversational communication can be challenging for them. As a result, people with dementia often experience difficulty in answering all parts of a compound question (see Extract 7).

Extract 7

008 DEM
01 DOC .hh Do you have any problems er with
02 reading or writing.
03 (0.6)
04 DOC Or spelling?
05 (1.2)
06 PAT .hh Er, (4.0) er, (0.2) tck What do you
07 mean, r-reading?
08 DOC Yeah. Can you read OK?
09 (0.3)
10 PAT Yeah, I can read.
11 DOC Can you write what you want?
12 PAT .hhh Er, well it takes me a lot longer.
13 I have to sit and think about it.= Er
14 when my pal's with me (0.5) he sometimes
15 b- b[etter with] what to do like.
16 DOC [Yeah.]
17 (1.4)
18 PAT D'you know what I mean?
19 DOC Can you spell OK?
20 (0.2)
21 PAT .hh Er, er, I'm n- not very good
22 speller.= But (0.4) sometimes, (1.4)
23 it- (1.0) an' it dun't come to me.

The patient is unable to respond to all three parts of this question and instead pursues clarification from the doctor (lines 6-7). The doctor prompts the patient by continuing to break the question down, asking each point individually, after which the patient responds. This suggests that the patient is unable to recall and process the three parts of the question simultaneously. The patient not only answers the questions with a relevant response when asked individually but he also displays an understanding of each of the functions through gestured actions. Such compound questions pose a greater difficulty for those with dementia than they do for people with FMD.

Patient delays in responding

There are a number of additional features that relate to the ones described above. Typically, in response to the neurologist's questions, people with dementia either do not respond (prompting further questioning from the neurologist, e.g. Extract 5, lines 1-5 and 11-15) or take longer to answer than people with FMD. In the

sequences from the consultations with FMD patients in previous extracts it is apparent that patients answers contiguously, with no intervening delay in responding. This pattern of no delay appears in Extract 1 (lines 1-2 – responding to the question about age); Extract 3 (lines 5-6 - responding to the neurologist’s questions about the reason for the patient’s visit to the clinic and what expectations they have); Extract 4 (lines 8-9 – the patient being asked to describe a typical migraine); and Extract 6 (lines 4-5 – responding to a request to background information). However, by contrast quite often the responses given by people with dementia are delayed (see Extract 5 lines 3, 5, 9, 21, 34, 41 and Extract 7 lines 3, 5, 9, 20). Often this delay is substantial, for example in Extract 5 (line 3) there is a delay of 6.8 seconds between the neurologist’s initial question and their follow up questions (a total delay of 9.8 seconds until the patient produces a response). Delay in interaction is therefore a further differentiating feature between FMD patients and those with dementia.

The elaboration of patient’s accounts through detailing

The detail people exhibit in their talk also has the potential to differentiate between the diagnoses. When people with FMD respond to the neurologist’s questions, their responses consist of multiple and extended turns containing detailed examples and additional often-unsolicited information (see Extract 8).

Extract 8

04_FMD
01_ DOC .hh So how long have you been running
02 the (Shop name),
03 PAT .h Twenty-five years.
04 (0.6)
05 PAT We had a twenty-fifth birthday party
06 last Saturday.
07 (2.4) ((Neurologist writing))
08 DOC .h So how many: (.) hours a week are
09 you working,
10 PAT tch Qoh: hh .hh I go to the wholesale
11 market at six o’clock in the
12 mo[rning th]ree days a week to buy
13 DOC [Mm hmm.]
14 PAT the fruit and veg,
15 (0.8)
16 PAT And then I go to the shop every morning
17 about t for about an hour and a half or
18 so, .hh
19 DOC Mm hmm.
20 PAT And then I bake a lot of cakes a couple
21 of times a week.=So I: must do: twenty
22 hours a week at lea[st.=Ye]ah.
23 DOC [Right.]

Here the patient answers each of the neurologist’s questions (lines 1-2 and 8-9), her responses being elaborated over more than one turn. She introduces new information, which has not been solicited by the neurologist in her questions. Following the first question, ‘So how long have you been running the (shop)’ (lines 1-2), that patient’s answer is contiguously, ‘Twenty-five years’ (line 3). She then volunteers more information about the celebration they had to mark the anniversary; detailing what day of this week this event took place (lines 5-6). An extended response was also produced following the neurologist’s second question enquiring about how many hours a week the patient works (lines 8-9). Although an answer, ‘Twenty hours a

week at least' (lines 21-22) was given, this came after a detailed description of the patient's weekly activities, including what time she goes to the market ('six o'clock in the morning', lines 10-11) and what she purchases ('fruit and veg', line 14). This extract illustrates a recurrent pattern in the consultations with FMD patients in that their talk is detailed, and this detail is often unsolicited by the neurologist. Furthermore, volunteering this detailed information displays episodic memory capabilities in the interaction.

When patients with dementia respond to the neurologist's questions, their responses often consist of admissions of forgetfulness or lack of knowledge/understanding (often attesting to a lack of episodic memory), for example 'Can't remember now' and 'Not off hand' (Extract 5, lines 6 and 35), and below in Extract 9 (line 4) in which the patient admits to 'not entirely' being able to describe his understanding about why he is at the clinic. These, as well as other responses, are typically very short, consisting of only single utterance or single turn units. These lack specificity and detail and rarely volunteer any additional information that falls outside the requirement of the question. The patient with dementia in the extract below (9) exhibits delayed and undetailed interaction.

Extract 9

033_DEM

01 DOC Could you describe what's: What you
 02 understand about why you're here?
 03 (0.3)
 04 PAT Erm:: (0.4) not entirely::.
 05 (2.0)
 06 DOC .hh Wha:t's been the problem?
 07 (0.2)
 08 PAT Erm: (0.2) memory sho:rtage.
 09 DOC Okay,
 10 (10.4 seconds omitted)
 11 DOC An' (.) could you::, give me an
 12 example of the last time your memory
 13 (0.2) let you down,
 14 (0.6)
 15 PAT tch .hh [Erm:. (2.4)
 16 PAT [((PAT turns to look at COM))]
 17 COM °In the car-° you've lost your sense
 18 of direction.

Following each of the neurologist's questions (lines 1-2, 6, 11-13) the patient delays responding (lines 3, 7 and 14). His responses lack any additional detail (of the kind seen in Extract 3, lines 6-24, Extract 6, lines 7-24 and Extract 8, lines 10-22), and are short, often consisting of just two words as in lines 4 and 8. In response to the third question the patient's only response is 'Erm', before turning his head to seek help from his companion in answering the question. Although the head turning sign (Fukui, 2011) appears to be a recurrent pattern in these consultations and is a characteristic feature of the communicative practice of people with dementia, it is not the focus here. Instead it is important to note that the responses given people with dementia are delayed and undetailed, often leaving the detail to be provided by their companion (in this instance this appears in line 17). These divergent patterns in the content and delay of interaction contribute to the different conversational profiles.

Discussion

In this exploratory stage of the project we have identified a number of interactional features that begin to form differential conversational profiles between people with dementia and those with other functional memory complaints.

The initial findings show that people with FMD are able to respond contiguously and in detail to questions, including those requesting personal information, often volunteering additional information (which is unsolicited). They can display working memory in interaction and engage in additional interactional work, marking self-repetitive talk using resources such as 'as I said'. In doing so they orient to a recollection that the information has been previously voiced and display an awareness of their repetition. They are also able to process and respond to compound questions. In our consultations we have observed distinct differences in the communicative practices of those with dementia, who are frequently unable produce accurate information about themselves such as how old they are or where they live. Their responses to questions are delayed, minimal (consisting of single utterances or single turns) and undetailed. People with dementia regularly display a lack of working memory in interaction by repeating themselves, providing information that is unmarked and produced as if for the first time. They also struggle to transact compound questions.

This paper has focused on some of the dyadic features of interaction, touching only briefly on some of the triadic patterns that have begun to emerge such as the head turning sign (Fukui, 2011). Further analysis of the data may demonstrate that triadic features can contribute to fuller differential conversational profiles. Taken independently these features do not hold significant diagnostic value but used collectively they have the potential to enable differential diagnosis based on communicative practices.

As demonstrated in the seizure research (Reuber et al, 2009), conversational profiling has potential clinical applications, in this case in the differential diagnosis of dementia and functional memory disorders. Practitioners would need actively to listen for the interactional practices outlined in the profiles. Additionally this research and the resulting profiles could underpin an interactional toolkit to aid practitioners in designing their consultation to maximise diagnostic potential, for instance they may wish to design appropriate compound questions and listen for the response. This contravenes much of the standard communications training on 'effective communication', which uses the KISS principle – keep it short and simple (Manthorpe et al, 2011). However, there is a diagnostic benefit to 'conversational testing' that is perceived to be less stressful than the battery of neuropsychological examinations the patients experience during the process (Lai et al, 2008; Gibson & Anderson, 2011).

Furthermore, these interactional features correspond with domains of cognition that are tested during neuropsychological examination, which employ standardized cognitive screening instruments such as the ACE-R. These domains include memory, attention/orientation, language and verbal fluency. Being able to identify features of cognition (such as aspects of memory); which are displayed in interaction, offers an alternative and complementary screening tool for dementia based on communication. Such a tool might help to reduce the unnecessary imposition on patients of the full battery of neuropsychological testing.

It is acknowledged here that both dementia and FMD are not homogenous categories and this pilot project does not offer an insight into the differentiating subtypes of dementia including Alzheimer's disease, fronto-temporal dementia, dementia with Lewy body and other Parkinsonian syndromes for example, as well as

into the spectrum of functional memory complaints including those caused by both behavioural and cognitive difficulties. The heterogeneous nature of these subtypes makes any group studies difficult due to the variability in the nature of the associated cognitive and communicative impairments (Gardner 1974; Bayles 1985). However, common symptoms of each subtype can be identified, and indeed similarities between subtypes can be recognized. Further studies may explore the differential diagnosis between subtypes of both dementia and of FMD.

With government initiatives seeking better ways to screen for dementia and to improve the patient experience during this screening process, practitioners in both primary care and secondary services are looking for new approaches to meet these goals. These interactional profiles have the potential to offer a method of assessment that is conversation in nature, providing further evidence for clinical judgments made during history taking. Using 'conversation as assessment' could be of significant diagnostic value and provide a useful screening tool for dementia as well as other functional memory disorders.

References

- Alzheimer's Society (2014) Dementia diagnosis rates. Available
>http://www.alzheimers.org.uk/site/scripts/documents_info.php?documentID=2165<
- Bayles, K.A. (1985) 'Communication in dementia', in H. Ulatowska (ed.) *The Aging Brain*. San Diego: College-Hill Press, pp 157-173.
- Boustani, M., Callahan, C.M., Unverzagt, F.W., Austrom, M.G., Perkins, A.J., Fultz, B.A., Hui, S.L. Hendrie, H.C. (2005) Implementing a Screening and Diagnosis Program for Dementia in Primary Care. *Journal of General Internal Medicine*, 20(7): 572-577.
- Department of Health (2012) Commissioning for Quality and Innovation (CQUIN) framework. Accessed 28th August 2013. Available
><https://www.gov.uk/government/publications/using-the-commissioning-for-quality-and-innovation-cquin-payment-framework-guidance-on-new-national-goals-for-2012-13><
- Drew, P., Chatwin, J., & Collins, S. (2001) Conversation analysis: a method for research into interactions between patients and health-care professionals. *Health Expectations*, 4:58-70.
- Drew, P. (2005) Conversation analysis. In K. L. Fitch, & R. E. Sanders (Eds.), *Handbook of language and social interaction* (pp. 71–102). London, UK: Lawrence Erlbaum Associates Publishers.
- Ekberg, K., McDermott, J., Moynihan, C., Brindle, L., Little, P. and Leydon, G.M. (2014) The role of helplines in cancer care: Intertwining emotional support with information or advice seeking needs. *Journal of Psychosocial Oncology*, 1-42.
- Fukui, T., Yamazaki, T. & Kinno, R. (2011) Can the 'Head-Turning Sign' Be a Clinical Marker of Alzheimer's Disease? *Dementia and Geriatric Cognitive Disorders Extra*, 1(1): 310–317.
- Gardner, H. (1974) *The Shattered Mind: The Person After Brain Damage*. New York: Random House.
- Gibson, A. & Anderson, K. (2011) Difficult Diagnoses: Family Caregivers' Experiences During and Following the Diagnostic Process for Dementia. *American Journal of Alzheimer's disease and other dementias*, 26(3):212-217.
- Heritage, J. & Stivers, T (1999) Online commentary in acture medical visits: a method of shaping patient expectations. *Social science & medicine*, 49:1501-1517.
- Heritage, J & Maynard, D.W. (2006) *Communication in Medical Care*. Cambridge University Press, Cambridge.

- Heritage, J. & Robinson, J. (2006) The structure of Patients' presenting concerns: Physicians Opening Questions. *Health Communication*, 19(2): 89-102.
- Heritage, J., Robinson, J., Elliott, M., Veckett, M. & Wilkes, M. (2007) Reducing Patients' Unmet Concerns in Primary Care: the Difference One Word Can Make. *Journal of General Internal Medicine*, 22(10):1429-33.
- Jefferson, G. (1983). Issues in the transcription of naturally-occurring talk: Caricature versus capturing pronunciation particulars. *Tilburg papers in language and literature*, Tilburg University, Tilburg. pp. 1-12.
- Jefferson, G. (2004). Glossary of transcript symbols with an introduction. In G. H. Lerner (Ed.), *Conversation analysis: Studies from the first generation* (pp. 13-23). Philadelphia: John Benjamins.
- Jones, D. (2012) Conversations with a person with Alzheimer's disease: A conversation analytic study. Unpublished PhD dissertation, University of York, York, UK.
- Jones, D. (2013) A family living with Alzheimer's disease: The communicative challenges. *Dementia: The International journal of social research and practice*, September 18, 2013, doi:10.1177/1471301213502213.
- Knopman, D.S., DeKosky, S.T., Cummings, L.L., Chui, H., Corey-Bloom, J., Relkin, N., Small, G.W., Miller, B. Stevens, J.C. (2001) Practice parameter: Diagnosis of dementia (an evidence-based review). *Neurology*, 59(9):1143-1153.
- Lai, J.M., Hawkins, K.A., Gross, C.P. & Karlawish, J.H. (2008) Self-reported distress after cognitive testing in patients with Alzheimer's disease. *Journal of Gerontology Series A*, 63(8): 855-859.
- Manthorpe, C., Young, T., Howells, D. (2011) DEMTEC: Dementia Toolkit for Effective Communication. Available at >http://www.demtalk.org.uk/wp-content/uploads/2012/11/DemTalk_Full_HealthCareProfessionals.pdf<
- NHS direct enhanced service (2013) Facilitating timely diagnosis and support for people with dementia. Available ><http://www.england.nhs.uk/wp-content/uploads/2013/03/ess-dementia.pdf><
- Reuber, M., Monzoni, C., Sharrack, B. & Plug, L (2009) Using interactional and linguistic analysis to distinguish between epileptic and psychogenic nonepileptic seizures: A prospective, blinded multirater study. *Epilepsy & Behavior*, 16(1):139-144 .
- Robson, C., Drew, P., Walker, T. & Reuber, M. (2012) Catastrophising and normalising in patient's accounts of their seizure experiences. *Seizure*, 21(10):795-801.
- Royal College of Psychiatrists (2013) English National Memory Clinic Audit Report. Available ><http://www.rcpsych.ac.uk/pdf/English%20National%20Memory%20Clinics%20Audit%20Report%202013.pdf><
- Schmidtke, K., Pohlmann, S. & Metternich, B (2008) The Syndrome of Functional Memory Disorder: Definition, Etiology, and Natural Course. *American Journal of Geriatric Psychiatry*, 16: 981-988.
- Schmidtke, K. & Metternich, B. (2009) Validation of two inventories for the diagnosis and monitoring of functional memory disorder. *Journal of Psychosomatic Research*, 67(3): 245-251.
- Schwabe, M., Howell, S.J. & Reuber, M. (2007) Differential diagnosis of seizure disorders: A conversation analytic approach. *Social Science & Medicine*, 65:712-724.
- Sidnell, J. (2010) *Conversation analysis: An introduction*. Sussex, UK: Wiley-Blackwell.
- Stivers, T (2002) Participating in decisions about treatment: overt parent pressure for antibiotic medication in pediatric encounters. *Social Science & Medicine*, 54: 1111-1130.
- Stivers, T. (2004) "No no no" and Other Types of Multiple Sayings in Social Interaction. *Human Communication Research*, 30(2): 260-293.

Appendix 1

Transcription symbols

DOC/PAT	Speaker labels (DOC = Doctor/Neurologist; PAT = Patient; COM Companion)
=	Links talk produced in close temporal proximity (latched talk)
> <	Talk between symbols is rushed or compressed
° °	Encloses talk which is produced quietly
<u>underline</u>	Underlining marks emphasis of some kind
CAPS	Words or parts of words spoken loudly marked in capital letters
s:::::	Sustained or stretched sound; the more colons, the longer the sound
. ? ,	Stop indicates falling intonation; a question mark indicates rising intonation over a word; a comma indicates a slight rising intonation at the end of word
.hhh	Inbreath, the number of 'h's representing the length of the inbreath
hhh.	Outbreath, the number of 'h's representing the length of the outbreath
[]	Encloses talk in overlap i.e. when more than one speaker is speaking
(word)	Parentheses indicate transcriber doubt
(this/that)	Alternative hearings
((description))	Description of what can be heard, rather than transcription e.g. ((shuffling papers))
cu-	Cut-off word or sound
(0.6)	Silence in seconds
(.)	Silence of less than two tenths of a second
^ or □	Indicates marked pitch rise
□	Indicates marked fall in pitch
(hhenhh)	Indicates laughter while speaking (aspiration)